

# NUMERICAL ANALOGUES OF THE KODAIRA DIMENSION AND THE ABUNDANCE CONJECTURE

THOMAS ECKL

ABSTRACT. We add further notions to Lehmann's list of numerical analogues to the Kodaira dimension of pseudo-effective divisors on smooth complex projective varieties, and show new relations between them. Then we use these notions and relations to prove that the Abundance Conjecture, as formulated in the context of the Minimal Model Program, and the Generalized Abundance Conjecture using these numerical analogues to the Kodaira dimension, are equivalent for non-uniruled complex projective varieties.

## 0. INTRODUCTION

During the last decade a plethora of numerical analogues to the Kodaira dimension for pseudoeffective divisors on (smooth) complex projective varieties was introduced, by Nakayama [Nak04], Demailly, Boucksom, Paun and Peternell [BDPP13], Siu [Siu11] and Lehmann [Leh13]. Lehmann furthermore clarified lots of relations between these numerical dimensions, adding some new notions, ordering them by the way how they are constructed and showing that most of them are equal. In this note we slightly extend his list and prove some more relations. We use these notions and results to show that the Abundance Conjecture as formulated in the context of the Minimal Model Program (see e.g. [Mat02, Conj.3-3-4]) is equivalent to a Generalised Abundance Conjecture introduced in [BDPP13]. On the way, we prove the birational equivalence of most of these notions of numerical dimension. In more details, we will discuss the following notions of numerical dimension, ordered according to their construction method as suggested by Lehmann, and postponing some technical definitions to section 1:

**Definition 0.1.** *Let  $X$  be a smooth complex projective variety and  $D$  a pseudoeffective  $\mathbb{R}$ -divisor on  $X$ . Then we define the following numerical dimensions using*

- *positive product conditions:*
  - (1)  $\nu_{\text{K\"{a}h}}(D) := \max \{k \in \mathbb{N} \mid \langle [D]^k \rangle_{\text{K\"{a}h}} \neq 0\}$ , where  $[D]$  denotes the  $(1, 1)$ -cohomology class of the integration current associated to  $D$ ;
  - (2)  $\nu_{\text{alg}}(D) := \max \{k \in \mathbb{N} \mid \langle D^k \rangle_{\text{alg}} \neq 0\}$ ;
  - (3)  $\nu_{\text{res}}(D) := \max \{\dim W \mid \langle D^{\dim W} \rangle_{X|W} > 0\}$  where  $W \subset X$  ranges over subvarieties not contained in the diminished base locus  $\mathbb{B}_-(D)$  (defined in 1.3);
- *volume conditions:*
  - (4)  $\nu_{\text{Vol}}(D) := \max \{k \in \mathbb{N} \mid \exists C > 0 : C \cdot t^{n-k} < \text{vol}(D + tA) \text{ for all } t > 0\}$ , where  $A$  is a sufficiently ample  $\mathbb{Z}$ -divisor on  $X$ ;

---

1991 *Mathematics Subject Classification.* 14C20, 14E05.

*Key words and phrases.* numerical Kodaira dimension, Abundance Conjecture.

- (5)  $\nu_{\text{Vol, res}}(D) := \max \{ \dim W \mid \lim_{\epsilon \rightarrow 0} \text{vol}_{X|W}(D + \epsilon A) > 0 \}$ , where  $W \subset X$  ranges over subvarieties not contained in  $\mathbb{B}_-(D)$  and  $A$  is a sufficiently ample  $\mathbb{Z}$ -divisor on  $X$ ;
- (6)  $\nu_{\text{Vol, Zar}} := \max \left\{ \dim W \mid \inf_{\phi} \text{vol}_{\widetilde{W}}(P_{\sigma}(\phi^* D)|_{\widetilde{W}}) > 0 \right\}$ , where  $W \subset X$  ranges over subvarieties not contained in  $\mathbb{B}_-(D)$ , the morphism  $\phi : (\widetilde{X}, \widetilde{W}) \rightarrow (X, W)$  ranges over all smooth  $W$ -birational models of  $(X, W)$  and  $P_{\sigma}(\phi^* D)$  is the positive part of the Zariski decomposition of  $\phi^* D$  (all defined in 1.6);
- *perturbed growth conditions:*

(7)  $\kappa_{\sigma}(D) := \max \{ k \in \mathbb{N} \mid \limsup_{m \rightarrow \infty} m^{-k} h^0(X, \mathcal{O}_X(A + \lfloor mD \rfloor)) > 0 \}$ , where  $A$  is a sufficiently ample  $\mathbb{Z}$ -divisor;

(8)  $\kappa_{\text{num}}(D) := \sup_{k \geq 1} \left\{ \limsup_{m \rightarrow \infty} \frac{\log h^0(X, \mathcal{O}_X(\lfloor mD \rfloor + kA))}{\log m} \right\}$ , where  $A$  is an ample  $\mathbb{Z}$ -divisor;
  - *Seshadri-type conditions:*

(9)  $\kappa_{\nu}(D) := \min \{ \dim W \mid D \not\preceq W \}$ , where  $D \succeq W$  means that  $D$  dominates  $W$  (defined in 1.9);

(10)  $\kappa_{\nu, \text{Leh}}(D) := \min \{ \dim W \mid \forall \epsilon > 0 : \phi_W^* D - \epsilon E_W \text{ not pseudoeffective} \}$ , where  $\phi_W : \widetilde{X} \rightarrow X$  is any birational morphism of smooth varieties such that  $\mathcal{O}_{\widetilde{X}}(E_W) = \phi^{-1} \mathcal{I}_W \cdot \mathcal{O}_{\widetilde{X}}$ .

For attributions of these definitions see also section 1.

These notions of numerical dimensions are related in the following way:

**Theorem 0.2.** *Let  $X$  be a smooth complex projective variety and  $D$  a pseudoeffective  $\mathbb{R}$ -divisor on  $X$ . Then:*

$$\begin{array}{ccc}
 \nu_{\text{Vol}}(D) & \kappa_{\text{num}}(D) & \equiv \equiv \kappa_{\text{num}}(D) \\
 \parallel & \vee & \wedge \\
 \nu_{\text{alg}}(D) = \nu_{\text{res}}(D) = \nu_{\text{Vol, res}}(D) = \nu_{\text{Vol, Zar}}(D) \leq \kappa_{\sigma}(D) \leq \kappa_{\nu}(D) \leq \nu_{\text{alg}}(D) & & \\
 \wedge & & \vee \\
 \nu_{\text{K\"ah}}(D) & & \kappa_{\nu, \text{Leh}}(D)
 \end{array}$$

In section 2 and 3 we will prove the inequalities  $\nu_{\text{alg}}(D) \leq \nu_{\text{K\"ah}}(D)$ ,  $\kappa_{\nu}(D) \leq \nu_{\text{alg}}(D)$ ,  $\nu_{\text{Vol, Zar}}(D) \leq \kappa_{\text{num}}(D)$ ,  $\kappa_{\text{num}}(D) \leq \kappa_{\nu}(D)$  and  $\kappa_{\nu, \text{Leh}}(D) \leq \kappa_{\nu}(D)$ , and we locate the proofs of the other inequalities in the works of Lehmann [Leh13] and Nakayama [Nak04]. Our proofs of  $\nu_{\text{Vol, Zar}}(D) \leq \kappa_{\text{num}}(D)$  and  $\kappa_{\text{num}}(D) \leq \kappa_{\nu}(D)$  also rely on Lehmann's ideas. But for the proof of  $\kappa_{\nu}(D) \leq \nu_{\text{alg}}(D)$  we need a new ingredient: the derivative of the restricted volume, generalizing Thm.A in [BFJ09] (see Thm 3.1).

The theorem shows that most of the notions in Def. 0.1 are equal. Therefore the following definition is justified:

**Definition 0.3.** *Let  $X$  be a smooth complex projective variety and  $D$  a pseudoeffective  $\mathbb{R}$ -divisor on  $X$ . Then the numerical dimension  $\nu_X(D)$  of  $D$  is defined as one of the equal numbers*

$$\nu_{\text{alg}}(D) = \nu_{\text{res}}(D) = \nu_{\text{Vol, res}}(D) = \nu_{\text{Vol, Zar}}(D) = \kappa_{\sigma}(D) = \kappa_{\nu}(D) = \kappa_{\text{num}}(D).$$

In section 4 we show that the numerical dimension of a pseudoeffective divisor behaves well under birational morphisms:

**Proposition 0.4 (= Proposition 4.1).** *Let  $f : \widetilde{X} \rightarrow X$  be a birational morphism between smooth complex projective varieties, let  $D$  be a pseudoeffective divisor on*

$X$  and  $\tilde{D}$  a pseudoeffective divisor on  $\tilde{X}$  such that  $\tilde{D} - f^*D$  is an  $f$ -exceptional divisor. Then:

$$\nu_X(D) = \nu_{\tilde{X}}(\tilde{D}).$$

In a celebrated theorem Boucksom, Demailly, Paun and Peternell show that the canonical divisor  $K_X$  of a non-uniruled smooth complex variety  $X$  is pseudoeffective [BDPP13, Cor.0.3]. Consequently, the numerical dimension of the canonical divisor can be used to state the Abundance Conjecture:

**Conjecture 0.5** (Abundance Conjecture). *Let  $X$  be a non-uniruled smooth complex projective variety. Then:*

$$\nu(X) := \nu_X(K_X) = \kappa(X).$$

Here  $\kappa(X) = \kappa_X(K_X)$  denotes the Kodaira dimension of the canonical divisor  $K_X$ , defined e.g. as

$$\kappa_X(K_X) := \limsup_{m \rightarrow \infty} \frac{\log h^0(X, \mathcal{O}_X(mK_X))}{\log m}.$$

Note that Boucksom, Demailly, Paun and Peternell refer in their Generalized Abundance Conjecture [BDPP13, Conj.3.8] to  $\nu_{\text{K\"ah}}(K_X)$  which is only conjecturally equal to  $\nu(X)$ , as discussed in 2.1.

In the context of the Minimal Model Program the Abundance Conjecture is formulated under the assumption that minimal models of smooth complex projective varieties exist (see Section 4 for definitions):

**Conjecture 0.6** (Abundance Conjecture, MMP version [Mat02, Conj.3-3-4]). *Let  $S$  be a minimal model of a non-uniruled smooth projective complex variety  $X$ . Then  $|mK_S|$  is base point free for sufficiently divisible and large  $m \in \mathbb{N}$  (that is,  $K_S$  is semi-ample).*

We use the birational invariance of the numerical dimension to show in section 4 that the two Abundance Conjectures as stated above are equivalent:

**Theorem 0.7** (= **Theorem 4.5**). *Let  $S$  be a minimal model of a non-uniruled smooth projective complex variety  $X$ . Then*

$$\nu_X(K_X) = \kappa_X(K_X) \iff K_S \text{ is semi-ample.}$$

Note that this equivalence is asserted in passing on several occasions (see e.g. [GL13], [DHP13]). However, the author still thinks that it is worth presenting the argument in detail, emphasizing in particular that not all the possible definitions of numerical dimension are easily shown to be birationally invariant.

## 1. NOTIONS OF NUMERICAL DIMENSION

In the following  $X$  is always a smooth  $n$ -dimensional complex projective variety and  $D$  a pseudoeffective  $\mathbb{R}$ -divisor on  $X$ .

1.1.  $\nu_{\text{K\"ah}}(D)$ . This notion is defined in [BDPP13, Def.3.6]. The *moving intersection product*  $\langle [D]^k \rangle_{\text{K\"ah}}$  of the  $(1, 1)$ -cohomology class of the integration current  $[D]$  is constructed in [BDPP13, Thm.3.5] following [Bou02]: For suitably chosen birational morphisms  $\mu_m : X_m \rightarrow X$  of smooth complex varieties, real numbers  $\delta_m \downarrow 0$ ,

closed semi-positive forms  $\beta_{i,m}$  representing big and nef classes in  $N^1(X_m)$ , effective  $\mu_m$ -exceptional  $\mathbb{Q}$ -divisors  $E_{i,m}$  on  $X_m$  and any ample class  $\omega$  on  $X$  such that  $[E_{i,m}] + \beta_{i,m}$  represents the  $(1,1)$ -class  $(\mu_m)^*([D] + \delta_m\omega)$  we can set

$$\langle [D]^k \rangle_{\text{K\"ah}} := \lim_{m \rightarrow \infty} (\mu_m)_*([\beta_{1,m} \wedge \dots \wedge \beta_{k,m}]),$$

where the limits are taken in  $H^{k,k}(X)$ . Note that other choices of  $\mu_m, \delta_m, \beta_{i,m}, E_{i,m}$  satisfying the properties above will yield "smaller"  $(k,k)$ -classes  $\alpha$ , that is,  $\langle [D]^k \rangle_{\text{K\"ah}} - \alpha$  is represented by a positive current.

1.2.  $\nu_{\text{alg}}(D)$ . This notion appears first in [Leh13] where it is nevertheless attributed to [BDPP13]. In fact, Lehmann uses the algebraic analogue of the moving intersection product  $\langle [D]^k \rangle_{\text{K\"ah}}$  as defined in [BFJ09]: To calculate  $\langle [D]^k \rangle_{\text{alg}}$  Boucksom, Favre and Jonsson replace the  $(k,k)$ -cohomology class  $[\beta_{1,m} \wedge \dots \wedge \beta_{k,m}] \in H^{k,k}(X_m)$  by the intersection  $k$ -cycle class  $[\beta_{1,m}] \cdots [\beta_{k,m}] \in N^k(X_m)$  and take the limit in  $N^k(X)$ . The connection to  $\langle [D]^k \rangle_{\text{K\"ah}}$  is discussed in 2.1.

Note that the moving intersection product is continuous and homogeneous on the cone spanned by the classes of big divisors ([BFJ09, Prop.2.9]). Furthermore it coincides with the usual intersection number if the numerical classes are represented by nef divisors [BFJ09, Prop.2.12].

1.3.  $\nu_{\text{res}}(D)$ . This notion is defined in [Leh13]. The *diminished* or *restricted base locus* of an  $\mathbb{R}$ -divisor

$$\mathbb{B}_-(D) := \bigcup_{A \text{ ample}} \mathbb{B}(D + A)$$

appears in [ELM<sup>+</sup>06, Def.1.12] and [Nak04, Def.III.2.6&p.168]. Here,

$$\mathbb{B}(D + A) := \bigcap_{m \geq 1} \text{Bs}(\lfloor m(D + A) \rfloor)$$

is the *stable base locus* of  $D + A$ . Later on, we also need the *augmented base locus*

$$\mathbb{B}_+(B) := \bigcap_{A \text{ ample}} \mathbb{B}(B - A)$$

of a big  $\mathbb{R}$ -divisor  $B$  (see [ELM<sup>+</sup>06, Def.1.2]).

The *restricted moving intersection*  $\langle D^k \rangle_{X|W}$  is constructed in [BFJ09] for divisors  $W$  and generalized to arbitrary subvarieties  $W \not\subset \mathbb{B}_-(D)$  of dimension at least  $k$  in [Leh13, Def.2.6] (then  $D$  is called a *W-pseudoeffective divisor*): Similar to 1.2,

$$\langle D^k \rangle_{X|W} := \lim_{m \rightarrow \infty} (\mu_m)_*([B_{1,m}] \cdots [B_{k,m}] \cdot \widetilde{W}),$$

where the  $B_{i,m}$  are suitably chosen big and nef divisors on the smooth variety  $X_m$  such that  $\mu_m : X_m \rightarrow X$  is a birational morphism whose center does not contain  $W$  (a so-called *W-birational model* of  $X$ ), the  $\mathbb{Q}$ -divisors  $\mu_m^*(D + \delta_m A) - B_{i,m}$  are effective and  $\mu_m$ -exceptional for a fixed ample divisor  $A$  on  $X$  and real numbers  $\delta_m \downarrow 0$ , and  $\widetilde{W}$  is the strict  $\mu_m$ -transform of  $W$ . On the cone spanned by classes of big divisors  $B$  such that  $W \not\subset \mathbb{B}_+(B)$  (then  $B$  is called a *W-big divisor*), the restricted product is continuous and homogeneous (see [BFJ09, Prop.2.9&Prop.4.6] resp. [Leh13, Prop.4.7]). This implies furthermore that

$$\langle D^k \rangle_{X|W} = \lim_{\delta \downarrow 0} \langle (D + B^{(\delta)})^k \rangle_{X|W},$$

for arbitrary  $W$ -big divisors  $B^{(\delta)}$  converging to 0 when  $\delta \downarrow 0$ .

Note also that by setting  $W := X$  the moving intersection cycle class  $\langle D^k \rangle_{\text{alg}}$  can be obtained as a special case of the restricted moving intersection product. Finally, in the calculation of  $\langle D^k \rangle_{X|W}$  one can choose  $B_{1,m} = \cdots = B_{k,m}$  (see the proof of [BFJ09, Lem.2.6]).

1.4.  $\nu_{\text{Vol}}(D)$ . This notion is defined in [Leh13]. Note that the volume of the big  $\mathbb{R}$ -divisor  $D + tA$  can be defined as

$$\text{vol}(D + tA) := \limsup_{m \rightarrow \infty} \frac{h^0(X, \mathcal{O}_X(\lfloor m(D + tA) \rfloor))}{m^n}$$

because this definition coincides with the one in [Laz04, §2.2.C] as the continuous extension of the volume function on  $\mathbb{Q}$ -divisors to the big cone.

Fujita's theorem [BFJ09, Thm.3.1] states that  $\text{vol}(D + tA) = \langle (D + tA)^n \rangle_{\text{alg}}$ .

1.5.  $\nu_{\text{Vol, res}}(D)$ . This notion is introduced in [Leh13] and uses the restricted volume investigated in [ELM<sup>+</sup>09] (see also [Leh13, Def.2.12] for the definition):

$$\text{vol}_{X|W}(D + \epsilon A) := \limsup_{m \rightarrow \infty} \frac{h^0(X|W, \mathcal{O}_X(\lfloor m(D + \epsilon A) \rfloor))}{m^{\dim W} / (\dim W)!},$$

where  $H^0(X|W, \mathcal{O}_X(\lfloor m(D + \epsilon A) \rfloor))$  is defined as

$$\text{Im}(H^0(X, \mathcal{O}_X(\lfloor m(D + \epsilon A) \rfloor)) \rightarrow H^0(W, \widetilde{\mathcal{O}_W}(\lfloor m(D + \epsilon A) \rfloor))).$$

By the Generalised Fujita Theorem [ELM<sup>+</sup>09, Prop.2.11 & Thm.2.13],

$$\langle (D + \epsilon A)^{\dim W} \rangle_{X|W} = \text{vol}_{X|W}(D + \epsilon A).$$

Consequently, the restricted volume is continuous and homogeneous on the cone spanned by the classes of  $W$ -big divisors  $B$ .

1.6.  $\nu_{\text{Vol, Zar}}(D)$ . Again this notion is introduced in [Leh13]. Note that morphisms  $\phi : (\widetilde{X}, \widetilde{W}) \rightarrow (X, W)$  are  $W$ -birational if the irreducible subvariety  $W \subset X$  is not contained in the center of the birational map  $\phi$ , and  $\widetilde{W}$  is the strict  $\phi$ -transform of  $W$ . The *divisorial Zariski decomposition* or  $\sigma$ -decomposition

$$\phi^*D = P_\sigma(\phi^*D) + N_\sigma(\phi^*D)$$

into a positive part  $P_\sigma$  and a negative part  $N_\sigma$  is constructed by Nakayama [Nak04, III.1] and [Bou04]. Lehmann characterized the negative part  $N_\sigma(\phi^*D)$  as the divisorial part of the diminished base locus  $\mathbb{B}_-(D)$  [Leh13, Prop.3.3(3)], whereas Nakayama [Nak04, Lem.III.1.14(1)] showed that the numerical class of  $P_\sigma(\phi^*D)$  lies in the closure of the movable cone  $\text{Mov}(X)$  spanned by fixed-part free divisors. For later purposes we need more details of Nakayama's construction of the  $\sigma$ -decomposition:

**Definition 1.1.** *Let  $X$  be a smooth projective complex variety,  $B$  a big  $\mathbb{R}$ -divisor and  $\Gamma$  a prime divisor on  $X$ . We set*

$$\sigma_\Gamma(B) := \inf\{\text{mult}_\Gamma \Delta \mid \Delta \equiv B, \Delta \geq 0\}.$$

*If  $D$  is a pseudoeffective  $\mathbb{R}$ -divisor and  $A$  an ample divisor on  $X$  we set*

$$\sigma_\Gamma(D) := \lim_{\epsilon \downarrow 0} \sigma_\Gamma(D + \epsilon A)$$

and define

$$N_\sigma(D) := \sum_{\Gamma} \sigma_\Gamma(D) \cdot \Gamma.$$

The well-definedness of  $\sigma_\Gamma(D)$  is shown in [Nak04, II.1.5].

1.7.  $\kappa_\sigma(D)$ . This notion is defined in [Nak04, Def.V.2.5].

1.8.  $\kappa_{\text{alg}}(D)$ . This notion is defined in [Siu11].

1.9.  $\kappa_\nu(D)$ . This notion is defined in [Nak04, Def.V.2.20], requiring the notion of *numerical dominance*:

**Definition 1.2** ([Nak04, Def.V.2.12&V.2.16]). *Let  $D$  be an  $\mathbb{R}$ -divisor on a smooth projective variety  $X$  and  $W \subset X$  an irreducible subvariety. We say that  $D$  dominates  $W$  numerically and write  $D \succeq W$  if there exists a birational morphism  $\phi : \tilde{X} \rightarrow X$  and an ample divisor  $A$  on  $\tilde{X}$  such that  $\phi^{-1}\mathcal{I}_W \cdot \mathcal{O}_{\tilde{X}} = \mathcal{O}_{\tilde{X}}(E_W)$  is the locally free sheaf of an effective divisor  $E_W$  on  $\tilde{X}$ , and for every real number  $b > 0$  there exist real numbers  $x > b, y > b$  such that*

$$x \cdot \phi^* D - y \cdot E_W + A$$

*is pseudoeffective.*

Note that the condition above is satisfied for any birational morphism  $\psi : Y \rightarrow X$  with  $\psi^{-1}\mathcal{I}_W \cdot \mathcal{O}_Y = \mathcal{O}_Y(F_W)$  for an effective divisor  $F_W$  and ample divisor  $B$  once it is satisfied for  $\phi$  and  $A$ .

1.10.  $\kappa_{\nu, \text{Leh}}(D)$ . This notion is introduced in [Leh13] using [Leh13, Def.5.1]. See the discussion of the inequality  $\kappa_{\nu, \text{Leh}}(D) \leq \kappa_\nu(D)$  in 2.9 for why the two invariants may be different.

## 2. INEQUALITIES BETWEEN NOTIONS OF NUMERICAL DIMENSION

2.1.  $\nu_{\text{alg}}(D) \leq \nu_{\text{K\"ah}}(D)$ . The inequality holds because  $k$ -cycles are numerically equivalent if the corresponding integration currents are cohomologically equivalent. Equality will hold if the converse is also true. However, a class map  $N^k(X)_{\mathbb{R}} \rightarrow H_{\mathbb{R}}^{k,k}(X)$  only exists conjecturally. In particular, its existence is a consequence (but not equivalent) to the Hodge Conjecture. For more details see [Ful84, 19.1, 19.3].

2.2.  $\nu_{\text{alg}}(D) \leq \nu_{\text{Vol}}(D)$ . This inequality is proven in [Leh13, Thm.6.2.(1)=(7)].

2.3.  $\nu_{\text{alg}}(D) \leq \nu_{\text{res}}(D)$ . This inequality is proven in [Leh13, Thm.6.2.(1)=(2)].

2.4.  $\nu_{\text{res}}(D) \leq \nu_{\text{Vol, res}}(D)$ . This inequality is proven in [Leh13, Thm.6.2.(2)=(3)].

2.5.  $\nu_{\text{Vol, res}}(D) \leq \nu_{\text{Vol, Zar}}(D)$ . This inequality is proven in [Leh13, Thm.6.2.(3)=(4)].

2.6.  $\nu_{\text{Vol, Zar}}(D) \leq \kappa_\sigma(D)$ . This inequality is proven in [Leh13, Thm.6.2.(4)≤(5)].

2.7.  $\kappa_\sigma(D) \leq \kappa_\nu(D)$ . This inequality is proven in [Nak04, Prop.V.2.22(1)].

2.8.  $\nu_{\text{Vol,Zar}}(D) \leq \kappa_{\text{num}}(D)$ . For a sufficiently ample divisor  $A$  Lehmann shows in [Leh13, Thm.6.2.(4)≤(5)] that there exists a constant  $C > 0$  so that for every sufficiently large  $m$

$$Cm^{\nu_{\text{Vol,Zar}}(D)} \leq h^0(X, \mathcal{O}_X(\lfloor mD \rfloor + A)).$$

Taking the logarithm, dividing by  $\log m$  and letting  $m$  tend to  $\infty$  shows the desired inequality.


2.9.  $\kappa_{\nu, \text{Leh}}(D) \leq \kappa_{\nu}(D)$ . Let  $W \subset X$  be an irreducible subvariety,  $\phi : \tilde{X} \rightarrow X$  a birational morphism of smooth varieties such that  $\mathcal{O}_{\tilde{X}}(E_W) = \phi^{-1}\mathcal{I}_W \cdot \mathcal{O}_{\tilde{X}}$  and  $A$  an ample divisor on  $\tilde{X}$ . If  $\phi^*D - \epsilon E_W$  is pseudoeffective for an  $\epsilon > 0$  then

$$\frac{b+1}{\epsilon} \phi^*D - (b+1)E_W + A$$

is also pseudoeffective, for any  $b > 0$ , hence  $D \succeq W$ . Consequently,

$$\kappa_{\nu, \text{Leh}}(D) \leq \kappa_{\nu}(D).$$

Note that the argument for equality in the proof of [Leh13, Prop.5.3] does not work because projections of finite-dimensional vector spaces are not closed maps. In particular equality could fail if  $\phi^*D$  sits on a non-polyhedral part of the boundary of the big cone  $\text{Big}(\tilde{X})$ , as illustrated in the following diagram of a cut through the big cone by the affine plane in  $\text{NS}(\tilde{X})_{\mathbb{R}}$  passing through  $E_W$ ,  $\phi^*D$  and  $E_W - \frac{1}{b+1}A$ , for arbitrary  $b > 0$ :



In this situation,  $\phi^*D - \epsilon E_W$  is not pseudoeffective for all  $\epsilon > 0$ , but  $\phi^*D - \epsilon E_W + \frac{\epsilon}{b+1}A$  is pseudoeffective for all sufficiently small  $\epsilon > 0$ . Consequently,  $\frac{b+1}{\epsilon} \phi^*D - (b+1)E_W + A$  is pseudoeffective, hence  $D \succeq W$ .

Note also that Nakayama's proof of  $\kappa_{\sigma}(D) \leq \kappa_{\nu}(D)$  does not work if we replace  $\kappa_{\nu}(D)$  with  $\kappa_{\nu, \text{Leh}}(D)$ : The definition of  $\kappa_{\sigma}(D)$  only allows to find sections of  $\mathcal{O}_X(\lfloor mD \rfloor + A)$ , with the ample divisor  $A$  on  $X$  added.

2.10.  $\kappa_{\text{num}}(D) \leq \kappa_{\nu}(D)$ . We adapt [Leh13, Thm.6.7(7)] and its proof to Nakayama's definition of  $\kappa_{\nu}(D)$ : Let  $A$  be a sufficiently ample divisor on  $X$ , and let  $W \subset X$  be a subvariety such that  $\dim W = \kappa_{\nu}(D)$  and  $D \not\succeq W$ . In particular, for a resolution  $\phi : \tilde{X} \rightarrow X$  of  $W$  and an ample divisor  $H$  on the smooth projective variety  $\tilde{X}$ , there exists  $b > 0$  such that  $x\phi^*D - yE_W + H$  is not pseudoeffective for any choice of  $x, y > b$ .

Choose  $q \in \mathbb{N}$  large enough so that  $qH - \phi^*A$  is pseudoeffective, and consider any sufficiently large  $m \in \mathbb{N}$ . Then the  $\mathbb{R}$ -divisor  $m\phi^*D - q[b+1]E_W + qH$  and hence

$$m\phi^*D - q[b+1]E_W + qH - (qH - \phi^*A) = \phi^*(mD + A) - q[b+1]E_W$$

is not pseudoeffective. Therefore  $\phi^*([mD] + A) - q[b+1]E_W$  is not effective, and we obtain

$$h^0(\tilde{X}, \mathcal{O}_{\tilde{X}}(\phi^*([mD] + A) - q[b+1]E_W)) = 0.$$

Consequently,  $h^0(X, \mathcal{O}_X([mD] + A) \otimes \mathcal{I}_W^{q[b+1]}) = 0$ . Set  $q' = q[b+1]$  and write  $W_{q'}$  for the subscheme defined by the ideal sheaf  $\mathcal{I}_W^{q'}$ . Then there is an injection

$$H^0(X, \mathcal{O}_X([mD] + A)) \rightarrow H^0(W_{q'}, \mathcal{O}_{W_{q'}}([mD] + A)).$$

Since  $[mD] + A \leq m[D + A]$  the rate of growth for the right hand side is bounded by a multiple of  $m^{\dim W_{q'}} = m^{\kappa_\nu(D)}$ . In particular, there exists a constant  $C > 0$  such that

$$h^0(X, \mathcal{O}_X([mD] + A)) \leq C \cdot m^{\kappa_\nu(D)}.$$

Taking the logarithm, dividing by  $\log m$  and letting  $m$  tend to  $\infty$  shows the desired inequality.

### 3. PROOF OF $\kappa_\nu(D) \leq \nu_{\text{alg}}(D)$

To show this inequality we cannot just adapt the proof of [Leh13, Thm.6.2(6)  $\leq$  (1)] to Nakayama's definition of  $\kappa_\nu(D)$  but need a new ingredient: the derivative of the restricted volume function. The following statement generalizes Thm.A in [BFJ09].

**Theorem 3.1.** *Let  $X$  be a  $n$ -dimensional smooth projective complex variety and  $V = H_1 \cap \dots \cap H_{n-k}$  a  $k$ -dimensional complete intersection variety cut out by very general very ample linearly equivalent divisors  $H_i$ . If  $\alpha$  is a  $V$ -big and  $\gamma$  an arbitrary divisor class then*

$$\frac{d}{dt} \Big|_{t=0} \text{vol}_{X|V}(\alpha + t\gamma) = k \cdot \langle \alpha^{k-1} \rangle_{X|V} \cdot \gamma.$$

To prove this theorem and the inequality we first need further facts on the restricted moving intersection product and volume.

**Lemma 3.2.** *Let  $X$  be a smooth projective complex variety,  $V \subset X$  a subvariety and  $D$  a  $V$ -pseudoeffective divisor on  $X$ . Furthermore, let  $F \subset X$  be a very general element of a free family of subvarieties, that is, a general element of the family intersects any given algebraic subset of  $X$  in the expected codimension. Then for  $k \leq \dim V \cap F$ :*

$$\langle D^k \rangle_{X|V} \cdot F = \langle D^k \rangle_{X|V \cap F}.$$

*Proof.* This is a generalisation of [Leh13, Lem.4.18(2)]: Consider a countable set of smooth  $V$ -birational models  $\phi_m : (\tilde{X}_m, \tilde{V}_m) \rightarrow (X, V)$  on which the restricted product can be calculated, as

$$\langle D^k \rangle_{X|V} = \lim_{m \rightarrow \infty} (\phi_m)_* ([B_{1,m}] \cdots [B_{k,m}])$$

for big and nef divisors  $B_{i,m}$  on  $\tilde{X}_m$ . Choose  $F$  sufficiently general so that it does not contain any of the  $\phi_m$ -exceptional centers and intersects  $V$  generically transversally. Then the strict transform  $\widetilde{V \cap F}$  of  $V \cap F$  on  $\tilde{X}_m$  will be a cycle representing the class  $[\phi_m^* F] \cdot [\tilde{V}_m]$ . Thus we can identify the classes

$$\begin{aligned} (\phi_m)_* \left( [B_{1,m}] \cdots [B_{k,m}] \cdot \tilde{V}_m \right) \cdot F &= (\phi_m)_* \left( [B_{1,m}] \cdots [B_{k,m}] \cdot [\phi_m^* F] \cdot \tilde{V}_m \right) = \\ &= (\phi_m)_* \left( [B_{1,m}] \cdots [B_{k,m}] \cdot [\widetilde{V \cap F}] \right), \end{aligned}$$



and that implies the claimed equality.  $\square$

**Lemma 3.3.** *Let  $X$  be a smooth projective complex variety,  $V \subset X$  a subvariety of dimension  $d$  and  $D$  a  $V$ -pseudoeffective divisor on  $X$ . If  $k \leq d$  and  $A$  is an ample divisor on  $X$  then*

$$\langle D^k \rangle_{X|V} \neq 0 \iff \langle D^k \rangle_{X|V} \cdot A^{d-k} > 0.$$

*Proof.* Intersection theory (as presented for example in [Ful84]) implies that the group  $N^k(X)$  of numerical classes of  $k$ -cycles on  $X$  is generated by Segre classes, that are classes represented by push forwards of complete intersections on  $\mathbb{P}_X(E)$  of the correct dimension and with respect to the tautological line bundle  $\mathcal{O}_{\mathbb{P}_X(E)}(1)$  associated to the vector bundle  $E$  on  $X$ . The idea is that Segre classes generate Chern classes of vector bundles which in turn generate all numerical classes of cycles, via resolutions of structure sheaves of subschemes.

Now a Segre class  $\sigma$  such that  $\langle D^k \rangle_{X|V} \cdot \sigma \neq 0$  is represented by a subscheme  $Y \subset X$ , say of codimension  $d - k$ . As every subscheme  $Y$  is a component of a complete intersection  $A_1 \cap \dots \cap A_{d-k}$  of hyperplane sections  $A_i \in |A|$  for some  $l \gg 0$ . If  $A_1 \cap \dots \cap A_{d-k} = Y \cup Y'$  we have

$$\begin{aligned} l^{d-k} \cdot \langle D^k \rangle_{X|V} \cdot A^{d-k} &= \langle D^k \rangle_{X|V} \cdot Y + \langle D^k \rangle_{X|V} \cdot Y' \\ &\geq \langle D^k \rangle_{X|V} \cdot Y = \langle D^k \rangle_{X|V} \cdot \sigma > 0 \end{aligned}$$

since the intersection of a restricted product with an effective cycle of the correct dimension is non-negative, by construction.

The opposite direction is obvious.  $\square$

**Proposition 3.4.** *Let  $X$  be an  $n$ -dimensional smooth projective complex variety,  $V = H_1 \cap \dots \cap H_{n-k} \subset X$  a  $k$ -dimensional complete intersection subvariety cut out by very general free big and nef divisors  $H_i$  linearly equivalent to  $H$  and  $A, B$   $V$ -big and nef  $\mathbb{R}$ -divisors. Then:*

$$\text{vol}_{X|V}(A - B) \geq A^k \cdot H^{n-k} - k \cdot A^{k-1} \cdot B \cdot H^{n-k}.$$

*Proof.* This is a generalisation of [Laz04, Thm.2.2.15]. By continuity of the usual intersection product it is enough to choose an ample divisor  $H'$  and prove the inequality for  $A + \epsilon H', B + \epsilon H'$ , that is for ample  $\mathbb{R}$ -divisors  $A, B$ . Since the restricted volume is continuous and homogeneous on the cone spanned by the classes of ample divisors, we can even assume that  $A, B$  are very ample divisors.

Let us fix  $m > 0$  and choose  $m$  general divisors  $B_1, \dots, B_m \in |B|$ . Then we have a commutative diagram

$$\begin{array}{ccccc} 0 & \longrightarrow & H^0(X, \mathcal{O}_X(mA - \sum_{i=1}^m B_i)) & \longrightarrow & H^0(X, \mathcal{O}_X(mA)) & \longrightarrow & \bigoplus_{i=1}^m H^0(B_i, \mathcal{O}_{B_i}(mA)) \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & H^0(X|V, \mathcal{O}_X(mA - \sum_{i=1}^m B_i)) & \longrightarrow & H^0(X|V, \mathcal{O}_X(mA)) & \longrightarrow & \bigoplus_{i=1}^m H^0(B_i|V \cap B_i, \mathcal{O}_{B_i}(mA)) \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & H^0(V, \mathcal{O}_V(mA - \sum_{i=1}^m B_i)) & \longrightarrow & H^0(V, \mathcal{O}_V(mA)) & \longrightarrow & \bigoplus_{i=1}^m H^0(V \cap B_i, \mathcal{O}_{V \cap B_i}(mA)) \end{array}$$

where in the upper row the vertical arrows correspond to surjective maps whereas in the lower row the vertical arrows correspond to inclusions. Consequently,

$$h^0(X|V, \mathcal{O}_X(m(A - B))) \geq h^0(X|V, \mathcal{O}_X(mA)) - \sum_{i=1}^m h^0(B_i|V \cap B_i, \mathcal{O}_{B_i}(mA)).$$

Dividing by  $\frac{m^k}{k!}$  and going to the limit  $m \rightarrow \infty$  we obtain

$$\begin{aligned}
\text{vol}_{X|V}(A - B) &\geq \text{vol}_{X|V}(A) - \sum_{i=1}^m \frac{k}{m} \text{vol}_{B_i|V \cap B_i}(A) \\
&= \langle A^k \rangle_{X|V} - \sum_{i=1}^m \frac{k}{m} \langle A^{k-1} \rangle_{B_i|V \cap B_i} \\
&= \langle A^k \rangle_X \cdot H^{n-k} - \sum_{i=1}^m \frac{k}{m} \langle A^{k-1} \rangle_{B_i} \cdot H^{n-k} \\
&= A^k \cdot H^{n-k} - k \cdot A^{k-1} \cdot B \cdot H^{n-k}
\end{aligned}$$

using the Generalised Fujita Theorem (see 1.5), Lemma 3.2 and the ampleness resp. freeness of  $A$ ,  $H$  and the  $B_i$ .  $\square$

In the following,  $D_1 \leq_V D_2$  means that the difference  $D_2 - D_1$  of the two  $\mathbb{R}$ -divisors  $D_1, D_2$  on  $X$  is effective and the support of  $D_1 - D_2$  does not contain the subvariety  $V \subset X$ .

**Proposition 3.5.** *Let  $X$  be an  $n$ -dimensional smooth projective complex variety,  $V = H_1 \cap \dots \cap H_{n-k} \subset X$  a  $k$ -dimensional complete intersection subvariety cut out by very general free big and nef divisors  $H_i$  linearly equivalent to  $H$  and  $B$  a big and nef  $\mathbb{R}$ -divisor such that  $B \leq_V H$ . If  $\gamma$  is an arbitrary divisor class such that  $H \pm \gamma$  is still nef then*

$$\text{vol}_{X|V}(B + t\gamma) \geq B^k \cdot H^{n-k} + k \cdot t \cdot B^{k-1} \cdot \gamma \cdot H^{n-k} - c \cdot t^2$$

for every  $0 \leq t \leq 1$  and some constant  $c > 0$  only depending on  $H^n$ .

*Proof.* This is a generalisation of [BFJ09, Cor.3.4]. As in [BFJ09, Cor.2.4] we can use the assumption that  $H \pm \gamma$  is nef to conclude that for  $0 \leq t \leq 1$  and some constant  $c' > 0$  only depending on  $H^n$ ,

$$(B + t\gamma)^k \cdot H^{n-k} \geq B^k \cdot H^{n-k} + k \cdot t \cdot B^{k-1} \cdot \gamma \cdot H^{n-k} - c' \cdot t^2,$$

by replacing  $\gamma$  with  $(H + \gamma) - H$  and using that  $H + \gamma \leq 2H$ . If we also write  $B + t\gamma$  as the difference of the two nef classes  $C := B + t(\gamma + H)$  and  $D := tH$  then we have furthermore

$$(B + t\gamma)^k \cdot H^{n-k} = (C - D)^k \cdot H^{n-k} \leq C^k \cdot H^{n-k} - k \cdot C^{k-1} \cdot D \cdot H^{n-k} + c'' \cdot t^2,$$

where  $c''$  once again only depends on  $H^n$ : Indeed,  $c''$  is controlled by  $C^i \cdot H^{n-i}$ ,  $0 \leq i \leq k-2$ , and we have  $C \leq 3H$ . Thus we have

$$C^k \cdot H^{n-k} - k \cdot C^{k-1} \cdot D \cdot H^{n-k} = B^k \cdot H^{n-k} + k \cdot t \cdot B^{k-1} \cdot \gamma \cdot H^{n-k} - (c' + c'')t^2.$$

The result follows by applying Prop. 3.4 to  $B + t\gamma = C - D$ .  $\square$

*Proof of Thm. 3.1.* Let  $H$  be a very general divisor linearly equivalent to the  $H_i$ , and assume that  $\alpha \leq_V H$  and  $H \pm \gamma$  is nef. If this is not the case replace  $\alpha, \gamma$  by multiples  $s\alpha, s\gamma$  with  $s > 0$  sufficiently small. The claim for  $\alpha, \gamma$  still follows, by homogeneity of restricted volumes and moving intersection numbers.

Let  $\beta$  be a nef divisor class on a  $V$ -birational model  $\phi : (\tilde{X}, \tilde{V}) \rightarrow (X, V)$  such that  $\beta \leq_{\tilde{V}} \phi^* \alpha$ , hence also  $\beta \leq_{\tilde{V}} \phi^* H$ . Since  $\tilde{V}$  is cut out by the big and nef divisors  $\phi^* H_i$  Prop. 3.5 shows

$$\text{vol}_{X|V}(\alpha + t\gamma) \geq \text{vol}_{\tilde{X}|\tilde{V}}(\beta + t\phi^* \gamma) \geq \beta^k \cdot (\phi^* H)^{n-k} + k \cdot t \cdot \beta^{k-1} \cdot \phi^* \gamma \cdot (\phi^* H)^{n-k} - c \cdot t^2$$

for every  $0 \leq t \leq 1$  and some constant  $c > 0$  only depending on  $H^n$ . Taking the supremum over all nef classes  $\beta \leq_{\tilde{\nu}} \phi^* \alpha$  yields

$$\text{vol}_{X|V}(\alpha + t\gamma) \geq \text{vol}_{X|V}(\alpha) + k \cdot t \cdot \langle \alpha^{k-1} \rangle_{X|V} \cdot \gamma - c \cdot t^2,$$

using Lem. 3.2 and the Generalised Fujita Theorem. This holds for every  $0 \leq t \leq 1$ , and in fact also for every  $-1 \leq t \leq 0$ , by replacing  $\gamma$  with  $-\gamma$ .

Exchanging the roles of  $\alpha + t\gamma \leq 2H$  and  $\alpha = (\alpha + t\gamma) + t \cdot (-\gamma)$  we obtain

$$\text{vol}_{X|V}(\alpha) \geq \text{vol}_{X|V}(\alpha + t\gamma) - k \cdot t \cdot \langle (\alpha + t\gamma)^{k-1} \rangle_{X|V} \cdot \gamma - c' \cdot t^2$$

for a constant  $c'$  possibly larger than  $c$  but still only depending on  $H^n$ . Combining the two inequalities shows that

$$\frac{d}{dt} \Big|_{t=0} \text{vol}_{X|V}(\alpha + t\gamma) = k \cdot \langle \alpha^{k-1} \rangle_{X|V} \cdot \gamma$$

as desired, since  $\langle (\alpha + t\gamma)^{k-1} \rangle_{X|V}$  converges to  $\langle \alpha^{k-1} \rangle_{X|V}$  if  $t \rightarrow 0$ .  $\square$

To prove  $\kappa_{\nu}(D) \leq \nu_{\text{alg}}(D)$  we finally need to connect divisorial Zariski decomposition and algebraic moving intersection product. For the Kähler intersection product this was done in [BDPP13, Thm.3.5].

**Proposition 3.6.** *Let  $X$  be a smooth projective complex variety and  $D$  a pseudo-effective  $\mathbb{R}$ -divisor. Then the negative part of the divisorial Zariski decomposition  $D = P_{\sigma}(D) + N_{\sigma}(D)$  can be calculated as*

$$N_{\sigma}(D) = D - \langle [D] \rangle_{\text{alg}}.$$

*Proof.* For an ample divisor  $A$ ,

$$\lim_{\epsilon \downarrow 0} (D + \epsilon A - \sum \sigma_{\Gamma}(D + \epsilon A)\Gamma) = D - \sum \sigma_{\Gamma}(D)\Gamma$$

by Def. 1.1 and

$$\lim_{\epsilon \downarrow 0} \langle [D + \epsilon A] \rangle_{\text{alg}} = \langle [D] \rangle_{\text{alg}}$$

by the definition of the moving intersection product in 1.2. Hence we only have to show the claimed equality for big  $\mathbb{R}$ -divisors.

By [Nak04, III.1.17(3)], for big  $\mathbb{R}$ -divisors  $B$  it is enough to show that

$$B - \langle [B] \rangle_{\text{alg}} = \lim_{m \rightarrow \infty} \frac{1}{m} |[mB]|_{\text{fix}}.$$

To this purpose let  $\pi_m : X_m \rightarrow X$  be a log resolution of  $|[mB]|$ . Then  $\pi_m^* B$  decomposes into

$$\pi_m^* B = \frac{1}{m} (|\pi_m^*([mB])|_{\text{fix}} + |\pi_m^*([mB])|_{\text{free}}) + \pi_m^* \Delta_m,$$

where  $\Delta_m := B - \frac{1}{m} [mB]$  is an effective  $\mathbb{R}$ -divisor. By construction of  $\langle [B] \rangle_{\text{alg}}$  (see 1.2) this implies

$$B - \langle [B] \rangle_{\text{alg}} \leq \pi_{m*} \left( \frac{1}{m} |\pi_m^*([mB])|_{\text{fix}} \right) + \Delta_m$$

and hence  $B - \langle [B] \rangle_{\text{alg}} \leq \lim_{m \rightarrow \infty} \frac{1}{m} |[mB]|_{\text{fix}}$  since  $\Delta_m \rightarrow 0$  if  $m \rightarrow \infty$ .

Vice versa, choose an ample divisor  $A$  on  $X$  and consider small enough  $\epsilon > 0$  such that  $B - \epsilon A$  is still big. As above, for all such  $\epsilon$  we can find birational morphisms  $\pi_{\epsilon} : X_{\epsilon} \rightarrow X$  between smooth projective varieties such that

$$\pi_{\epsilon}^*(B - \epsilon A) \equiv D_{\epsilon} + B_{\epsilon},$$

where  $D_\epsilon$  is an effective  $\mathbb{Q}$ -divisor and  $B_\epsilon$  a nef  $\mathbb{R}$ -divisor, and

$$\lim_{\epsilon \downarrow 0} \pi_{\epsilon*}(D_\epsilon) = B - \langle [B] \rangle_{\text{alg}}.$$

Next, we use that for these  $\epsilon$  there exist ample  $\mathbb{R}$ -divisors  $A_\epsilon$  on  $X_\epsilon$  such that  $\pi_\epsilon^* A - A_\epsilon$  is an effective  $\mathbb{Q}$ -divisor supported on  $\pi_\epsilon$ -exceptional divisors. Then

$$\pi_\epsilon^* B \equiv D_\epsilon + B_\epsilon + \epsilon \pi_\epsilon^* A = D'_\epsilon + B_\epsilon + \epsilon A_\epsilon$$

for an effective  $\mathbb{Q}$ -divisor  $D'_\epsilon$  such that  $\pi_{\epsilon*}(D_\epsilon) = \pi_{\epsilon*}(D'_\epsilon)$ . Since the difference  $(1 + \epsilon)\pi_\epsilon^* B - D'_\epsilon - (B_\epsilon + \epsilon A_\epsilon)$  is big there exist effective  $\mathbb{R}$ -divisors  $\Delta_{m,\epsilon} \equiv \epsilon \pi_\epsilon^* B$  such that

$$m(1 + \epsilon)\pi_\epsilon^* B \sim mD'_\epsilon + m(B_\epsilon + \epsilon A_\epsilon) + m\Delta_{m,\epsilon},$$

for sufficiently large  $m$ . Since  $B_\epsilon + \epsilon A_\epsilon$  is ample this implies

$$\frac{1}{\lceil m(1 + \epsilon) \rceil} \lceil [m(1 + \epsilon)] \pi_\epsilon^* B \rceil|_{\text{fix}} \leq \frac{1}{m} \lceil [m(1 + \epsilon)] \pi_\epsilon^* B \rceil|_{\text{fix}} \leq D'_\epsilon + \Delta_{m,\epsilon}$$

for sufficiently large  $m$ . This implies  $\lim_{m \rightarrow \infty} \frac{1}{m} \lceil [m\pi_\epsilon^* B] \rceil|_{\text{fix}} \leq D'_\epsilon + \Delta_{m,\epsilon}$ , hence  $\lim_{m \rightarrow \infty} \frac{1}{m} \lceil [mB] \rceil|_{\text{fix}} \leq \pi_{\epsilon*}(D'_\epsilon)$  because  $\pi_\epsilon^*(\lceil [mB] \rceil) \leq \lceil [m\pi_\epsilon^* B] \rceil$ . Consequently,

$$\lim_{m \rightarrow \infty} \frac{1}{m} \lceil [mB] \rceil|_{\text{fix}} \leq B - \langle [B] \rangle_{\text{alg}}.$$

□

*Proof of  $\kappa_\nu(D) \leq \nu_{\text{alg}}(D)$ .* First assume that  $\nu_{\text{alg}}(D) = 0$ . By definition this means that the positive product  $\langle D \rangle_{\text{alg}} = 0$ , hence  $P_\sigma(D) \equiv 0$  and  $D \equiv N_\sigma(D)$  by Prop. 3.6. Consequently,  $\kappa_\nu(D) = 0$  by [Nak04, V.2.22(2)].

So from now on we assume  $1 \leq k := \nu_{\text{alg}}(D) < \kappa_\nu(D) \leq n := \dim X$  and derive a contradiction: Let  $W$  be a  $k$ -dimensional intersection of very general very ample divisors. Set  $\phi : Y \rightarrow X$  to be the blow up of  $X$  along  $W$ , with exceptional divisor  $E$ . Fix a very ample divisor  $H$  on  $Y$ . By [Nak04, V.2.21]  $k < \kappa_\nu(D)$  implies that  $D \succeq W$ , that is, for each sufficiently small  $\epsilon > 0$  there exists a  $\tau > 0$  such that  $\phi^* D - \tau E + \epsilon H$  is pseudoeffective and  $\frac{\tau}{\epsilon} \rightarrow \infty$  when  $\epsilon \rightarrow 0$ .

Fix  $\epsilon > 0$ . Choose birational models  $\psi_i : (\tilde{Y}_i, \tilde{E}_i) \rightarrow (Y, E)$  on which both the restricted product  $\langle (\phi^* D + \epsilon H)^k \rangle_{Y|E}$  and the product  $\langle (\phi^* D + \epsilon H)^{k+1} \rangle_Y$  can be computed. Choose big and nef divisors

$$A_i \leq_{\tilde{E}_i} \psi_i^*(\phi^* D + \epsilon H)$$

on  $\tilde{Y}_i$  such that the limit of the push forwards  $\psi_{i*}(A_i^k \cdot \tilde{E}_i)$  calculates the first product.

As in the proof of [Leh13, Prop.5.5] we can conclude that  $P_\sigma(\phi^* D + \epsilon H) - \tau E$  and hence  $\psi_i^* P_\sigma(\phi^* D + \epsilon H) - \tau \psi_i^* E$  are also pseudoeffective. Since by [Nak04, III.5.16]

$$\psi_i^* N_\sigma(\phi^* D + \epsilon H) \geq N_\sigma(\psi_i^* P_\sigma(\phi^* D + \epsilon H))$$

we also have that  $P_\sigma(\psi_i^*(\phi^* D + \epsilon H)) - \tau \psi_i^* E$  is pseudoeffective. [Leh13, Prop.3.5&3.7] tell us that for a suitable effective divisor  $G$  on  $Y$  we can further assume that the big and nef divisors  $A_i$  from above satisfy

$$A_i \leq_{\tilde{E}_i} P_\sigma(\psi_i^*(\phi^* D + \epsilon H)) \leq_{\tilde{E}_i} A_i + \frac{1}{i} \psi_i^* G.$$

In particular,  $A_i + \frac{1}{i}\psi_i^*G - \tau\tilde{E}_i$  is pseudoeffective. Therefore,

$$\begin{aligned} 0 &\leq (A_i + \frac{1}{i}\psi_i^*G - \tau\tilde{E}_i) \cdot A_i^k \cdot \psi_i^*H^{n-k-1} = \\ &= A_i^{k+1} \cdot \psi_i^*H^{n-k-1} + \frac{1}{i}\psi_i^*G \cdot A_i^k \cdot \psi_i^*H^{n-k-1} - \tau\tilde{E}_i \cdot A_i^k \cdot \psi_i^*H^{n-k-1}. \end{aligned}$$

By definition,

$$0 \leq A_i^{k+1} \cdot \psi_i^*H^{n-k-1} \leq \langle (\phi^*D + \epsilon H)^{k+1} \rangle_Y \cdot H^{n-k-1}$$

and

$$0 \leq A_i^k \cdot \psi_i^*G \cdot \psi_i^*H^{n-k-1} \leq \langle (\phi^*D + \epsilon H)^k \rangle_Y \cdot G \cdot H^{n-k-1}.$$

So taking the limit over all models  $\tilde{Y}_i$  we obtain

$$(3.1) \quad 0 \leq \langle (\phi^*D + \epsilon H)^{k+1} \rangle_Y \cdot H^{n-k-1} - \tau \langle (\phi^*D + \epsilon H)^k \rangle_{Y|E} \cdot H^{n-k-1}.$$

If  $V = H_1 \cap \dots \cap H_{n-k-1} \subset Y$  is a  $(k+1)$ -dimensional complete intersection subvariety cut out by  $n-k-1$  very general very ample divisors  $H_i \in |H|$ , Thm. 3.1 and Lem. 3.2 imply that

$$\begin{aligned} \frac{d}{dt} \Big|_{t=0} \langle (\phi^*D + (\epsilon + t)H)^{k+1} \rangle_{Y|V} &= (k+1) \cdot \langle (\phi^*D + \epsilon H)^k \rangle_{Y|V} \cdot H = \\ &= (k+1) \cdot \langle (\phi^*D + \epsilon H)^k \rangle_Y \cdot H^{n-k}. \end{aligned}$$

Furthermore, by definition  $\lim_{\epsilon \downarrow 0} \langle (\phi^*D + \epsilon H)^k \rangle_Y = \langle (\phi^*D)^k \rangle_Y$ , and the assumption  $\nu_{\text{alg}}(D) = k$  implies  $\langle (\phi^*D)^k \rangle_Y \cdot H^{n-k} > 0$  by Lem. 3.3 whereas  $\langle (\phi^*D)^{k+1} \rangle_Y = 0$ . Consequently, there exists  $c > 0$  such that  $\langle (\phi^*D + \epsilon H)^{k+1} \rangle_Y \cdot H^{n-k-1} \leq c \cdot \epsilon$ . Then (3.2) implies that

$$\tau \leq \frac{\langle (\phi^*D + \epsilon H)^{k+1} \rangle_Y \cdot H^{n-k-1}}{\langle (\phi^*D + \epsilon H)^k \rangle_{Y|E} \cdot H^{n-k-1}} \leq \epsilon \cdot \frac{c}{\langle (\phi^*D + \epsilon H)^k \rangle_{Y|E} \cdot H^{n-k-1}}.$$

The denominator of the right-hand side fraction tends to  $\langle (\phi^*D)^k \rangle_{Y|E} \cdot H^{n-k-1}$  if  $\epsilon \rightarrow 0$ . By choosing sufficiently general elements  $H_1, \dots, H_{n-k-1} \in |H|$  we may assume that  $\phi$  restricted to  $E \cap H_1 \dots \cap H_{n-k-1}$  is a finite morphism onto  $W$ . If  $A_1, \dots, A_{n-k}$  denote the very ample divisors on  $X$  cutting out  $W$  there exists  $C > 0$  such that

$$\begin{aligned} \langle (\phi^*D)^k \rangle_{Y|E} \cdot H^{n-k-1} &= \langle (\phi^*D)^k \rangle_{Y|E \cap H_1 \dots \cap H_{n-k-1}} = C \cdot \langle D^k \rangle_{X|W} = \\ &= C \cdot \langle D^k \rangle_X \cdot A_1 \cdots A_{n-k} \end{aligned}$$

where the first and the last equality follow from Lem. 3.2 and the middle equality from [Leh13, Prop.4.20]. By assumption and Lem. 3.3 this last product is positive, contradicting the unboundedness of  $\frac{\tau}{\epsilon}$  for  $\epsilon \rightarrow 0$ .  $\square$

#### 4. BIRATIONAL INVARIANCE AND ABUNDANCE CONJECTURE

To prove that the Abundance Conjecture 0.5 is equivalent to the MMP-version of the Abundance Conjecture 0.6 we need the birational invariance of the numerical dimension of the canonical bundle:

**Proposition 4.1.** *Let  $X$  be a smooth projective complex variety and  $D$  a pseudoeffective divisor on  $X$ . Let  $f : Y \rightarrow X$  be a birational morphism of smooth projective varieties and  $E$  an  $f$ -exceptional effective  $\mathbb{R}$ -divisor on  $Y$ . Then:*

$$\nu_X(D) = \nu_Y(f^*D + E).$$

*Proof.* Let  $E_1, \dots, E_k$  be the prime components of  $E = \sum_{i=1}^k x_i E_i$ ,  $x_i > 0$ . Assume first that  $D$  is big. Let  $\Delta$  be an effective  $\mathbb{R}$ -divisor  $\equiv f^*D + E$  on  $Y$ .

*Claim.*  $\text{mult}_{E_i} \Delta \geq \text{mult}_{E_i}(E) = x_i$  for all  $i = 1, \dots, k$ .

*Proof of Claim.* If  $\text{mult}_{E_j} \Delta < \text{mult}_{E_j}(E)$  for a  $j \in \{1, \dots, k\}$  we subtract a multiple of  $E_j$  from  $\Delta$  and  $E$  to obtain  $\Delta' \geq 0$ ,  $E' \geq 0$  such that  $\Delta' \equiv f^*D + E'$  and  $0 = \text{mult}_{E_j} \Delta' < \text{mult}_{E_j} E'$ . Pushing forward  $\Delta'$  we have  $f_* \Delta' \equiv D$ . Hence for real numbers  $y_i$  such that  $y_j \geq 0$ ,

$$f^*D \equiv f^* f_* \Delta' = \Delta' + \sum_{i=1}^k y_i E_i \equiv f^*D + \sum_{i=1}^k y_i E_i + E'.$$

Thus,  $\sum_{i=1}^k y_i E_i + E'$  is a non-trivial linear combination of the  $E_i$  numerically equivalent to 0. But this is impossible as numerical classes of  $f$ -exceptional prime divisors are always linearly independent: On  $X$ , sufficiently general complete intersection curves  $C$  avoid all centers  $f(E_i)$  but one, hence the strict transform  $\bar{C} \subset Y$  intersects the corresponding prime divisor on  $Y$  but none else.  $\square$

The claim implies that  $\sigma_{E_i}(f^*D + E) \geq \text{mult}_{E_i}(E)$ . Taking the limit this also holds when  $D$  is only pseudoeffective. Hence  $E \leq N_\sigma(f^*D + E)$ , this implies  $N_\sigma(f^*D + E) - E = N_\sigma(f^*D)$ , and

$$P_\sigma(f^*D + E) = (f^*D + E) - N_\sigma(f^*D + E) = f^*D - N_\sigma(f^*D) = P_\sigma(f^*D).$$

The same holds when  $\phi : \tilde{Y} \rightarrow Y$  is a further birational morphism between smooth projective varieties:

$$P_\sigma(\phi^*(f^*D + E)) = P_\sigma(\phi^*f^*D + \phi^*E) = P_\sigma(\phi^*f^*D).$$

Using that the numerical dimension can be defined by  $\nu_{\text{Vol,Zar}}$  (see 1.6 and Def. 0.3) this implies  $\nu_Y(f^*D + E) = \nu_Y(f^*D)$ . Defining the numerical dimension via positive intersection products as  $\nu_{\text{alg}}$  shows that  $\nu_Y(f^*D) = \nu_X(D)$ , together with the projection formula and the fact that  $f^*$  defines a homomorphism on the intersection rings.  $\square$

*Remark 4.2.* The proof above also shows that  $\nu_Y(f^*D) = \nu_Y(f^*D + E)$  for a pseudoeffective  $\mathbb{Q}$ -divisor  $D$  on  $X$  and an effective  $f$ -exceptional divisor  $E$  on  $Y$  even when  $X$  is not smooth but only  $\mathbb{Q}$ -factorial.

**Corollary 4.3.** *Let  $X$  be a non-uniruled smooth projective complex variety and  $f : Y \rightarrow X$  a birational morphism between smooth projective varieties. Then:*

$$\nu_X(K_X) = \nu_Y(K_Y).$$

*Proof.* By [BDPP13, Cor.0.3] the canonical divisors  $K_X$  and  $K_Y$  are pseudoeffective on the non-uniruled varieties  $X$  and  $Y$ . Hence it is possible to calculate their numerical dimension. Since the pullback of canonical forms through a birational morphism is again a canonical form, there exists an effective  $f$ -exceptional divisor  $E$  such that  $K_Y = f^*K_X + E$ . The corollary follows from Prop. 4.1.  $\square$

A *minimal model* of a non-uniruled smooth projective complex variety  $X$  is a normal variety  $S$  such that there exists a sequence of divisorial contractions and flips

$$X = X_0 \dashrightarrow X_1 \dashrightarrow \dots \dashrightarrow X_n = S$$

and  $K_S$  is nef (see e.g. [Mat02, Def.3-3-1] and passim for further definitions). In particular,  $S$  is  $\mathbb{Q}$ -factorial and has only terminal singularities, that is, every Weil

divisor on  $S$  is a  $\mathbb{Q}$ -Cartier divisor and if  $f : Y \rightarrow S$  is a birational morphism from a smooth projective variety  $Y$  then in the ramification formula

$$K_Y = f^*K_S + \sum a_i E_i,$$

the coefficients  $a_i$  of all the  $f$ -exceptional prime divisors  $E_i$  are  $> 0$ .

Note that on a minimal model  $S$  it is possible to construct intersection products of ( $\mathbb{Q}$ -)Cartier divisors and to define the numerical triviality of the resulting (rational) cycles (see [Ful84, 19.1]). Hence it makes sense to set the numerical dimension of a nef ( $\mathbb{Q}$ -)Cartier divisor  $D$  on  $S$  equal to

$$\nu_S(D) := \max\{k : D^k \neq 0\}.$$

If  $S$  is smooth this numerical dimension coincides with the one defined in Def. 0.3, by construction of positive intersection products (see 1.2).

The following result of Kawamata [Kaw85] sits at the core of the proof that the two versions of the Abundance Conjecture are equivalent:


**Theorem 4.4** (Kawamata). *On a minimal model  $S$ ,  $\kappa_S(K_S) = \nu_S(K_S)$  if and only if  $K_S$  is semi-ample.*

**Theorem 4.5.** *Let  $S$  be a minimal model of a non-uniruled smooth projective complex variety  $X$ . Then*

$$\nu_X(K_X) = \kappa_X(K_X) \iff K_S \text{ is semi-ample.}$$

*Proof.* By Kawamata's Theorem we only need to prove that  $\kappa_X(K_X) = \kappa_S(K_S)$  and  $\nu_X(K_X) = \nu_S(K_S)$ .

The first equality follows from using a common resolution



of  $X$  and  $S$  such that  $K_Y = f^*K_S + E_S = \phi^*K_X + E_X$  where  $E_S$  and  $E_X$  are  $f$ - resp.  $\phi$ -exceptional effective divisors. Since the sections in  $H^0(Y, \mathcal{O}_Y(mK_Y)) = H^0(Y, \mathcal{O}_Y(m\phi^*K_X + mE_X))$  can be interpreted both as rational functions on  $X$  and  $Y$ , we have  $H^0(Y, \mathcal{O}_Y(mK_Y)) \subset H^0(X, \mathcal{O}_X(mK_X))$ , and since  $E_X$  is effective the inverse inclusion also holds. Similarly on  $S$ , and the equality follows.

For  $\nu_X(K_X) = \nu_S(K_S)$  we use Cor. 4.3 and Rem. 4.2 to deduce the chain of equalities

$$\nu_X(K_X) = \nu_Y(K_Y) = \nu_Y(f^*K_S + E_S) = \nu_Y(f^*K_S) = \nu(K_S)$$

where the last equality follows from the projection formula and the fact that  $f^*$  defines a homomorphism on the intersection rings.  $\square$

REFERENCES

[BDPP13] Sébastien Boucksom, Jean-Pierre Demailly, Mihai Păun, and Thomas Peternell. The pseudo-effective cone of a compact Kähler manifold and varieties of negative Kodaira dimension. *J. Algebraic Geom.*, 22(2):201–248, 2013.

[BFJ09] Sébastien Boucksom, Charles Favre, and Mattias Jonsson. Differentiability of volumes of divisors and a problem of Teissier. *J. Algebraic Geom.*, 18(2):279–308, 2009.

[Bou02] S. Boucksom. *Cônes positifs des variétés complexes compactes*. PhD thesis, Grenoble, 2002.

- [Bou04] Sébastien Boucksom. Divisorial Zariski decompositions on compact complex manifolds. *Ann. Sci. École Norm. Sup. (4)*, 37(1):45–76, 2004.
- [DHP13] Jean-Pierre Demailly, Christopher D. Hacon, and Mihai Păun. Extension theorems, non-vanishing and the existence of good minimal models. *Acta Math.*, 210(2):203–259, 2013.
- [ELM<sup>+</sup>06] Lawrence Ein, Robert Lazarsfeld, Mircea Mustăţă, Michael Nakamaye, and Mihnea Popa. Asymptotic invariants of base loci. *Ann. Inst. Fourier (Grenoble)*, 56(6):1701–1734, 2006.
- [ELM<sup>+</sup>09] Lawrence Ein, Robert Lazarsfeld, Mircea Mustăţă, Michael Nakamaye, and Mihnea Popa. Restricted volumes and base loci of linear series. *Amer. J. Math.*, 131(3):607–651, 2009.
- [Ful84] W. Fulton. *Intersection Theory*. Ergebnisse der Mathematik und ihrer Grenzgebiete 3/2. Springer, Heidelberg, 1984.
- [GL13] Y. Gongyo and B. Lehmann. Reduction maps and minimal model theory. *Compos. Math.*, 149(2):295–308, 2013.
- [Kaw85] Y. Kawamata. Pluricanonical systems on minimal algebraic varieties. *Invent. Math.*, 79(3):567–588, 1985.
- [Laz04] R. Lazarsfeld. *Positivity in Algebraic Geometry I*. Erg. Math. 48. Springer, New York, 2004.
- [Leh13] Brian Lehmann. Comparing numerical dimensions. *Algebra Number Theory*, 7(5):1065–1100, 2013.
- [Mat02] K. Matsuki. *Introduction to the Mori program*. Springer Verlag, 2002.
- [Nak04] N. Nakayama. *Zariski-decomposition and abundance*, volume 14 of *MSJ Memoirs*. Mathematical Society of Japan, Tokyo, 2004.
- [Siu11] Yum-Tong Siu. Abundance conjecture. In *Geometry and analysis. No. 2*, volume 18 of *Adv. Lect. Math. (ALM)*, pages 271–317. Int. Press, Somerville, MA, 2011.

THOMAS ECKL, DEPARTMENT OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF LIVERPOOL,  
 MATHEMATICAL SCIENCES BUILDING, LIVERPOOL, L69 7ZL, ENGLAND, U.K.

*E-mail address:* thomas.eckl@liv.ac.uk

*URL:* <http://pcwww.liv.ac.uk/~eckl/>