

Article (refereed) - postprint

Smart, Simon M.; Ellison, Aaron M.; Bunce, Robert G.H.; Marrs, Robert H.; Kirby, Keith J.; Kimberley, Adam; Scott, Andy W.; Foster, David R.. 2014.

Quantifying the impact of an extreme climate event on species diversity in fragmented temperate forests: the effect of the October 1987 storm on British broadleaved woodlands. *Journal of Ecology*, 102 (5). 1273-1287.

[10.1111/1365-2745.12291](https://doi.org/10.1111/1365-2745.12291)

© 2014 The Authors. *Journal of Ecology* © 2014 British Ecological Society

This version available <http://nora.nerc.ac.uk/509702/>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the rights owners. Users should read the terms and conditions of use of this material at

<http://nora.nerc.ac.uk/policies.html#access>

This document is the author's final manuscript version of the journal article, incorporating any revisions agreed during the peer review process. Some differences between this and the publisher's version remain. You are advised to consult the publisher's version if you wish to cite from this article.

The definitive version is available at <http://onlinelibrary.wiley.com>

Contact CEH NORA team at
noraceh@ceh.ac.uk

1 **Quantifying the impact of an extreme climate event on species diversity in fragmented**
2 **temperate forests: the effect of the October 1987 storm on British broadleaved woodlands**

3
4 Smart, SM^{1*}, Ellison, AM², Bunce, RGH³, Marrs, RH⁴, Kirby, KJ⁵, Kimberley, A¹, Scott, WA¹
5 and Foster, DR²

6
7 1 NERC Centre for Ecology & Hydrology, Library Avenue, Bailrigg, Lancaster LA1 4AP UK

8 2 Harvard Forest, Harvard University, 324 North Main St, Petersham, MA 01366, USA

9 3 Estonian University of Life Sciences, Freidrich Reinhold Kreutzwaldi 1, 51014 Tartu, Estonia

10 4 Ecology and Ocean Sciences, Nicholson Building, University of Liverpool, L69 3GP UK

11 5 Department of Plant Sciences, South Parks Road, Oxford, OX1 3RB, UK

12
13 *Correspondence author. E-mail: ssma@ceh.ac.uk

14
15 **SUMMARY**

- 16 1. We report the impact of an extreme weather event, the October 1987 severe storm, on
17 fragmented woodlands in southern Britain. We analysed ecological changes between
18 1971 and 2002 in 143 200-m² plots in 10 woodland sites exposed to the storm with an
19 ecologically equivalent sample of 150 plots in 16 non-exposed sites. In both years,
20 understorey species-richness, species composition, soil pH and woody basal area of the
21 tree and shrub canopy were measured.
- 22 2. We tested the hypothesis that the storm had deflected sites from the wider national
23 trajectory of an increase in woody basal area and reduced understorey species-richness
24 associated with ageing canopies and declining woodland management. We also expected
25 storm disturbance to amplify the background trend of increasing soil pH, a UK-wide
26 response to reduced atmospheric sulphur deposition. Path analysis was used to quantify
27 indirect effects of storm exposure on understorey species richness via changes in woody
28 basal area and soil pH.
- 29 3. By 2002, storm exposure was estimated to have increased mean species richness per 200
30 m² by 32%. Woody basal area changes were highly variable and did not significantly
31 differ with storm exposure.

- 1 4. Increasing soil pH was associated with a 7% increase in richness. There was no evidence
2 that soil pH increased more as a function of storm exposure. Changes in species richness
3 and basal area were negatively correlated: a 3.4% decrease in richness occurred for every
4 0.1-m² increase in woody basal area per plot.
- 5 5. Despite all sites substantially exceeding the empirical critical load for nitrogen
6 deposition, there was no evidence that in the 15 years since the storm, disturbance had
7 triggered a eutrophication effect associated with dominance of gaps by nitrophilous
8 species.
- 9 6. *Synthesis*: Although the impacts of the 1987 storm were spatially variable in terms of
10 impacts on woody basal area, the storm had a positive effect on understorey species
11 richness. There was no evidence that disturbance had increased dominance of gaps by
12 invasive species. This could change if recovery from acidification results in a soil pH
13 regime associated with greater macronutrient availability.

14

15 **Key-words:** Bayesian Structural Equation Modelling, biodiversity, extreme weather, global
16 change, land-use, mixed models, path analysis, plant traits, resilience

17

18

1 **INTRODUCTION**

2

3 Ecosystems embedded in densely populated landscapes are increasingly exposed to novel
4 combinations of stressors, including pollutant deposition, land-use and climatic change (Foster *et*
5 *al.* 1997; Verheyen *et al.* 2012). Predicting the impacts of these changes on biodiversity and
6 ecosystem function requires quantification of the responses of ecosystems to these anthropogenic
7 press disturbance regimes and how they interact with pulse disturbances such as extreme weather
8 impacts, to generate potentially novel outcomes (Smith *et al.* 2009).

9 Important insights can come from analysis of natural perturbations that have operated in
10 combination with other factors (e.g., Bruelheide & Luginbühl 2009; Romme *et al.* 2011). This
11 depends on the serendipitous availability of data before and after the event in control and
12 impacted areas and where like-with-like contrasts can be constructed (Flinn & Vellend 2005). In
13 Britain, an ideal example is provided by the October 1987 storm. Its impact on broadleaved
14 woodland ecosystems was partially captured by a detailed national ecological survey of 16 200-
15 m² plots in each of 103 woodlands carried out in 1971 and repeated in 2002 (Kirby *et al.* 2005a;
16 Corney *et al.* 2006). Ten of the 103 woodlands were exposed to the October 1987 storm (Fig. 1).
17 The availability of a regional series of sites not exposed to the storm allowed the selection of
18 reference woodlands for comparison. We focus on the response of the understorey because in
19 temperate forests this is where most of the plant diversity is concentrated (Flinn & Vellend
20 2005). The biodiversity of the understorey in ancient woodlands also is of high conservation
21 value and is often different from secondary woodlands of more recent origins (Peterken & Game
22 1984; Motzkin *et al.* 1999; Kimberley *et al.* 2014).

23 The 1987 storm event was typified by wind speeds thought only to be likely every 200
24 years and locally gusting to 160 kph (Burt & Mansfield 1988). An estimated 15 million trees

1 were blown down across southeast England. Damage was locally severe; however wind speeds
2 and the extent of damage to trees were variable within the storm-track, resulting from context
3 dependent interactions between topography, tree species, form, age, substrate and variation in
4 wind strength (Whitbread 1991; Hopkins 1994; Harmer 2012). The variation in impact and the
5 difficulty in explaining its source within the storm-track was summarized by Peterken (1996) “..
6 storm damage generally appeared to be random and patchy at all scales. Some districts were
7 devastated, whilst others within the storm-track were virtually untouched. Some very exposed
8 stands escaped with little more than superficial branch-break. It was rarely possible to find a
9 reason why one tree within a wood fell while its neighbors survived.”

10 Previous analysis of all 103 woodland sites showed a widespread suppressive effect of
11 increased shading on understorey plant species density as woodland canopies aged following
12 intensive timber removal across many British forests during and just after the end of World War
13 II (Kirby *et al.* 2005a). This pattern was associated with a mean loss of eight species per plot
14 across the national sample (Kirby *et al.* 2005a).

15 Analysis also showed a national increase in woodland soil pH (from a mean of 4.98 to
16 5.31 between the 1971 and 2002 surveys (Kirby *et al.* 2005a) consistent with recovery of soils
17 following reductions in atmospheric sulphur deposition since the early 1970s (Kirby *et al.* 2005a;
18 Kirk *et al.* 2006). Soil pH increased less where woody basal area had increased the most, a
19 pattern consistent with the build up of soil organic matter with shading and succession and a
20 proportionally greater input from higher C:N tree leaf litter and woody debris.

21 These large-scale changes in soils, land-use, and atmospheric deposition in British
22 woodlands define the ecological context against which we test our primary hypothesis: that the
23 October 1987 storm changed sites in the storm-track away from the national trajectories of

1 canopy growth and reduction in understorey species-richness. Since we were interested in how
2 storm exposure impacted species richness and species and trait composition via changes in soil
3 and canopy, a path analysis was constructed and tested (Fig. 2).

4 We also tested whether the interaction between storm disturbance, the chronic effects of
5 long term increases in nitrogen deposition and the reduction in acidifying sulphur deposition
6 since the early 1970s in Britain and other parts of Europe (RoTAP 2012) had driven changes in
7 understorey species composition and whether this had tended to homogenize the flora between
8 storm-impacted sites. For example, had gap creation within the storm track triggered a
9 eutrophication effect leading to dominance by rapidly growing generalist species with high
10 specific leaf areas (SLA), including non-natives (Gilliam 2006; Verheyen *et al.* 2012). This
11 follows from the novelty of patch conditions and forest context following the 1987 storm relative
12 to the landscape of pre-industrial Britain. Small forests surrounded by intensive land-use make it
13 more likely that gaps will be colonised by edge species favoured by exposure to macronutrient
14 surpluses and more suited to modern land-use in addition to residual vegetation typical of older
15 forest (Smart *et al.* 2005; Smart *et al.* 2006a,b; Kimberley *et al.* 2013). Nitrogen limitation in the
16 understorey is also likely to have been alleviated by atmospheric deposition, although its impact
17 on the vegetation will depend upon the pH of the substrate and phosphorus availability (Stevens
18 *et al.* 2011; Verstraeten *et al.* 2013). Enrichment impacts on the understorey were analysed by
19 quantifying and interpreting differences in the species composition of the understorey in 1971
20 versus 2002 between storm exposed plots and those outside the storm-track and by quantifying
21 changes in cover-weighted SLA given the positive association between this trait and more
22 productive soils (Kimberley *et al.* 2014; Laughlin 2011).

1 Our hypotheses find support from within three conceptual frameworks. Together they
2 describe the outcomes of natural disturbance factors interacting with global change drivers to
3 impact understorey species composition in fragmented forests embedded in the often intensively
4 farmed landscapes of the temperate zone. The model of Roberts (2004) considers the response of
5 the woodland understorey as a function of the severity of disturbance to soil, canopy and
6 herbaceous vegetation. The partitioning of effects among these three ecosystem compartments
7 aligns well with the driving variables and their postulated linkage via regression equations in our
8 path model (Fig 2). The storm event is the key exogenous disturbance whose impacts we
9 hypothesise to be propagated through to change in understorey species richness and composition.
10 The principal effect of the storm is expected to be via canopy damage leading to a reduction in
11 woody basal area (Fig 2, β_3 and β_4) but direct residual effects are also possible where for
12 example canopy removal or damage to individuals is not detectable via basal area change (Fig 2,
13 β_2). Soil disturbance and its effect on the understorey is considered in terms of the relationships
14 between changing soil pH between surveys and change in species richness or cover-weighted
15 Specific Leaf Area (Fig 2, β_6). Soil pH change is then modelled as a function of background
16 variation in pH (Fig 2, β_7), the impact of storm disturbance (Fig 2, β_9) and change in woody
17 basal area (Fig 2, β_3 and β_5).

18 The Roberts (2004) model provides a foundation for understanding the impact of natural
19 disturbance agents within human-dominated landscapes but does not explicitly consider global
20 change drivers. Their impacts on the woodland ecosystem are considered in terms of the
21 Hierarchical Response Framework (HRF) of Smith *et al.* (2009). The critical insight here is that
22 ecosystems experience pulsed changes in resources within natural limits to which the biota is
23 adapted and where local ecosystem feedbacks can exert control over these changes in resource

1 availability. Anthropogenic activity introduces press rather than pulse regimes involving
2 directional, chronic changes in resource availability from land-use, population growth and
3 atmospheric pollutants (Smith *et al.* 2009). Local ecosystem feedbacks cannot moderate these
4 changes in resource availability because the origins of these resource inputs are geographically
5 distant from the impacted ecosystem. The consequence is biotic change which maybe rapid and
6 stepwise when triggered by interaction with a pulse of disturbance. In forest ecosystems for
7 example, storm events remove the canopy temporarily reducing the influence of the dominant
8 plant species on ecosystem processes. In our sample of woodlands, understorey development in
9 post-storm gaps could reflect an interaction with high cumulative N deposition and recently
10 reduced deposition of acidifying pollutants. We speculate that a small number of generalist
11 winners more typical of the surrounding farmed landscape could have increased in abundance at
12 the expense of forest specialist ‘losers’ and that non-random filtering has occurred preferentially
13 on storm-exposed sites as a result of the alleviation of light and nutrient limitation. This
14 conjecture includes aspects of the *biodiversity* and *biogeochemistry hypotheses* in Smith *et al.*
15 (2009). If the same pattern is seen across sites then this would also be consistent with the notion
16 of biotic homogenisation (McKinney & Lockwood, 1999; Baeten *et al.* 2012).

17 Lastly, we test hypotheses that concern the influence of the background species pool and
18 within-site beta diversity on change in local species richness. Sites with greater beta diversity
19 and a larger species pool might be expected to provide a greater pool of potential colonists for
20 exploiting the mosaic of abiotic conditions resulting from storm disturbance thus leading to
21 higher richness in individual plots post-storm. In non-storm sites, higher beta diversity in 1971
22 could result in a larger reduction in mean richness per plot if shading and lack of management
23 filters the understorey favouring a smaller number of residual shade tolerators. Thus the

1 relationship between beta diversity and changing species richness in plots is likely to be
2 dependent on storm disturbance and the extent to which within-site heterogeneity correlates with
3 species pool size. For example, in European forests changes in beta diversity tend to have been
4 driven more by reduction in species pool size rather than turnover of existing forest species or
5 the spread of immigrants into more locations within each site (Baeten *et al.* 2012; 2014). We
6 therefore included within-site beta diversity in 1971 in our path model (Fig. 2, β_8) and also
7 tested whether change in beta diversity had occurred between surveys, whether directions of
8 change differed depending on storm exposure and whether the influence of beta diversity was
9 largely due to differences in site species pool size.

10

11 **MATERIALS AND METHODS**

12 **Study region**

13 We analysed data from 26 woodlands all located in southern England between 48m and 198m
14 above sea level (Fig. 1). Regional climate is temperate maritime (Peel *et al.* 2007) with annual
15 precipitation of 600 - 800 mm, a mean January temperature of 3 - 4.5 °C and mean monthly July
16 temperature of 16.5 - 17.5 °C (Goudie & Burden 1994). The region is densely populated (401
17 persons per km² in England in 2012 - www.ons.gov.uk) and has seen widespread intensification
18 of agriculture since the end of World War II, including drainage and improvement of land,
19 mechanization of agriculture, and increases in agricultural productivity (North 2000;
20 Chamberlain *et al.* 2000). However, the study area also has the highest proportional cover of
21 broadleaved woodland in the British Isles (Morton *et al.* 2011). Woodland sites within the storm
22 track were selected as those coinciding with wind speeds above 144 kph in the peak gust wind

1 footprint defined by the RMS Europe Windstorm Model for the October 1987 storm (Risk
2 Management Solutions 2007).

3 The floristic affinities of the sample reflected the soils of southeastern England: base poor
4 brown earths and podzols (Rodwell 1991). Forty percent of plots were referable to the W10
5 *Quercus robur* – *Pteridium aquilinum* – *Rubus fruticosus* woodland and 20% to the W16
6 *Quercus* spp – *Betula* spp – *Deschampsia flexuosa* woodland. The major woodland community
7 type of calcareous to neutral soils in south east England, the W8 *Fraxinus excelsior* – *Acer*
8 *campestre* – *Mercurialis perennis* woodland, was represented by 11% of plots (see Fig. S1 in
9 Supporting Information). Canopy dominants comprised *Quercus robur* (in 77% of plots),
10 *Corylus avellana* (62%), *Fraxinus excelsior* (61%), *Betula pubescens* & *pendula* (48%), *Acer*
11 *pseudoplatanus* (39%) and *Fagus sylvatica* (38%). Nomenclature for plants follows Stace
12 (1997).

13

14 **Survey design and data collection**

15 Full details of the sampling design and sampling methods were published by Kirby *et al.*
16 (2005a,b) and Corney *et al.* (2006). In summary, the Great Britain (GB) Woodland Survey was
17 based on sites representing woodland types as determined from an earlier multivariate
18 classification of 2453 British woodlands (Ratcliffe 1977; Bunce 1982). Woodland sites were
19 visited in 1971 and again in 2002. Indicators of woodland management and surrounding land-use
20 were recorded at site level and from 16 random 200-m² square plots located in each woodland.
21 Cover and presence of all vascular and selected non-vascular plants were also recorded in each
22 plot. Individual trees and shrubs were identified and counted and the diameter at 1.3 m above-
23 ground (DBH) was measured for stems > 1 cm diameter. Understorey species richness comprises

1 the count of vascular plants, common bryophytes and species of trees and shrubs but only where
2 these were present as seedlings or saplings (individuals <1.3m in height and <1cm DBH). Fresh
3 soil pH was measured in deionized water on a homogenized 15cm topsoil sample taken from
4 each plot. A Quality Assurance (QA) survey of a subset of the sites was carried out in 2002
5 (Kirby *et al.* 2005b). QA procedures were also applied to the soils analysis including a partial re-
6 analysis of the stored 1971 samples (see Kirby *et al.* 2005a,b for details). The repeat visit in
7 2002, as far as possible, recorded all data from the same plot based on 1:10,000 site maps
8 prepared at the time of the first survey. A quantitative analysis of plot relocation error was
9 carried out by comparing mean similarity coefficients between temporal pairs of plots assumed
10 to have been recorded in the same locations in 1971 and 2002 versus similarity coefficients for
11 randomized pairings of plots within the same site. Results are reported in Kirby *et al.* (2005b)
12 and showed that, on average, attempts to re-find the 1971 plot resulted in greater similarity
13 between temporal pairs than randomized pairs.

14

15 **Datasets**

16 Data from the 1971 survey of the 10 sites situated inside the October 1987 storm-track were
17 matched with a dataset of plots from 1971 in sites outside the storm-track (Fig. 1). The floristic
18 composition of the storm sites was used to stratify non-storm exposed sites. These plots were
19 then randomly sampled to identify a dataset that was floristically equivalent to the storm-exposed
20 sites and for which plots and sites had equivalent average levels of pre-storm soil pH,
21 understorey species richness, mean woody basal area and mean cover-weighted SLA (Fig. S2).
22 This selection process yielded 150 plots in 16 non-storm exposed sites and 143 plots in the 10
23 sites inside the storm-track.

1
2 All sites were exposed to roughly equivalent levels of other potential driving variables, including
3 changes in atmospheric deposition of sulphur and nitrogen compounds, and intensive land-use
4 surrounding each woodland in 1971 (Fig. S2). All storm-exposed and non-storm sites showed
5 substantial reductions in modeled S deposition between 1970 and 2000 (Fig. S2). Deposition of
6 N at all sites was substantially above the empirical critical load (10-20 kg N ha⁻¹ yr⁻¹; Tipping *et al.*
7 *al.* 2013), whether inside the storm track or not (Fig. S2), but storm-track sites were exposed to
8 lower modeled N deposition in 2000.

9
10
11 **Path analysis**
12 The path model specified in Fig. 2 was implemented in OpenBUGS version 3.2.1
13 (<http://www.openbugs.info/w/>) (Grace *et al.* 2012). The separate regression models that made up
14 the path model were initially run within SAS (Little *et al.* 2000). This was done to derive
15 parameter estimates against which to help check the outputs from the OpenBUGS model. The
16 hierarchical structure of the data was specified by implementing a random-intercepts model in
17 OpenBUGS on all covariates that were measured on plots within sites (Kéry 2010).

18 To produce standardized regression coefficients and path coefficients, one version of the
19 path model was run with all covariates centred and standardized to zero mean and unit variance –
20 see Supplementary Material. A second path analysis was run to generate regression coefficients
21 and residual variances for variation partitioning (Gajewski *et al.* 2006; Shipley 2000). In the
22 latter, covariates were neither centred nor standardized. The regression coefficients in the latter
23 model also were used to interpret the estimated effect of a unit change in hypothesized
24 explanatory variables on proportional change in understory species-richness between 1971 and

1 2002. Since species-richness change between surveys was transformed to $\ln[(\text{richness } 2002 + 1)/$
2 $(\text{richness } 1971 + 1)]$, the regression coefficients involving species richness were back-
3 transformed by exponentiating them in order to derive an estimate of the effect of the
4 explanatory variable on the proportional change in species richness from 1971 to 2002. For
5 example if the average richness in 1971 were twice the 2002 value then the raw ratio would be
6 0.5 and the parameter estimate approximately -0.69. Storm impact was coded as 1 (storm) and 0
7 (no-storm) in the data.

8 Two Monte Carlo Markov chains were initialized for each path analysis with varying
9 starting values. Convergence of all posterior distributions was monitored with trace plots and by
10 the Gelman-Rubin statistic (Kéry 2010). After a 20,000 iteration burn-in, a subsequent 20,000
11 iterations were summarized to describe posterior distributions for all parameters of interest.

12

13 **Testing hypothesised paths**

14 All response variables could be approximated by normal error distributions. Thus, path
15 coefficients were calculated by sampling from the posterior distributions of the products of
16 standardized regression coefficients (Grace 2006). The following path coefficients were
17 specified to test particular hypotheses as follows:

18

19 $\beta_3^* \beta_4$: Exposure to the 1987 storm was associated with increased species-richness or reduced
20 loss of richness via the effect of reduced woody basal area and hence greater light availability at
21 ground level (Kirby 1988). Thus the storm was expected to have deflected the wider national
22 trend for canopies to age and close (Kirby *et al.* 2005a).

23

1 $\beta_3^* \beta_5^* \beta_6$: Storm-driven reductions in woody basal area increased species-richness by driving
2 increased soil pH in forest gaps. Mechanisms include reduced input of lower pH, higher C:N leaf
3 litter and woody debris compared to sites not exposed to the storm (van Oijen *et al.* 2005) and
4 increased input of higher pH litter from early successional trees, shrubs and gap-phase herbs
5 (Sydes & Grime 1981; Nordén 1994; Cooper-Ellis *et al.* 1999; Borschenius *et al.* 2004), or
6 increasing pH via soil disturbance (Guo *et al.* 2004; Strandberg *et al.* 2005; van Oijen *et al.*
7 2005).

8

9 $\beta_7^* \beta_6$: Given that lower pH soils have been more susceptible to historical acidification impacts
10 in the UK, these soils should show a greater recovery than higher pH soils (Norton *et al.* 2012;
11 Kirk *et al.* 2006). Since changes in species richness are expected to respond positively to
12 increased pH, lower pH soils should have increased most in species richness because of changes
13 in soil pH (Borschenius *et al.* 2004; De Keersmaeker *et al.* 2004). However, if increased soil pH
14 covaries with, or results from, eutrophication, then this could drive understorey dominance and
15 reduced richness (Kirby 1998; Erjnaes *et al.* 2003; Brewer *et al.* 2012). The mean of the soil pH
16 in 1971 and 2002 was used as the predictor to circumvent any regression to the mean artefact
17 associated with plot relocation error (Kirk *et al.* 2006).

18

19 All direct and indirect effects were also tested taking into account that, on average, woodland
20 sites were surveyed 38 days earlier in 2002 than in 1971 and closer to the height of the growing
21 season (Fig 2, β_1) We therefore expected to detect more species in each plot in 2002 than in
22 1971.

23

1 **Variance partitioning**

2 In path analysis, covariates can be response variables as well as explanatory variables. The
3 covariate at the terminal node of the path diagram – in this case change in species richness – was
4 subject to explanation by the largest number of preceding explanatory variables, some of which
5 are conditional on intermediate variables. This results in variation in species-richness change
6 being broken down into the largest number of variance components. Other covariates in the path
7 diagram are explained by progressively fewer effects, whilst exogenous variables such as storm
8 exposure and difference in date of site survey are not subject to explanation by any preceding
9 variables (Fig. 2). Variance explained was also decomposed to the site and plot level since
10 predictors such as storm exposure and within-site beta diversity in 1971, were only measured at
11 the site scale while others were measured within plots and so can potentially explain between
12 plot and between site variation.

13

14 **Changes in species composition**

15 A binomial test (Zar 1984) was used to calculate the cumulative probability of the observed
16 number of presences of each species in either 1971 or 2002, assuming a 50% chance of occurring
17 in either year. The results convey inequality in the distribution of records between years in the
18 sample plots as a basis for interpreting whether there appears to have been an increased chance
19 of recording species associated with nutrient enrichment in 2002 in the storm-impacted plots
20 versus the non-storm plots.

21

1 The impact of enrichment on changes in understorey species composition also was quantified by
2 analysis of differences in cover-weighted SLA between years and within storm versus non-storm
3 sites. Cover-weighted SLA was calculated as follows:

$$4 \quad cSLA_j = \text{sum}(SLA_{ij} \times (cov_{ij}))/\text{sum}(cov_{ij}), \quad 1)$$

6
7 where (cov_{ij}) was the square root transformed percentage cover value for species i in each
8 sample plot j . All calculations of cSLA used a single published value per species. SLA values
9 were taken from Grime *et al.* (1995), Kleyer *et al.* (2008) and Wright *et al.* (2004). All vascular
10 plant species had SLA values. cSLA was also analysed in another version of the path analysis
11 model in Fig. 2 and subjected to the same hypothesized drivers of change to determine whether
12 there was any relationship with soil pH change, woody basal area change, site beta diversity and
13 the direct effect of storm exposure.

14 15 *Beta diversity change and effects on species richness*

16 Beta diversity within each site in each year of survey and change between surveys was calculated
17 based on the ΣDi community heterogeneity metric using the the *rDev* function in R provided by
18 Baeten *et al.* (2014). This metric sums the binomial deviances associated with variation in
19 species frequency across plots in a site. Deviance is greatest for species occupying 50% of plots
20 and so the metric attains lower values as species increase in commonness or rarity. Because we
21 were only interested in impacts on the understorey, the metric was calculated after excluding
22 records for all canopy trees and shrubs.

1 Beta diversity values for the understorey across each site in 1971 were entered into the
2 path model as predictors of change in plot-level species richness (Fig 2). The regression was re-
3 run with the ΣDi metric standardized by species pool size to remove the effect of differences in
4 site species richness. We simply divided each site value of the metric by the size of the site pool.
5 Comparing the residual variation between a regression model based on standardized versus
6 unstandardized values of ΣDi indicated how much of the explanatory power of ΣDi was due to
7 differences in site richness rather than between-plot frequency.

8

9 **RESULTS**

10 *Species richness*

11 Plots inside the storm-track had a lower loss of understorey species richness or increased in
12 richness (Figs. 3, 4a and Table 1, $\beta 2$). Increased woody basal area was associated with decreased
13 species richness (Table 1, $\beta 4$) and increasing soil pH was associated with increased species
14 richness, yet change in woody basal area and soil pH did not differ significantly between storm
15 and non-storm sites (Figs. 3, 4b, 4d and Table 1, $\beta 6$).

16 Interpretation of the regression coefficients based on path analysis of uncentred and
17 unstandardised data (Table 1) indicated that storm exposure resulted in an estimated mean 32%
18 increase in species-richness by 2002; estimates ranged from 8% to 61% (Table 1, 95% credible
19 interval on $\beta 2$). A 3.4% reduction in species richness was estimated to occur for every 0.1 m²
20 increase in woody basal area and a 12% increase in species richness was estimated to occur for
21 every one pH unit increase (Table 1). Species richness on average increased by 2% for every 10
22 days earlier plots were sampled in 2002 relative to 1971.

23

1 In total, 39% of the observed between-plot variation and 87% of the between-site variation in
2 species-richness was explained by the path model (Fig. 5 and Table 1). The best predictors at the
3 site level were within-site beta diversity of the ground flora across each site in 1971 (20%) and
4 storm exposure (60%). Collinearity between difference in survey date and 1971 beta diversity
5 was observed; when both predictors were included in the path model the variation explained by
6 difference in survey date dropped from 16% to 3.6%. Change in mean soil pH across sites
7 explained 2% of the mean change in species richness. Despite the expectedly influential role of
8 woody basal area change as a driver of gap creation and changing light regime, only 0.8% of
9 mean change in species richness among sites and 5.2% of change in species richness among plots
10 within sites, was explained by mean woody basal area change (Fig. 5). Species-richness change
11 among plots within sites was highly variable and weakly correlated with the variation in mean
12 species-richness change across sites. Consequently only 3.5% of the between-plot variation in
13 species-richness change was explained by between-site variation in species-richness change (Fig.
14 5).

15 Within-site beta diversity (ΣDi) in 1971 was a stronger predictor of change in species
16 richness, explaining 20% of the mean change in richness among sites (Fig 5). The relationship
17 was negative indicating that higher beta diversity in 1971 was associated with a greater loss of
18 species richness over time or smaller gains in richness for sites that increased in species richness.
19 The interaction with storm exposure was not significant. When standardized ΣDi values were
20 regressed against species richness change, variance explained dropped to 4% so that the majority
21 of the explanatory power of the beta diversity measure was due to differences in the size of the
22 species pool between sites and not turnover of species between plots.

23

1 *Soil pH*

2 Soil pH increased between 1971 and 2002 across all sites (Fig. 4b), changes that were
3 significantly correlated with an increase in understorey species richness (Table 1, β_6). Contrary
4 to expectation soil pH was more likely to increase significantly in plots with a higher initial pH
5 (Table 1, β_7). Mean site-level soil pH explained 30% of the between-site change in soil pH but
6 mean plot-level soil pH explained only 2.4% of the plot-level change in soil pH (Fig. 5).
7 Variation in mean site pH was able to explain 78% of the variation between plots indicating that
8 most of the variation was between sites with much less between plots within sites. Change in
9 woody basal area at plot level only explained 0.2% of the change in soil pH at plot level.
10 Explanatory power at site level was an order of magnitude less. Soil pH change did not
11 significantly differ between storm and non-storm sites (Table 1, β_9).

12

13 *Woody basal area*

14 There was no evidence of a difference in basal area change between plots inside or outside the
15 storm-track (Fig 4d). However, storm exposure explained 29% of the variation in mean woody
16 basal area change among sites (Fig. 5). An average 1.2% decrease in soil pH was estimated to
17 occur with every 0.1-m² increase in woody basal area per 200 m², but this estimate ranged
18 between a 3.2% decrease and a 0.7% increase and was therefore not significant (Table 1, 95%
19 credible interval on β_5).

20

21 *Indirect effects*

22 Mean soil pH in 1971 and 2002 had a significant positive effect on change in species richness via
23 soil pH change (Table 1, β_6), however the effect size, as measured by the path coefficient, was

1 very small (Table 1, β_7 β_6). No other significant indirect relationships were detected where
2 storm exposure impacted species-richness change via impacts on mediating variables (Table 1,
3 path coefficients).

4 *Changes in species composition*

5 Five species were more frequent by 2002 in both storm and non-storm sites of which two,
6 *Ranunculus repens* and *Galium aparine*, are widespread generalist species extremely common in
7 lowland Britain. Thirty-three species were more frequent in storm-impacted sites by 2002 but
8 were not significantly different in frequency in non-storm sites (Table 2). These included the
9 alien shrub *Prunus laurocerasus* and the widespread weeds *Senecio jacobaea* and *Cirsium*
10 *vulgare*. However, the majority of species that increased in frequency in storm sites but not in
11 non-storm sites were more typical of base-poor, low productivity substrates, such as *Carex*
12 *binervis*, *C. pilulifera*, *Juncus effusus*, *Holcus mollis*, *Digitalis purpurea* and *Teucrium*
13 *scorodonia* (Table 2). Five of the species that increased only on storm sites are considered
14 ancient woodland specialists in south eastern England; *Anemone nemorosa*, *Hyacinthoides non-*
15 *scripta*, *Lysimachia nemorum*, *Hypericum pulchrum* and *Chrysosplenium oppositifolium* (Table
16 2). Moreover, all increasing generalist species that would be favoured by nutrient enrichment
17 were still much less frequent in 2002 than typical woodland species (Table 2).

18
19 Cover-weighted SLA did not change significantly between surveys either in storm-
20 impacted or non-storm sites (Fig. 4c) and was not significantly explained by any of the path
21 model relationships – Supplementary Material. Thus, fifteen years after the storm, differences in
22 species frequency and plant trait contribution between storm and non-storm plots showed no
23 evidence of a widespread shift toward assemblages that would indicate eutrophic conditions.

24

1 *Changes in understorey beta diversity*

2 Eighteen of the 26 sites showed significant changes in beta diversity between 1971 and 2002. On
3 storm exposed sites the only significant changes were increases, while on non-storm sites five
4 decreased and seven increased - see Supplementary Material.

5

6 **DISCUSSION**

7

8 *The effects of the October 87 storm on understorey species richness*

9 By 2002 exposure to the 1987 storm had significantly offset the reduction in species richness
10 associated with non-impacted sites and typical of the long-term trend in woodlands across
11 Britain (Fig. 4a). Among the range of predictors tested, storm exposure had the strongest effect
12 on the change in mean species richness across woodland sites; it had 40 times the explanatory
13 power of mean soil pH change and 75 times the explanatory power of mean woody basal area
14 change. Despite this apparently strong effect, both woody basal area change and the species-
15 richness response were highly variable across plots and forest sites. Contrary to expectation
16 storm exposure explained only 29% of the change in mean woody basal area across sites, which
17 in turn explained a miniscule 0.8% of mean site-level species richness change. Yet storm
18 exposure directly and uniquely explained 60% of the change in mean site-level species richness.
19 The mechanism whereby storm exposure impacted species richness but independently of change
20 in woody basal area must comprise a range of other disturbance effects. These include gap
21 creation by blowdown of trees with stems outside plots but whose canopies shaded plots. Also
22 moderate damage to trees could have resulted in additional light penetration at ground level but
23 where trees continued to grow. If not killed, most of the broadleaved canopy species can re-
24 sprout and re-leaf quickly. Thus change in basal area may not be strongly correlated with post-

1 disturbance changes in canopy cover that alter light availability and impact species richness
2 (Clinton & Baker 2000; Brewer *et al.* 2012; Barker-Plotkin *et al.* 2013). Whilst all 10 sites were
3 exposed to the storm, the extent of disturbance reported by surveyors varied from none to
4 widespread (Kirby *et al.* 2005b). For example, many fallen trees were reported as still alive and
5 vigorously regrowing in 2002. Indeed a recent assessment indicated that the majority of timber
6 was not damaged. In the two counties completely within the storm track 24% of standing timber
7 volume was blown down in East Sussex and 18% in Kent (Harmer 2012). However, the apparent
8 absence of significant change in woody basal area in the storm-impacted sites seems at odds with
9 the likely effects of such exposure. A possible explanation is that the interval of 15 years
10 between the storm and the 2002 survey was sufficient for regrowth to have achieved woody
11 basal area values similar to those in the first survey in 1971. Previous analysis has shown that
12 substantially younger cohorts of stems were present across the survey sites in 1971 than in 2002
13 (Kirby *et al.* 2005a). Such an explanation assumes that widespread reduction in basal area of an
14 equivalent magnitude to the storm must have occurred around **the mid-1950s. This could reflect**
15 **the culmination of severe post-WWII timber extraction but in the absence of historical**
16 **management information for the sites involved this is a matter for speculation and further**
17 **enquiry.**

18 Since storm salvage operations were also apparent in the aftermath of the storm, it is
19 unclear how these may have altered woody basal area and impacted the structure and species
20 composition of gaps (cf. Cooper-Ellis *et al.* 1999; Brewer *et al.* 2012; Barker-Plotkin *et al.*
21 2013). During the 2002 resurvey, land owners often reported how destructive the storm had
22 been, but also indicated that it had been the stimulus for interventions, including clearing out
23 fallen timber and dead wood and then restocking. However, surveyors' reports indicated how the

1 apparent effects of post-storm tidying varied greatly; in some places these effects were linked to
2 further suppression of understorey species-richness following dense restocking, whilst in others
3 they were associated with greater light penetration and herbaceous regrowth and the removal of
4 dead wood (also see Whitbread 1991).

5 The severity of canopy damage and the dynamics of recovery depends on a range of other
6 scale-dependent factors, including legacy effects of management, slope, tree species and age,
7 ground wetness, nutrient availability, litter inputs and plant traits (Foster & Boose 1992;
8 Whitbread & Montgomery 1994; Peterson & Pickett 1995; Cooper-Ellis *et al.* 1999; Clinton &
9 Baker 2000). Even if canopy gaps are created, the subsequent timing and direction of change in
10 species richness depends upon propagule availability from nearby populations and the
11 persistence of vegetative material in and around gaps (Whitney & Foster 1988; Vellend 2003;
12 Roberts 2004). Post-storm salvage operations as well as these other factors will have
13 undoubtedly contributed to the large amounts of residual variance not explainable by the few
14 predictors applied in the analysis. However, despite the chaotic nature of the storm's impact at
15 multiple scales, our cross-site study explained 87% of the change in mean species richness at
16 site-level and provides a novel estimate that exposure of woodland plots to the storm increased
17 understorey species-richness per 200m² by an average of 32% of their starting values in the
18 following 15 years.

19

20 *Changes in species diversity; was there evidence of a eutrophication effect?*

21 In four of the storm-exposed sites, surveyors reported locally vigorous colonization of canopy
22 gaps by species whose consolidation was associated with low species-richness. Colonising
23 dominants included the non-native, invasive shrubs *Prunus laurocerasus* and *Rhododendron* spp,

1 the native rhizomatous fern *Pteridium aquilinum* and dense juvenile stems of the native tree
2 *Betula pubescens*. Rapid gap colonization by a small number of dominants is consistent with
3 other observations on storm-affected sites (Parker 1994; Cole & Weltzin 2005). However, the
4 identity of the species that were more frequent on storm-exposed sites by 2002 did not indicate a
5 widespread increase in nitrophilous species. Whilst a small number of such species were more
6 frequent by 2002, a larger number of woodland specialists and species more typical of low
7 productivity substrates had increased even more. Moreover, cover-weighted SLA did not change
8 significantly indicating no average increase in abundance of species favoured by more enriched
9 conditions. These changes are also consistent with the average increase in plot species richness
10 in storm-exposed sites rather than suppression of species richness by a small number of
11 dominants.

12 The lack of a eutrophication effect could be attributable to a number of factors. Fifteen
13 years might be too short a time for the expression of a cross-site pattern of competitive
14 suppression in the understorey. However, on storm-exposed sites total atmospheric N deposition
15 ranged from an estimated 31 to 43 kg ha⁻¹ yr⁻¹, well in excess of the current European empirical
16 critical load for nitrogen. Significant species compositional change in the herbaceous understorey
17 has been found elsewhere following addition of lower N loads than this over shorter time
18 intervals (reviewed in Gilliam 2006). Yet, in other experiments and observational studies,
19 changes in dominance and diversity either have been much slower or have not been observed and
20 appear to depend upon the soil chemistry of the study system and the presence of responsive
21 species at the start (De Schrijver *et al.* 2011; Verstraeten *et al.* 2013). It is therefore possible that
22 eutrophication effects have yet to influence understorey dominance hierarchies and may only do
23 so dependent on the biogeochemical susceptibility of different locations.

1 An additional constraint is soil pH and its influence on macro-nutrient availability
2 (Schaffers 2002; Falkengren-Grerup & Diekmann, 2003). The non-significant path from storm
3 exposure to soil pH change via woody basal area change means that changes in soil pH were
4 independent of both factors even though significant soil pH change did occur between 1971 and
5 2002 (Fig. 4b). The overall increase in soil pH is therefore consistent with recovery from
6 acidification following reduced atmospheric sulphur deposition since the mid-1970s (Norton *et*
7 *al.* 2012; Kirk *et al.* 2006) but there is no evidence that this background change in pH was
8 amplified on storm-exposed sites. As more woodlands recover from acidification, those moving
9 into a pH window of between ≈ 5.5 and 7.0, (Schaffers 2002; Stevens *et al.* 2011), are expected to
10 show increasing dominance by nitrophiles unless continued lack of disturbance and increased
11 shading prevents such light-demanding species from becoming abundant (De Keersmaeker *et al.*
12 2004; Verheyen *et al.* 2012; Baeten *et al.* 2009).

13 A significant positive relationship also was found between soil pH change and species-
14 richness change between 1971 and 2002. A positive spatial relationship between soil pH and
15 species richness is consistent with other datasets for temperate forests (Borschenius *et al.* 2004;
16 De Keersmaeker *et al.* 2004; Corney *et al.* 2006) but it is interesting to find such a clear coupling
17 between species-richness change and soil pH change over time. This suggests a responsive
18 woodland species pool and a signal detectable despite apparent storm-driven changes in species
19 richness within the same dataset. The mean soil pH changed from 4.7 to 5.3 in the 30 years, a
20 change estimated to have driven an average increase of 7% of the 1971 starting species-richness
21 or an addition of 1 species per plot given that the mean richness in 1971 was 14 per 200m². This
22 change moved the average woodland to just below the threshold where macro-nutrient
23 availability confers susceptibility to dominance by nitrophilous species in the understorey and

1 reduced species richness (Stevens *et al.* 2011). Moreover, higher pH soils tended to have shown
2 the greatest increase in pH between 1971 and 2002 (Fig. 6). Soil and vegetation responses to
3 changing pollutant deposition differ depending upon the biogeochemical starting point and
4 whether the starting pH was previously reduced by historical acidification (Verstraeten *et al.*
5 2013; Baeten *et al.* 2009). Where pollutant deposition drives pH down to below about 4.2-4.3
6 species richness typically declines. Mechanisms include toxicity of aluminium and H⁺ and the
7 loss of species unable to effectively utilise NH₄⁺ (Stevens *et al.* 2011; Stevens *et al.* 2009).
8 Within the sample of 293 woodland plots, 20 plots moved into the pH_{>=5.5} window while 54
9 remained below a pH of 4.2 and 75 moved from below pH 4.2 to between 4.2 and 5.5. These
10 movements between critical pH windows should predict increases or decreases in diversity
11 reflecting recovery from acidification and then the onset of eutrophication. However, we do not
12 know whether these pH changes are a consequence of pollutant deposition driving down pH
13 earlier in the 20th century with recovery since the late 1970s. Analysis along crossed gradients of
14 sulphur and nitrogen deposition history and soil pH is not possible for the small sample of paired
15 storm and non-storm woodlands because all sites saw reduced acidification and all were subject
16 to high N loads (Fig. S2). With no gradient of effects along which to analyse change no signal
17 can be attributed (Smart *et al.* 2012). Analysis of the full set of 103 sites offers a chance of
18 further characterizing the effects of pollutant deposition on soil pH and understorey diversity but
19 in the absence of storm disturbance effects.

20

21 *Did the understorey species composition become more homogenous?*

22 Different mechanisms could result in homogenization of the woodland understorey depending
23 upon exposure to the storm. On storm-disturbed sites homogenization would result where

1 regenerating understoreys were dominated by a small number of widespread generalists
2 associated with the wider farmed landscapes of lowland Britain, at the expense of a larger pool
3 of typical woodland species. On non-storm exposed sites, suppression of the understorey by
4 continued shading and lack of management would lead to greater homogeneity if the same
5 smaller pool of shade-tolerant plant species persists across sites. The latter scenario appears
6 broadly typical of recent changes in many European forests (Baeten *et al.* 2014). Our results
7 clearly indicated increased differentiation and heterogeneity of the understorey on storm-exposed
8 sites with no evidence of the release of suppressive nitrophiles. On non-storm sites, within-site
9 beta diversity increased or decreased in roughly equal measure. However, analysis of the wider
10 site series has shown a marked loss of species richness within British broadleaved woodlands
11 with a species-compositional shift toward a more shade-tolerant flora (Kirby *et al.* 2005a).
12 Ongoing lack of disturbance is not necessarily a counsel of despair since shaded undisturbed
13 woods may be poor in plant species per unit area but richer in groups of invertebrates, fungi and
14 bryophytes that prefer dead wood, low light, humidity and shade (Townsend 2006; Hambler &
15 Speight 1995). However, these specialist taxa may also be increasingly rare given the
16 fragmentation of woodlands and the negative effects of pollutant deposition.

17

18 *Developing the conceptual framework of forest responses to global change*

19 A combination of the three conceptual frameworks provided a useful basis for hypothesis
20 generation and testing. Roberts (2004) provides a convenient separation of disturbance effects
21 along three axes that align well with the effect of storm disturbance as a natural pulsed changed
22 in resource availability on soil, understorey and canopy. Roberts (2004) however does not
23 explicitly predict ecosystem dynamics in response to global change drivers. The HRF does so

1 and in particular makes predictions about the outcome of interactions between natural pulsed
2 disturbances and externally sourced chronic changes in resources whose scale means that
3 resource supplies cannot be modified by local ecosystem feedbacks. Finally, the biotic
4 homogenisation framework has been widely applied to woodland change (Wiegmann & Waller,
5 2006; Baeten *et al.* 2012) and increasingly valuable insights are likely to arise from focussing on
6 how plant traits that are known to drive feedbacks on ecosystem functioning become more
7 widely represented as global change drivers non-randomly select winning versus losing taxa
8 from the local and regional species pools (Suding *et al.* 2008). The likelihood that new colonists
9 will include widespread generalists typical of human-modified landscapes is increased where
10 woodland patches are smaller and less buffered by existing older woodland (Kimberley *et al.*
11 2014). These spatial effects probably need to be more explicitly factored in when extending the
12 Roberts (2004) framework to fragmented, small woodlands typical in northern Europe and in
13 other parts of the temperate zone.

14

15 *Synthesis*

16 In summary, analysis of this unique dataset has shown that storm events can drive a reversal in
17 the direction of change in plant species richness resulting from at least 40 years of reduced
18 canopy disturbance. Direct storm impacts on the understorey were detectable, but quantifying the
19 links between storm exposure, different types and severities of canopy damage and the impact of
20 these on soil and understorey vegetation will likely require a wider range of measurements.

21 Many temperate woodlands are now embedded in intensively managed landscapes and
22 subject to legacy effects of elevated atmospheric nitrogen deposition but reduced sulphur
23 deposition. Hence the future consequence of disturbance, whether from storm events or

1 reinstated management, could be the development of very different herbaceous understoreys
2 dominated by rapidly growing species more typical of nutrient rich conditions. More frequent
3 recording at the impacted and unimpacted woodland sites would be highly desirable to determine
4 the ongoing course of post-disturbance trajectories and the extent to which these help us
5 understand the resilience of temperate woodlands to the interacting effects of future stressors
6 (Bruehlheide & Luginbühl 2009).

7

8 **Acknowledgements**

9 SMS thanks the NERC Centre for Ecology & Hydrology, Mark Bailey, Alan Jenkins, Terry Parr
10 and colleagues in the Land Use Group for supporting a sabbatical leave at the Harvard Forest
11 funded by the Charles Bullard fellowship scheme during which this work was initiated. Work by
12 AME and DRF on this project was supported by the Harvard Forest LTER site, through NSF
13 grants 0620443 and 12-37491. Comments from the journal editors Lauren Sandhu, Frank Gilliam
14 and David Gibson, and two anonymous referees greatly improved the original manuscript.
15 Andrew Titman is thanked for statistical advice. The woodland resurvey was funded by DEFRA,
16 Natural England, Forestry Commission and the Woodland Trust.

17

18 **Data accessibility**

19 The woodland survey database is freely available via the following DOI:

- 20 - doi.org/10.5285/fb1e474d-456b-42a9-9a10-a02c35af10d2
- 21 - doi.org/10.5285/d6409d40-58fe-4fa7-b7c8-71a105b965b4
- 22 - doi.org/10.5285/2d023ce9-6dbe-4b4f-a0cd-34768e1455ae
- 23 - doi.org/10.5285/4d93f9ac-68e3-49cf-8a41-4d02a7ead81a

24

25

26 **SUPPORTING INFORMATION**

- 1 **Figure S1** Phytosociological composition of the study plots in the baseline year of 1971.
- 2 **Figure S2** Distributions of measured variables in storm (1) and non-storm (0) sites.
- 3 **Table S1** Percentage of the variation in response variables explained by hypothesized predictor
- 4 variables.
- 5 **Table S2** Path analysis of change in cover-weighted Specific Leaf Area (cSLA).
- 6 **Figure S4** Path analysis diagram for change in cover-weighted SLA between 1971 and 2002.
- 7 **Table S3** Significance tests of change in understorey community heterogeneity (ΣDi).
- 8 **Text S1** Notes on variation partitioning.
- 9 **Text S2** OpenBUGS code.
- 10 **Text S3** Notes on construction of the path analysis in OpenBUGS.
- 11

1 **REFERENCES**

2 Baeten, L., Bauwens, B., De Schrijver, A., De Keersmaecker, L., Van Calster, H.,
3 Vanderkerkhove, K., Roelandt, B., Beeckmann, H. & Verheyen, K. (2009) Herb layer changes
4 (1954-2000) related to the conversion of coppice-with-standards forest and soil acidification.
5 *Journal of Vegetation Science*, **12**, 187-197.

6
7 Baeten, L., Vangansbeke, P., Hermy, M., Peterken, G., Vanhuyse, K. & Verheyen, K. (2012)
8 Distinguishing between turnover and nestedness in the quantification of biotic homogenization.
9 *Biodiversity & Conservation* **21**, 1399-1409.

10
11 Baeten, L., Warton, D.I., Van Calster, H., De Frenne, P., Verstraeten, G., Bonte, D., Bernhardt-
12 Römermann, M., Cornelis, J., Decocq, G., Eriksson, O., Hédli, R., Heinken, T., Hermy, M.,
13 Hommel, P., Kirby, K., Naaf, T., Petřík, P., Walther, G-R., Wulf, M. & Verheyen, K. (2014) A
14 model-based approach to studying changes in compositional heterogeneity. *Methods in Ecology*
15 *& Evolution* **5**, 156-164.

16
17 Barker-Plotkin, A., Foster, D.R., Carlson, J. & Magill, A. (2013) Survivors, not invaders, control
18 forest development following simulated hurricane. *Ecology*, **94**, 414-423.

19
20 Borschenius, F., Nielsen, P.K. & Lawesson, J.E. (2004) Vegetation structure and diversity of an
21 ancient temperate deciduous forest in SW Denmark. *Plant Ecology*, **175**, 121-135.

22

- 1 Brewer, J.S., Bertz, C.A., Cannon, J.B., Chesser, J.D. & Maynard, E.E. (2012) Do natural
2 disturbances or the forestry practices that follow them convert forests to early successional
3 communities? *Ecological Applications*, **22**, 442-458.
- 4
- 5 Bruelheide, H. & Luginbühl, U. (2009) Peeking at ecosystem stability: making use of a natural
6 disturbance experiment to analyze resistance and resilience. *Ecology*, **90**, 1314-1325.
- 7
- 8 Bunce, R.G.H. (1982) *A field key for classifying British woodland vegetation. Parts 1 & 2.*
9 Cambridge: Institute of Terrestrial Ecology, UK.
- 10
- 11 Burt, S.D. & Mansfield, D.A. (1988) The great storm of 15-16 October 1987. *Weather*, **43**, 90-
12 108.
- 13
- 14 Chamberlain, D.E., Fuller, R.J., Bunce, R.G.H., Duckworth, J.C. & Shrubbs, M. (2000) Changes
15 in the abundance of farmland birds in relation to the timing of agricultural intensification in
16 England and Wales. *Journal of Applied Ecology*, **37**, 771-788.
- 17
- 18 Clinton, B.D. & Baker, C.R. (2000) Catastrophic windthrow in the southern Appalachians:
19 characteristics of pits and mounds and initial vegetation responses. *Forest Ecology and*
20 *Management*, **126**, 51-60.
- 21
- 22 Cole, P.G. & Weltzin, J.F. (2005) Light limitation creates patchy distribution of an invasive grass
23 in eastern deciduous forests. *Biological Invasions*, **7**, 477-488.

1
2 Cooper-Ellis, S., Foster, D.R., Carlton, G. & Lezberg, A. (1999) Forest response to catastrophic
3 wind: results from an experimental hurricane. *Ecology*, **80**, 2683-2696.
4
5 Corney, P.M., Le Duc, M.G., Smart, S.M., Kirby, K.J., Bunce, R.G.H. & Marrs, R.H. (2006)
6 Relationships between the species composition of forest field-layer vegetation and environmental
7 drivers, assessed using a national scale survey. *Journal of Ecology*, **94**, 383-401.
8
9 De Keersmaeker, L., Martens, L., Verheyen, K., Hermy, M., De Schrijver, A. & Lust, N. (2004)
10 Impact of soil fertility and insolation on diversity of herbaceous woodland species colonizing
11 afforestations in Muizen forest (Belgium). *Forest Ecology and Management*, **188**, 291-304.
12
13 De Schrijver, A., De Frenne, P., Ampoorter, A., Van Nevel, L., Demey, A., Wuyts, K.,
14 Verheyen, K. (2011) Cumulative nitrogen input drives species loss in terrestrial ecosystems.
15 *Global Ecology & Biogeography*, **20**, 803-816.
16
17 Erjnaes, R., Hansen, D.N. & Aude, E. (2003) Changing course of secondary succession in
18 abandoned sandy fields. *Biological Conservation* **109**, 343-350.
19
20 Falkengren-Grerup, U. & Diekmann, M. (2003) Use of a gradient of N deposition
21 to calculate effect-related soil and vegetation measures in deciduous forests. *Forest Ecology*
22 *and Management*, **180**, 113-124.
23

- 1 Flinn, K.M. & Vellend, M. (2005) Recovery of forest plant communities in post-agricultural
2 landscapes. *Frontiers in Ecology and the Environment*, **3**, 243-250.
- 3
- 4 Foster, D.A. & Boose, E. (1992) Patterns of damage resulting from catastrophic wind in Central
5 New England, USA. *Journal of Ecology*, **80**, 79-98.
- 6
- 7 Foster, D.R., Aber, J.D., Melillo, J.M., Bowden, R.D. & Bazzaz, F.A. (1997) Forest response to
8 disturbance and anthropogenic stress. *BioScience*, **47**, 437-445.
- 9
- 10 Foster, D.R., Swanson, F., Aber, J.D., Burge, I., Brokaw, N., Tilman, D. & Knapp, A. (2003)
11 The importance of land-use legacies to ecology and conservation. *BioScience*, **53**, 77-88.
- 12
- 13 Garnier, E., Cortez, J., Billes, G., Navas, M.L., Ruomet, C., Debussche, M., Laurent, G.,
14 Blanchard, A., Aubry, D., Bellmann, A., Neill, C. & Toussaint, J.P. (2004) Plant functional
15 markers capture ecosystem properties during secondary succession. *Ecology*, **85**, 2630-2637.
- 16
- 17 Gajewski, B.J., Lee, R., Thompson, S., Dunton, N., Becker, A. & Wells, V. (2006) Non-normal
18 path analysis in the presence of measurement error and missing data: a Bayesian analysis of
19 nursing homes' structure and outcomes. *Statistics in Medicine*, **25**, 3632-3647.
- 20
- 21 Gilliam, F.S. (2006) Response of the herbaceous layer of forest ecosystems to excess nitrogen
22 deposition. *Journal of Ecology*, **94**, 1176-1191.
- 23

1 Goudie, A.S. & Burden, P. (1994) *The Environment of the British Isles: an atlas*. Oxford
2 University Press, Oxford, UK.
3
4 Grace, J.B. (2006) *Structural Equation Modelling and Natural Systems*. Cambridge University
5 Press, Cambridge, UK.
6
7 Grace, J.B., Schoolmaster, D.R., Guntenspergen, G.R., Little, A.M., Mitchell, B.R., Miller, K.R.,
8 Schweiger, E.W. (2012) Guidelines for a graph-theoretic implementation of structural equation
9 modelling. *Ecosphere* **3**, Article 73.
10
11 Grime, J.P., Hodgson, J.G., Hunt, R. & Thompson, K. (1995) *The Electronic Comparative Plant*
12 *Ecology*. London. Chapman & Hall, UK.
13
14 Guo, D., Mou, P., Jones, R.H. & Mitchell, R.J. (2004) Spatio-temporal patterns of soil available
15 nutrients following experimental disturbance in a pine forest. *Oecologia*, **138**, 613-621.
16
17 Hambler, C. & Speight, M.R. (1995) Biodiversity conservation in Britain: science replacing
18 tradition. *British Wildlife*, **6**, 137-148.
19
20 Harmer, R. (2012) What happened in the woods? *Weather*, **67**, 261-265.
21

- 1 Hopkins, B (1994) The storm. *In: Ecological Responses to the 1987 Great Storm in the woods of*
2 *south-east England* (eds. K.J. Kirby, G.P. Buckley). English Nature Science series (23).
3 Peterborough, UK.
- 4
- 5 Kéry, M (2010) *Introduction to WinBUGS for Ecologists*. Elsevier, Academic Press, Amsterdam,
6 Netherlands.
- 7
- 8 Kimberley, A., Blackburn, G.A., Whyatt, J.D. & Smart, S.M (2013) Identifying the trait
9 syndromes of conservation indicator species: how distinct are British ancient woodland indicator
10 plants from other woodland species? *Applied Vegetation Science* **16**, 667-675.
- 11
- 12 Kimberley, A., Blackburn, G.A., Whyatt, J.D. & Smart, S.M (2014) Traits of plant communities
13 in fragmented forests: the relative influence of habitat spatial configuration and local abiotic
14 conditions. *Journal of Ecology* **102**, 632-640.
- 15
- 16 Kirby, K.J. (1998) The distribution and growth of bramble (*Rubus fruticosus*) in British semi-
17 natural woodland and the implications for nature conservation. *Journal of Practical Ecology and*
18 *Conservation*, **2**, 31-41.
- 19
- 20 Kirby, K.J. (1988) Changes in the ground flora under plantations on Ancient Woodland sites.
21 *Forestry*, **61**, 317-338.
- 22

1 Kirby, K.J. (2006) Ancient Woodland Indicator (AWI) plants. In: Rose, F. *The Wildflower Key*.
2 Pp. 558-561. Penguin Group, London, UK.
3
4 Kirby, K.J., Smart, S.M., Black, H.I.J., Bunce, R.G.H., Corney, P.M. & Smithers, R.J. (2005a)
5 *Long-term ecological changes in British broadleaved woodland 1971-2002*. English Nature,
6 Research Report. 653, part 1. Peterborough, UK. On-line at
7 <http://publications.naturalengland.org.uk/publication/94019?category=550043>
8
9 Kirby, K.J., Smart, S.M., Black, H.I.J., Bunce, R.G.H., Corney, P.M. & Smithers, R.J. (2005b)
10 *Long-term ecological changes in British broadleaved woodland 1971-2002. Appendices*. English
11 Nature, Research Report. 653, part 2. Peterborough, UK. On-line at
12 <http://publications.naturalengland.org.uk/publication/94019?category=550043>
13
14 Kirk, G.J.D., Bellamy, P.H. & Lark, R.M. (2006) Changes in soil pH across England and Wales
15 in response to decreased acid deposition. *Global Change Biology*, **16**, 3111-3119.
16
17 Kleyer, M., Bekker, R.M., Knevel, I.C., Bakker, J.P., Thompson, K., Sonnenschein, M.,
18 Poschlod, P., van Groenendael, J.M., Klimes, L., Klimesova, J., Klotz, S., Rusch, G.M., Hermy,
19 M., Adriaens, D., Boedeltje, G., Bossuyt, B., Danneman, A., Edndels, P., Gotzenberger, L.,
20 Hodgsno, J.G., Jackel, A.K., Kuhn, I., Kunzmann, D., Ozinga, W.A., Romermann, C., Stadler,
21 M., Schlegelmilch, J., Steendham, H.J., Tackenberg, O., Wilmann, B., Cornelissen, J.H.C.,
22 Eriksson, O., Granier, E. & Peco, B. (2008) The LEDA traitbase: a database of life-history traits
23 of the NW European flora. *Journal of Ecology*, **96**, 1266-1274.

1
2 Laughlin, D.C. (2011) Nitrification is linked to dominant leaf traits rather than functional
3 diversity. *Journal of Ecology*, **99**, 1091-1099.
4
5 Little, R.C., Milliken, G.A., Stroup, W.W. & Wolfinger, R.D. (2000) *SAS System for Mixed*
6 *Models*. 4th edn. Pp.437-432 Cary, NC: SAS Institute Inc., USA.
7
8 McKinney, M.L. & Lockwood, J.L. (1999) Biotic homogenisation: a few winners replacing
9 many losers in the next mass extinction. *Trends in Ecology & Evolution*, **14**, 450-453.
10
11 Morton, D, Rowland, C, Wood, C., Meek, L., Marston, C., Smith, G., Wadsworth, R. &
12 Simpson, I.C. (2011) *Countryside Survey: Final Report for LCM2007 – the new UK Land Cover*
13 *Map*. NERC Centre for Ecology & Hydrology, Lancaster UK. Pg 83.
14
15 Motzkin, G., Wilson, P., Foster, D.R. & Allen, A. (1999) Vegetation patterns in heterogenous
16 landscapes: the importance of history and environment. *Journal of Vegetation Science*, **10**, 903-
17 920.
18
19 North, J. (2000) Changes in agricultural structures in the 20th century. *Journal of the Royal*
20 *Agricultural Society of England. The Millenium Journal*, 29-40.
21
22 Norton, L.R., Maskell, L.C., Smart, S.M., Dunbar, M.J., Emmett, B.A., Carey, P.D., Williams,
23 P., Crowe, A., Chandler, K., Scott, W.A. & Wood, C.M. (2013) Measuring stock and change in

1 the GB countryside for policy - Key findings and developments from the Countryside Survey
2 2007 field survey. *Journal of Environmental Management*, **113**, 117-127.

3

4 van Oijen, D., Feijen, M., Hommel, P., den Ouden, J. & de Waal, R. (2005) Effects of tree
5 species composition on within-forest distribution of understorey species. *Applied Vegetation
6 Science*, **8**, 155-166.

7

8 Peel, M.C., Finlayson, B.L. & McMahon, T.A. (2007) Updated world map of the Köppen-Geiger
9 climate classification. *Hydrology & Earth System Sciences*, **11**, 1633-1644.

10

11 Peterken, G.F. (1981) *Woodland Conservation and Management*. London: Chapman &
12 Hall, UK.

13

14 Peterken, G.F. (1996) *Natural Woodland. Ecology and conservation in northern temperate
15 regions*. Cambridge University Press, UK.

16

17 Peterken, G.F. & Game, M. (1984) Historical factors affecting the number and distribution of
18 vascular plants in the woodlands of central Lincolnshire. *Journal of Ecology*, **72**, 155-182.

19

20 Peterken, G.F. & Jones, E.W. (1987) Forty years of change in Lady Park Wood: the
21 old growth stands. *Journal of Ecology*, **75**, 479-512.

22

- 1 Peterson, C.J. & Pickett, S.T.A. (1995) A case study in an old-growth forest catastrophic
2 blowdown. *Ecology*, **76**, 763-774.
- 3
- 4 Ratcliffe, D.A. (1977) *A Nature Conservation Review; the selection of biological sites of*
5 *national importance to nature conservation in Britain*. Cambridge: Cambridge University
6 Press, UK.
- 7
- 8 Risk Management Solutions, Inc. (2007) *The Great Storm of 1987: a 20-year retrospective. RMS*
9 *Special Report*. Risk Management Solutions, Inc., London, UK.
- 10
- 11 Roberts, M.R. (2004) Response of the herbaceous layer to natural disturbance in North American
12 forests. *Canadian Journal of Botany* **82**, 1273-1283.
- 13
- 14 Rodwell, J.S. (1991) *Woodland & Scrub*. British Plant Communities. Vol 1. Cambridge
15 University Press, Cambridge, UK.
- 16
- 17 Schaffers, A.P. (2002) Soil, biomass and management of semi-natural vegetation. *Plant*
18 *Ecology*, **158**, 229-246.
- 19
- 20 Shipley, B. (2000) *Cause and Correlation in Biology: a user's guide to path analysis, structural*
21 *equations and causal inference*. Cambridge University Press, Cambridge, UK.
- 22

1 Smart, S.M., Bunce, R.G.H., Marrs, R.H., Le Duc, M., Firbank, L.G., Maskell, L.C., Scott,
2 W.A., Thompson, K. & Walker, K.J. (2005) Large-scale changes in the abundance of common
3 higher plants across Britain between 1978, 1990 and 1998; tests of hypothesised changes in traits
4 linked to human activity. *Biological Conservation*, **124**, 355-371.

5

6 Smart, S.M., Thompson, K., Marrs, R.H., Le Duc, M.G., Maskell, L.C., Firbank, L.G. (2006a)
7 Biotic homogenization and changes in species diversity across human-modified ecosystems.
8 *Proceedings of the Royal Society. London. B.* **263**, 2659-2665.

9

10 Smart, S.M., Le Duc, M., Marrs, R.H., Rossall, M.J., Bunce, R.G.H., Thompson, K. & Firbank,
11 L.G. (2006b) Spatial relationships between intensive land cover and residual plant species
12 diversity in temperate, farmed landscapes. *Journal of Applied Ecology*, **43**, 1128-1137.

13

14 Smart, S.M., Henrys, P., Purse, B.V., Murphy, J., Bailey, M.J. & Marrs, R.H. (2012) Clarity or
15 confusion? – problems in attributing large-scale ecological changes to anthropogenic drivers.
16 *Ecological Indicators*, **20**, 51-56.

17

18 Smith, M.D., Knapp, A.K. & Collins, S.L. (2009) A framework for assessing ecosystem
19 dynamics in response to chronic resource alterations induced by global change. *Ecology*, **90**,
20 3279-3289.

21

22 Stace, C. (1997) *New Flora of the British Isles*. 2nd edn. Cambridge University Press, Cambridge,
23 UK.

1

2 Stevens, C.J., Dise, N.B., Gowing, D.J. (2009) Regional trends in soil acidification and metal
3 mobilisation related to acid deposition. *Environmental Pollution*, **157**, 313-319.

4

5 Stevens, C.J., Manning, P., Van den Berg, L.J.L., De Graaf, M.C.C., Wamelink, G.W., Weiger,
6 B., Andries, W., Bleeker, A., Vergeer, P., Arroniz-Crespo, M., Limpens, J., Lamers, L.P.M.,
7 Bobbink, R. & Dorland, E. (2011) Ecosystem responses to reduced and oxidized nitrogen
8 impacts in European terrestrial habitats. *Environmental Pollution*, **159**, 665-676.

9

10 Strandberg, B., Kristiansen, S.M. & Tybirk, K. (2005) Dynamic oak-scrub to forest succession:
11 effects of management on understorey vegetation, humus forms and soils. *Forest Ecology and*
12 *Management*, **211**, 318-328.

13

14 Suding, K.N., Lavorel, S., Chapin, F.S., Cornelissen, J.H.C., Díaz, S., Garnier, E., Goldberg, D.,
15 Hooper, D.U., Jackson, S.T. & Navas, M.L. (2008) Scaling environmental change through the
16 community-level; a trait-based response-and-effect framework for plants. *Global Change*
17 *Biology*, **14**, 1125-1140.

18

19 Sydes, C. & Grime, J.P. (1981) Effects of tree leaf litter on herbaceous vegetation in deciduous
20 woodland. II. An experimental investigation. *Journal of Ecology*, **69**, 249-262.

21

1 Romme, W.H., Boyce, M.S., Gresswell, R., Merrill, E.H., Minshall, G.W., Whitlock, C. &
2 Turner, M.G. (2011) Twenty years after the 1988 Yellowstone Fires: Lessons about disturbance
3 and ecosystems. *Ecosystems*, **14**, 1196-1215.
4
5 RoTAP (2012) *Review of Transboundary Air Pollution: Acidification, Eutrophication,*
6 *Ground Level Ozone and Heavy Metals in the UK*. Contract Report to the Department for
7 Environment, Food and Rural Affairs. Centre for Ecology & Hydrology. www.rotap.ceh.ac.uk
8
9 Taverna, K, Peet, R.K. & Phillips, L.C. (2005) Long-term change in ground-layer vegetation of
10 deciduous forests of the North Carolina Piedmont, USA. *Journal of Ecology*, **93**, 202-213.
11
12 Tipping, E., Henrys, P.A., Maskell, L.C. & Smart, S.M. (2013) Nitrogen deposition effects on
13 plant species diversity: threshold loads from field data. *Environmental Pollution* **179**, 218-223.
14
15 Townsend, M. (2006) Long-term change in British woodland (1971-2001) – Lies, damn lies and
16 statistics. *Ecos*, **27**, 90-95.
17
18 Vellend, M. (2003) Habitat loss inhibits recovery of plant diversity as forests regrow. *Ecology*,
19 **84**, 1158-1164.
20
21 Verheyen, K., Baeten, L, De Frenne, P., Bernhardt-Römermann, M., Brunet, J., Cornelis, J,
22 Decocq, G., Dierschke, H., Eriksson, O., Hédli, R., Heinken, T., Hermy, M., Hommel, P., Kirby,
23 K., Naaf, T., Peterken, G., Petřík, P., Pfandenauer, J., Van Calster, H., Walther, G-R, Wulf, M. &

1 Verstraeten, G. (2012) Driving factors behind the eutrophication signal in understory plant
2 communities of deciduous temperate forests. *Journal of Ecology*, **100**, 352-365.
3

4 Verstraeten, G., Baeten, L., Van den Broeck, T., De Frenne, P., Demey, A., Tack, W., Muys, B.
5 & Verheyen, K. (2013) Temporal changes in forest plant communities at different site types.
6 *Applied Vegetation Science*, **16**, 237-247.
7

8 Wiegmann, S.M. & Waller, D.M. (2006) Fifty years of change in northern upland forest
9 understories: Identity and traits of “winner” and “loser” plant species. *Biological Conservation*,
10 **129**, 109-123.
11

12 Whitbread, T. (1991) *When the wind blew: life in our woods after the great storm of*
13 *1987*. Royal Society for Nature Conservation, The Wildlife Trusts Partnership, Lincoln, UK.
14

15 Whitbread, T. & Montgomery, H.A.C. (1994) Storm damage in The Mens. *In:*
16 *Ecological Responses to the 1987 Great Storm in the woods of south-east England* (eds K.J.
17 Kirby, G.P. Buckley). Peterborough: English Nature Science series (23), UK.
18

19 Wright, I.J., Reich, P.B., Westoby, M., Ackerly, D.D., Baruch, Z., Bongers, F., Cavender-Bares,
20 J., Chapin, T., Cornelissen, J.H.C., Diemer, M., Flexas, J., Garnier, E., Groom, P.K., Gulias, J.,
21 Hikosaka, K., Lamont, B.B., Lee, T., Lee, W., Lusk, C., Midgely, J.J., Navas, M.L., Niinemets,
22 U., Oleksyn, J., Osada, N., Poorter, H., Poot, P., Prior, L., Pyankov, V.I., Roumet, C., Thomas,

- 1 S.C, Tjoelker, M.G., Veneklaas, E.J. & Villar, R. (2004) The world-wide leaf economics
- 2 spectrum. *Nature*, **428**, 821-827.
- 3
- 4 Zar, J.H. (1984) *Biostatistical Analysis*. 2nd edn. Prentice Hall, New Jersey, USA.

Table 1: Summary statistics for all model parameters from the Bayesian path analysis of October 1987 storm impacts on British broadleaved woodlands. Posterior distributions of path coefficients (β s) were estimated from analysis of centred and standardized data. See text and Figs. 1 and 2. Significant effects by Bayes P value are emboldened. Species richness change was analysed as $\ln[(\text{richness } 2002 + 1) / (\text{richness } 1971 + 1)]$

Description	Parameters	Mean	St dev	Monte Carlo SE	2.5%tile	median	97.5%tile	Bayes P value
<i>Regression coefficients</i>								
Species richness change given difference in survey date	β_1	0.001572	0.001418	0.00002	-0.00112	0.001536	0.004463	0.1280
Species richness change given storm exposure	β_2	0.2791	0.1003	0.00137	0.07788	0.2795	0.4765	0.0045
Species richness change given within-site beta diversity in 1971	β_8	-0.1862	0.05196	0.00048	-0.2914	-0.1856	-0.08451	0.0003
Woody basal area change given storm exposure	β_3	0.06268	0.07264	0.00088	-0.08131	0.06193	0.2063	0.1859
pH change given storm exposure	β_9	-0.23	0.2228	0.00481	-0.6668	-0.231	0.2155	0.1130
pH change given woody basal area change	β_5	-0.1243	0.1025	0.00059	-0.3224	-0.1253	0.07724	0.1446
pH change given mean soil pH across survey years	β_7	0.2971	0.06795	0.00283	0.1648	0.2972	0.4309	0
Species richness change given woody basal area change	β_4	-0.4076	0.07978	0.00064	-0.5644	-0.4075	-0.2523	0
Species richness change given pH change	β_6	0.1094	0.04278	0.00030	0.02605	0.1093	0.1937	0.0062
<i>Path coefficients</i>								

Storm effect on richness change via pH change via Basal area change	$\beta s3^* \beta s5^* \beta s6$	-0.00119	0.00221	1.83E-05	-0.00685	-6.47E-04	0.002045	0.2549
Storm effect on richness change via basal area change	$\beta s3^* \beta s4$	-0.03584	0.04174	4.59E-04	-0.1223	-0.03419	0.04349	0.1802
Effect of mean pH across survey years on species richness change via pH change	$\beta s7^* \beta s6$	0.05665	0.02602	2.00E-04	0.01182	0.05454	0.1131	0.0058
<i>Residual standard deviations</i>								
SITE level woody basal area change		0.09801	0.04765	0.00145	0.008076	0.0985	0.1933	
Random intercepts for (pH 71+02)/2		1.148	0.1776	0.00102	0.862	1.127	1.551	
SITE-level pH change		0.4734	0.1066	0.00189	0.2938	0.4633	0.7114	
SITE-level species richness change		0.1521	0.06665	0.00199	0.01575	0.151	0.2884	
Difference in date of survey		39.99	5.894	0.03106	30.4	39.3	53.44	
within-SITE beta diversity in 1971		1.032	0.1532	0.00085	0.7842	1.014	1.382	
PLOT-level woody basal area change		0.4735	0.02047	0.00017	0.4351	0.4728	0.5157	
PLOT-level pH change		0.803	0.03534	0.00027	0.7377	0.8016	0.8761	
PLOT-level mean pH (71+02)/2		0.6154	0.02687	0.00015	0.5654	0.6143	0.6707	
PLOT-level species richness change		0.6217	0.027	0.00024	0.5717	0.6209	0.6771	

Table 2: Differences in species frequency between 1971 and 2002 in paired sample plots (no storm; n=150, storm; n=143). Species are excluded if the cumulative probability of finding the smaller of the two counts, 1971 versus 2002, was >0.05 in both storm and non-storm plots. I = Ancient Woodland Indicators for south east England (Kirby 2006). 'Dir' indicates increased or decreased frequency between surveys where binomial $P \leq 0.05$.

Species	No storm					Storm				
	1971	2002	Change	Bin P	Dir	1971	2002	Change	Bin P	Dir
<i>Solidago virgaurea(I)</i>	0	5	5	0.063		23	2	-21	0.000	down
<i>Oxalis acetosella (I)</i>	42	23	-19	0.025	down	33	14	-19	0.008	down
<i>Deschampsia flexuosa</i>	27	2	-25	0.000	down	21	6	-15	0.006	down
<i>Sanicula europaea (I)</i>	4	2	-2	0.688		17	3	-14	0.003	down
<i>Rubus idaeus</i>	14	4	-10	0.031	down	14	4	-10	0.031	down
<i>Ranunculus acris</i>	5	1	-4	0.219		7	0	-7	0.016	down
<i>Hypericum pulchrum (I)</i>	5	2	-3	0.453		0	6	6	0.031	up
<i>Chrysosplenium oppositifolium(I)</i>	5	5	0	1.000		0	6	6	0.031	up
<i>Carex binervis</i>	0	0	0			0	6	6	0.031	up
<i>Carex pilulifera</i>	3	1	-2	0.625		0	7	7	0.016	up
<i>Luzula campestris/multiflora</i>	4	2	-2	0.688		0	7	7	0.016	up
<i>Cirsium vulgare</i>	1	2	1	1.000		0	7	7	0.016	up
<i>Hypericum tetrapterum</i>	0	0	0			0	7	7	0.016	up
<i>Senecio jacobaea</i>	0	4	4	0.125		0	8	8	0.008	up
<i>Betula pubescens</i>	5	10	5	0.302		2	10	8	0.039	up
<i>Carex remota (I)</i>	2	15	13	0.002	up	3	12	9	0.035	up
<i>Galium saxatile</i>	5	0	-5	0.063		0	9	9	0.004	up
<i>Anthriscus sylvestris</i>	1	4	3	0.375		1	10	9	0.012	up
<i>Anthoxanthum odoratum</i>	5	3	-2	0.727		2	11	9	0.022	up
<i>Teucrium scorodonia</i>	0	0	0			0	9	9	0.004	up
<i>Prunus avium</i>	5	21	16	0.002	up	0	10	10	0.002	up
<i>Ranunculus repens</i>	7	20	13	0.019	up	5	15	10	0.041	up
<i>Arum maculatum</i>	8	25	17	0.005	up	12	23	11	0.090	
<i>Salix caprea</i>	4	5	1	1.000		1	12	11	0.003	up

<i>Digitalis purpurea</i>	4	10	6	0.180		7	19	12	0.029	up
<i>Prunus laurocerasus</i>	3	2	-1	1.000		2	14	12	0.004	up
<i>Cardamine hirsuta/flexuosa</i>	6	7	1	1.000		1	13	12	0.002	up
<i>Lysimachia nemorum (I)</i>	8	10	2	0.815		11	25	14	0.029	up
<i>Athyrium filix-femina</i>	12	13	1	1.000		3	17	14	0.003	up
<i>Holcus mollis</i>	30	22	-8	0.332		16	32	16	0.029	up
<i>Juncus effusus</i>	22	15	-7	0.324		13	30	17	0.014	up
<i>Glechoma hederacea</i>	28	26	-2	0.892		17	34	17	0.024	up
<i>Acer pseudoplatanus</i>	73	80	7	0.628		35	55	20	0.045	up
<i>Dryopteris dilatata</i>	62	64	2	0.929		30	51	21	0.026	up
<i>Carpinus betulus</i>	0	2	2	0.500		20	42	22	0.007	up
<i>Galium aparine</i>	19	40	21	0.009	up	7	31	24	0.000	up
<i>Anemone nemorosa (I)</i>	11	20	9	0.150		3	33	30	0.000	up
<i>Hyacinthoides non-scripta (I)</i>	50	54	4	0.769		52	82	30	0.012	up
<i>Poa nemoralis/trivialis</i>	42	51	9	0.407		35	67	32	0.002	up
<i>Luzula pilosa (I)</i>	46	13	-33	0.000	down	21	13	-8	0.229	
<i>Chamerion angustifolium</i>	34	7	-27	0.000	down	16	9	-7	0.230	
<i>Potentilla sterilis</i>	20	6	-14	0.009	down	13	6	-7	0.167	
<i>Ligustrum vulgare</i>	16	5	-11	0.027	down	8	2	-6	0.109	
<i>Hieracium 'indeterminate'</i>	11	0	-11	0.001	down	3	0	-3	0.250	
<i>Ribes nigrum (I)</i>	10	1	-9	0.012	down	5	2	-3	0.453	
<i>Abies sp.</i>	0	6	6	0.031	up	2	0	-2	0.500	
<i>Epilobium montanum</i>	19	2	-17	0.000	down	14	13	-1	1.000	
<i>Polypodium vulgare sens.lat. (I)</i>	6	0	-6	0.031	down	0	0	0		
<i>Ranunculus ficaria</i>	0	9	9	0.004	up	0	2	2	0.500	
<i>Arrhenatherum elatius</i>	7	0	-7	0.016	down	3	5	2	0.727	
<i>Pinus nigra</i>	0	6	6	0.031	up	0	2	2	0.500	
<i>Cirsium arvense</i>	1	8	7	0.039	up	5	7	2	0.774	
<i>Brachypodium sylvaticum</i>	20	42	22	0.007	up	9	14	5	0.405	
<i>Geranium robertianum</i>	13	31	18	0.010	up	9	15	6	0.307	

<i>Stachys sylvatica</i>	1	8	7	0.039	up	13	19	6	0.377
<i>Arctium agg.</i>	13	2	-11	0.007	down	5	12	7	0.143
<i>Ilex aquifolium</i>	66	107	41	0.002	up	72	90	18	0.181

Fig 1. The woodlands included in this study showing sites inside and outside the region of south east England associated with the track of the October 1987 storm (grey shaded).



Fig 2: Path diagram depicting hypothesized relationships between ecosystem changes and understorey species-richness change measured in 26 broadleaved woodland sites in lowland Britain in 1971 and again in 2002. Ten of the sites were inside the October 1987 storm track. Expected relationships are indicated by arrows each associated with a numbered regression coefficient linked to hypotheses described in the text. Ellipses indicate covariates recorded at the level of plots within woodland sites. Rectangles indicate effects recorded at the site level only.




Fig 3: Results of analysis of October 1987 storm impacts on changes in woodland attributes between 1971 and 2002 in England. Squares indicate site-level covariates. Ovals indicate plot-level covariates. Thick arrows indicate significant paths. Dashed arrows indicate a negative regression relationship. Numbers are regression coefficients based on uncentred and unstandardized data. Understorey species-richness change was analysed as $\ln((\text{richness } 2002 + 1)/(\text{richness } 1971 + 1))$.




Fig 5: Percentage variation explained given hypothesized relationships. Only variables hypothesized to be causally influenced by another variable are on the vertical axis. Therefore, since between-plot changes in species richness were expected to be impacted by all variables its variance was decomposed into the largest number of factors. The height of each bar indicates the total amount of variation in each variable that was explained.




Fig 6: Change in soil pH between 1971 and 2002 versus mean pH in the two survey years. Mean pH is plotted on the axis to avoid regression to the mean artefacts where sampling error results in extreme values switching back to near average values thus causing a strong negative slope. Random sampling of heterogenous soils within plots results in individually large changes in pH. N=293 plots across 26 woodland sites in SE England.




Fig 4: Means \pm 95% confidence intervals for: a) Understorey species-richness, b) soil pH, c) cover-weighted Specific Leaf Area (cSLA), d) woody basal area (m^2) per 200m^2 plot, and e) beta diversity of the understorey in each site (ΣDi). Data from 293 woodland plots within 26 woodland sites in 1971, recorded again in 2002. 10 sites were exposed to the October 1987 storm.

