

LETTERS TO THE EDITOR

To the Editor:

The pathophysiology of menopause related vasomotor symptoms remains an enigma. Recently, authors have hypothesized about the effect of endogenous peripheral nitric oxide (NO) and these symptoms.

Alison J. Huang et al¹ postulate that continuous use of nitroglycerin rapidly leads to tolerance to the drug's vasodilatory effects and cross-tolerance to endogenous nitrates, as a result of enhanced NO degradation.²⁻⁷ Although this tolerance limits the usefulness of nitroglycerin for chest pain, it offers a potentially innovative approach to treating hot flashes, because women who develop cross-tolerance should experience a marked reduction in hot flashes as a result of suppression of NO-mediated peripheral vasodilation.

On the other hand, Bailey et al,⁸ explain that exercise training reduces thermoregulatory dysfunction via stabilization of central thermoregulatory control, that is, lowering core body temperature and improving heat dissipation thresholds, alongside improvements in peripheral mechanisms that allow for greater heat dissipation (sweating sensitivity). These adaptations likely include increases in the number of sweat expulsions per minute, sweat gland hypertrophy, increased NO availability, and others.⁹ The reduction in estrogen associated with menopause causes endothelial dysfunction via decreased NO bioavailability¹⁰ and/or increased reactive oxygen species scavenging NO.¹¹ Exercise increases endothelial NO synthase expression via similar mechanisms of transcriptional regulation to that of estrogen¹² and augments NO-mediated vasodilation.¹³

It appears that these hypotheses are in conflict. Further discussion by the authors may clarify the existing knowledge base regarding the role of nitric oxide and vasodilatation. This information may aid menopause practitioners in educating and caring for their patients.

Financial disclosure/conflicts of interest: None reported.

Ann Steiner, MD
Department of Obstetrics and Gynecology
Perelman School of Medicine
University of Pennsylvania
Radnor, PA

REFERENCES

1. Huang AJ, Cummings SR, Schembri M, Vittinghoff E, Ganz P, Grady D. Continuous transdermal nitroglycerin therapy for menopausal hot flashes: a single-arm, dose-escalation trial. *Menopause* 2016;23:330-334.
2. Chen Z, Foster MW, Zhang J, et al. An essential role for mitochondrial aldehyde dehydrogenase in nitroglycerin bioactivation. *Proc Natl Acad Sci U S A* 2005;102:12159-12164.
3. Daiber A, Mulsch A, Hink U, et al. The oxidative stress concept of nitrate tolerance and the antioxidant properties of hydralazine. *Am J Cardiol* 2005;96:25i-36i.
4. Laursen JB, Boesgaard S, Poulsen HE, Aldershvile J. Nitrate tolerance impairs nitric oxide-mediated vasodilation in vivo. *Cardiovasc Res* 1996;31:814-819.
5. Laursen JB, Mulsch A, Boesgaard S, et al. In vivo nitrate tolerance is not associated with reduced bioconversion of nitroglycerin to nitric oxide. *Circulation* 1996;94:2241-2247.
6. Munzel T, Sayegh H, Freeman BA, Tarpey MM, Harrison DG. Evidence for enhanced vascular superoxide anion production in nitrate tolerance. A novel mechanism underlying tolerance and cross-tolerance. *J Clin Invest* 1995;95:187-194.
7. Thomas GR, DiFabio JM, Gori T, Parker JD. Once daily therapy with isosorbide-5-mononitrate causes endothelial dysfunction in humans: evidence of a free-radical-mediated mechanism. *J Am Coll Cardiol* 2007;49:1289-1295.
8. Bailey TG, Cable N Timothy, et al. Exercise training reduces the frequency of menopausal hot flashes by improving thermoregulatory control. *Menopause* 2016;23:708-718.
9. Shibasaki M, Wilson TE, Crandall CG. Neural control and mechanisms of eccrine sweating during heat stress and exercise. *J Appl Physiol* 2006;100:1692-1701.
10. Virdis A, Ghiadoni L, Pinto S, et al. Mechanisms responsible for endothelial dysfunction associated with acute estrogen deprivation in normotensive women. *Circulation* 2000;101:2258-2263.
11. Kojda G, Harrison D. Interactions between NO and reactive oxygen species: pathophysiological importance in atherosclerosis, hypertension, diabetes and heart failure. *Cardiovasc Res* 1999;43:562-571.
12. Chambliss KL, Shaul PW. Estrogen modulation of endothelial nitric oxide synthase. *Endocr Rev* 2002;23:665-686.
13. Green DJ, Maiorana A, O'Driscoll G, Taylor R. Effect of exercise training on endothelium-derived nitric oxide function in humans. *J Physiol* 2004;561:1-25.

In Reply:

We thank Professor Steiner for her interest in our recent publication and the opportunity to discuss the role of nitric oxide (NO) in menopausal hot flush mechanisms and treatment.

Initially, it is important to highlight that while the two studies investigate the effects of "treatment" via different interventions (exercise training¹ and nitroglycerin therapy²) to improve the frequency and severity of self-reported hot flashes in postmenopausal women; both interventions alter NO availability. In the following text, we have outlined how the two interventions potentially impact hot flashes, potentially via NO pathways.

During a bout of exercise the endothelium releases NO in response to episodic increases in blood flow, and subsequently shear stress along the artery wall. NO diffuses into the smooth muscle cells causing relaxation and artery vasodilation, which

is endothelium-dependent. Exercise training, as performed in our study,¹ causes regular, episodic increases in vascular shear stress that causes NO release during each exercise session³ and thus has been shown to improve both local³ and systemic^{4,5} NO-mediated endothelial function. Nitroglycerin is an NO donor that causes increased systemic NO availability that acts directly on smooth muscle cells causing endothelium-independent vasodilation when administered acutely. Huang et al,² administered a daily dose of nitroglycerin for 6 weeks and inferred that cross-tolerance to organic nitrates occurred with the continuous nitroglycerin administration that suppressed NO-mediated vasodilation. Endothelial function, microvascular function (eg, cutaneous vasodilation) or other thermoregulatory variables (e.g. core body temperature) were not measured by Huang,² nor was the frequency and severity of hot flushes prior to the onset of cross-tolerance where NO availability would have been elevated.

There is evidence that acute alterations in NO have a role in cutaneous vasodilation during an individual hot flush,⁶ but it is currently unknown whether NO is involved in the trigger for hot flushes. Nevertheless, the self-report data from ours¹ and Huang's² studies suggest that both interventions reduce the frequency of hot flushes, which may provide evidence that NO availability is involved in the trigger for hot flushes; the specific mechanism(s) are unclear however it is plausible that NO impacts on the thermoregulatory system,⁷ the sympathetic nervous system⁸ or neurotransmitters in the brain^{9,10}; all of which have been shown to be involved in the hot flush trigger.

The self-report data from ours¹ and Huang's² studies also suggest that both interventions reduce the severity of hot flushes. Physiological severity was not measured in the Huang study, but we have direct evidence that exercise training reduces the physiological severity of hot flushes by reducing the amount of cutaneous vasodilation and sweating during a hot flush.¹¹ We suggest that this is due to more efficient thermoregulation, including endothelial dependent NO-mediated vasodilation. Huang infers that cross tolerance to NO would reduce cutaneous vasodilation during a hot flush.

There are a number of distinct differences between the two interventions that are noteworthy. Exercise training enhances endothelial function,¹ which has been shown to be impaired in postmenopausal women and especially in symptomatic women,¹² whereas chronic daily administration of nitroglycerin can cause "cross tolerance" to organic nitrates and possibly negate the requirement for endothelial dependent vasodilation, both of which may lead to endothelial dysfunction, that might have implications for those with coronary artery disease.^{13,14} Furthermore, exercise does not cause headaches like nitroglycerin, and exercise training has additional benefits to reducing hot flush frequency and severity (eg, improved thermoregulation, neural activity¹⁵) as well as several well-known multifaceted benefits for improving health in postmenopausal women.¹⁶

It is also important to emphasize that both of these studies^{1,2} require confirmation with full randomized control trials

to determine efficacy, ideally which include outcomes measurements of both self-reported and physiologically measured hot flushes that include cutaneous vasodilation, sweating, and vascular function.¹¹

Financial disclosure/conflicts of interest: None reported.

Tom G. Bailey, PhD
School of Health and Sport Sciences
University of the Sunshine Coast
Maroochydore DC, Australia

Tim Cable, PhD
Research Institute for Sport and Exercise Sciences
Liverpool John Moores University
Liverpool, UK
Department of Sports Science
Aspire Academy
Doha, Qatar

Nabil Aziz, MD
Department of Gynecology and Reproductive Medicine
Liverpool Women's Hospital
Liverpool, UK

Rebecca Dobson, MD
Victoria S. Sprung, PhD
Department of Obesity and Endocrinology
University of Liverpool
Liverpool, UK

David A. Low, PhD
Helen Jones, PhD
Research Institute for Sport and Exercise Sciences
Liverpool John Moores University
Liverpool, UK

REFERENCES

1. Bailey TG, Cable NT, Dobson RL, et al. Exercise training reduces the frequency of menopausal hot flushes by improving thermoregulatory control. *Menopause* 2016;23:708-718.
2. Huang AJ, Cummings SR, Schembri M, Vittinghoff E, Ganz P, Grady D. Continuous transdermal nitroglycerin therapy for menopausal hot flushes: a single-arm, dose-escalation trial. *Menopause* 2016;23:330-334.
3. Atkinson CL, Carter HH, Dawson EA, Naylor LH, Thijssen DH, Green DJ. Impact of handgrip exercise intensity on brachial artery flow-mediated dilation. *Eur J Appl Physiol* 2015;115:1705-1713.
4. Birk GK, Dawson EA, Atkinson C, et al. Brachial artery adaptation to lower limb exercise training: role of shear stress. *J Appl Physiol* 2012;112:1653-1658.
5. Hambrecht R, Adams V, Erbs S, et al. Regular physical activity improves endothelial function in patients with coronary artery disease by increasing phosphorylation of endothelial nitric oxide synthase. *Circulation* 2003;107:3152-3158.
6. Hubing KA, Wingo JE, Brothers RM, Del Coso J, Low DA, Crandall CG. Nitric oxide synthase inhibition attenuates cutaneous vasodilation during postmenopausal hot flash episodes. *Menopause* 2010;17:978-982.
7. Freedman RR. Menopausal hot flushes: mechanisms, endocrinology, treatment. *J Steroid Biochem Molec Biol* 2014;142c:115-120.
8. Low D, Hubing K, Del Coso J, Crandall C. Mechanisms of cutaneous vasodilation during the menopausal hot flash. *Menopause* 2011;18:359-365.
9. Mittelman-Smith MA, Williams H, Krajewski-Hall SJ, McMullen NT, Rance NE. Role for kisspeptin/neurokinin B/dynorphin (KNDy) neurons in cutaneous vasodilation and the estrogen modulation of body temperature. *Proc Natl Acad Sci U S A* 2012;109:846-851.
10. Jayasena CN, Comminos AN, Stefanopoulou E, et al. Neurokinin B administration induces hot flushes in women. *Sci Rep* 2015;5:8466.
11. Bailey TG, Cable NT, Aziz N, et al. Exercise training reduces the acute physiological severity of post-menopausal hot flushes. *J Physiol* 2016;594:657-667.
12. Bechlioulis A, Kalantaridou SN, Naka KK, et al. Endothelial function, but not carotid intima-media thickness, is affected early in menopause and is

- associated with severity of hot flashes. *J Clin Endocrinol Metab* 2010;95:1199-1206.
13. Nakamura Y, Moss AJ, Brown MW, Kinoshita M, Kawai C. Long-term nitrate use may be deleterious in ischemic heart disease: A study using the databases from two large-scale postinfarction studies. Multicenter Myocardial Ischemia Research Group. *Am Heart J* 1999;138(3 pt 1): 577-585.
 14. Parker JD. Nitrate tolerance, oxidative stress, and mitochondrial function: another worrisome chapter on the effects of organic nitrates. *J Clin Invest* 2004;113:352-354.
 15. Oneda B, Cardoso CG Jr, Forjaz CL, et al. Effects of estrogen therapy and aerobic training on sympathetic activity and hemodynamics in healthy postmenopausal women: a double-blind randomized trial. *Menopause* 2014;21:369-375.
 16. Nunes PR, Barcelos LC, Oliveira AA, et al. Effect of resistance training on muscular strength and indicators of abdominal adiposity, metabolic risk, and inflammation in postmenopausal women: controlled and randomized clinical trial of efficacy of training volume. *Age (Dordr)* 2016;38:40.

In Reply:

We agree that the pathophysiology of menopause-related vasomotor symptoms remains frustratingly unclear. However, our exploration of continuous nitroglycerin as a treatment strategy for hot flashes is not predicated on any particular hypothesis about the causal trigger for these symptoms.¹ We only assume that, once triggered, hot flashes manifest in the form of cutaneous vasodilation that is likely mediated by nitric oxide (NO) activation of vascular smooth muscle.² If this is true, then treatment strategies directed at dampening NO-mediated vasodilation (such as induction of nitrate cross-tolerance) may suppress the primary physical manifestations of hot flashes, regardless of the original stimulus for these symptoms.

We appreciate that menopause may be associated with changes in vascular endothelial function that are in turn associated with decreased NO bioavailability, and that a prior study has suggested that changes in endothelial-dependent flow-mediated dilation are associated with more severe vasomotor symptoms.³ However, evidence is lacking that these changes are specifically responsible for development of menopausal vasomotor symptoms. Similar changes in flow-mediated dilation have also been observed in other populations such as aging men⁴ or cardiac patients using long-term oral nitrates,⁵ and yet high rates of vasomotor symptoms are not reported in these populations. Even if decreased endothelial response to endogenous NO is associated with vasomotor symptoms, it is unlikely that it is the primary cause of these symptoms.

With regard to the exercise study by Bailey et al,⁶ larger-scale randomized trials of exercise-based interventions have unfortunately failed to demonstrate a reduction in vasomotor

symptoms relative to controls.⁷ But even if we assume that exercise is effective in reducing hot flashes, it does not necessarily follow that this is mediated by exercise-induced increase in NO availability.⁶ Regular exercise has a multitude of effects—not just at the tissue-specific level, but also at the level of the organ system and the individual woman and her perception of vasomotor symptoms. Reported reductions in hot flashes among women who exercise may (and probably do) arise from other mechanisms.

Two things are certain—hot flashes involve vasodilation, and NO is a vasodilator. If we go one step further and accept laboratory-based evidence that NO is the primary vasodilator implicated in hot flashes,² then interventions that target NO-mediated vasodilation may offer promise for treatment. Of course, this does not lessen the need for a better understanding of the upstream stimuli that give rise to these perplexing symptoms in the first place.

Financial disclosure/conflicts of interest: None reported.

Allison J. Huang, MD, MAS
Peter Ganz, MD
Deborah Grady, MD, MPH
Department of Medicine
University of California
San Francisco, CA

Steven R. Cummings, MD, MPH
San Francisco Coordinating Center
California Pacific Medical Center Research Institute
San Francisco, CA

REFERENCES

1. Huang AJ, Cummings SR, Schembri M, Vittinghoff E, Ganz P, Grady D. Continuous transdermal nitroglycerin therapy for menopausal hot flashes: a single-arm dose-escalation trial. *Menopause* 2016;23:330-334.
2. Hubing KA, Wingo JE, Brothers RM, Del Coso J, Low DA, Crandall CG. Nitric oxide synthase inhibition attenuates cutaneous vasodilation during postmenopausal hot flash episodes. *Menopause* 2010;17:978-982.
3. Bechlioulis A, Kalantaridou SN, Naka KK, et al. Endothelial function, but not carotid intima-media thickness, is affected early in menopause and is associated with severity of hot flashes. *J Clin Endocrinol Metab* 2010;95:1199-1206.
4. Celermajer DS, Sorensen KE, Spiegelhalter DJ, Georgakopoulos D, Robinson J, Deanfield JE. Aging is associated with endothelial dysfunction in healthy men years before the age-related decline in women. *J Am Coll Cardiol* 1994;24:471-476.
5. Lai J, Wu B, Sun J, Shang Y, Zhu J. Long-term isosorbide mononitrate treatment impairs endothelial function in patients with coronary artery disease. *Coronary Artery Dis* 2013;24:566-571.
6. Bailey TG, Cable NT, Aziz N, et al. Exercise training reduces the frequency of menopausal hot flashes by improving thermoregulatory control. *Menopause* 2016;23:708-718.
7. Daley A, Stokes-Lampard H, Thomas A, MacArthur C. Exercise for vasomotor menopausal symptoms. *Cochrane Database Syst Rev* 2014; 4:CD006108.