

## Modulation of the Passive Membrane Properties of a Pair of Strongly Electrically Coupled Neurons by Anaesthetics

Munir M Qazzaz<sup>1</sup> and William Winlow<sup>2\*</sup>

<sup>1</sup>Faculty of Pharmacy, Nursing and Health Professions, Birzeit University, Birzeit, West Bank, Palestine

<sup>2</sup>Department of Biology, Università degli Studi di Napoli, Federico II, Via Cintia 26, Napoli, Italia and Institute of Ageing and Chronic Diseases, The APEX Building, 6 West Derby Street, University of Liverpool, Liverpool L7 8TX, UK

\*Corresponding Author: William Winlow, Department of Biology, University of Naples, Federico II, Italy and Institute of Ageing and Chronic Diseases, University of Liverpool, Liverpool, UK.

Received: May 02, 2017; Published: May 26, 2017

### Abstract

Differential effects of volatile general anaesthetics and sodium pentobarbital on input resistance [ $R_m$ ], membrane time constant [ $\tau_m$ ] and capacitance [ $C_m$ ] of VD1 and RPD2 neurons of *Lymnaea* are reported in this paper. The volatile anaesthetics halothane and isoflurane caused a significant reduction in the input resistance, membrane time constant and capacitance in a dose dependent manner and halothane was more potent than isoflurane. The clinical concentration [100  $\mu$ M] of pentobarbitone caused no change in  $R_m$ ,  $\tau_m$  and  $C_m$  of VD1 and RPD2, whereas 2.0 and 0.5 mM pentobarbitone increased them significantly during phase I but caused no change in these parameters during phase II.

**Keywords:** General Anaesthetics; Electrical Coupling; Identified Neurons; Input Resistance; Time Constant; Capacitance

### Abbreviations

$C_m$ : Membrane Capacitance;  $R_m$ : Membrane Input Resistance;  $\tau_m$ : Membrane Time Constant; PDS: Paroxysmal Depolarizing Shift; RPD2: Right Parietal Dorsal Neuron 2; VD1: Visceral Dorsal Neuron 1

### Introduction

Electrical synapses [gap junctions] are now known to be important in promoting synchronicity in many regions of the mammalian brain [1], in the brains of lower vertebrates [2] and also in invertebrate preparations [3] as first described in the crayfish by Furshpan and Potter [4,5]. In the pond snail *Lymnaea stagnalis* there are numerous examples of electrically coupled neurons and neural clusters [6-9]. Recent evidence has shown that gap junctions may exhibit synaptic plasticity, being dynamically regulated by neuromodulators such as dopamine [10]. It is therefore of some importance that the actions of volatile and systemic anaesthetics on electrical synapses should be understood, although a previous study suggested that electrical synapses are less sensitive to most anaesthetics than chemical synapses [11].


In a previous paper [12] we demonstrated that barbiturates and volatile anaesthetics have markedly different effects on electrically coupled neurons. For example, volatile anaesthetics cause a significant reduction in the coupling coefficient between the giant VD1 and RPD2 neurons of *Lymnaea stagnalis* [12] and also between the smaller neurons of the left and right pedal A clusters [9]. However, sodium pentobarbitone, which is widely used as a veterinary anaesthetic [13,14] had very different actions. Pentobarbitone, within the clinical concentration range, caused no change in the coupling coefficient though the action potential was abolished and the cells became quiescent. In contrast, 0.5 and 2.0 mM pentobarbitone first generated paroxysmal depolarising shifts [PDS] while increasing the coupling coefficient but it then returned to control values as the neurons became quiescent.

The effects described above are may be partly caused by alterations in the passive membrane properties of the nerve cells for instance by altering the membrane input resistance  $[R_m]$  and capacitance  $[C_m]$  which will also have direct effects on the membrane time constant  $[\tau_m]$ . The details of these passive properties of nerve cell membranes are well described elsewhere [15,16], but although subthreshold changes in passive membrane properties may reflect changes in the lipid membrane, other components of the membrane will also be affected. Here we describe the actions of the same concentrations of these anaesthetics as we used in our previous experiments [12] on the passive membrane properties of VD1 and RPD2: i.e.  $R_m$ ,  $\tau_m$  and  $C_m$ . Preliminary data have appeared elsewhere [17].

**Methods**

Intact *Lymnaea* brain preparations were prepared and intracellular recordings of neuronal electrical activity were made from target neurons according to the methods of Benjamin and Winlow [6] and Qazzaz and Winlow [12]. Sub-threshold hyperpolarizing current pulses [0.1 nA for 1 sec] were injected into VD1 or RPD2 neurons via the recording microelectrode using Neurolog NL102 bridge balance preamplifiers. Data was captured using a CED 1401 data acquisition interface and a personal computer with Spike2 data capture and analysis software installed. All data were statistically analyzed using the paired Student’s t-test and expressed as mean ± standard error of the mean [SEM]. Values were considered statistically significant at a confidence level of 95%. All recordings were made in the summer months (June to August) since date collected at other times of year were variable and inconsistent.

When a square current pulse is injected into a neuron two individual and separate stages are seen before it reaches the final potential value. First, there is a very fast stage followed by a slower exponential stage (Figure 1). This is due to the membrane acting as a capacitor, where the membrane represents a good insulator between two good electrolytic conductors. Being a capacitor means that it stores charge across its surface opposing any changes in its potential. Membrane capacitance can be defined as the ratio between the charge separated by the membrane and the change in its potential. Furthermore, membrane capacitance is directly proportional to its surface area is and inversely proportional to its thickness.




**Figure 1:** Calculating input resistance - Injection of a hyperpolarizing current pulse into a neuron cell body causes a drop in membrane potential for the duration of the pulse where  $\Delta V$  [change in membrane voltage] can be calculated from Ohm’s law  $[\Delta V = I.R_m]$  where  $I$  is the applied current and  $R_m$  is the calculated input resistance of the cell membrane.

To calculate membrane capacitance, two passive properties are required; the input resistance and membrane time constant. The former can be determined by passing constant current pulse through the neuronal membrane to earth as shown in Figure 1. Knowing both the applied current and the induced voltage change the input resistance of the membrane can be calculated according to Ohm’s Law [voltage [V] = current [I] x input resistance [Rm]] The membrane time constant is defined as the time needed for a current pulse to charge the membrane capacitance to 63% of its final value [15]. Figure 2 illustrates how to measure the time constant  $[\tau_m]$  and the equation below demonstrates the relationship between the membrane time constant  $[\tau_m]$ , input resistance  $[R_m]$  and capacitance  $[C_m]$ :

$$\tau_m = R_m \times C_m$$

Membrane capacitance [ $C_m$ ] was calculated using input resistance and the time constant data, according to this equation.



**Figure 2:** Measuring membrane time constant - A diagram showing the change in membrane potential during the exponential stage as a function of time after the injection of a hyperpolarising current pulse. Membrane time constant [ $\tau_m$ ] is calculated by measuring the time at 63% of the potential change. Adapted from Kandel [15].

## Results


### Input Resistance

The input resistance of VD1 and RPD2 were measured separately in the presence and absence of the anaesthetics. Because VD1 and RPD2 are strongly electrically connected, in 10 preparations we either crushed the commissure between the visceral and right parietal ganglia to isolate the neurons from one another or destroyed one of the pair of neurons. There was no change in the input resistance of either neuron under these conditions.

**Effects of halothane:** Both neurons became quiescent in 1% and 2% halothane as previously reported [12]. The control input resistance of VD1 [ $285 \pm 19 \text{ M}\Omega$ ;  $n = 10$ ] and RPD2 [ $344 \pm 11 \text{ M}\Omega$ ;  $n = 8$ ] significantly decreased [ $193 \pm 6 \text{ M}\Omega$  and  $243 \pm 7 \text{ M}\Omega$  respectively] after application of 1% halothane. 2% halothane further decreased the input resistance of both neurons [ $143 \pm 6 \text{ M}\Omega$  and  $163 \pm 7 \text{ M}\Omega$  respectively] and these effects were completely reversed by washing in anaesthetic free saline as illustrated in Figure 3 for RPD2.


**Effects of isoflurane:** Isoflurane was applied as for halothane and at the same percentage concentrations, again resulting in quiescence of both neurons. The control input resistances of VD1 and RPD2 were  $277 \pm 10 \text{ M}\Omega$  [ $n = 10$ ] and  $344 \pm 11 \text{ M}\Omega$  [ $n = 8$ ] respectively. As with halothane, isoflurane decreased the input resistance reversibly in a dose dependent manner, as illustrated in Figure 3. The input resistances of VD1 and RPD2 in 1% halothane were  $245 \pm 7 \text{ M}\Omega$  and  $290 \pm 12 \text{ M}\Omega$  respectively and in 2% isoflurane the respective values were  $219 \pm 7 \text{ M}\Omega$  and  $205 \pm 7 \text{ M}\Omega$ .

**Effects of sodium pentobarbitone:** Three concentrations of pentobarbitone were applied,  $100 \mu\text{M}$  [within the clinical range],  $0.5 \text{ mM}$  and  $2 \text{ mM}$ , but the order of application was varied and in some cases, they were applied after halothane and/or isoflurane with a 30 min delay between each concentration and/or drug. No differences in the responses to pentobarbitone were noted as a consequence of these different application regimes.



**Figure 3:** Effects of halothane and isoflurane on RPD2 – The effects on VD1 [not shown] were similar. The actions of the anaesthetics on both neurons were dose dependent and both neurons became quiescent in the presence of the anaesthetics. Halothane was more potent in reducing the input resistance than isoflurane as demonstrated in Figures 5 and 6.

**Application of 2mM pentobarbitone:** 2mM pentobarbitone produced two distinct phases of activity in VD1/RPD2 (Figure 4a) as previously reported [12]. Phase I started within a few seconds of the application of the drug and action potential activity was modified into paroxysmal depolarizing shifts [PDS] within a few minutes. This was followed by quiescence of the neurons in phase II. There was a clear and distinctive difference in the input resistances of VD1 and RPD2 in these two phases. In phase I, from control values of  $298 \pm 7 \text{ M}\Omega$  [n = 12] and  $355 \pm 9 \text{ M}\Omega$  [n = 14] for VD1 and RPD2 respectively, the input resistances significantly increased to  $373 \pm 8 \text{ M}\Omega$  and  $410 \pm 14 \text{ M}\Omega$  respectively. However, the input resistances were not significantly different from control values in phase II [ $288 \pm 10 \text{ M}\Omega$  and  $356 \pm 9 \text{ M}\Omega$  respectively].




**Figure 4:** Effects of sodium pentobarbitone on VD1 on input resistance – The effects on RPD2 [not shown] were similar. High concentrations of sodium pentobarbitone caused paroxysmal depolarizing shifts [PDS] [phase 1] followed a few minutes later by quiescence [phase 2]. During phase 1,  $R_m$  increased significantly, but this was not dose-dependent and then declined to control values in phase 2, again not dose dependent at these higher concentrations. At clinical concentrations of  $100 \mu\text{M}$ , the neurons simply became quiescent without any intervening PDS, but  $R_m$  did not differ from control values.

**Application of 0.5mM pentobarbitone:** Very similar results were obtained as compared with 2mM pentobarbitone, with apparently no dose-dependency at these concentrations outside the clinical range, but their effects were completely reversible (Figure 4b). Control values for input resistance of  $286 \pm 7 \text{ M}\Omega$  [ $n = 12$ ] and  $369 \pm 11 \text{ M}\Omega$  [ $n = 14$ ] were obtained for VD1 and RPD2 respectively. During phase I these values increased to  $362 \pm 13 \text{ M}\Omega$  and  $427 \pm 13 \text{ M}\Omega$  respectively, while during the quiescence of phase II there was no significant difference from control values [ $287 \pm 12 \pm 9 \text{ M}\Omega$  and  $376 \pm 12 \text{ M}\Omega$  respectively].

**Application of 100 $\mu\text{M}$  pentobarbitone:** 100 $\mu\text{M}$  pentobarbitone, which is in the clinical range, had little effect on input resistance of VD1/RPD2, although both cells became quiescent (Figure 4c). Control values were  $295 \pm 13 \text{ M}\Omega$  [ $n = 8$ ] and  $322 \pm 8 \text{ M}\Omega$  [ $n = 6$ ] respectively, which were little different from experimental values of  $280 \pm 15 \text{ M}\Omega$  and  $329 \pm 10 \text{ M}\Omega$  respectively.

The data for pentobarbitone, isoflurane and halothane are summarized in Figure 5.




**Figure 5:** Summary of the actions of sodium pentobarbitone, halothane and isoflurane on input resistance of VD1 [upper panel] and RPD2 [lower panel]. The  $R_m$  of VD1 was significantly increased during the PDS phase [I] induced by both 0.5 mM and 2.0 mM pentobarbitone [ $n = 10$ ,  $P \leq 0.01$ ] but there was no significant change in phase II in comparison with control when the cells became quiescent.  $R_m$  was significantly reduced by both 1% and 2% isoflurane in a dose dependent manner [ $n = 10$ ,  $P \leq 0.01$ ]. 1% and 2% halothane induced a greater reduction in  $R_m$  in comparison with isoflurane [ $\leq 0.01$ ] again in a dose dependent manner. The input resistance of RPD2 was affected by the two volatile anaesthetics [ $n = 12$ ,  $P \leq 0.01$ ] and barbiturates [ $n = 12$ ,  $P \leq 0.01$ ] in a similar way to VD1. Error bars represent mean  $\pm$  standard error.

### Time Constant and Membrane Capacitance


The time constant can be affected differently depending upon the drug or anaesthetic agent as shown in figure 6. Clearly the changes in time constant and input resistance also determine changes in membrane capacitance and changes in both  $\tau_m$  and  $C_m$  are summarized as percentages of control values for VD1 and RPD2 in Figures 7 and 8.

The calculated  $C_m$  of VD1 and RPD2, showed that control values of  $0.43 \pm 0.036 \text{ nF}$  [ $n = 10$ ] and  $0.57 \pm 0.042 \text{ nF}$  [ $n = 8$ ] decreased to  $0.27 \pm 0.017 \text{ nF}$  and  $0.32 \pm 0.021 \text{ nF}$  with 1% halothane and  $0.17 \pm 0.015 \text{ nF}$  and  $0.22 \pm 0.019 \text{ nF}$  with 2% halothane respectively (Figure 8).

**Halothane:** Perfusion of the CNS with HBS-containing halothane [1% and 2%] produced a significant reduction of  $\tau_m$  in both neurons VD1 and RPD2. Control values of  $121 \pm 11 \text{ msec}$  and  $202 \pm 14 \text{ msec}$  respectively were decreased to  $54 \pm 7 \text{ msec}$  and  $77 \pm 6 \text{ msec}$  after the application of 1% halothane and  $29 \pm 05 \text{ msec}$  and  $36 \pm 09 \text{ msec}$  with 2% halothane. In all experiments, the control values were regained when the CNS was washed with HBS (Figure 7).



**Figure 6:** An example to demonstrate the differential effects of general anaesthesia on membrane time constant. The exponential curves in the diagrams a and b represent the second stage of an injected hyperpolarizing current pulse into a single VD1 neuron. [a] Shows the effect of 2 mM pentobarbitone on the control [ $\tau = 121 \pm 11$  msec]. After application, the time constant increased significantly [ $\tau = 201 \pm 08$  msec;  $n = 12$ ]. [b] In sharp contrast 2% halothane significantly reduced the time constant [ $\tau = 29 \pm 05$  msec;  $n = 10$ ].



**Figure 7:** Effects of volatile anaesthetics and barbiturates on time constant of VD1 and RPD2 neurons. The different anaesthetics [halothane, isoflurane and pentobarbitone; at different concentrations] were applied to the CNS in different order. These anaesthetics did not alter the effect of each other. Volatile anaesthetics decreased the time constant in a dose dependent manner and the effect of halothane was greater than that of isoflurane. However, there was no significant change in their values from the control value when 100 µM pentobarbitone was applied. Furthermore, 2.0 mM and 0.5 mM pentobarbitone significantly increased their time constant during phase I, but they showed no significant difference from their control value in phase II. Note that the bars were calculated as a percentage of the halothane control. Bars are mean  $\pm$  SEM.

**Isoflurane:** Isoflurane produced similar results to those produced by halothane but its effects were slightly weaker. 1% isoflurane significantly reduced the  $\tau_m$  of VD1 and RPD2 from the control value [ $0.126 \pm 12$  msec,  $n = 10$  and  $205 \pm 13$  msec,  $n = 8$ ] to  $75 \pm 07$  msec and  $111 \pm 08$  msec respectively. A greater reduction was demonstrated with 2% isoflurane [ $45 \pm 07$  msec and  $50 \pm 04$  msec] (Figure 7).


When  $C_m$  of VD1 and RPD2 was calculated, it was found to be reduced in a dose dependent manner after perfusing the CNS with HBS-containing 1% and 2% isoflurane. Control values of  $0.45 \pm 0.42$  nF [ $n = 10$ ] and  $0.55 \pm 0.034$  nF [ $n = 8$ ] were decreased to  $0.31 \pm 0.023$  nF and  $0.39 \pm 0.033$  nF by 1% and  $0.20 \pm 0.019$  nF and  $0.24 \pm 0.014$  nF with 2% (Figure 8).

**Sodium pentobarbitone:** Pentobarbitone affected the measured parameters of the VD1 and RPD2 neurons differently from that produced by volatile anaesthetics.

Effects of 2 mM pentobarbitone: During phase I of the effect of 2 mM pentobarbitone,  $\tau_m$  of VD1 and RPD2 was found to increase significantly from the control [ $125 \pm 04$  msec [ $n = 12$ ] and  $209 \pm 08$  msec [ $n = 14$ ]] to  $201 \pm 08$  msec and  $315 \pm 09$  msec respectively. No significant difference from the control values in phase II was noticed [ $128 \pm 06$  msec and  $211 \pm 07$  msec respectively] (Figure 7).

2 mM pentobarbitone produced an increase in the  $C_m$  of VD1 and RPD2 during phase I [ $0.58 \pm 0.022$  nF [ $n = 12$ ] and  $0.71 \pm 0.028$  nF [ $n = 14$ ]] compared to controls of  $0.44 \pm 0.018$  nF and  $0.57 \pm 0.035$  nF respectively. However, no significant change was noticed during phase II [ $0.43 \pm 0.015$  nF and  $0.57 \pm 0.019$  nF respectively] (Figure 8).

Effects of 0.5 mM pentobarbitone - After the application of 0.5 mM pentobarbitone on the neurons VD1 and RPD2, their  $\tau_m$  was found to increase significantly from their controls [ $126 \pm 06$  msec,  $n = 12$  and  $207 \pm 06$  msec,  $n = 14$ ] to  $217 \pm 08$  msec and  $302 \pm 08$  msec respectively (Figure 7). These values returned almost to control values during phase II [ $125 \pm 07$  msec and  $213 \pm 05$  msec respectively].



**Figure 8:** Effects of volatile anaesthetics and barbiturates on membrane capacitance of VD1 and RPD2 neurons. The different general anaesthetic agents [halothane, isoflurane and pentobarbitone] was tested at different concentrations and their application was in different order. They did not alter the effect of each other on membrane capacitance. Volatile anaesthetics decreased the membrane capacitance in a dose dependent manner with the effect of halothane being greater than that of isoflurane. However, there was no significant change in its value, from its control value, when 100  $\mu$ M pentobarbitone was applied. Furthermore, 2.0 mM and 0.5 mM pentobarbitone significantly increased the membrane capacitance of VD1 [a] and RPD2 [b] neurons during phase I, but they showed no significant change in their capacitance value in phase II. Note that the bars were calculated as a percentage of the halothane control. Bars are mean  $\pm$  SEM.

Similar results to those with 2 mM pentobarbitone were gained after the application of 0.5 mM pentobarbitone. Control values of  $C_m$  for VD1 and RPD2 were  $0.43 \pm 0.016$  nF [ $n = 12$ ] and  $0.59 \pm 0.025$  nF [ $n = 14$ ] respectively. There was a significant increase in their  $C_m$  during phase I [ $0.56 \pm 0.017$  nF and  $0.72 \pm 0.032$  nF] but no significant difference from the control values during phase II [ $0.44 \pm 0.023$  nF and  $0.59 \pm 0.013$  nF] respectively (Figure 8).

**Effects of 100  $\mu$ M pentobarbitone:** After the application of 100  $\mu$ M pentobarbitone, the effects on  $\tau_m$  and  $C_m$  of VD1 and RPD2 were completely different from the effect of the other concentrations. There was no significant difference on either of these parameters from the control [ $\tau_m = 133 \pm 06$  msec [ $n = 8$ ] and  $209 \pm 07$  msec [ $n = 6$ ] which became  $138 \pm 09$  msec and  $212 \pm 07$  msec (Figure 7) and [ $C_m$ ] were equal to  $0.45 \pm 0.021$  nF [ $n = 8$ ] and  $0.59 \pm 0.031$  nF [ $n = 6$ ] and became equal to  $0.44 \pm 0.017$  nF and  $0.59 \pm 0.036$  nF respectively (Figure 8).

## Discussion

It is already known that the effects of general anaesthetics are very complex and, furthermore, the effects on one neuron cannot be generalized to other neurons [18-20]. Over the years, a variety of effects on ionic currents, intracellular components and on chemical and electrical synapses have been reported [12,21-29]. Here we have shown that there are distinct differences between the effects of volatile anaesthetics and barbiturates on the same neuron. The data we report clearly demonstrate that the application of the volatile anaesthetics isoflurane and halothane reduce  $R_m$ ,  $C_m$  and  $\tau_m$  in a dose dependent manner, whereas the responses to sodium pentobarbitone were quite different with a significant rise in all parameters in phase I, followed by a fall to control values in quiescent phase II.

**Variability between nerve cells:** The effects of anaesthetics on neuronal function are very diverse, for example halothane caused different effects on different neurons in the same and in different species. It hyper-activates some *Lymnaea* neurons, such as in the CGCs, J cells, right parietal A group and visceral M group causing them to exhibit PDS [19]. This response was seen in the intact brain [20,30] as well as in isolated cultured neurons [31]. In sharp contrast, other *Lymnaea* neurons became hyperpolarized and consequently quiescent [20,26,32,33]. Indeed, the spontaneous firing activity of RPeD1, RPD1 and VV1/2 [20] and VD1 and RPD2 [26] decreased, eventually leading to quiescence after the application of halothane or isoflurane. In a study on the small pedal A cluster neurons of *Lymnaea* [9] we discovered that they became quiescent after the application of volatile anaesthetics, with no observed change in the resting membrane potential. This indicates that general anaesthetics shut off the voltage-gated channels involved in action potential generation confirming the results of Winlow and Yar [34] in pedal I cluster neurons, who recently demonstrated that a high-voltage activated Ca current was reversibly depressed in a dose dependent manner by the exposure of cultured *Lymnaea* neurons to clinical concentrations of halothane, again in pedal I-cluster neurons [35]. Similarly, Winlow, et al. [36] found that general anaesthetics blocked voltage-gated calcium and potassium currents [25,36]. The likely molecular targets of general anaesthetics are discussed in detail elsewhere [28,29].

## What is happening in the neuronal membrane?

### Time constant and Membrane capacitance

It could be argued that the effects of anaesthetics on the measured  $\tau_m$  and the calculated  $C_m$  are entirely due to interactions with membrane lipids, since under normal circumstances  $C_m$  is not expected to alter very much, but small changes could occur due to changes in the physical properties of the cell membrane [specifically in the lipid bilayer], i.e. surface area or thickness [15]. If the neuronal membrane was a true capacitor any increase in  $\tau_m$  and  $C_m$  would suggest that membrane had become thinner while a decrease in  $\tau_m$  and  $C_m$  would suggest membrane thickening. Thus, the volatile anaesthetics which decrease  $C_m$  would thicken the membrane while the barbiturate would be expected to thin it during phase I. However, cell membranes consist of both lipids and proteins and surround complex intracellular constituents which may have effects on the membrane and may themselves be modified by applied anaesthetics. Since the now outdated lipid hypothesis of general anaesthesia has been replaced by the membrane protein hypothesis of general anaesthetic action [29,37] it is more probable that the changes in the passive membrane properties induced by anaesthetics are due largely to effects on membrane proteins. In particular, these same results could also be produced by an increases or decreases in membrane conductance [ $\equiv$  increases or decreases in input resistance respectively] i.e. the membrane becomes more or less leaky, perhaps due to the direct actions of anaesthetics on membrane proteins.

### Input resistance

**Actions of volatile anaesthetics:** Our data clearly demonstrate that volatile anaesthetics and sodium pentobarbitone have differential effects on input resistance and we suspect that they are due to modifications of membrane proteins by the anaesthetics. Volatile anaesthetics are known to inhibit or potentiate ligand-gated and voltage-gated ion channels at clinically relevant concentrations [29]. Franks and Lieb [33] reported that, in the CNS of *Lymnaea* there are two different types of neurons; sensitive and insensitive neurons to anaesthetics. They suggested the presence of a novel potassium current in the sensitive neurons which they called the anaesthetic-activated potassium current [ $I_{K[an]}$ ]. They proposed that this current was activated by the application of halothane, but the activation of  $I_{K[an]}$  caused the membrane potential to hyperpolarize far below the threshold that is necessary to generate an action potential. We previously demonstrated that the membrane potentials of VD1 and RPD2 were significantly hyperpolarised after the application of halothane or isoflurane and during phase II of 0.5 or 2.0 mM pentobarbitone [9,27] in agreement with of Franks and Lieb [27] and Nicoll and Madison [38]. Thus, a reduced input resistance indicates an increased neuronal conductance implying the presence of either  $I_{K[an]}$  or other anaesthetic potentiated currents in VD1/RPD2.

**Actions of pentobarbitone:** The application of pentobarbitone [100  $\mu$ M] abolished the action potentials and slightly hyperpolarised the membrane of VD1 and RPD2 neurons with no change in their input resistance. Calcium-activated potassium current is one possible pathway through which general anaesthetics hyperpolarise neuronal membranes. Alternatively, activation of a chloride current is another possibility.  $\gamma$ -aminobutyric acid [GABA] is the principle inhibitory amino acid neurotransmitter known to induce a chloride current [33,38-40]. Nicoll and Madison [38] demonstrated that a reduction in the extracellular concentration of chloride in frog motoneurons had no effect on the hyperpolarisation caused by anaesthetics. They concluded that general anaesthetics do not produce their effects by increasing chloride conductance. However, sodium pentobarbitone also causes hyperpolarization of VD1/RPD2 and this is mimicked by 100 $\mu$ M GABA which causes hyperpolarization (from -45.5mV to -53.4 mV, n = 7; P = 0.002 – Moccia., *et al.* [40]) of the giant dopamine containing, multiaction neuron RPeD1 [42] in *Lymnaea* via GABA<sub>A</sub> receptors. We therefore suggest that a GABA<sub>A</sub> chloride current is responsible for the pentobarbitone induced hyperpolarization in VD1/RPD2.

Moody., *et al.* [43] found that inhalational anaesthetics increase the chloride conductance in rat cortical synaptoneurosome and this was supported by findings from Nakahiro., *et al.* [44] on cultured rat dorsal root ganglion neurons. They concluded that inhalational anaesthetics augmented the sustained current evoked by GABA and this work was supported by several further studies [39,40] and Zimmerman., *et al.* [45] found that the responses to the application of GABA to rat hippocampal neurons were prolonged by many general anaesthetics. Thus, the actions on GABA receptors are not limited to barbiturate anaesthetics.

**Paroxysmal depolarising shifts [PDS]:** VD1 and RPD2 neurons generated PDS after the application of 0.5 and 2.0 mM pentobarbitone before becoming quiescent. Furthermore, during which the input resistance was significantly increased. PDS is a kind of neuronal epileptiform activity that can be produced under pathological conditions in different species, which “typically consists of an abrupt depolarisation of the cell membrane to a level at which the cell is unable to generate a full-sized action potential, usually followed by repolarisation for a brief period before the membrane depolarises again, producing square-wave membrane potential oscillation” [46]. A full explanation of PDS is given elsewhere [47]. However it is believed to be connected to rising intracellular calcium concentration [ $Ca^{2+}$ ] [46-49] and Moghadam and Winlow [50] have demonstrated that pentobarbitone raises [ $Ca^{2+}$ ] while blocking both calcium and potassium channels. An increase in Na<sup>+</sup> conductance may also explain the induction of PDS [47,51,52] by high concentrations of pentobarbitone but it does not explain the accompanying increase in the membrane input resistance of VD1 and RPD2 and this needs further attention, unless the decreased conductance of sodium and calcium channels is much greater than the presumed increase in sodium conductance.

### Does electrical coupling vary on a seasonal basis?

**Assessment of  $R_m$**  in VD1/RPD2 appears to vary significantly between different laboratories. Here we have shown the control  $R_m$  to be in the order of 300 M $\Omega$  for VD1 and 350 M $\Omega$  for RPD2 [see Figure 5]. This is in the normal range for molluscan neurons [ $10^5$  to  $10^8 \Omega$  – [15]], but was found to be rather variable, in the region of 14 - 80 M $\Omega$  by Benjamin and Pilkington [53]. Further studies [54] suggested that  $R_m$

for VD1 was 100 M $\Omega$  and for RPD2  $R_m$  was 50 M $\Omega$ . The explanation of these differences may be due to the well-known seasonal plasticity in the nervous system of *Lymnaea* [55-59]. Furthermore, we were only able to obtain consistent data on the passive membrane properties of VD1/RPD2 during the summer months. The chemical synaptic connections between the giant dopamine cell and its follower cells fail in winter snails [56], which disables the respiratory central pattern generator [60,61] and Wood and Winlow [62] reported a difference in the frequency-half width correlation in specific neurons between summer and winter. Furthermore, Qazzaz [58] demonstrated seasonal variation in the weak electrical coupling between small pedal A cluster neurons [9]. During the summer months (June to August), the electrical connections between these cells were present and very clear, but disappeared completely during the winter season (October to January). Further studies on the electrical connections of VD1-RPD2 to determine whether any changes in  $R_m$  are related to seasonal factors are therefore required. Another possible cause of variation in  $R_m$  may be due to the ageing process which is known to reduce  $R_m$  VD1/RPD2 [63] and in some *Aplysia* neurons [64]. This is largely due to the increasing size of the neurons and may disturb the synchrony of firing of neuron discharges in the VD1-RPD2 pairing and may also reduce their sensitivity to synaptic inputs [63]. According to Benjamin and Pilkington [53] there is a low resistance/high conductance pathway between VD1 and RPD2 of about 20M $\Omega$  when recorded in the cell bodies. The cell membranes have much higher resistance which ensures access of currents between the cells which is sufficient to keep them synchronized. Although we now know the effects of the anaesthetics on the soma membranes, their effects on the junctional resistance remain to be clarified.

## Conclusions

It is now appreciated that electrical synapses are highly plastic, modifiable by a variety of agents, including neurotransmitters [27,29], and subject to block by a wide variety of agents, many of which are non-specific [64].

Here we have demonstrated that the volatile anaesthetics isoflurane and halothane have similar actions on electrically coupled synapses between the identified neurons VD1 and RPD2 in *Lymnaea*, where they both cause a dose dependent decrease in electrical coupling between the cells. These actions are in contrast to the effects of sodium pentobarbitone which at low doses [100  $\mu$ M] causes quiescence with no alteration in coupling coefficient, but at higher doses [0.5 and 2.00 mM] causes an increase in coupling coefficient and PDS [phase I], followed by quiescence and a reduction of coupling coefficient to control values [phase II].

Given that both the volatile anaesthetics and sodium pentobarbitone raise  $[Ca^{2+}]_i$ , further studies to elucidate the modulation of membrane potential and input resistance by anaesthetics, need to be performed while monitoring  $[Ca^{2+}]_i$  during application of each class of anaesthetic. Furthermore, the effects of anaesthetics on the junctional resistance and the junctional proteins need to be understood.

## Bibliography

1. Curti, *et al.* "Synergy between electrical coupling and membrane properties promotes strong synchronization of neurons of the mesencephalic trigeminal nucleus". *Journal of Neuroscience* 32.13 (2012): 4341-4359.
2. Michael J Hull, *et al.* "Modelling the Effects of Electrical Coupling between Unmyelinated Axons of Brainstem Neurons Controlling Rhythmic Activity". *PLOS Computational Biology* 11.5 (2015): e1004240.
3. Carew TJ and Kandel ER. "Two functional effects of decreased conductance EPSP's: synaptic augmentation and increased electrotonic coupling". *Science* 192.4235 (1976): 150-153.
4. Furshpan EJ and Potter DD. "Transmission at the giant motor synapses of the crayfish". *Journal of Physiology* 45.2 (1959) 289-325.
5. Furshpan EJ and Potter DD. "Mechanism of nerve impulse transmission at a crayfish synapse". *Nature* 180 (1957): 342-343.
6. Benjamin P.R. and Winlow W. "The distribution of three wide-acting synaptic inputs to identified neurones in the isolated brain of *Lymnaea stagnalis* (L)". *Comparative Biochemistry and Physiology* 70A (1981): 293-307.

7. Haydon PG and Winlow W. "Multipolar neurones of *Lymnaea stagnalis*. I. Multiple spike initiation sites and propagation failure allow neuronal compartmentalisation". *Journal of Comparative Physiology* 147 (1982): 503-510.
8. Syed NI and Winlow W. "Coordination of locomotor and cardiorespiratory networks of *Lymnaea stagnalis* by a pair of identified interneurons". *Journal of Experimental Biology* 158 (1991): 37-62.
9. Qazzaz M and Winlow W. "Effects of volatile anaesthetics on the electrical activity and coupling coefficient of weakly electrically coupled neurones". *Acta Biologica Hungarica* 50.1-3 (1999): 199-213.
10. Pereda AE, *et al.* "Gap junction-mediated electrical transmission: regulatory mechanisms and plasticity". *Biochimica et Biophysica Acta* 1828.1 (2013): 134-146.
11. Johnston MF, *et al.* "Interaction of anaesthetics with electrical synapses". *Nature* 286 (1980): 498-500.
12. Qazzaz MM and Winlow W. "Differential Actions of Volatile Anaesthetics and a Systemic Barbiturate on Strongly Electrically Coupled Neurons". *EC Neurology* 2.4 (2015): 188-204.
13. "Barbiturates drug profile". European Monitoring Centre for Drugs and Drug addiction (2015).
14. "LARC Formulary - Anesthesia and Analgesia in Laboratory Animals". Oregon State University (2011): 1-29.
15. Kandel ER. "Cellular Basis of Behavior – an introduction to behavioral neurobiology". Freeman, San Francisco (1976).
16. Nicholls JG., *et al.* "From Neuron to Brain". 4<sup>th</sup> edition. Sinauer, Sunderland, Mass (2001).
17. Qazzaz M and Winlow W. "Volatile anaesthetics and barbiturates have differential effects on time constant and input capacitance of identified molluscan neurones". *Journal of Physiology* 504 (1997): 16P-17P.
18. Haydon PG., *et al.* "The effects of menthol on central neurones of the pond-snail, *Lymnaea stagnalis* (L)". *Comparative Biochemistry and Physiology* 73C (1982): 95-100.
19. Girdlestone D and Winlow W. "A relationship between neuronal topography and responses to anaesthetics in *Lymnaea*". *Journal of Physiology* 386 (1987): 33.
20. Winlow W., *et al.* "Differential effects of general anaesthetics on identified molluscan neurones in vivo and in culture". *General Pharmacology* 23.6 (1992): 985-992.
21. Richards CD. "Actions of general anaesthetics on synaptic transmission in the CNS". *British Journal of Anaesthesia* 55.3 (1978): 201-207.
22. Richards CD. "Actions of general anaesthetics on synaptic transmission". *General Pharmacology* 9 (1983): 287-293.
23. Girdlestone D., *et al.* "The actions of halothane on spontaneous activity, action potential shape and synaptic connections of the giant serotonin-containing neurone of *Lymnaea stagnalis* (L)". *Comparative Biochemistry and Physiology* 93.2 (1989): 333-339.
24. Frenkel C., *et al.* "Molecular actions of pentobarbital isomers on sodium channels from human brain cortex". *Anesthesiology* 72.4 (1990): 640-649.

25. Moghadam HF and Winlow W. "Actions of isoflurane and halothane on whole cell  $\text{Ca}^{2+}$  and  $\text{K}^+$  currents in cultured, identified molluscan neurones". *British Journal of Anaesthesia* 75 (1995): 661-662.
26. Qazzaz M and Winlow W. "Differential effects of volatile anaesthetics and barbiturates on electrical coupling between identified molluscan neurones". *Journal of Physiology* 489 (1995): 21.
27. Qazzaz M and Winlow W. "Volatile anaesthetics and barbiturates have different actions on the input resistance of an identified molluscan neurone". *British Journal of Anaesthesia* 75 (1995): 656P-657P.
28. Hemmings HC. "Molecular targets of general anesthetics in the nervous system". In "Suppressing the mind, Contemporary clinical neuroscience". Humana Press (2010): 11-31.
29. Hudson AE., et al. "Pharmacology of inhaled anesthetics". In ClinicalKey.com (2016): 159-179.
30. Walcourt-Ambakederemo A and Winlow W. "Halothane induces paroxysmal depolarization shifts in synaptically isolated Lymnaea neurones in situ and in culture". *Journal of Physiology* 473 (1993): 187.
31. Winlow W., et al. "Studies on mechanisms underlying general anaesthesia using cultured molluscan neurones". *Molecular and Cellular Mechanisms of Alcohol and Anaesthetics* 625 (1991): 269-272.
32. Winlow W and Girdlestone D. "Modulation of nervous activity in Lymnaea by general anaesthetics". *Symposia Biologica Hungarica* 36 (1988): 451-461.
33. Franks NP and Lieb WR. "Selective actions of volatile general anaesthetics on identified neurons". *Annals of the New York Academy of Sciences* 625 (1991): 54-70.
34. Winlow W and Yar T. "Effect of halothane on cultured neurones". *British Journal of Anaesthesia* 69 (1992): 530P-531P.
35. Yar T and Winlow W. "Effects of halothane on whole-cell calcium channel currents in cultured Lymnaea neurones". *Neurology* 4.1 (2016): 3-22.
36. Winlow W., et al. "Multiple cellular and subcellular actions of general anaesthetics on cultured molluscan neurones". *Acta Biologica Hungarica* 46.2-4 (1995): 381-393.
37. Franks NP and Lieb WR. "Molecular and cellular mechanisms of general anaesthesia". *Nature* 367.6464 (1994): 607-614.
38. Nicoll RA and Madison DV. "General anaesthetics hyperpolarise neurones in the vertebrates central nervous system". *Science* 217.4564 (1982): 1055-1057.
39. Wakamori M., et al. "Effects of two volatile anaesthetics and a volatile convulsant on the excitatory and inhibitory amino acid responses in dissociated CNS neurones of rat". *Journal of Neurophysiology* 66.6 (1991): 2014-2021.
40. Mody I., et al. "Halothane enhances tonic neuronal inhibition by elevating intracellular calcium". *Brain Research* 538.2 (1991): 319-323.
41. Moccia F., et al. "GABA (A)- and AMPA-like receptors modulate the Activity of an identified neuron within the central pattern generator of the pond snail Lymnaea stagnalis". *Invertebrate Neuroscience* 9.1 (2009): 29-41.

42. Winlow W and Benjamin PR. "Postsynaptic effects of a multi-action giant interneurone on identified snail neurones". *Nature* 268.5617 (1977): 263-265.
43. Moody E., *et al.* "Modulation benzodiazepine/-amino butyric acid receptor chloride channel complex by inhalational anaesthetics". *Journal of Neurochemistry* 51.5 (1988):1386-1393.
44. Nakahiro M., *et al.* "General anaesthetics modulate GABA receptor channels complex in rat dorsal root ganglion neurones". *FASEB Journal* 3.7 (1989): 1850-1854.
45. Zimmerman SA., *et al.* "Potentiation of -aminobutyric acid A receptor Cl- current correlates with in vivo anaesthetic potency". *Journal of Pharmacology and Experimental Therapeutics* 270 (1994): 987-991.
46. Brown AM and McCrohan C. "Differential responses of two identified neurones of the pond snail *Lymnaea stagnalis* to the convulsant drug pentelenetetrazol". *Brain Research* 565.2 (1991): 247-253.
47. Maxim B., *et al.* "Cellular and network mechanisms of electrographic seizures". *Drug Discovery Today: Disease Models* 5.1 (2008): 45-57.
48. Walden J., *et al.* "Epileptogenesis: Contribution of calcium ions and antiepileptic calcium antagonists". *Acta Neurologica Scandinavica* 86 (1992): 41-46.
49. Angstadt JD and Choo JJ. "Sodium-dependent plateau potentials in cultured Retzius cells of the medicinal leech". *Journal of Neurophysiology* 76.3 (1996): 1491-1502.
50. Moghadam HF and Winlow W. "Pentobarbitone alters intracellular calcium concentration in cultured, identified, molluscan neurones". *Journal of Physiology* 526 (2000): 62.
51. McCrohan CR and Gillette R. "Cyclic AMP-stimulated sodium current in identified feeding neurones of *Lymnaea stagnalis*". *Brain Research* 438.1-2 (1988): 115-123.
52. McCrohan CR and Gillette R. "Enhancement of cyclic AMP-dependent sodium current by the convulsant drug pentylenetetrazol". *Brain Research* 452.1-2 (1988): 21-27.
53. Benjamin PR and Pilkington JB. "The electrotonic location of low-resistance intercellular junctions between a pair of giant neurones in the snail *Lymnaea*". *Journal of Physiology* 370 (1986):111-126.
54. Beekharry CC., *et al.* "Role for electrical synapses in shaping the output of coupled peptidergic neurons from *Lymnaea*". *Brain Research* 1603 (2015): 8-21.
55. Winlow W. "The plastic nature of action potentials". In: "Cellular Basis of Neuronal Plasticity", (ed.) AGM Bulloch, Manchester University Press, Manchester (1989): 3-27.
56. Hetherington MS., *et al.* "A quantitative analysis of the biogenic amines in the central ganglia of the pond snail *Lymnaea stagnalis* (L.)". *Comparative Biochemistry and Physiology* 107.1 (1994): 83-93.
57. Copping J., *et al.* "Seasonal plasticity of synaptic connections between identified neurones in *Lymnaea*". In: "Neurobiology of Invertebrates. Membranes, Chemical Signalling and Systems Approach". (eds) K. Elekes and J. Salanki (2000): 205-210.

58. Winlow W and Polese G "A Neuroplastic Network Underlying Behaviour and Seasonal Change in *Lymnaea stagnalis*: A Neuroecological Standpoint". In: "Neuroecology and Neuroethology in Molluscs: the interface between behaviour and environment". Nova Science Publishers, Inc, New York (2014): 145-176.
59. Qazzaz M. "Actions of general anaesthetics on electrical synapses". PhD Thesis, University of Leeds (1997).
60. Syed NI, *et al.* "In vitro reconstruction of the respiratory central pattern generator of the mollusc *Lymnaea*". *Science* 250.4978 (1990): 282-285.
61. Syed NI and Winlow W. "Respiratory behaviour in the pond snail *Lymnaea stagnalis*. II. Neural elements of the central pattern generator". *Journal of Comparative Physiology* 169.5 (1991): 557-568.
62. Wood G and Winlow W. "Seasonal variation of spike width in identified neurones of *Lymnaea stagnalis*". *Journal of Physiology* 495 (1996): 38P.
63. Janse C., *et al.* "Peptidergic neurons in the ageing molluscan brain". In: "Comparative Aspects of Neuropeptide Function" (Ed) Ernst Florey and George B Stefano. Manchester University Press (1991): 128-139.
64. Peretz B. "Neurobiology of ageing: An hypothesis of neuron viability". In: "The Cellular Basis of neuronal Plasticity: physiology, morphology and biochemistry of molluscan neurons" (Ed) Andrew Bulloch. Manchester University Press (1989): 200-220.

**Volume 6 Issue 4 May 2017**

**© All rights reserved by Munir M Qazzaz and William Winlow.**