

Manuscript Number: SCHRES-D-17-00387R1

Title: Birth experiences, trauma responses and self-concept in postpartum psychotic-like experiences

Article Type: Full Length Article

Keywords: Psychosis; postnatal; trauma; self-concept; adjustment.

Corresponding Author: Dr. Lyndsey Holt, DClinPsy

Corresponding Author's Institution: Mersey Care NHS Foundation Trust

First Author: Lyndsey Holt, DClinPsy

Order of Authors: Lyndsey Holt, DClinPsy; Pauline Slade; William Sellwood

Abstract: The frequency of psychotic-like experiences (PLEs) amongst new mothers is beginning to be explored but the mechanisms underlying such experiences are yet to be understood. First time mothers (N = 10,000) receiving maternity care via the UK National Health Service were contacted postnatally via Emma's Diary, an online resource for mothers. Measures assessed birth experience, trauma appraisals, post-traumatic stress symptoms, adjustment to motherhood, self-concept clarity and PLEs (in the form of hallucinations and delusions). There was a 13.9% response rate (N = 1,393) and 1,303 participants reported experiencing at least one PLE (93.5%). Three competing nested path models were analysed.

A more negative birth experience directly predicted delusions, but not hallucinations. Trauma appraisals and poorer adjustment to motherhood indirectly predicted PLEs, via disturbed self-concept clarity. Post-traumatic stress symptoms directly predicted the occurrence of all PLEs.

PLEs in first time mothers may be more common than previously thought. A key new understanding is that where new mothers have experienced birth as traumatic and are struggling with adjustment to their new role, this can link to disturbances in a coherent sense of self (self-concept clarity) and be an important predictor of PLEs. Understanding the development of PLEs in new mothers may be helpful in postnatal care, as would public health interventions aimed at reducing the sense of abnormality or stigma surrounding such experiences.

**Cover letter**

Dear Editor,

Please find attached the revised manuscript entitled 'Birth experiences, trauma responses and self-concept in postpartum psychotic-like experiences'.

Also enclosed is a document stating the comments made by the reviewer, along with responses and amendments made to the manuscript.

Thank you for considering this article for publication.

We look forward to hearing from you.

Yours sincerely,

Dr. Lyndsey Holt  
Clinical Psychologist

# Birth experiences, trauma responses and self-concept in postpartum psychotic-like experiences

Lyndsey Holt<sup>a\*</sup>

Pauline Slade<sup>b</sup>

William Sellwood<sup>c</sup>

<sup>a</sup>Mersey Care NHS Foundation Trust, Park Lodge CMHT, Orphan Drive, Liverpool, L6 7UN, UK. E-mail address: [lyndsey.holt@merseycare.nhs.uk](mailto:lyndsey.holt@merseycare.nhs.uk). Telephone: +44 (0151) 330 8942. Fax: 0151 3308990

<sup>b</sup>Institute of Psychology, Health & Society, Brownlow Hill, University of Liverpool, Liverpool LG9 3GB, UK. E-mail address: [Pauline.Slade@liverpool.ac.uk](mailto:Pauline.Slade@liverpool.ac.uk)

<sup>c</sup>Division of Health Research, University of Lancaster, Bailrigg, LA1 4YW, UK. E-mail address: [b.sellwood@lancaster.ac.uk](mailto:b.sellwood@lancaster.ac.uk)

\*Correspondence

## Abstract

The frequency of psychotic-like experiences (PLEs) amongst new mothers is beginning to be explored but the mechanisms underlying such experiences are yet to be understood. First time mothers ( $N = 10,000$ ) receiving maternity care via the UK National Health Service were contacted postnatally via *Emma's Diary*, an online resource for mothers. Measures assessed birth experience, trauma appraisals, post-traumatic stress symptoms, adjustment to motherhood, self-concept clarity and PLEs (in the form of hallucinations and delusions). There was a 13.9% response rate ( $N = 1,393$ ) and 1,303 participants reported experiencing at least one PLE (93.5%). Three competing nested path models were analysed.

A more negative birth experience directly predicted delusions, but not hallucinations. Trauma appraisals and poorer adjustment to motherhood indirectly predicted PLEs, via disturbed self-concept clarity. Post-traumatic stress symptoms directly predicted the occurrence of all PLEs.

PLEs in first time mothers may be more common than previously thought. A key new understanding is that where new mothers have experienced birth as traumatic and are struggling with adjustment to their new role, this can link to disturbances in a coherent sense of self (self-concept clarity) and be an important predictor of PLEs. Understanding the development of PLEs in new mothers may be helpful in postnatal care, as would public health interventions aimed at reducing the sense of abnormality or stigma surrounding such experiences.

Keywords: Psychosis; postnatal; trauma; self-concept; adjustment.

## 1. Introduction

One to two women per thousand is diagnosed with postpartum psychosis (Valdimarsdóttir et al., 2009; VanderKruik et al., 2017), with onset in the first postnatal month (Sit et al., 2006). However, the occurrence of psychotic-like experiences (PLEs), non-clinical experiences of hallucinations or delusions similar to psychosis but in a diminished form (Cicero et al., 2013), are common (Mannion and Slade, 2014). The mechanisms underlying the development of such experiences are still unclear.

The continuity hypotheses (Johns and Van Os, 2001) postulates that psychosis exists on a spectrum within the population, from no psychotic experiences or symptoms through to clinical levels of psychosis. Postpartum psychosis is associated with distress, illness related behaviours, impairment of functioning and help-seeking behaviour (Van Os et al., 2009; *The Diagnostic and Statistical Manual of Mental Disorders*, DSM-5, American Psychiatric Association, 2013). However, PLEs are often not associated with distress or illness-related behaviour, and maybe fleeting in duration. Longitudinal studies indicate that people who experience PLEs are at increased risk of clinical psychosis (Poulton et al., 2000). Five percent of the population experience PLEs during their lifetime (Van Os et al., 2009). Attempts have been made to understand the development of PLEs and what may cause a transition to psychosis. Childhood traumas are associated with PLEs and psychosis in adulthood (Peters et al., 2016; Read et al., 2008; Varese et al., 2012b). Certain traumas are related to specific psychotic symptoms (Bentall et al., 2014) and these relationships are mediated by factors including dissociation (Varese et al., 2012a), and attachment (Sitko et al., 2014). A further potential mediator is self-concept clarity (SCC) i.e. the extent to which a person's beliefs about themselves are well-defined, confidently held, internally coherent, stable and cognitively accessible (Campbell et al., 1996). SCC is reduced in schizophrenia (Cicero et al., 2015). SCC mediates the relationship between childhood traumas and psychosis (Evans et al., 2015). If PLEs and psychosis can be regarded as on the same continuum (Van Os et al., 2009), then difficult life events and low SCC may play key roles in the development of both.

PLEs may occur in mothers as result of childbirth (Barratt et al. 2012; Mannion and Slade, 2014). Despite childbirth being a normal event, up to half of women experience birth as traumatic

(O'Donovan et al., 2014), and three to fifteen percent meet the criteria for post-traumatic stress disorder (PTSD) (Ayers et al., 2016; Czarnocka and Slade, 2000). Both fearful birth experiences (Mannion and Slade, 2014) and a traumatic birth (Barratt, 2012) are associated with increased risk of PLEs. Given that birth can be experienced as traumatic and lead to post-traumatic stress symptoms (PTSS) and becoming a mother entails substantial life adjustments, which in turn may threaten self-concept, the relevance of self-concept to the occurrence of PLEs warrants investigation in new mothers.

This study aimed to explore the relationships between birth experience and trauma, adjustment to motherhood, SCC and PLEs in new mothers. We aimed to test three competing hypotheses (Figure 1). Each model hypothesised that birth experience would directly predict the occurrence of PLEs, as has been found previously (Mannion and Slade, 2014). We hypothesised that birth experience would predict trauma appraisals, PTSS and adjustment to motherhood. Birth experiences have been found to predict PTSS (Edworthy et al., 2008) and trauma appraisals and PTSS are associated (Ehlers and Clark, 2000), whilst the relationship between birth experience and adjustment to motherhood is yet to be established. We also hypothesized that trauma appraisals, PTSS and adjustment to motherhood would in turn predict SCC, as they are part of the cognitive affect system that corresponds to self-concept (Markus and Wurf, 1987). SCC would then predict PLEs, in accordance with the general psychosis literature (Cicero et al., 2013). As the association between PTSS and PLEs has been established (Alsawy et al., 2015; Hamner et al., 1999) we included this relationship throughout the models. However, we were uncertain whether trauma appraisals and adjustment to motherhood would directly predict PLEs, therefore these paths were removed in two of the models. We also wished to test whether birth experience directly predicted SCC in one of the models, as this relationship is yet to be established. In accordance with the defined onset of postpartum psychosis in the DSM-5, it is hypothesised that PLEs occur in the first month postpartum. However, in order to encapsulate the entire theorised period of risk we extended this time frame to two months.

## 2. Method

### 2.1 Participants

Women aged 16-50, proficient in English, who gave birth to their first child 2-6 months before recruitment, were included. This time frame was chosen for consistency, and to ensure all participants had passed the hypothesised risk period for PLEs. To ensure our estimates were conservative, and not confounded by participants with known mental health problems receiving treatment, participants with a history of psychosis or having had input from perinatal mental health teams (from self-report prior to questionnaire completion) were excluded.

### 2.2 Procedure

University of Liverpool ethics committee provided approval. Recruitment took place via *Emma's Diary* ([www.emmasdiary.co.uk](http://www.emmasdiary.co.uk)), an online resource for mothers. Information about Emma's Diary is routinely supplied to pregnant women by their NHS general practitioner or midwife. An emailed invitation was sent to 10,000 website registrants meeting the inclusion criteria. Scales were administered via Qualtrics ([www.qualtrics.com](http://www.qualtrics.com)); 2,870 participants commenced the survey, 77 participants were excluded (35 past history of psychosis, 42 input from perinatal mental health teams, 9 both) and 1,400 did not complete the data set. 1,393 participants completed the survey (13.93% - Figure 2).

Participants retrospectively reported on the first two months postpartum (the theorised phase of risk), apart from depressive symptoms which participants reported concurrently to control for current levels of depressive symptoms in analyses.

### 2.3 Measures

Demographic and obstetric details were recorded.

*2.3.1 Birth experience: The Wijma Delivery Experience Questionnaire B (Version B; Wijma et al., 1998).*

This scale targets appraisals of the birth experience, with higher scores indicating higher fear. This scale demonstrated excellent internal consistency ( $\alpha = .94$ ).

### 2.3.2 Trauma appraisals

Questions were developed based upon the *The Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV-TR; American Psychiatric Association 2000). Participants were asked if at any time during childbirth they (a) thought that their or their babies' life was at risk or at risk of serious injury (a)(i) if so, was this for themselves, their baby or both of them (b) experienced intense fear, helplessness or horror. Responses were scored on a binary scale (0 = *no*, 1 = *yes*). The internal consistency was acceptable ( $\alpha = .69$ ).

### 2.3.3 Post-traumatic stress symptoms: *The Impact of Event Scale - Revised* (Weiss, 2007).

Three subscales cover arousal, intrusions and avoidance, and provide a total. The clinical cut off is  $\geq 33$  (Creamer et al., 2003). Cronbach's alpha for the current sample was .93.

### 2.3.4 Adjustment to motherhood: *The Being a Mother Scale (BaM-13; Matthey, 2011)*.

This measures social isolation, regret, coping and guilt, with higher scores indicating greater difficulty adjusting to motherhood. Good internal consistency ( $\alpha = .87$ ) was found in the present sample.

### 2.3.5 Self-concept clarity: *The Self-Concept Clarity Scale* (Campbell et al., 1996)

This measures the extent to which participants' self-concepts are clear, confidently held, internally consistent, stable and cognitively accessible with higher scores indicating higher clarity. The scale showed excellent internal consistency in this sample,  $\alpha = .90$ .

### 2.3.6 Hallucination experiences: *Launay Slade Hallucination Scale (LSHS; Launay and Slade, 1981)*.

Measures hallucinations on 12 binary scale (0 = *false*, 1 = *true*) items, with higher scores indicating more hallucinations. There are 4 subscales; vivid thoughts, vivid day dreams, auditory hallucinations and visual hallucinations. Full scale internal consistency was acceptable ( $\alpha = .66$ ).

### 2.3.7 Delusional experiences: *Peters Delusion Inventory (PDI-21; Peters et al., 2004)*

This measures delusions across 21 binary items, with higher scores indicating more delusions. If participants answer yes further questions ask how distressing and intrusive the delusions are, along with how much they believe them to be true. Only data on occurrence of delusions were included in analyses. Internal consistency was good,  $\alpha = .73$ .

### 2.3.8 Depressive symptoms: *The Edinburgh Postnatal Depression Scale (EPDS; Cox et al., 1987).*

Depression is assessed using 10 items. The clinical cut off is a score  $\geq 13$  (Matthey et al., 2006). The scale is widely used and showed good internal consistency,  $\alpha = .86$ .

### 2.3.9 Sleep: *Pittsburgh Sleep Quality Index (PSQI; Buysse et al., 1989).*

Participants answered 18 questions on 4-point Likert scales (0 = *not at all*, 3 = *three or more times a week*). The overall scale has a binary outcome (0 = *poor sleep*, 1 = *good sleep*) and showed good internal consistency,  $\alpha = .76$ .

## 2.4 Data analysis

Descriptive and correlational analyses were undertaken in SPSS v.21 (IBM Corporation., Armonk, NY, USA) and path analysis in AMOS v. 22 (IBM Corporation., Armonk, NY, USA). Path analysis allows for: (i) estimation of direct and indirect (mediation) effects, (ii) multiple endogenous (dependent) variables to be modelled simultaneously, (iii) interdependence of endogenous variables to be accounted for, and (iv) analysis of several hypothesised models. To test the three competing models, the regression weights on the paths not included in each model were constrained to zero. We controlled for variables potentially having significant relationships with relevant variables: age, ethnicity, marital status, employment, mode of birth, education, depression and sleep.

Data for both dependent variables (hallucinations and delusions) were positively skewed. Therefore bias-corrected bootstrapping (Hayes and Scharkow, 2013) and 10,000 resamples were used (Mallinckrodt et al., 2006). To assess model fit the Bollen and Stine (1992) bootstrap adjusted  $p$  value was used in relation to the Chi-square statistic, along with the CFI and RMSEA. After the model was fitted, separate indirect effects via a single mediator were obtained separately, by constraining both the correlation between the alternative mediators, and path from the predictor to alternative mediators, to 0 (MacKinnon, 2008).

### 2.4.1 Missing and incomplete data

Due to text input being used for certain questions sleep data on the PSQI were missing for 223 participants. Other missing data were minimal (marital status only, Table 1). To minimise data loss, the PSQI was recoded into three binary contrast variables (unknown sleep, poor sleep and good

sleep). Only unknown sleep and good sleep were included in the path models, allowing poor sleep to be the reference category.

### **3. Results**

#### **3.1 Descriptive statistics and correlations**

See Table 1 for demographic characteristics and mode of birth.

##### *3.1.1 Prevalence of psychotic like experiences*

One thousand and eighty-five participants (77.9%) experienced at least one hallucination, whilst 1,217 participants experienced at least one delusion (87.4%). When considering both hallucinations and delusions, 999 participants experienced at least one hallucination and at least one delusion (71.7 %). To understand the distribution of PLEs as a whole, the scales were combined; a total of 1,303 participants experienced at least one PLE (93.58%). Table 4 presents the items of the LSHS and PDI-21 in order of frequency of endorsement. Descriptive data are presented in Table 2 and correlations between demographic details and symptom measures in Table 3.

#### **3.2 Path models**

We analysed three competing theoretical path models (Figure 1). In all hypothesised models, birth experiences were the exogenous predictor variable and there were two stages of mediation (i) trauma appraisals, PTSS and adjustment to motherhood (ii) SCC. Hallucination and delusional experiences were the outcome variables.

Table 5 outlines the goodness-of-fit statistics for the competing hypothesised models.

#### **3.3 Best fitting model**

Model 2 (Figure 3), supporting hypotheses 2, was the best fit. It did not have a significantly worse fit in comparison to model 1,  $\Delta\chi^2(3) = 0.36, p = .95$ , and was a good fit for the data;  $\chi^2(4) = 4.49, p = .34$ ; CFI = 1.00, RMSEA = 0.01, 90% CI [0.00, 0.03]  $p = .98$ . Path and associated maximum likelihood and bootstrap SEs/CIs are presented in Table 6.

### **4. Discussion**

This is the first study to explore the role of predictive and mediating variables, in relation to PLEs in new mothers ( $N = 1,393$ ). There was a high prevalence of PLEs (93.58%). Assuming that the

86.07% of participants who did not complete the survey did not experience any PLEs, 13.03% of all the invited participants experienced at least one PLE over the short postnatal time frame specified. This is substantially higher than the five percent lifetime prevalence rate in the general population (Van Os et al., 2009). This emphasises that PLEs are not uncommon postnatally.

Our findings supported model 2. Birth experience predicted delusions, but not hallucinations. Trauma appraisals of the birth and struggling in adjustment to motherhood indirectly predicted PLEs, a relationship dependent on SCC. PTSS predicted PLEs.

The intra and inter-personal transitions for new mothers theoretically correspond to changes in self-concept (Markus and Wurf, 1987), and SCC has been shown to be in flux at this time (Smith, 1999). The finding that SCC predicted PLEs is consistent with studies in other populations (Cicero et al., 2013; Evans et al., 2015).

It was surprising that birth experience did not predict hallucinations, as Mannion and Slade (2014) found that fear during childbirth predicted both hallucinations and delusions. As expected, PTSS predicted all PLEs, which is comparable to general population research (Morrison et al., 2003).

The finding that trauma appraisals and adjustment to motherhood were mediating factors between birth experience and SCC, but PTSS were not, is noteworthy. This suggests that traumatic birth or difficulties with adjustment, rather than symptom experience, are important in disruption of SCC. As expected, PTSS mediated the relationship between birth experience and both hallucinations and delusions. However, given that this was full mediation for hallucinations and partial mediation for delusions, there may be other important factors in the birth experience and delusion relationship that we did not investigate. It also suggests that the mechanisms underlying these PLEs may differ.

In comparison to both Barratt (2012) and Mannion and Slade (2014) studies, the reported occurrence of PLEs were higher. Possibly, this is due to data being collected anonymously online, as the women could honestly reflect on their experiences without fear of disclosure. Mothers can potentially experience feelings of shame surrounding disclosure of psychological difficulties (Slade et al., 2010). Given that psychosis in the general population (Johns and Van Os, 2001) and postnatal psychosis (Valdimarsdóttir et al., 2009) have been found to be associated with lower socio-economic

status, we may have found higher prevalence rates if the sample were more representative, so our estimates may be conservative.

The significant two stage mediation pathways highlight the importance of indirect relationships between birth experience and PLEs. It was unexpected that PTSS did not predict SCC, and the relationship between fear during childbirth and SCC is a novel finding. This emphasises the importance of experience (e.g. fear, appraisals of trauma, difficulties with adjusting) over and above PTSS in the prediction of SCC. Clinically this is important as women whose birth does not go as planned are more likely to experience birth as traumatic (Soet et al., 2003) and 34% of first time mothers birth experiences are worse than expected (National Perinatal Epidemiology Unit, 2014). In line with self-concept theories (Markus and Wurf, 1987) we can infer that experiences, incongruent with expectations, can impact upon a women's SCC, in terms of fundamental assumptions about themselves. For example, women who prior to birth believe themselves to be competent, organised and in control may have difficulties making sense of an experience where they feel out of control, inept and a failure. This is noteworthy as first time mothers are more likely to feel out of control than multiparous women (Green and Baston, 2003).

#### **4.1 Limitations**

Mediation models imply a sequence of time (Maxwell and Cole, 2007); however this was a cross-sectional study and therefore may not reflect true causal mediation pathways. We imposed theoretical models, which were found to be a good fit to the data. However, it could be possible that disrupted SCC is a precursor to a traumatic birth, rather than the consequence of birth trauma.

Duration from birth was not evaluated, thus we could not control for this. We cannot be certain that recall from participants two months postpartum did not differ from those six months postpartum. However, memory surrounding childbirth is unaffected during the first year postpartum (Waldenström, 2003).

#### **4.2 Implications**

PLEs in new mothers are more common than previously known. There are probably multiple pathways, that predict the occurrence of PLEs with a variety of underlying mechanisms. The model

suggests that PTSS predict the occurrence of PLEs, whilst SCC is critical for women who have appraised birth as traumatic and have difficulties adjusting to motherhood. Indeed, the process of incorporating intra and inter-personal changes into the self-concept postpartum may be much more important than previously thought. Therefore, SCC should, perhaps, be considered by health-care professionals. The same is true for trauma appraisals and maternal-adjustment. This is especially pertinent given that these factors are less overt than other difficulties (e.g. PTSS), and many distressed women may be overlooked.

#### **4.3 Future research**

Longitudinal studies, beginning antenatally, may be most helpful in understanding the development of PLEs in new mothers, especially given that Mannion and Slade (2014) found the occurrence of PLEs were higher during pregnancy than postpartum. Given that relationships were found between psycho-social factors and PLEs (Table 3.), and obstetric trauma has been found to increase the risk of postnatal psychosis (Nager et al., 2008), further investigation of these areas would be pertinent. Additionally, longitudinal studies investigating the transition from PLEs to psychosis are indicated.

#### **4.4 Conclusion**


PLEs in first time mothers are not uncommon and can be associated with distress. PTSS directly predicted the occurrence of PLEs. However, other pathways exist and there maybe women who are less likely to present with clinical symptoms as their difficulties surround trauma appraisals, adjustment to motherhood or disruptions in sense of self-concept. SCC seems to be an important factor previously unrecognised. It is imperative for health-care professionals to recognise the development of PLEs and understand these experiences. Public health interventions that reduce the stigma of PLEs, whilst also improving knowledge and understanding may be important.

## References

- American Psychiatric Association, 2000. Diagnostic and statistical manual of mental disorders DSM IV-TR. American Psychiatric Publishing, Washington.
- American Psychiatric Association, 2013. Diagnostic and statistical manual of mental disorders DSM 5. American Psychiatric Publishing, Washington.
- Alsawy, S., Wood, L., Taylor, P., Morrison, A., 2015. Psychotic experiences and PTSD: exploring associations in a population survey. *Psychological Medicine* 45(13), 2849-2859.
- Ayers, S., Bond, R., Bertullies, S., Wijma, K., 2016. The aetiology of post-traumatic stress following childbirth: a meta-analysis and theoretical framework. *Psychological medicine* 46(06), 1121-1134.
- Barratt, S., 2012. Transient psychotic symptoms in new mothers: an unrecognized and potentially frequent problem. Dissertation, University of Liverpool.
- Bentall, R.P., de Sousa, P., Varese, F., Wickham, S., Sitko, K., Haarmans, M., Read, J., 2014. From adversity to psychosis: pathways and mechanisms from specific adversities to specific symptoms. *Social Psychiatry and Psychiatric Epidemiology* 49(7), 1011-1022.
- Bollen, K.A., Stine, R.A., 1992. Bootstrapping goodness-of-fit measures in structural equation models. *Sociological Methods & Research* 21(2), 205-229.
- Buysse, D.J., Reynolds, C.F., Monk, T.H., Berman, S.R., Kupfer, D.J., 1989. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Research* 28(2), 193-213.
- Campbell, J.D., Trapnell, P.D., Heine, S.J., Katz, I.M., Lavallee, L.F., Lehman, D.R., 1996. Self-concept clarity: Measurement, personality correlates, and cultural boundaries. *Journal of Personality and Social Psychology* 70(1), 141.
- Cicero, D.C., Becker, T.M., Martin, E.A., Docherty, A.R., Kerns, J.G., 2013. The role of aberrant salience and self-concept clarity in psychotic-like experiences. *Personality Disorders: Theory, Research, and Treatment* 4(1), 33.
- Cicero, D.C., Martin, E.A., Becker, T.M., Kerns, J.G., 2015. Decreased self-concept clarity in people with schizophrenia. *The Journal of Nervous and Mental Disease*. 204(2),142-147.
- Cox, J.L., Holden, J.M., Sagovsky, R., 1987. Detection of postnatal depression. Development of the 10-item Edinburgh Postnatal Depression Scale. *The British Journal of Psychiatry* 150(6), 782-786.
- Creamer, M., Bell, R., Failla, S., 2003. Psychometric properties of the impact of event scale - revised. *Behaviour Research and Therapy* 41:1489-1496
- Czarnocka, J., Slade, P., 2000. Prevalence and predictors of post-traumatic stress symptoms following childbirth. *British Journal of Clinical Psychology* 39(1), 35-51.
- Edworthy, Z., Chasey, R., Williams, H., 2008. The role of schema and appraisals in the development of post-traumatic stress symptoms following birth. *Journal of Reproductive and Infant Psychology* 26(2), 123-138.
- Ehlers, A., Clark, D.M., 2000. A cognitive model of posttraumatic stress disorder. *Behaviour Research and Therapy* 38(4), 319-345.
- Evans, G.J., Reid, G., Preston, P., Palmier-Claus, J., Sellwood, W., 2015. Trauma and psychosis: The mediating role of self-concept clarity and dissociation. *Psychiatry Research* 228(3), 626-632.
- Green, J.M., Baston, H.A., 2003. Feeling in control during labor: concepts, correlates, and consequences. *Birth* 30(4), 235-247.
- Hamner, M.B., Frueh, B.C., Ulmer, H.G., Arana, G.W., 1999. Psychotic features and illness severity in combat veterans with chronic posttraumatic stress disorder. *Biological Psychiatry* 45(7), 846-852.
- Hayes, A.F., Scharkow, M., 2013. The relative trustworthiness of inferential tests of the indirect effect in statistical mediation analysis does method really matter? *Psychological Science*, 0956797613480187.
- Johns, L.C., Van Os, J., 2001. The continuity of psychotic experiences in the general population. *Clinical Psychology Review* 21(8), 1125-1141.
- Launay, G., Slade, P., 1981. The measurement of hallucinatory predisposition in male and female prisoners. *Personality and Individual Differences* 2(3), 221-234.
- MacKinnon, D., 2008. Introduction to statistical mediation analysis. Routledge.


- Mallinckrodt, B., Abraham, W.T., Wei, M., Russell, D.W., 2006. Advances in testing the statistical significance of mediation effects. *Journal of Counseling Psychology* 53(3), 372.
- Mannion, A., Slade, P., 2014. Psychotic-like experiences in pregnant and postpartum women without a history of psychosis. *Schizophrenia Research* 160(1), 118-123.
- Markus, H., Wurf, E., 1987. The dynamic self-concept: A social psychological perspective. *Annual Review of Psychology* 38(1), 299-337.
- Matthey, S., 2011. Assessing the experience of motherhood: the Being a Mother Scale (BaM-13). *Journal of Affective Disorders* 128(1), 142-152.
- Maxwell, S.E., Cole, D.A., 2007. Bias in cross-sectional analyses of longitudinal mediation. *Psychological Methods* 12(1), 23.
- Morrison, A.P., Frame, L., Larkin, W., 2003. Relationships between trauma and psychosis: A review and integration. *British Journal of Clinical Psychology* 42(4), 331-353.
- Nager, A., Sundquist, K., Ramírez-León, V. and Johansson, L.M., 2008. Obstetric complications and postpartum psychosis: a follow-up study of 1.1 million first-time mothers between 1975 and 2003 in Sweden. *Acta Psychiatrica Scandinavica*, 117(1), 12-19.
- National Perinatal Epidemiology Unit, 2014. Safely delivered: A national survey of women's experience of maternity care. National Perinatal Epidemiology Unit, Oxford
- O'Donovan, A., Alcorn, K.L., Patrick, J.C., Creed, D.K., Dawe, S., Devilly, G.J., 2014. Predicting posttraumatic stress disorder after childbirth. *Midwifery* 30(8), 935-941.
- Peters, E., Joseph, S., Day, S., Garety, P., 2004. Measuring delusional ideation: the 21-item Peters et al. Delusions Inventory (PDI). *Schizophrenia Bulletin* 30(4), 1005.
- Peters, E., Ward, T., Jackson, M., Morgan, C., Charalambides, M., McGuire, P., Woodruff, P., Jacobsen, P., Chadwick, P., Garety, P.A., 2016. Clinical, socio-demographic and psychological characteristics in individuals with persistent psychotic experiences with and without a "need for care". *World Psychiatry* 15(1), 41-52.
- Poulton, R., Caspi, A., Moffitt, T.E., Cannon, M., Murray, R., Harrington, H., 2000. Children's self-reported psychotic symptoms and adult schizophreniform disorder: a 15-year longitudinal study. *Archives of General Psychiatry* 57(11), 1053-1058.
- Read, J., Fink, P., Rudegeair, T., Felitti, V., Whitfield, C., 2008. Child maltreatment and psychosis: a return to a genuinely integrated bio-psycho-social model. *Clinical Schizophrenia & Related Psychoses* 2(3), 235-254.
- Sit, D., Rothschild, A.J., Wisner, K.L., 2006. A review of postpartum psychosis. *Journal of Women's Health* 15(4), 352-368.
- Sitko, K., Bentall, R.P., Shevlin, M., Sellwood, W., 2014. Associations between specific psychotic symptoms and specific childhood adversities are mediated by attachment styles: an analysis of the National Comorbidity Survey. *Psychiatry Research* 217(3), 202-209.
- Slade, P., Morrell, C.J., Rigby, A., Ricci, K., Spittlehouse, J., Brugha, T.S., 2010. Postnatal women's experiences of management of depressive symptoms: a qualitative study. *Br J Gen Pract* 60(580), e440-e448.
- Smith, J., 1999. Identity development during the transition to motherhood: An interpretative phenomenological analysis. *Journal of Reproductive and Infant Psychology* 17(3), 281-299.
- Soet, J.E., Brack, G.A., DiIorio, C., 2003. Prevalence and predictors of women's experience of psychological trauma during childbirth. *Birth* 30(1), 36-46.
- Valdimarsdóttir, U., Hultman, C.M., Harlow, B., Cnattingius, S., Sparén, P., 2009. Psychotic illness in first-time mothers with no previous psychiatric hospitalizations: a population-based study. *PLoS Med* 6(2), e1000013.
- VanderKruik, R., Barreix, M., Chou, D., Allen, T., Say, L. and Cohen, L.S., 2017. The global prevalence of postpartum psychosis: a systematic review. *BMC psychiatry*, 17(1), p.272.
- Van Os, J., Linscott, R.J., Myin-Germeys, I., Delespaul, P., Krabbendam, L., 2009. A systematic review and meta-analysis of the psychosis continuum: evidence for a psychosis proneness–persistence–impairment model of psychotic disorder. *Psychological Medicine* 39(02), 179-195.
- Varese, F., Barkus, E., Bentall, R., 2012a. Dissociation mediates the relationship between childhood trauma and hallucination-proneness. *Psychological Medicine* 42(05), 1025-1036.

- Varese, F., Smeets, F., Drukker, M., Lieveise, R., Lataster, T., Viechtbauer, W., Read, J., van Os, J., Bentall, R.P., 2012b. Childhood adversities increase the risk of psychosis: a meta-analysis of patient-control, prospective-and cross-sectional cohort studies. *Schizophrenia Bulletin* 38(4), 661-671.
- Waldenström, U., 2003. Women's memory of childbirth at two months and one year after the birth. *Birth* 30(4), 248-254.
- Weiss, D.S., 2007. The impact of event scale: revised, Cross-cultural assessment of psychological trauma and PTSD. Springer 219-238.
- Wijma, K., Wijma, B., Zar, M., 1998. Psychometric aspects of the W-DEQ; a new questionnaire for the measurement of fear of childbirth. *Journal of Psychosomatic Obstetrics & Gynecology* 19(2), 84-97.



**Figure 1** Three nested competing path models exploring the relationship between birth experience and psychotic like experiences (hallucination and delusional experiences) through two stages of mediating variables (i) trauma appraisals, post-traumatic stress symptoms and adjustment to motherhood (ii) self-concept clarity

- - - - Pathways included in Model 1 only
- ..... Pathway included in Model 2 only
- Pathways included in Model 1, 2 and 3



**Figure 2** Participant recruitment flowchart

**Table 1** Participant characteristics

	Range	Mean ( <i>SD</i> )
Age (years)	16-48	28.9 (5.51)
		Total n (%)
Marital status		
Married		732 (52.5)
Not married		661 (47.5)
Missing data		7 (0.50)
Educational attainment		
Low education (no qualifications/GCSE's)		184 (13.2)
Mid education (A Levels/vocational qualifications)		500 (35.9)
High education (graduate/post graduate)		709 (50.9)
Pre-pregnancy employment		
Employed (full time/part time/self employed)		1263 (90.6)
Unemployed (out of work/voluntary work/student)		130 (9.4)
Ethnic origin		
White		1254 (90)
Other		139 (10)
Number of babies delivered		
Singleton		1357 (98.7)
Multiple		18 (1.3)
Mode of birth		
Normal vaginal delivery		707 (50.8)
Assisted delivery (e.g. Kiwi, Forceps or Ventouse)		330 (23.7)
Caesarean section		356 (25.6)

*N* = 1393.

**Table 2** Descriptive statistics

Variable	Mean ( <i>SD</i> )	Possible ranges of scores
Birth experience	54.25 (28.88)	0-149
Post-traumatic stress symptoms		
Arousal	0.65 (0.72)	0-4
Intrusion	0.62 (0.67)	0-4
Avoidance	0.77 (0.81)	0-4
Total score	14.98 (15.1)	0-88
Trauma appraisal	0.77 (0.85)	0-2
Adjustment to motherhood		
Child	6.03 (3.88)	0-18
Adult	4.60 (3.36)	0-15
Emotional closeness	0.57 (0.95)	0-6
Total score	11.2 (7.02)	0-39
Self-concept clarity	42.36 (9.52)	12-60
Hallucination experiences		
Auditory hallucinations	0.69 (0.84)	0-4
Visual hallucinations	0.04 (0.20)	0-1
Vivid thoughts	0.58 (0.78)	0-2
Vivid day dreams	0.46 (0.81)	0-3
Total score	2.20 (2.03)	0-12
Delusion experiences		
Distress <sup>a</sup>	9.44 (8.86)	0-105
Preoccupation <sup>b</sup>	9.35 (8.61)	0-105
Conviction <sup>c</sup>	12.08 (9.92)	0-105
Total score	3.49 (2.84)	0-21
Composite score	30.46 (29.46)	0-336
Depressive symptoms	7.06 (4.72)	0-30
Average hours of sleep	5.83 (1.61)	1-16
	Frequency (%)	
Quality of sleep		
Good	335 (24.00)	-
Poor	835 (60.00)	-
Unknown	223 (16.00)	-
Trauma appraisals		
Experienced danger	524 (37.60)	-
Danger for self	27 (5.20)	-
Danger for baby	269 (51.30)	-
Danger for self and baby	228 (16.40)	-
Experienced intense fear	557 (40.00)	-
Both criteria met	384 (27.60)	-
Above clinical cut off for PTSD <sup>d</sup>	167 (12.00)	-
Above clinical cut off for depression <sup>e</sup>	172 (12.30)	-

*N* = 1,393.

<sup>a, b, c</sup> *n* = 1217, corresponding to number of participants reporting at least one delusion. <sup>d</sup> Score  $\geq$  33, recommended by Creamer et al. (2003). <sup>e</sup> Score  $\geq$  13, recommended by Matthey et al. (2006)

**Table 3** Inter-correlations of study variables

	1	2	3	4	5	6	7	8	9	10	11	12	13	14
1. Birth experience	-													
2. Trauma appraisals	.64**	-												
3. Post-traumatic stress symptoms	.56**	.47**	-											
4. Adjustment to motherhood	.44**	.28**	.47**	-										
5. Self-concept clarity	-.37**	-.27**	-.41**	-.59**	-									
6. Hallucination experiences	.22**	.21**	.39**	.30**	-.40**	-								
7. Delusional experiences	.24**	.22**	.42**	.36**	-.46**	.52**	-							
8. Depression	.36**	.25**	.44**	.61**	-.63**	.38**	.46**	-						
9. Good sleep	-.22**	-.17**	-.26**	-.30**	.20**	-.13**	-.15**	-.29**	-					
10. Age	.02	-.04	-.06*	-.01	.19**	-.21**	-.20**	-.13**	-.11**	-				
11. Ethnicity	.01	.00	.01	.06*	-.13**	-.00	-.16**	-.03	.00	-.05	-			
12. Employment status	-.02	.00	-.07*	.01	.05*	-.12**	-.15**	-.08**	.00	.23**	.16**	-		
13. Mode of Birth	.33**	.28**	.13**	.03	-.01	-.02	-.03	.03	-.07*	.22**	.00	.06*	-	
14. Marital Status	-.03	-.04	-.06*	.01	.13**	-.13**	-.09**	-.08**	-.03	.33**	-.16**	.11**	.04	-
15. Education	.02	.00	.02	.08**	.12**	-.12**	-.09**	-.06*	-.07*	.37**	-.15**	.11**	.03	.31**

$N = 1393$ . Correlations represent Spearman's  $r$  or phi ( $r\phi$ ) coefficients.

\*  $p < .05$ ; \*\*  $p < .01$

Note: In order to undertake correlational analyses demographic data were simplified and recoded as appropriate: Ethnicity (0 = *not White British*, 1 = *White British*), marital status (0 = *not married*, 1 = *married*), employment status (0 = *not employed*, 1 = *employed*), mode of birth (coded in terms of increasing intervention; 1 = *normal delivery*, 2 = *assisted delivery*, 3 = *caesarean section*) and education level (1 = *low education*, 2 = *mid education*, 3 = *high education*).

**Table 4** Endorsement of items on the Launay Slade Hallucination Scale and the 21-item Peters Delusion Inventory

Scales, subscales and items	Frequency (%)
<b>Launay Slade Hallucination Scale</b>	
No matter how much I try to concentrate on my work unrelated thoughts always creep into my mind	594 (42.6)
Sometimes a passing thought will seem so real that it frightens me	439 (31.5)
I have never been troubled by hearing voices in my head <sub>r</sub>	379 (27.2)
Sometimes my thoughts seem as real as actual events in my life	370 (26.6)
The sounds I hear in my daydreams are generally clear and distinct	272 (19.5)
In my daydreams I can hear the sound of a tune almost as clearly as if I were actually listening to it	249 (17.9)
I have never heard the voice of the Devil <sub>r</sub>	215 (15.4)
In the past I have had the experience of hearing a person's voice and then found that there was no-one there	159 (11.4)
I often hear a voice speaking my thoughts aloud	141 (10.1)
The people in my daydreams seem so true to life that I sometimes think they are	125 (9)
In the past I have heard the voice of God speaking to me	63 (4.5)
On occasions I have seen a person's face in front of me when no-one was in fact there	58 (4.2)
<b>21-item Peters Delusion Inventory</b>	
Do you ever feel as if some people are not what they seem to be	941 (67.6)
Do you ever feel as if people seem to drop hints about you or say things with a double meaning?	553 (39.7)
Are you often worried that your partner may be unfaithful?	413 (29.6)
Do you ever feel that people look at you oddly because of your appearance?	317 (22.8)
Do you ever feel as if electrical devices such as computers can influence the way you think?	297 (21.3)
Do you ever feel that you are a very special or unusual person?	276 (19.8)
Do you ever feel as if things in magazines or on TV were written for you?	213 (15.3)
Do you ever think people can communicate telepathically?	205 (14.7)
Do you ever feel as if you are, or destined to be someone very important?	186 (13.4)
Do you ever feel that you are especially close to God?	179 (12.8)
Do you ever feel that you have sinned more than the average person?	167 (12.0)
Do you ever feel as if you are being persecuted in some way?	163 (11.7)
Do you believe in the power of witchcraft, voodoo or the occult?	163 (11.7)
Do you ever feel as if you had no thoughts in your head at all?	160 (11.5)
Do you ever feel as if you have been chosen by God in some way?	123 (8.8)
Do your thoughts ever feel alien to you in some way?	113 (8.1)
Do you ever feel as if your own thoughts were being echoed back to you?	83 (6.0)
Do you ever feel as if the world is about to end?	82 (5.9)
Have your thoughts ever been so vivid that you were worried other people would hear them?	76 (5.5)
Do you ever feel as if there is a conspiracy against you?	76 (5.5)
Do you ever feel as if you are a robot or zombie without a will of your own?	71 (5.1)


*N* = 1,393

<sub>r</sub> Reverse scored item

**Table 5** Goodness-of-fit statistics for hypothesised models

Model	Comparative model	$\chi^2$	<i>df</i>	<i>p</i> value	$\chi^2$	CFI	RMSEA	$\Delta\chi^2$	$\Delta df$	<i>p</i> value $\Delta\chi^2$
Model 1	-	4.13	1	.042	.999	.047	-	-	-	-
Model 2	Model 1	4.49	4	.344	1.000	.009	0.36	3		.948
Model 3	Model 2	8.71	5	.121	.999	.023	4.22	1		.040

$\Delta\chi^2$  = difference in  $\chi^2$  values between models;  $\Delta df$  = difference in number of degrees of freedom between models.



**Figure 3 Model 2 exploring the relationship between birth experience and psychotic like experiences** (hallucination and delusional experiences) through two stages of mediating variables (i) trauma appraisals, post-traumatic stress symptoms and adjustment to motherhood (ii) self-concept clarity. The regression coefficients for paths trauma appraisals\*hallucinations, trauma appraisals\*delusions, adjustment to motherhood\*hallucinations and adjustment to motherhood\*delusions were constrained to 0. For simplicity error terms and control variables have been omitted. Estimates on the endogenous variables were controlled for by age, birth method, education level, employment status, ethnicity, marital status, depression and sleep quality. Error terms were correlated as appropriate, with all stage one mediator error terms (trauma appraisals, post-traumatic stress symptoms and adjustment to motherhood) correlated with each other and outcome variable error terms (hallucination and delusional experiences) correlated with each other. Estimates shown are standardised betas ( $\beta$ ). Significance levels were established via bootstrapped CIs (10,000 resamples). A bootstrapped Chi square test indicated a good model fit,  $\chi^2(4) = 4.49, p = .34$ ; CFI = 1.00, RMSEA = 0.01, 90% CI [0.00, 0.03]  $p = .98$ .

———— Significant path                      ———— Non-significant path

**Table 6** Path estimates for model 2 (Figure 3)

Path	Estimate		SE of $b$		Bootstrap 95% CIs $b$		Bootstrap 95% CIs $\beta$	
	$b$	$\beta$	ML	Bootstrap	Lower	Upper	Lower	Upper
Direct path estimates								
$a_1$	0.27***	.52	.01	.02	.24	.30	.47	.57
$a_2$	0.02***	.60	.00	.00	.02	.02	.56	.64
$a_3$	0.07***	.28	.01	.01	.06	.08	.23	.32
$b_1$	- 0.02	-.03	.02	.02	-.05	.01	-.08	.02
$b_2$	- 0.56*	-.05	.28	.28	- 1.11	.00	-.10	.00
$b_3$	- 0.42***	-.31	.04	.04	-.49	-.35	-.36	-.28
$c_1$	- 0.04***	-.17	.01	.01	-.05	-.02	-.23	-.11
$c_2$	- 0.07***	-.24	.01	.01	-.09	-.05	-.30	-.18
$d_1$	0.00	-.05	.00	.00	-.01	.00	-.11	.13
$d_2$	- 0.01*	-.06	.00	.00	-.01	.00	-.12	.00
$e_1$	0.04***	.28	.00	.00	.03	.05	.21	.34
$e_2$	0.05***	.24	.01	.01	.03	.06	.18	.30
$f$	- 0.02*	-.06	.01	.01	-.04	.00	-.12	.00
Indirect path estimates								
$a_1b_1$	.00	.00	-	.00	-.01	.00	-.04	.01
$a_1e_1$	.01***	.16	-	.00	.01	.01	.12	.19
$a_1e_2$	.01***	.14	-	.00	.01	.02	.11	.18
$a_2b_2$	-.01*	-.03	-	.01	-.02	.00	-.06	.00
$a_3b_3$	-.03***	-.09	-	.00	-.04	-.02	-.11	-.07
$b_1c_1$	.00	.01	-	.00	.00	.00	.00	.01
$b_1c_2$	.00	.01	-	.00	.00	.00	-.01	.02
$b_2c_1$	.02*	.01	-	.01	.00	.05	.00	.02
$b_2c_2$	.03*	.01	-	.02	.00	.08	.00	.02
$b_3c_1$	.02***	.05	-	.00	.01	.02	.03	.07
$b_3c_2$	.03***	.07	-	.01	.02	.04	.05	.10
$a_1b_1c_1$	.00*	.01	-	.00	.00	.00	.00	.01
$a_1b_1c_2$	.00*	.02	-	.00	.00	.00	.01	.02
$a_2b_2c_1$	.00**	.01	-	.00	.00	.00	.01	.01
$a_2b_2c_2$	.00**	.01	-	.00	.00	.00	.01	.02
$a_3b_3c_1$	.00***	.02	-	.00	.00	.00	.01	.02
$a_3b_3c_2$	.00***	.02	-	.00	.00	.01	.02	.03
$fc_1$	.01*	.01	-	.00	.01	.02	.14	.02
$fc_2$	.02*	.01	-	.00	.01	.02	.14	.03

$N = 1393$ . ML = maximum likelihood estimation. Probability values determined on bootstrapped CIs (10,000 resamples)

\*  $p < .05$ ; \*\*  $p < .01$ ; \*\*\*  $p < .001$

**Responses to reviewer's comments: re paper entitled 'Birth experiences, trauma responses and self-concept in postpartum psychotic-like experiences'**

**Comment 1:** *"In the introduction the authors define PLEs as "...non-clinical experiences similar to psychosis but in a diminished form" but this requires elaboration. The authors should clarify to readers from psychiatry how PLEs postpartum differ from DSM-V's characterisation of postpartum psychosis. The authors should clarify how they bridge their work with DSM-V's conceptualisation of postpartum psychosis (which only requires 1 symptom to be present and could actually meet the authors' definition of PLEs if the disturbance is for just 1 day). Why/when do PLEs not meet the criteria in terms of their duration, frequency and severity?"*

**Response:** Van Os et al. (2009) highlight that experiencing hallucinations/delusions is not always associated with the presence of a disorder. In their review, they found there to be a distinction between psychotic experiences without distress or help seeking behaviour (prevalence approximately 8%), subclinical psychotic symptoms which have a degree of distress and some help seeking behaviour (prevalence approximately 4%), and psychotic disorder associated with significant distress and help seeking behaviour (prevalence approximately 3%). In our study, we aimed to investigate both the psychotic experiences/subclinical psychotic symptoms.

Therefore, in accordance with DSM-V's criteria, postpartum psychosis would be assumed to be associated with distress, illness related behaviours, and impairment of some kind. However, we would assume that some PLEs would not necessarily be associated with distress or illness related behaviour. Additionally, we would assume PLEs to be more prevalent in the population than postpartum psychosis.

Finally, we would argue that PLEs could be fleeting in duration, lasting less than 1 day.

**Amendment made:** "Postpartum psychosis is associated with distress, illness related behaviours, impairment of functioning and help-seeking behaviour (Van Os et al., 2009; *The Diagnostic and Statistical Manual of Mental Disorders*, DSM-5, American Psychiatric Association, 2013). However, PLEs are usually not associated with distress or illness-related behaviour, and maybe fleeting in duration." (Page 3, lines 9-13)

**Comment 2:** *"In the introduction the authors should clarify whether they define PLE's as strictly one (or more) of the 4 symptoms of brief psychotic disorder (delusions, hallucinations, disorganised speech and catatonic behaviour). If not, the authors should explain what additional sorts of symptoms PLE's can include with some examples."*

**Response:** In this paper PLEs were defined as either delusions or hallucination experiences.

**Amendment made:** "of hallucinations or delusions" (Page 3, line 4)

**Comment 3:** *"In the introduction the authors should clarify how they define PLE's in duration in order to support their definition of PLEs as non-clinical experiences. In brief psychotic disorder the disturbances can occur for as little as 1 day. Do the authors argue that PLEs occur for less than the 1 day, and/or do they argue for a shorter or longer maximum than the 1 month in brief psychotic disorder? The authors need to address potential criticism that PLEs are actually clinically significant"*

*and actually meet the criteria for brief psychotic disorder.”*

**Response:** The reviewer has made a really helpful point here. As highlighted in point 1, we argue that PLEs may occur over short, possibly fleeting, lengths of time, and *may* be present for less than one day. Some participants may well have met criteria for brief psychotic disorder in terms of DSM. Indeed, due to the definition of PLEs (as highlighted in point 1), the criteria are associated with lack of help seeking behaviour/distress rather than absolute duration

**Amendment made:** “Postpartum psychosis is associated with distress, illness related behaviours, impairment of functioning and help-seeking behaviour (Van Os et al., 2009; *The Diagnostic and Statistical Manual of Mental Disorders*, DSM–5, American Psychiatric Association, 2013). However, PLEs are not necessarily associated with distress or illness-related behaviour, and maybe fleeting in duration.” (Page 3, lines 9-13).

**Comment 4:** *“In the introduction, with regards to defining PLEs postpartum, what is the authors' cut-off point (in weeks or months) after childbirth? In the method they later present a 2 month cut-off point but it is not clear whether there is support for this in previous literature.”*

**Response:** As only two studies to date have investigated PLE’s postpartum there are limited data surrounding the risk period. Mannion and Slade (2014), investigated PLEs 7 days postpartum, whilst Barratt (2012) also examined the first two months post-partum.

By definition the onset of post-partum psychosis occurs up to a month, but often within days, of giving birth. Therefore, as we are looking at a continuum model, we hypothesised that PLEs would occur within the same time frame. However, in order to encapsulate the entire theorised period of risk we extended this time frame.

**Amendment made:** “In accordance with the defined onset of post-partum psychosis in the DSM-5, it is hypothesised that PLEs occur in the first month postpartum. However, in order to encapsulate the entire theorised period of risk we extended this time frame to two months.” (Page 4, lines 23-25)

**Comment 5:** *“In the introduction the authors should elaborate on sources of trauma during childbirth that increase the risk of postpartum psychotic symptoms. The authors should give examples (with references) of (a) obstetric sources of trauma such as pre-eclampsia; (b) non-obstetric sources of trauma such as emergency transfers from birth centres to hospital. The authors should cite systematic reviews (where available) showing the level of risk associated with different sorts of obstetric and non-obstetric factors. The authors' method focuses on e.g. fear but they need to justify why - given previous literature about the huge role of obstetric trauma - they elected to focus on fearfulness etc.”*

**Response:** It has been established that appraisal predicts whether birth is experienced as traumatic, and indeed PTSD symptoms, rather than the experience, type of physical trauma or physical intervention.

For example, Devilly et al. (2014), in their longitudinal prospective study, found that subjective reaction to childbirth explained variance in distress over and above just experiencing the event. Additionally, experiencing fear, helplessness, terror, or horror during childbirth significantly predicted progression distress related to PTSD symptoms over time. Furthermore, subjective appraisal of threat during or immediately after childbirth was a better predictor of longer term distress related to PTSD

symptoms, than an event specific measure of childbirth trauma. O'Donovan et al. (2014) found that psychological factors such as perceived lack of control and discrepancy in expectations surrounding childbirth, rather than medical procedures, were predictive of a traumatic birth experience. Indeed, PTSD symptoms were predicted predominantly by past trauma, rather than factors during birth such as pain.

Qualitative studies have found the focus to also be on appraisal of the experience, rather than medical interventions. Reed et al. (2017), in their mixed-methods study exploring women's descriptions of trauma, found that the majority (66.7%) of participants described care provider actions as the traumatic component of their experience, as apposed to physical interventions/trauma events per-se (e.g. haemorrhage). Also, one of the studies major themes was entitled 'violation', highlighting the appraisal of the birthing experience to be particularly important.

In a study related specifically to PLE's postpartum, Mannion and Slade (2014) found that 'fear' during childbirth (specifically measured by one question: during labour, were you fearful for your life or your baby's life, or fearful of serious injury or permanent damage?) was significantly associated with PLEs postnatally.

Therefore, given the evidence to indicate that 'fear' during labour is an important factor in the predictor of PLE's postpartum, and the evidence demonstrating that appraisal is a more important factor in the predictor of postpartum distress related to PTSD symptoms, we felt it imperative that we focus on this areas as opposed to sources of obstetric trauma.

**Amendment made:** None.

**Comment 6:** *"In the method the authors should justify their selection criteria - why did they exclude mothers who had given birth 0 to 8 weeks before the study? The authors mention that 2 months is their theorised phase of risk but this is problematic unless the authors present strong evidence for this cut-off point in the introduction. In DSM-V the cut-off point is 4 weeks postpartum, which is arguably too short. Eight weeks is, also, arguably questionable - there is no clear justification for this from the authors"*

**Response:** This was chosen for consistency. If we had included women within our time frame of reference, i.e. 0-8 weeks, they would have had variable exposure to the time frame. Therefore, it would have been difficult to comment on prevalence, as some women with negative responses may have had subsequent experience of PLEs.

**Amendment made:** "This time frame was chosen for consistency, and to ensure all participants had passed the hypothesised risk period for PLEs." (Page 5, lines 3-4)

**Comment 7:** *"Why did the authors exclude "participants disclosing a history of psychosis or having had input from perinatal mental health teams"?"*

**Response:** The aim of the study was to investigate experiences among women without a significant history of psychosis. The exclusion criteria were therefore chosen to ensure our estimates were conservative and not inflated by reports from women with known serious and enduring mental health problems. In addition, frequencies of difficulties and indeed issues of coherence of self-concept might be confounded by ongoing medication or other intervention. The choice was made therefore to attempt to investigate these aspects in a sample where such confounding factors had been excluded.

**Amendment made:** "To ensure our estimates were conservative, and not confounded by participants

with known mental health problems receiving treatment” (Page 5, lines 4-5)

**Comment 8:** “How did the authors ascertain the women's psychiatric history? Presumably this was self-reported but how did they word the questions about psychotic history etc. - did they ask the women if they had a history of a specific disorder?”

**Response:** Participants were asked to respond, via self-report, to two specific questions to ascertain participant's psychiatric history:

- a) Have you ever been treated for psychosis prior, during or after your pregnancy?
- b) Are you under the care of the perinatal mental health team?

This approach was thought to be the best method to capture those who would meet point 7.

**Amendment made:** “participants with a history of psychosis or having had input from perinatal mental health teams (from self-report prior to questionnaire completion) were excluded.” (Page 5, lines 7-8)

**Comment 9:** *“The authors imply that their measurement of hallucinations captured just proneness - rather than actual experiences. In the context of postpartum psychosis (or postpartum PLEs) the authors cannot assume that proneness is the same as actually having had hallucinations. Later they present data about the proportion of respondents who had at least 1 hallucination therefore the method should clarify whether this is what was analysed in their results (not proneness) - as they clarify with delusions.”*

**Response:** The reviewer is correct in highlighting that the way the data are presented could be confusing to the reader. The word ‘proneness’ was used to highlight that we thought these experiences were PLEs, rather than being necessarily clinical forms of hallucinations/delusions that could meet diagnostic criteria. However, by using the LSHS and PDI-21 we did measure self-reported experience of hallucinations and delusions. Therefore the word ‘proneness’ has been removed entirely from the paper, in order to clarify to the reader that we did measure self-reported occurrence of hallucinations/delusions. However, these were perceived to be representative of PLEs, as the participants in the study were presumed to have not engaged in any help seeking behaviour/had not been diagnosed with psychosis (in line with the exclusion criteria).

**Amendment made:** Removal of the word proneness, and experience where appropriate, throughout the paper. Grammar has also been corrected accordingly.

**Comment 10:** *“In the discussion the authors ponder on the reasons why their measure of childbirth trauma (fear) did not predict hallucinations; the authors should include a comment on whether the results would have been different had they measured obstetric trauma in the way that the literature about risk levels does - see odds ratios associated with different types of obstetric complications, and also transfers. I appreciate that the previous data mainly concern postpartum depression - but there is some previous literature on odds ratios and postpartum psychosis. The authors can comment on the obstetric factors that should be measured in future work. Likewise, the authors should comment on the role of additional factors (e.g. a history of mental illness, spousal support and socioeconomic status) in increasing the risk of postpartum psychosis - are these risk factors likely to be relevant to PLEs?”*

**Response:** Despite the evidence cited in point 5 we acknowledge that our focus on fear and appraisal may have limited the findings. The reviewer has helpfully noted that other factors, such as obstetric trauma, may be relevant and should be investigated in future studies. Indeed, Nager et al. (2008) found the risk of postnatal psychosis in first time mothers increased in relation to respiratory disorder in the neonate, severe birth asphyxia, preterm birth, caesarean section, perinatal death and giving birth to a SGA infant. However, only pre-term birth and acute caesarean section were significant risk factors for psychosis, over and above previous psychiatric disorder with hazard ratios of 1.2 and 1.31 respectively. We explored the impact of mode of birth, and found this was not associated with PLEs, however we did not distinguish between elective and acute caesarean section. Therefore if we had made this distinction, and included data surrounding pre-term birth, it is possible that we would have found there to be some variance in the model accounted for by these factors.

The reviewer has helpfully highlighted that other factors, such as history of low-socio economic status and spousal support, may also be relevant to both postnatal psychosis and PLEs. The literature in this area is currently sparse; however Nager et al. (2005) found that both older age and being a single mother increased the risk of being hospitalised due to postnatal psychosis. In our study (see Table 3.) we found that PLEs (in terms of both hallucination and delusions) increased amongst participants who were older, not married, not employed or with lower educational attainment. PLEs were also significantly correlated with depressive symptoms, and negatively correlated with good sleep. Delusions were also significantly correlated with ethnicity, with higher delusional experiences being associated with participants who were not White.

It would be recommended that future studies investigate the role of psycho-social factors in relation to PLEs in new mothers.

**Amendment made:** “Given that relationships were found between psycho-social factors and PLEs (Table 3.), and obstetric trauma has been found to increase the risk of postnatal psychosis (Nager et al., 2008), further investigation of these areas would be pertinent.” (Page 11, lines 21-24).

**Comment 11:** *“The authors should make more precise recommendations about the practical implications of their research. First, the authors should address the question of whether or not PLEs are ever clinically significant. For example, is there a risk that - though isolated - PLEs are early warning signs of preponderance to a psychotic disorder? The authors should inform readers whether there is evidence showing that e.g. a history of PLEs in new mothers is associated with a higher incidence of schizophrenia. That way, clinician readers will be aware of whether new mothers presenting with PLEs are at greater risk of a psychotic disorder in the long-term and therefore whether PLEs ever require any early intervention.”*

**Response:** Unfortunately, no papers have yet investigated the link between PLEs in new mothers and post-partum psychosis. Additionally, longitudinal studies investigating the transition from PLEs to psychosis are yet to be undertaken. However, there is evidence that PLEs in the general population and clinical psychosis have similar co-morbidities, including depressive symptoms and blunting of affect (Van Os et al, 2000). Additionally, as stated in the introduction, both PLEs and clinical psychosis have been found to be associated with childhood traumas (Peters et al., 2016; Read et al., 2008; Varese et al., 2012b).

We would recommend that the relationship between PLEs in new mothers, and future psychotic conditions, be investigated in future studies.

**Amendment made:** “Additionally, longitudinal studies investigating the transition from PLEs to psychosis are indicated.” (Page 11, line 25)

**Additional amendments:** An additional reference has been added, due to a recent paper being published, reviewing the onset of post-partum psychosis: VanderKruik, R., Barreix, M., Chou, D., Allen, T., Say, L. and Cohen, L.S., 2017. The global prevalence of postpartum psychosis: a systematic review. *BMC Psychiatry*, 17(1), p.272.

Due to the upper word limit being met sections of the paper have been amended.

## References

- Creedy, D.K., Shochet, I.M. and Horsfall, J., 2000. Childbirth and the development of acute trauma symptoms: incidence and contributing factors. *Birth*, 27(2), 104-111.
- Devilly, G.J., Gullo, M.J., Alcorn, K.L. and O'Donovan, A., 2014. Subjective appraisal of threat (criterion A2) as a predictor of distress in childbearing women. *The Journal of Nervous and Mental Disease*, 202(12), 877-882.
- Kendler, K.S., Gallagher, T.J., Abelson, J.M., Kessler, R.C., 1996. Lifetime prevalence, demographic risk factors, and diagnostic validity of nonaffective psychosis as assessed in a US community sample. The National Comorbidity Survey. *Archives of General Psychiatry* 53, 1022–1031.
- Nager, A., Johansson, L.M. and Sundquist, K., 2005. Are sociodemographic factors and year of delivery associated with hospital admission for postpartum psychosis? A study of 500 000 first-time mothers. *Acta Psychiatrica Scandinavica*, 112(1), 47-53.
- Nager, A., Sundquist, K., Ramírez-León, V. and Johansson, L.M., 2008. Obstetric complications and postpartum psychosis: a follow-up study of 1.1 million first-time mothers between 1975 and 2003 in Sweden. *Acta Psychiatrica Scandinavica*, 117(1), 12-19.
- Peters, E., Ward, T., Jackson, M., Morgan, C., Charalambides, M., McGuire, P., Woodruff, P., Jacobsen, P., Chadwick, P. and Garety, P.A., 2016. Clinical, socio-demographic and psychological characteristics in individuals with persistent psychotic experiences with and without a “need for care”. *World Psychiatry*, 15(1), 41-52.
- Read, J., Fink, P., Rudegeair, T., Felitti, V. and Whitfield, C., 2008. Child maltreatment and psychosis: a return to a genuinely integrated bio-psycho-social model. *Clinical Schizophrenia & related Psychoses*, 2(3), 235-254.
- Reed, R., Sharman, R. and Inglis, C., 2017. Women's descriptions of childbirth trauma relating to care provider actions and interactions. *BMC Pregnancy and Childbirth*, 17(1), 21.
- Slade, P., Morrell, C.J., Rigby, A., Ricci, K., Spittlehouse, J. and Brugha, T.S., 2010. Postnatal women's experiences of management of depressive symptoms: a qualitative study. *British Journal of General Practice*, 60(580), 440-448.
- Varese, F., Smeets, F., Drukker, M., Lieveerse, R., Lataster, T., Viechtbauer, W., Read, J., van Os, J. and Bentall, R.P., 2012. Childhood adversities increase the risk of psychosis: a meta-analysis of patient-control, prospective-and cross-sectional cohort studies. *Schizophrenia Bulletin*, 38(4), 661-67.

Wisner, K.L., Sit, D.K., McShea, M.C., Rizzo, D.M., Zoretich, R.A., Hughes, C.L., Eng, H.F., Luther, J.F., Wisniewski, S.R., Costantino, M.L. and Confer, A.L., 2013. Onset timing, thoughts of self-harm, and diagnoses in postpartum women with screen-positive depression findings. *JAMA Psychiatry*, 70(5), 490-498.

**Acknowledgments**

Thank you to Dr. Philip Powell, who kindly provided advice and assistance in regards to the statistical analysis.

**Conflict of interest**

All authors declare they have no conflicts of interest.


## **Contributors**

Dr. Lyndsey Holt, Professor Pauline Slade and Professor William Sellwood designed the study. Dr. Lyndsey Holt wrote the protocol, managed the literature searches and undertook the statistical analysis. Dr. Lyndsey Holt wrote the first draft of the manuscript. All authors contributed to and have approved the final manuscript.

**Funding body arrangements**


This project was funded by The University of Liverpool, as part of Dr. Lyndsey Holt's Doctorate in Clinical Psychology.

Figures




**Figure 1** Three nested competing path models exploring the relationship between birth experience and psychotic like experiences (hallucination and delusional experiences) through two stages of mediating variables (i) trauma appraisals, post-traumatic stress symptoms and adjustment to motherhood (ii) self-concept clarity

- Pathways included in Model 1 only
- ..... Pathway included in Model 2 only
- Pathways included in Model 1, 2 and 3



**Figure 2** Participant recruitment flowchart



**Figure 3 Model 2 exploring the relationship between birth experience and psychotic like experiences** (hallucination and delusional experiences) through two stages of mediating variables (i) trauma appraisals, post-traumatic stress symptoms and adjustment to motherhood (ii) self-concept clarity. The regression coefficients for paths trauma appraisals\*hallucinations, trauma appraisals\*delusions, adjustment to motherhood\*hallucinations and adjustment to motherhood\*delusions were constrained to 0. For simplicity error terms and control variables have been omitted. Estimates on the endogenous variables were controlled for by age, birth method, education level, employment status, ethnicity, marital status, depression and sleep quality. Error terms were correlated as appropriate, with all stage one mediator error terms (trauma appraisals, post-traumatic stress symptoms and adjustment to motherhood) correlated with each other and outcome variable error terms (hallucination and delusional experiences) correlated with each other. Estimates shown are standardised betas ( $\beta$ ). Significance levels were established via bootstrapped CIs (10,000 resamples). A bootstrapped Chi square test indicated a good model fit,  $\chi^2(4) = 4.49, p = .34$ ; CFI = 1.00, RMSEA = 0.01, 90% CI [0.00, 0.03]  $p = .98$ .

## Tables

**Table 1** Participant characteristics

	Range	Mean ( <i>SD</i> )
Age (years)	16-48	28.9 (5.51)
		Total n (%)
Marital status		
Married		732 (52.5)
Not married		661 (47.5)
Missing data		7 (0.50)
Educational attainment		
Low education (no qualifications/GCSE's)		184 (13.2)
Mid education (A Levels/vocational qualifications)		500 (35.9)
High education (graduate/post graduate)		709 (50.9)
Pre-pregnancy employment		
Employed (full time/part time/self employed)		1263 (90.6)
Unemployed (out of work/voluntary work/student)		130 (9.4)
Ethnic origin		
White		1254 (90)
Other		139 (10)
Number of babies delivered		
Singleton		1357 (98.7)
Multiple		18 (1.3)
Mode of birth		
Normal vaginal delivery		707 (50.8)
Assisted delivery (e.g. Kiwi, Forceps or Ventouse)		330 (23.7)
Caesarean section		356 (25.6)

*N* = 1393.

**Table 2** Descriptive statistics

Variable	Mean ( <i>SD</i> )	Possible ranges of scores
Birth experience	54.25 (28.88)	0-149
Post-traumatic stress symptoms		
Arousal	0.65 (0.72)	0-4
Intrusion	0.62 (0.67)	0-4
Avoidance	0.77 (0.81)	0-4
Total score	14.98 (15.1)	0-88
Trauma appraisal	0.77 (0.85)	0-2
Adjustment to motherhood		
Child	6.03 (3.88)	0-18
Adult	4.60 (3.36)	0-15
Emotional closeness	0.57 (0.95)	0-6
Total score	11.2 (7.02)	0-39
Self-concept clarity	42.36 (9.52)	12-60
Hallucination experiences		
Auditory hallucinations	0.69 (0.84)	0-4
Visual hallucinations	0.04 (0.20)	0-1
Vivid thoughts	0.58 (0.78)	0-2
Vivid day dreams	0.46 (0.81)	0-3
Total score	2.20 (2.03)	0-12
Delusion experiences		
Distress <sup>a</sup>	9.44 (8.86)	0-105
Preoccupation <sup>b</sup>	9.35 (8.61)	0-105
Conviction <sup>c</sup>	12.08 (9.92)	0-105
Total score	3.49 (2.84)	0-21
Composite score	30.46 (29.46)	0-336
Depressive symptoms	7.06 (4.72)	0-30
Average hours of sleep	5.83 (1.61)	1-16
	Frequency (%)	
Quality of sleep		
Good	335 (24.00)	-
Poor	835 (60.00)	-
Unknown	223 (16.00)	-
Trauma appraisals		
Experienced danger	524 (37.60)	-
Danger for self	27 (5.20)	-
Danger for baby	269 (51.30)	-
Danger for self and baby	228 (16.40)	-
Experienced intense fear	557 (40.00)	-
Both criteria met	384 (27.60)	-
Above clinical cut off for PTSD <sup>d</sup>	167 (12.00)	-
Above clinical cut off for depression <sup>e</sup>	172 (12.30)	-

*N* = 1,393.

<sup>a, b, c</sup> *n* = 1217, corresponding to number of participants reporting at least one delusion. <sup>d</sup> Score  $\geq$  33, recommended by Creamer et al. (2003). <sup>e</sup> Score  $\geq$  13, recommended by Matthey et al. (2006)

**Table 3** Inter-correlations of study variables

	1	2	3	4	5	6	7	8	9	10	11	12	13	14
1. Birth experience	-													
2. Trauma appraisals	.64**	-												
3. Post-traumatic stress symptoms	.56**	.47**	-											
4. Adjustment to motherhood	.44**	.28**	.47**	-										
5. Self-concept clarity	-.37**	-.27**	-.41**	-.59**	-									
6. Hallucination experiences	.22**	.21**	.39**	.30**	-.40**	-								
7. Delusional experiences	.24**	.22**	.42**	.36**	-.46**	.52**	-							
8. Depression	.36**	.25**	.44**	.61**	-.63**	.38**	.46**	-						
9. Good sleep	-.22**	-.17**	-.26**	-.30**	.20**	-.13**	-.15**	-.29**	-					
10. Age	.02	-.04	-.06*	-.01	.19**	-.21**	-.20**	-.13**	-.11**	-				
11. Ethnicity	.01	.00	.01	.06*	-.13**	-.00	-.16**	-.03	.00	-.05	-			
12. Employment status	-.02	.00	-.07*	.01	.05*	-.12**	-.15**	-.08**	.00	.23**	.16**	-		
13. Mode of Birth	.33**	.28**	.13**	.03	-.01	-.02	-.03	.03	-.07*	.22**	.00	.06*	-	
14. Marital Status	-.03	-.04	-.06*	.01	.13**	-.13**	-.09**	-.08**	-.03	.33**	-.16**	.11**	.04	-
15. Education	.02	.00	.02	.08**	.12**	-.12**	-.09**	-.06*	-.07*	.37**	-.15**	.11**	.03	.31**

$N = 1393$ . Correlations represent Spearman's  $r$  or phi ( $r\phi$ ) coefficients.

\*  $p < .05$ ; \*\*  $p < .01$

*Note:* In order to undertake correlational analyses demographic data were simplified and recoded as appropriate: Ethnicity (0 = *not White British*, 1 = *White British*), marital status (0 = *not married*, 1 = *married*), employment status (0 = *not employed*, 1 = *employed*), mode of birth (coded in terms of increasing intervention; 1 = *normal delivery*, 2 = *assisted delivery*, 3 = *caesarean section*) and education level (1 = *low education*, 2 = *mid education*, 3 = *high education*).

**Table 4** Endorsement of items on the Launay Slade Hallucination Scale and the 21-item Peters Delusion Inventory

Scales, subscales and items	Frequency (%)
<b>Launay Slade Hallucination Scale</b>	
No matter how much I try to concentrate on my work unrelated thoughts always creep into my mind	594 (42.6)
Sometimes a passing thought will seem so real that it frightens me	439 (31.5)
I have never been troubled by hearing voices in my head <sub>r</sub>	379 (27.2)
Sometimes my thoughts seem as real as actual events in my life	370 (26.6)
The sounds I hear in my daydreams are generally clear and distinct	272 (19.5)
In my daydreams I can hear the sound of a tune almost as clearly as if I were actually listening to it	249 (17.9)
I have never heard the voice of the Devil <sub>r</sub>	215 (15.4)
In the past I have had the experience of hearing a person's voice and then found that there was no-one there	159 (11.4)
I often hear a voice speaking my thoughts aloud	141 (10.1)
The people in my daydreams seem so true to life that I sometimes think they are	125 (9)
In the past I have heard the voice of God speaking to me	63 (4.5)
On occasions I have seen a person's face in front of me when no-one was in fact there	58 (4.2)
<b>21-item Peters Delusion Inventory</b>	
Do you ever feel as if some people are not what they seem to be	941 (67.6)
Do you ever feel as if people seem to drop hints about you or say things with a double meaning?	553 (39.7)
Are you often worried that your partner may be unfaithful?	413 (29.6)
Do you ever feel that people look at you oddly because of your appearance?	317 (22.8)
Do you ever feel as if electrical devices such as computers can influence the way you think?	297 (21.3)
Do you ever feel that you are a very special or unusual person?	276 (19.8)
Do you ever feel as if things in magazines or on TV were written for you?	213 (15.3)
Do you ever think people can communicate telepathically?	205 (14.7)
Do you ever feel as if you are, or destined to be someone very important?	186 (13.4)
Do you ever feel that you are especially close to God?	179 (12.8)
Do you ever feel that you have sinned more than the average person?	167 (12.0)
Do you ever feel as if you are being persecuted in some way?	163 (11.7)
Do you believe in the power of witchcraft, voodoo or the occult?	163 (11.7)
Do you ever feel as if you had no thoughts in your head at all?	160 (11.5)
Do you ever feel as if you have been chosen by God in some way?	123 (8.8)
Do your thoughts ever feel alien to you in some way?	113 (8.1)
Do you ever feel as if your own thoughts were being echoed back to you?	83 (6.0)
Do you ever feel as if the world is about to end?	82 (5.9)
Have your thoughts ever been so vivid that you were worried other people would hear them?	76 (5.5)
Do you ever feel as if there is a conspiracy against you?	76 (5.5)
Do you ever feel as if you are a robot or zombie without a will of your own?	71 (5.1)

*N* = 1,393

<sub>r</sub> Reverse scored item

**Table 5** Goodness-of-fit statistics for hypothesised models

Model	Comparative model	$\chi^2$	$df$	$p$ value $\chi^2$	CFI	RMSEA	$\Delta\chi^2$	$\Delta df$	$p$ value $\Delta\chi^2$
Model 1	-	4.13	1	.042	.999	.047	-	-	-
Model 2	Model 1	4.49	4	.344	1.000	.009	0.36	3	.948
Model 3	Model 2	8.71	5	.121	.999	.023	4.22	1	.040

$\Delta\chi^2$  = difference in  $\chi^2$  values between models;  $\Delta df$  = difference in number of degrees of freedom between models.

**Table 6** Path estimates for model 2 (Figure 3)

Path	Estimate	SE of $b$	Bootstrap 95% CIs $b$	Bootstrap 95% CIs $\beta$
------	----------	-----------	-----------------------	---------------------------

	<i>b</i>	$\beta$	ML	Bootstrap	Lower	Upper	Lower	Upper
Direct path estimates								
a <sub>1</sub>	0.27***	.52	.01	.02	.24	.30	.47	.57
a <sub>2</sub>	0.02***	.60	.00	.00	.02	.02	.56	.64
a <sub>3</sub>	0.07***	.28	.01	.01	.06	.08	.23	.32
b <sub>1</sub>	- 0.02	-.03	.02	.02	-.05	.01	-.08	.02
b <sub>2</sub>	- 0.56*	-.05	.28	.28	- 1.11	.00	-.10	.00
b <sub>3</sub>	- 0.42***	-.31	.04	.04	-.49	-.35	-.36	-.28
c <sub>1</sub>	- 0.04***	-.17	.01	.01	-.05	-.02	-.23	-.11
c <sub>2</sub>	- 0.07***	-.24	.01	.01	-.09	-.05	-.30	-.18
d <sub>1</sub>	0.00	-.05	.00	.00	-.01	.00	-.11	.13
d <sub>2</sub>	- 0.01*	-.06	.00	.00	-.01	.00	-.12	.00
e <sub>1</sub>	0.04***	.28	.00	.00	.03	.05	.21	.34
e <sub>2</sub>	0.05***	.24	.01	.01	.03	.06	.18	.30
f	- 0.02*	-.06	.01	.01	-.04	.00	-.12	.00
Indirect path estimates								
a <sub>1</sub> b <sub>1</sub>	.00	.00	-	.00	-.01	.00	-.04	.01
a <sub>1</sub> e <sub>1</sub>	.01***	.16	-	.00	.01	.01	.12	.19
a <sub>1</sub> e <sub>2</sub>	.01***	.14	-	.00	.01	.02	.11	.18
a <sub>2</sub> b <sub>2</sub>	-.01*	-.03	-	.01	-.02	.00	-.06	.00
a <sub>3</sub> b <sub>3</sub>	-.03***	-.09	-	.00	-.04	-.02	-.11	-.07
b <sub>1</sub> c <sub>1</sub>	.00	.01	-	.00	.00	.00	.00	.01
b <sub>1</sub> c <sub>2</sub>	.00	.01	-	.00	.00	.00	-.01	.02
b <sub>2</sub> c <sub>1</sub>	.02*	.01	-	.01	.00	.05	.00	.02
b <sub>2</sub> c <sub>2</sub>	.03*	.01	-	.02	.00	.08	.00	.02
b <sub>3</sub> c <sub>1</sub>	.02***	.05	-	.00	.01	.02	.03	.07
b <sub>3</sub> c <sub>2</sub>	.03***	.07	-	.01	.02	.04	.05	.10
a <sub>1</sub> b <sub>1</sub> c <sub>1</sub>	.00*	.01	-	.00	.00	.00	.00	.01
a <sub>1</sub> b <sub>1</sub> c <sub>2</sub>	.00*	.02	-	.00	.00	.00	.01	.02
a <sub>2</sub> b <sub>2</sub> c <sub>1</sub>	.00**	.01	-	.00	.00	.00	.01	.01
a <sub>2</sub> b <sub>2</sub> c <sub>2</sub>	.00**	.01	-	.00	.00	.00	.01	.02
a <sub>3</sub> b <sub>3</sub> c <sub>1</sub>	.00***	.02	-	.00	.00	.00	.01	.02
a <sub>3</sub> b <sub>3</sub> c <sub>2</sub>	.00***	.02	-	.00	.00	.01	.02	.03
fc <sub>1</sub>	.01*	.01	-	.00	.01	.02	.14	.02
fc <sub>2</sub>	.02*	.01	-	.00	.01	.02	.14	.03

*N* = 1393. ML = maximum likelihood estimation. Probability values determined on bootstrapped CIs (10,000 resamples)

\* *p* < .05; \*\* *p* < .01; \*\*\* *p* < .001