

Manuscript Number: OO-D-17-710

Title: Head and Neck Reconstruction with Free Flaps Based on the Thoracodorsal System

Article Type: VSI: Head & Neck Reconstruct

Section/Category: Clinical

Keywords: Thoracodorsal; free flap; head and neck cancer; reconstruction

Corresponding Author: Mr. John Edward O'Connell, MB, BDS, FFD RCSI

Corresponding Author's Institution: Aintree University Hospitals NHS Foundation Trust

First Author: John Edward O'Connell, MB, BDS, FFD RCSI

Order of Authors: John Edward O'Connell, MB, BDS, FFD RCSI; Mandeep U Bajwa, MB, BDS, MSc, FRCS; Andrew G Schache, PhD FDSRCS FRCS(OMFS); Richard J Shaw, MD FDS FRCS(OMFS)

Abstract: The advent of micro-vascular free tissue transfer has facilitated the reconstruction of increasingly complex Head and Neck defects. There are multiple donor sites available, each with its own advantages and disadvantages. However, the subscapular system, including the thoracodorsal system, provides the widest array of soft and hard tissue flaps, as well as chimeric options. Its advantages include a long pedicle, independently mobile tissue components, relative sparing from atherosclerosis, and minimal donor site morbidity. The soft tissue flaps available from the thoracodorsal system include the Latissimus Dorsi, and Thoracodorsal Artery perforator flaps, while the tip of scapula provides the hard tissue component. This review paper outlines the anatomical basis for these flaps, as well as describing their utility in head and neck reconstruction.

To the editor of Oral Oncology

Dear Editor,

Please find attached a review manuscript, entitled '*Head and Neck Reconstruction with Free Flaps Based on the Thoracodorsal System*' that I would like you to consider for publication in your Journal.

Please note that there was no funding for this research, and that the authors do not have commercial associations, current and within the past five years, that might pose a potential, perceived or real conflict of interest.

Please let me know if you need any further information.

Yours sincerely,

John Edward O'Connell*

*Corresponding Author: Specialist Registrar, Department of Oral & Maxillofacial / Head and Neck Surgery
Aintree University Hospitals NHS Foundation Trust, Lower lane, Fazakerly, Liverpool. L9 7AL
Fax: 0151 529 5288; Tel: +44 7429666670; Email: johnedoconnell@yahoo.com

Oral Oncology Author Form

All manuscripts submitted to *Oral Oncology* must be accompanied by this form. Please scan the form and transmit it to the Editorial Office via EES with the manuscript. If you are unable to do this, please contact the Editorial Office at ooncology@elsevier.com to organise an alternative way of sending the form to the *Oral Oncology*.

Title of Manuscript:

..... Head and neck reconstruction with Free Flaps based on the Thoracodorsal System

Contribution

Author(s)

Study concepts:

..... Prof RJ Shaw, Mr AG Schache

Study design:

..... Prof RJ Shaw, Mr AG Schache, Mr JE O'Connell, Mr M U Bajwa

Data acquisition:

..... Mr JE O'Connell, Mr M U Bajwa

Quality control of data and algorithms:

..... Mr JE O'Connell, Mr M U Bajwa

Data analysis and interpretation:

..... Mr JE O'Connell, Mr M U Bajwa

Statistical analysis:

..... Prof R J Shaw, Mr A G Schache

Manuscript preparation:

..... Prof R J Shaw, Mr A G Schache

Manuscript editing:

Manuscript review:

Ethical Approval for Research: No / Yes N.A.

External Funding: No / Yes

Source of Funding: NA

Name of Principal Investigator:

(If funded, please include a statement as to the role of the study sponsor at end of manuscript under a heading 'Role of the Funding Source')

Possible Conflict of Interest: No / Yes

(Please ensure that a 'Conflict of Interest' statement is included in your manuscript)

Number of Tables: 0

Number of Figures: 7

Name and Title of Corresponding Author: Mr John Edward O'Connell

Address: Department of Oral and Maxillofacial/Head and Neck Surgery

Address: Aintree University Hospital, Lower Lane, Fazakerly


Postcode and country: L9 7AL, United Kingdom

Tel No: +44 7429656670

Fax No:

Email: johnedoconnell@yahoo.com

"I confirm that all the authors have made a significant contribution to this manuscript, have seen and approved the final manuscript, and have agreed to its submission to the *Oral Oncology*".

Signed (corresponding author): 

Date: 22/08/2017

Highlights

- The Thoracodorsal system (forms part of the sub-scapular system) provides a wide range of soft and hard tissue free flaps, as well as chimeric options, for head and neck reconstruction
- Latissimus Dorsi, Thoracodorsal Artery Perforator, and Tip of Scapula are main flaps available
- Particularly useful in vessel deplete necks, multi-surface defects, elderly patients
- Its use has been well described in mandibular, maxillary, mid-face, and scalp defects
- Technically challenging, but low morbidity relative to other free flap options

Title

Head and Neck Reconstruction with Free Flaps Based on the Thoracodorsal System

Authors and Affiliations

John Edward O'Connell¹

Mandeep U Bajwa^{1,2}

Andrew G Schache^{1,2}

Richard J Shaw^{1,2}

1. Regional Maxillofacial Unit, Aintree University Hospital, Liverpool, UK.

2. Department of Molecular and Clinical Cancer Medicine, Northwest Cancer Research Centre, University of Liverpool, Liverpool, UK.

Corresponding Author

John Edward O'Connell MB, BDS, FFD, MRCS

Specialist Registrar

Department of Oral & Maxillofacial / Head and Neck Surgery

Aintree University Hospitals NHS Foundation Trust

Lower lane, Fazakerly,

Liverpool. L9 7AL

Fax: 0151 529 5288

Tel: +44 7429666670

Email: johnedoconnell@yahoo.com

Work count

4,362

Key-words

Thoracodorsal; free flap; head and neck cancer; reconstruction;

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Conflict of Interest

None declared

Head and Neck Reconstruction with Free Flaps Based on the Thoracodorsal System

Abstract

The advent of micro-vascular free tissue transfer has facilitated the reconstruction of increasingly complex Head and Neck defects. There are multiple donor sites available, each with its own advantages and disadvantages. However, the subscapular system, including the thoracodorsal system, provides the widest array of soft and hard tissue flaps, as well as chimeric options. Its advantages include a long pedicle, independently mobile tissue components, relative sparing from atherosclerosis, and minimal donor site morbidity. The soft tissue flaps available from the thoracodorsal system include the Latissimus Dorsi, and Thoracodorsal Artery perforator flaps, while the tip of scapula provides the hard tissue component. This review paper outlines the anatomical basis for these flaps, as well as describing their utility in head and neck reconstruction.

Introduction

Evolution in the range of free flaps available for head and neck cancer has facilitated successful reconstructive surgery in increasingly demanding defects. Patients presenting with extensive primary or recurrent tumours, as well as cases of osteoradionecrosis, form a significant part of the head and neck reconstructive surgeon's practice. These cases often incorporate complex three-dimensional defects that include both hard and soft tissue components as well as more than one epithelial surface (e.g. mucosa and skin). Microvascular free tissue transfer is considered the gold standard in these complex cases and the choice of flap is varied and dependent on factors including the site, size, complexity of the defect as well as patient co-morbidities and indeed surgeon training and preference. The Thoracodorsal system offers both soft and hard tissue flaps as well as chimeric options. The common soft tissue flaps are the Latissimus Dorsi (LD) and Thoracodorsal Artery Perforator (TDAP) flaps, whilst the Tip of Scapula (TSCAP) remains the predominant hard tissue component, with or without additional soft tissue elements. We, herein, provide a review of the anatomical basis of these flaps and their relevance to reconstruction within head and neck practice.

Overview of Thoracodorsal Vascular Anatomy

Figure 1 summarises critical aspects of thoracodorsal arterial anatomy relevant to the reconstructive surgeon. The subscapular artery is the largest branch of the axillary artery with a mean diameter of 6mm (range 4 – 8mm)¹. Both it and the subscapular vein arise from the posterior surface of the third part of the axillary artery and vein, following the inferior margin of the subscapularis muscle for 2.2cm (range 2 – 3cm), before dividing into two terminal branches, the circumflex scapular and thoracodorsal arteries and veins respectively. The subscapular artery and vein arise in close proximity in the vast majority of cases. In those cases where the artery and vein originate separately, the subscapular artery lies proximally in the axilla by an average of 4.2cm^{2,3}. In addition, it should be noted that the

thoracodorsal artery arises directly from the axillary artery in 0.8 – 3% of cases^{1,4,5}. In some instances it may even arise from the lateral thoracic artery.

The thoracodorsal vessels travel cephalo-caudally along the lateral border of the scapula, through the fatty tissue of the axilla, before entering the hilum of the Latissimus Dorsi muscle. The mean length of the thoracodorsal pedicle between origin and muscular hilum is 9.3cm (range 6 – 16.5cm). At its origin, the mean diameter of the artery is 2.7mm (range 1.5 – 4mm) and the vein is 3.4mm (range 1.5 – 4.5mm)². Along its course to the muscle the thoracodorsal artery supplies named branches to the tip of scapula (angular branch), and to a number of muscles including subscapularis, teres major, and serratus anterior. The angular branch is reported as being consistent and has a mean pedicle length of 6.83 cm (range 2.5 – 8cm).⁶ Anecdotally however we have found it to be missing or diminutive in a small minority of cases (5%). It is of considerable relevance that the thoracodorsal system, relative to other vessels (particularly those in the lower limb and pelvis used to harvest vascularized bone⁷), is spared from atherosclerotic changes^{1,4}.

Cadaveric dissections have demonstrated that in 92% of cases, the thoracodorsal artery and vein cross the axilla together. In the remaining cases, the artery and vein may not join until the takeoff of branches to the serratus anterior. The thoracodorsal artery sends just one branch to the serratus anterior in 54% of cases, with 2 or 3 branches less frequently¹. In approximately 1% of cases, the thoracodorsal artery does not send a branch to the serratus anterior muscle⁸.

Latissimus Dorsi (LD)

First described by Tansini in 1896⁹ as a pedicled skin rotation flap and subsequently as a musculocutaneous flap, the LD has a long history in thoracic and limb reconstruction. Although described by Quillen¹⁰ in 1978 as a pedicled flap in head and neck reconstruction, Maxwell¹¹ first reported its use in free tissue transfer, also in 1978. Ease of dissection, large surface area, and length of the vascular pedicle are some of reasons that account for the popularity of the free LD flap.

The LD is a fan-like muscle originating medially and extending from the dorsal iliac crest caudally and travelling cranially via the sacral, lumbar, and thoracic (lower 6) vertebrae. The fibres insert into the humerus via a space between teres major and pectoralis major. Along with the teres major, it forms the posterior axillary fold^{12, 13}.

Within the LD, the thoracodorsal artery divides into transverse and vertical branches in 94% of cases, thus forming the anatomical basis for splitting the muscle longitudinally in order to harvest two separately vascularised muscular or musculocutaneous units (the thoracodorsal nerve parallels the vessel). In the remaining 6%, the vessel divides into 3 or 4 parallel vessels, which traverse the muscle¹⁴. The transverse branch usually runs parallel to the free upper border of the LD muscle, 3.5cm from the edge. The vertical branch usually takes a cranio-caudal course towards the iliac crest, 2.1cm

from the lateral edge of the muscle. These branches then become intra-muscular. These intra-muscular branches of the thoracodorsal artery supply the musculocutaneous perforators on which the TDAP flap is based. The anatomical basis for the TDAP flap will be outlined in more detail below.

Based on the work of Taylor and Palmer¹⁵, the LD is divided into three angiosomes. The most proximal segment is supplied predominantly by the thoracodorsal artery, with the middle portion by the posterior intercostal arteries, and the distal segment is supplied by the lumbar arteries. Therefore, the skin paddles overlying the distal angiosome are less reliable when based on the thoracodorsal pedicle only.

Owing to the high success rate with microvascular free tissue transfer, and increased versatility in flap design, the LD is now more commonly used as a free rather than pedicled flap. A failure rate of 0-8% has been described for the free LD used in head and neck reconstruction^{16, 17, 18}. The ability to utilise the vascular anatomy, and split the muscle longitudinally with 2 separate skin paddles based on the transverse and vertical branches of the thoracodorsal artery respectively, allows the reconstruction of multi-surface oral defects^{19, 20}. In our practice the main indication for this broad flap is in the reconstruction of large scalp, mid-face and skull base defects either in isolation or as a chimeric flap including bone sourced from the scapula. This is in keeping with other units^{21, 22, 23}. **Figure 2** demonstrates an example of LD-TSCAP chimeric flap from our own series.

In 2014, Lee and Mun²⁴ performed a systematic review of functional donor-site morbidity after LD muscle transfer and found that while functional impairment of the shoulder could occur, there was little evidence for interference with activities of daily living. As one would expect, the muscle-sparing LD and TDAP flaps showed lower functional morbidity.

While the focus of this review is primarily on free tissue transfer, the pedicled LD does have a potential role in salvage surgery and/or in patients with vessel depleted neck, and is included for completeness. Relative to other pedicled regional flaps, for example the pectoralis major, the pedicled LD has advantages including a broader surface area and a less obvious scar. The reported success rate is 80-90%^{25, 26, 27, 28}. Like all flaps the pedicled LD is not without its disadvantages²⁹ including a requirement to turn the patient (precluding simultaneous tumour ablation and flap raising), prolonged wound drainage and seroma formation, and risk of injury to the brachial plexus³⁰. In addition to these reported disadvantages, we have found that the pedicled LD has a propensity to pull away from defects in the head and neck, presumably due to the weight of the flap, and also carries a significant risk of vascular compromise as a result of venous occlusion at the apex of pedicle rotation.

Thoracodorsal Artery Perforator (TDAP) Flap

Angrigiana et al³¹ and Spinelli et al³², in 1995 and 1996 respectively, described the TDAP flap, which is based on perforators arising from the thoracodorsal artery and utilises the overlying skin but not the underlying LD muscle. The TDAP flap therefore shares the benefits of long pedicle length and broad large surface area, yet has the additional advantages of reduced thickness and decreased morbidity when comparison is made to the LD flap³³.

As described previously, the terminal branches of the thoracodorsal artery become intra-muscular within the Latissimus Dorsi. A single perforator is sufficient to supply a skin paddle of up to 15cm by 25cm³¹. Perforators originating in the vertical branch have a shorter intra-muscular course and are therefore preferred over those derived from the transverse branch. An average of more than three musculocutaneous perforators, predominantly from the vertical branch, have been demonstrated^{34, 35, 36}. The thoracodorsal artery may, in fact, supply direct cutaneous perforators before it enters the LD muscle¹³ but their use potentially comes at the expense of pedicle length.

Two consistent anatomical landmarks have been described for localising perforators intraoperatively. The first, published by Angrigiani³¹, describes the main perforators arising from the vertical branch as it leaves the muscle in the subcutaneous tissue. This point lies 8cm inferior to the axilla in the posterior axillary line, and 2cm medial to the lateral border of LD. The second relates to the bifurcation of the thoracodorsal artery, into the vertical and transverse branches. Heitman et al³⁴ suggested that all perforators were in close proximity to a point 4cm inferior to the tip of scapula, and 2.5cm medial to the free border of LD. Perhaps the most useful description is that of Schaverien et al³⁷ who found the greatest concentration of perforators (> 0.5mm in diameter) 9.5 – 15.4cm from the posterior axillary fold within 4.3 cm of the lateral border of LD. Therefore, siting a skin paddle over this area increases the chance of having a perforator with a diameter >0.5mm. **Figure 3** demonstrates the anatomical location of the perforators derived from the paper by Schaverien et al³⁷.

The TDAP flap possesses many of the qualities that are required for a soft tissue flap in head and neck reconstruction. It provides a large and relatively thin hairless skin paddle, and has a mean pedicle length of 15cm^{38, 39}. As with the LD, this makes it particularly useful in vessel-depleted necks, where the contra-lateral neck may be used for anastomosis, and in reconstruction of the mid-face, skull base and scalp. The main difference between the TDAP and LD is the volume of soft tissue, the TDAP being considerably thinner. The TDAP has been shown to be a reliable flap by Hamdi et al⁴⁰, who in a retrospective review of 90 TDAP flaps for breast reconstruction report only one case of flap loss. Similarly, Lee and Mun⁴¹, report the loss of only one TDAP flap in a review of 31 flaps.

Relative to the radial forearm (RFFF) and Antero-lateral thigh (ALT) free flaps, the TDAP flap, in our experience, is more technically challenging to raise. **Figure 4** shows a TDAP flap being raised with a large skin paddle and demonstrates the length of pedicle achievable. The intra-muscular dissection can be difficult and unforgiving due to the relatively small perforators and propensity for multiple

branches. However, the resultant scar below the posterior axillary fold is easily hidden and donor site morbidity is reported as being minimal when compared to the RFFF⁴². Importantly, the TDAP flap spares both the LD muscle and its motor nerve, thereby preserving muscle strength and shoulder function. While some authors⁴² have claimed it is possible to harvest the flap in the supine position and therefore facilitate two-team operating, harvesting in a lateral decubitus position is more common but as a result includes a time consuming element.

Tip of Scapula

Swartz et al⁴³, in what was the first published series of cases where the scapula was used as a donor site for head and neck reconstruction; found that when the tip of scapula was included, its blood supply was tenuous. Deraemaeker et al⁴⁴ subsequently identified the angular branch of the thoracodorsal artery as an additional or indeed independent blood supply to the caudal portion of the lateral scapula. Coleman and Sultan⁴⁵ consequently described a free flap with the scapular tip and Latissimus Dorsi as a single unit. In addition, they highlighted that use of the angular artery enabled the harvesting of 2 separate bony segments based on 2 separate branches of the subscapular artery, namely the circumflex scapular artery and the angular branch of thoracodorsal. However, it should be noted that up to 8% of patients do not have a subscapular artery; in such instances the circumflex and thoracodorsal arteries arise independently from the axillary artery^{46, 47}.

As previously mentioned Deraemaeker et al⁴⁴ identified the angular branch as an independent blood supply to the tip of scapula. Seitz et al⁴⁸ reported the mean combined pedicle length of the angular branch and thoracodorsal artery as being 14.8cm. This increases to 16.7cm when the subscapular artery is included. Chepeha et al⁴⁹ reported a mean length of bone in addition to pedicle as an impressive 27cm (range 23 – 32cm).

Regarding the amount of bone that can be safely harvested on the angular branch, Yoo et al⁵⁰ have reported the average length as 6.2cm with the longest segment measuring 8cm in their series. This compares similarly to the average length of 5.2cm (range 2.5 – 9cm) described by Chepeha et al⁴⁹. Seneviratne et al⁵¹, in 81 cadaveric dissections, found that the angular branch can supply up to 20cm of bone from both the medial and lateral scapula, however this has not yet been reproduced in a clinical series.

The Tip of Scapula can be harvested as a stand alone myo-osseous flap (with teres major), or as part of a chimeric flap. Its application as part of a chimeric flap will be discussed further below. Relative to other composite bone flaps, this flap has a number of advantages; namely a long pedicle, independently mobile chimeric options, and the scope for a variety of osseous shapes (albeit limited in terms of scapula thickness). However, the volume of bone available is limited by comparison to the iliac crest and fibula. The Tip of Scapula has been described for reconstruction of short posterior mandibular defects, including those of the angle (Brown Class 1)^{49, 52, 53, 54}. Its utility is exemplified in both the

description of a condyle-ramus unit by Yoo et al⁴⁹ and its usage for reconstruction of 39 maxillary defects by Miles & Gilbert⁵⁵.

Figure 6 demonstrates a Tip of Scapula flap being used for a Brown Class V maxillary defect requiring reconstruction of the orbit. In this case the flap was positioned vertically.

Chimeric Flaps

As previously mentioned, one of the greatest advantages of the thoracodorsal system is versatility of the chimeric flap options on offer. In addition, and owing to the fact that the circumflex scapular and thoracodorsal arteries are commonly derived from the subscapular artery, both scapular and parascapular flaps can be incorporated to form even reconstructive flexibility. **Figure 5** shows a complex chimeric option based on the TDAP and a composite parascapular flaps. Although not included within this free flap it is also possible to include the TSCAP as a further bony element if required. We have previously published the use of the TDAP-Scap in situations calling for both hard and soft tissue complex reconstruction, often in vessel depleted areas⁵⁶. The subscapular system is unique in its ability to address all of the reconstructive requirements in these cases. A long pedicle, along with independently mobile tissues can often negate the need for harvesting a second free flap.

While the need for 2 flaps to reconstruct a defect is rare, those defects that do require more than one flap are challenging. Wei et al⁵⁷ suggested that composite resections not adequately reconstructed with one flap, for example large intra-oral defects and complex 3 dimensional defects are some of those that may require chimeric flaps, the alternative being two separate free flaps. In our practice we have found that separate free flaps have an increased risk of failure and it would be our preference to utilise the subscapular system in these situations. **Figure 6** shows a photographic series of a case of complex mandibular reconstruction with the LD-TSCAP chimeric flap for an extensive through-and-through defect. This series demonstrates the versatility of the soft tissue and length of pedicle reaching the contra-lateral neck.

Uglesic et al⁵⁸ were the first group to publish a large series of cases where radical maxillectomy defects were reconstructed using a LD-TSCAP flap (similar to that shown in **Figure 2**). Several other authors have described similar reconstructions^{13, 59, 60}. Chepeha et al⁴⁹ highlighted the potential benefits of the chimeric TSCAP flap; including a long pedicle, independently mobile soft tissue components, and the 3-dimensional nature of the scapular tip. L'Heureux-Lebeau et al⁶¹, in a retrospective review, reported the outcomes of 16 patients in whom oro-mandibular defects were reconstructed with the LD-TSCAP flap. They reported no flap losses, and concluded that this chimeric reconstruction is both safe and reliable.

Our group⁶² previously detailed the use of a LD-TSCAP flap for management of 9 patients following midface resections, of which 8 were extended maxillectomies (Brown Class 4)⁶³ with orbital exenteration. While there were no micro-vascular failures in this series, the bony component of one flap proved not to have a viable blood supply was ultimately debrided.

Midface Reconstruction

The thoracodorsal system, with its variety of free flaps and pedicle length, is useful in maxillary/midface reconstruction. The 3-dimensional nature of these defects asks unique questions of the reconstructive surgeon. Not only may vascularised bone be required to replace the palate, facial profile or orbital wall, but soft tissue may also be required to line the oral and/or nasal cavities, as well as cutaneous defects. While **Figure 7** demonstrates how the TSCAP can be positioned in the vertical plane to reconstruct orbital defects, the TSCAP can also be positioned in the horizontal plane to reconstruct low-level defects (Brown Class 2 or 3). The TSCAP is particularly suited to this reconstruction as shown by Pagedar et al⁶⁴ who found remarkable similarities in both size and shape in a study based on CT imaging. However, as mentioned before, the thickness of bone is not ideal for endosseous implants and as such the potential for complete oral rehabilitation is compromised. Miles and Gilbert⁵⁵ reported a 46% complication rate in their series of 39 TSCAP flaps used to reconstruct maxillectomy defects, although most were relatively minor. Perhaps the most note-worthy complications were firstly that 36% of patients had an inadequately reconstructed gingivo-buccal sulcus (further reducing the chance of oral rehabilitation) and that 21% developed a palatal fistula (albeit half of which closed spontaneously).

While the traditional scapula flap, based on the circumflex scapular artery, has been used for maxillary reconstruction, it is limited by short pedicle length and more frequent necessity for vein grafts, and the inherent increase risk of microvascular compromise. For this reason, if a decision is made to utilise the scapula donor site for a patient requiring midface reconstruction, it is our preference to harvest the Tip of Scapula.

Mandibular Reconstruction

Use of the fibula, iliac crest, scapula, and composite radial free flaps have been well described for mandibular reconstruction⁶⁵. The Tip of Scapula, notwithstanding the decreased ability to place endosseous implants⁶⁶, and a lack of bone length, has some potential advantages. Its unique shape is ideal for defects of the mandibular angle and for those defects that extend into the ramus or body (Brown Class I). This shape may obviate the need for an osteotomy. In addition, this flap should also be considered where a short linear segment is required in combination with a large soft tissue defect. Yoo et al⁵⁰, describe the successful reconstruction of a condyle-ramus unit where the fibrous tip of scapula was used to replace the condylar head. As discussed previously for midface reconstruction, the pedicle length is advantageous, particularly in cases where the contra-lateral or inferior neck is used for anastomosis. However the TSCAP is not suitable for mandibular defects greater than 10cm in length, and in those requiring osteotomies.

For mandibular reconstruction, it is our practice to reserve the Tip of Scapula for predominantly Class I, and some Class II defects that require either contralateral neck access or ipsilateral transverse cervical vessels in patients not suitable for fibula flap harvest (e.g. peripheral vascular disease). In

addition to this, if there is an associated complex soft tissue defect(s) then the use of the scapula tip is favoured above all other flaps⁵⁴.

Donor Site Morbidity

While some of the advantages of the Thoracodorsal system of free flaps have been described in this paper, it is not without its problems. Harvesting these flaps simultaneously with tumour ablation is difficult, and consequent patient repositioning inevitably results in increased operative time. Care must be taken to stabilise the contralateral shoulder while bringing the patient into the lateral decubitus position, as injury to the brachial plexus can occur⁶⁷.

However, it seems that relative to other donor sites, there is less morbidity. Bach et al⁴² compared the cosmetic results of radial forearm free flaps with those of Thoracodorsal artery perforator flaps (TDAP) using three scar-scoring systems, and found greater patient satisfaction with the TDAP scar. In addition, they demonstrated minimal donor site morbidity associated with the TDAP donor site. Of note, they also state that while flap-harvesting time was greater for the TDAP flap, it was never longer than the cancer resection time. This is relevant, given that they harvest the TDAP in a supine position concurrent with tumour resection.

We have found that, as a result of taking *terres major* with tip of scapula flaps, a discreet hollowing/deformity of the skin can result. This, however, does not seem to be of concern to patients.

Hamdi et al⁶⁸, looked at shoulder function post TDAP flap harvesting and demonstrated that sparing of the motor nerve to *Latissimus Dorsi* preserves muscle strength and a full range of shoulder movement. In addition, they report that the absence of dead space in the axilla, and preservation of *Latissimus Dorsi* function are associated with decreased incidence of seroma.

Miles and Gilbert⁵⁵, in a retrospective review of 39 post-maxillectomy patients reconstructed with scapular tip free flaps, reported that mild post-operative seroma was a common occurrence, but that less than 8% required intervention. They stated further that shoulder morbidity was not a significant problem. Chepeha et al⁴⁹ reported on 21 patients reconstructed with a scapular tip free flap, and found that donor site complication rates were low. Two of their 21 patients had minor early complications (haematoma, and wound breakdown). Again, they found no significant shoulder dysfunction.

Shoulder morbidity may be minimised by suturing the remaining serratus anterior to the remaining scapula in combination with fastidious post-operative physiotherapy^{19, 69}. Ferrari et al⁷⁰, in a paper looking solely at donor site morbidity, found a very low rate of shoulder morbidity, which did not interfere with activities of daily living, in patients with scapular tip free flaps. In addition, they reported a mean time to ambulation of 2.7 days. Other authors, albeit reporting on morbidity following parascapular and scapular free flap harvesting, described similar results. Roll et al⁷¹, in a series of 20 patients reconstructed with a parascapular flap, reported that 3 had limited shoulder function post-operatively. Gibber et al⁴⁷, in their previously mentioned retrospective review of 105 patients, reported no post-operative shoulder dysfunction.

Anecdotally, we have found that there is less morbidity associated with flaps harvested from the subscapular system relative to, for example, the iliac crest or fibula. This is significant in elderly patients for whom early mobilisation is hugely important. In our practice this, along with the

aforementioned benefit of protection from atherosclerosis, means we often choose this flap in elderly or frail patients.

Discussion

There is no doubt that the advent of microvascular free tissue transfer has allowed the predictable reconstruction of large and complex head and neck defects. There are numerous donor sites available, each with their own advantages and disadvantages. The thoracodorsal system has a range of hard and soft tissue flaps that can be raised independently or as chimeric flaps with independently mobile components. These features in combination with a substantial pedicle length, that predictably negates the need for interpositional grafts or multiple flaps, are the main advantages of the thoracodorsal system. In addition, for patients in whom early post-operative mobility is critical, the thoracodorsal system has obvious advantages relative to the iliac crest⁷² and fibula. Further, the thoracodorsal system is particularly useful in patients for whom peripheral vascular disease precludes the use of other donor sites⁷. Finally it is easier, relative to other donor sites, to camouflage the resultant scar⁷³.

In conclusion, free flaps based on the thoracodorsal system are suited to complex defects requiring extensive soft and hard tissue reconstruction as well as those requiring long pedicle length (e.g. vessel deplete necks). In these complex cases, the use of thoracodorsal system-based free flaps may negate the need for interpositional vein grafts and/or the harvesting a second flap, both of which are factors that add a layer of complexity and are associated with increased potential for complications.

References:

- ¹ Rowsell AR, Davies DM, Eisenberg N, Taylor GI. The anatomy of the subscapular-thoracodorsal arterial system: study of 100 cadaver dissections. *Br J Plast Surg* 1984;37:574–576.
- ² Bartlett SP, May JW, Yaremchuk MJ: The latissimus dorsi muscle. A fresh cadaver study of the primary neurovascular pedicle. *Plast Reconstr Surg* 1981;67:631.
- ³ Urken M, Cheney M, Futran N et al. *Atlas of Regional and Free Flaps for Head and Neck Reconstruction: Flap Harvest and Insetting*. 2nd Edition, Lippincott Williams & Wilkin, 2012.
- ⁴ DeGaris C, Swartz W: The axillary artery in white and Negro stocks. *Am J Anat* 1928;4:353.
- ⁵ Kawamura K, Yajima H, Kobata Y, Shigematsu K, Takakura Y. Anatomy of Y-shaped configurations in the subscapular arterial system and clinical application to harvesting flow-through flaps. *Plast Reconstr Surg* 2005; 116:1082–1089.
- ⁶ Coeugnet E, Harchaoui A, Malka G, Pellerin P, Danino A. The osteomuscular latissimus dorsi scapula flap: anatomical study and 3 cases. *Ann Chir Plast Esthet* 2007;52:108-13.
- ⁷ Barry CP, Brown J, Hanlon R, Shaw R. Tailored approach to oromandibular reconstruction in patients with compromised lower limb vessels. *Head Neck*. 2017 May;39(5):916-920.
- ⁸ Halim AS, Wan Z: Anomalous arterial supply to the muscles in a combined latissimus dorsi and serratus anterior flap. *ClinAnat* 2004;17:358-359.
- ⁹ Tansini I. Sopra il mio nuovo processo di amputazione della mammella. *Gazz Med Ital* 1906;57:141.
- ¹⁰ Quillen CG, Shearin JC, Georgiade NG: Use of the latissimus dorsi myocutaneous island flap for reconstruction in the head and neck area, *Plast Reconstr Surg* 62:113-117, 1978.
- ¹¹ Maxwell GP, Stueber K, Hoopes JE: A free latissimus dorsi myocutaneous flap, *Plast Reconstr Surg* 62:462-466, 1978.
- ¹² Ong HS, Ji T, Zhang CP. The pedicled latissimus dorsi myocutaneous flap in head and neck reconstruction. *Oral Maxillofac Surg Clin North Am*. 2014 Aug;26(3):427-34.
- ¹³ Aviv JE, Urken ML, Vickery C, et al. The combined latissimus dorsi-scapular free flap in head and neck reconstruction. *Arch Otolaryngol Head Neck Surg* 1991;117:1242–50.
- ¹⁴ Tobin GR, Schusterman M, Peterson GH, Nichols G, Bland KI. The intramuscular neurovascular anatomy of the latissimus dorsi muscle: the basis for splitting the flap. *Plast Reconstr Surg*. 1981 May;67(5):637-41.
- ¹⁵ Taylor GI, Palmer J: The vascular territories (angiosomes) of the body; experimental study and clinical applications. *Br J Plast Surg* 1987;40: 113.
- ¹⁶ Girod A, Boissonnet H, Jouffroy T, Rodrigue J: Latissimus dorsi free flap reconstruction of anterior skull base defects. *J Craniomaxillofac Surg*, 2012, Feb;40(2):177-9.
- ¹⁷ Li BH, Jung HJ, Choi SW, Kim SM, Kim MJ, Lee JH. Latissimus dorsi (LD) free flap and reconstruction plate used for extensive maxillo-mandibular reconstruction after tumour ablation. *J Craniomaxillofac Surg*. 2012 Dec;40(8):e293-300.
- ¹⁸ Pohlenz P, Klatt J, Schon G, et al. Microvascular free flaps in head and neck surgery: complications and outcome of 1000 flaps. *Int J Oral Maxillofac Surg* 2012;41:739–743
- ¹⁹ Baker SR. Closure of large orbital-maxillary defects with free latissimus dorsi myocutaneous flaps. *Head Neck Surg*. 1984; 6:828–835.
- ²⁰ Maruyama Y, Iwahira Y, Hashimura C et al. One stage total cheek reconstruction with double folded extended latissimus dorsi musculocutaneous flap. *Acta Chir Plast*. 1986; 28:159–166.
- ²¹ Robson MC, Zachary LS, Schmidt DR et al. Reconstruction of large cranial defects in the presence of heavy radiation damage and infection utilizing tissue transferred by microvascular anastomoses. *Plast Reconstr Surg*, 1989; 83:438–442.
- ²² Girod A, Boissonnet H, Jouffroy T, Rodrigue J: Latissimus dorsi free flap reconstruction of anterior skull base defects. *J Craniomaxillofac Surg*, 2012, Feb;40(2):177-9.
- ²³ Hierner R, van Loon J, Goffin J, et al. Free latissimus dorsi flap transfer for subtotal scalp and cranium defect reconstruction: report of 7 cases. *Microsurgery* 2007;27:425–428
- ²⁴ Lee KT, Mun GH. A systematic review of functional donor-site morbidity after latissimus dorsi muscle transfer. *Plast Reconstr Surg*. 2014 Aug;134(2):303-14.
- ²⁵ Hayden RE, Kirby SD, Deschler DG. Technical modifications of the latissimus dorsi pedicled flap to increase versatility and viability. *Laryngoscope* 2000; 110:352–7.
- ²⁶ Barton FE Jr, Spicer TE, Byrd HS. Head and neck reconstruction with the latissimus dorsi myocutaneous flap: anatomic observations and report of 60 cases. *Plast Reconstr Surg* 1983;71: 199–204.
- ²⁷ Chowdhury CR, McLean NR, Harrop-Griffiths K, et al. The repair of defects in the head and neck region with the latissimus dorsi myocutaneous flap. *J Laryngol Otol* 1988;102:1127–32.
- ²⁸ Haughey BH, Fredrickson JM. The latissimus dorsi donor site. Current use in head and neck reconstruction. *Arch Otolaryngol Head Neck Surg* 1991; 117:1129–34.
- ²⁹ Germann G, Ohlbauer M. Latissimus dorsi flap. In: Wei FC, Mardini S, editors. *Flap and reconstructive surgery*. Philadelphia: Saunders Inc; 2009. p. 287–319.
- ³⁰ Sabatier RE, Bakamjian VY. Transaxillary Latissimus dorsi flap reconstruction in head and neck cancer. Limitations and refinements in 56 cases. *Am J Surg* 1985;150:427–34.

-
- ³¹ Angrigiani C, Grilli D, Siebert J. Latissimus dorsi musculocutaneous flap without muscle. *Plast Reconstr Surg*. 1995 Dec; 96 (7):1608-14.
- ³² Spinelli HM, Fink JA, The latissimus dorsi perforator-based fasciocutaneous flap. Muzaffar AR. *Ann Plast Surg*. 1996 Nov;37(5):500-6.
- ³³ Yang LC, Wang XC, Bentz ML, Fang BR, Xiong X, Li XF, Lu Q, Wu Q, Wang RF, Feng W, He JY, Wang YY. Clinical application of the thoracodorsal artery perforator flaps. *J Plast Reconstr Aesthet Surg*. 2013 Feb;66(2):193-200.
- ³⁴ Heitmann C, Guerra A, Metzinger SW, et al. The thoracodorsal artery perforator flap: anatomic basis and clinical application. *Ann Plast Surg* 2003;51:23—9.
- ³⁵ Mun GH, Lee SJ, Jeon BJ. Perforator topography of the thoracodorsal artery perforator flap. *Plast Reconstr Surg* 2008;121:497—504.
- ³⁶ Thomas BP, Geddes CR, Tang M, et al. The vascular basis of the thoracodorsal artery perforator flap. *Plast Reconstr Surg* 2005;116:818—22.
- ³⁷ Schaverien M, Wong C, Bailey S, Saint-Cyr M. Thoracodorsal artery perforator flap and Latissimus dorsi myocutaneous flap--anatomical study of the constant skin paddle perforator locations. *J Plast Reconstr Aesthet Surg*. 2010 Dec;63(12):2123-7.
- ³⁸ Thomas BP, Geddes CR, Tang M, et al. The vascular basis of the thoracodorsal artery perforator flap. *Plast Reconstr Surg* 2005;116:818—22.
- ³⁹ Guerra AB, Metzinger SE, Lund KM, et al. The thoracodorsal artery perforator flap: clinical experience and anatomic study with emphasis on harvest techniques. *Plast Reconstr Surg* 2004;114:32—41.
- ⁴⁰ Hamdi M, Van Landuyt K, Hijawi JB, Roche N, Blondeel P, Monstrey S. Surgical technique in pedicled thoracodorsal artery perforator flaps: a clinical experience with 99 patients. *Plast Reconstr Surg*. 2008 May;121(5):1632-41.
- ⁴¹ Lee SH, Mun GH. Transverse thoracodorsal artery perforator flaps: experience with 31 free flaps. *J Plast Reconstr Aesthet Surg*. 2008;61(4):372-9.
- ⁴² Bach CA, Dreyfus JF, Wagner I, Lachiver X, Guth A, Chabolle F. Comparison of radial forearm flap and thoracodorsal artery perforator flap donor site morbidity for reconstruction of oral and oropharyngeal defects in head and neck cancer. *Eur Ann Otorhinolaryngol Head Neck Dis*. 2015 Sep;132(4):185-9.
- ⁴³ Swartz WM, Banis JC, Newton ED, Ramasastry SS, Jones NF, Acland R. The osteocutaneous scapular flap for mandibular and maxillary reconstruction. *Plast Reconstr Surg* 1986;77:530—545.
- ⁴⁴ Deraemaeker R, Thienen C, Lejour M, Dor P. The serratus anterior scapular free flap: a new osteomuscular unit for reconstruction after radical head and neck surgery. *Proceedings of the Second International Conference on Head and Neck Cancer*; 1988. pp 133—137.
- ⁴⁵ Coleman JJ III, Sultan MR. The bipedicled osteocutaneous scapula flap: a new subscapular system free flap. *Plast Reconstr Surg* 1991;87:682—692.
- ⁴⁶ Patnaik V, Kalsey G, Singla Rajan K. Branching pattern of brachial artery— a morphological study. *J Anat Soc India* 2002;51:176—186.
- ⁴⁷ Gibber MJ, Clain JB, Jacobson AS, Buchbinder D, Scherl S, Zevallos JP, Mehra S, Urken ML. Subscapular system of flaps: An 8-year experience with 105 patients. *Head Neck*. 2015 Aug;37(8):1200-6.
- ⁴⁸ Seitz A, Papp S, Papp C, Maurer H. The anatomy of the angular branch of the thoracodorsal artery. *Cells Tissues Organs*. 1999;164(4):227-236
- ⁴⁹ Chepeha DB, Khariwala SS, Chanowski EJ, et al. Thoracodorsal artery scapular tip autogenous transplant: vascularized bone with a long pedicle and flexible soft tissue. *Arch Otolaryngol Head Neck Surg*. 2010 Oct;136(10):958-64.
- ⁵⁰ Yoo J, Dowthwaite SA, Fung K, Franklin J, Nichols A. A new angle to mandibular reconstruction: the scapular tip free flap. *Head Neck*. 2013 Jul;35(7):980-6.
- ⁵¹ Seneviratne S, Duong C, Taylor GI. The angular branch of the thoracodorsal artery and its blood supply to the inferior angle of the scapula: an anatomical study. *Plast Reconstr Surg* 1999; 104:85-8.
- ⁵² Brown JS, Barry C, Ho M, Shaw R. A new classification for mandibular defects after oncological resection. *Lancet Oncol* 2016; 17, e23-30.
- ⁵³ Dowthwaite SA, Theurer J, Belzile M, Fung K, Franklin J, Nichols A, Yoo J. Comparison of fibular and scapular osseous free flaps for oromandibular reconstruction: a patient-centered approach to flap selection. *JAMA Otolaryngol Head Neck Surg*. 2013 Mar;139(3):285-92.
- ⁵⁴ Ho MW, Brown JS, Shaw RJ. Refining the indications for scapula tip in mandibular reconstruction. *Int J Oral Maxillofac Surg*. 2017 Jun; 46(6):712-715.
- ⁵⁵ Miles BA, Gilbert RW. Maxillary reconstruction with the scapular angle osteomyogenous free flap. *Arch Otolaryngol Head Neck Surg*. 2011 Nov;137(11):1130-5.
- ⁵⁶ Shaw RJ, Ho MW, Brown JS. Thoracodorsal artery perforator - scapular flap in oromandibular reconstruction with associated large facial skin defects. *Br J Oral Maxillofac Surg*. 2015 Jul;53(6):569-71.
- ⁵⁷ Wei FC, Yazar S, Lin CH, Cheng MH, Tsao CK, Chiang YC. Double free flaps in head and neck reconstruction. *Clin Plast Surg*. 2005 Jul;32(3):303-8.
- ⁵⁸ Uglesic V, Virag M, Varga S, Knezevic P, Milenovic A. Reconstruction following radical maxillectomy with flaps supplied by the subscapular artery. *J Craniomaxillofac Surg*. 2000;28(3):153-160.
- ⁵⁹ Kakibuchi M, Fujikawa M, Hosokawa K, et al. Functional reconstruction of maxilla with free latissimus dorsi-scapular osteomusculocutaneous flap. *Plast Reconstr Surg*. 2002;109(4):1238-1245.
- ⁶⁰ Jones JW. Reconstruction of a complex hemifacial deformity with multiple simultaneous free-flap transfers:

case report. *J Reconstr Microsurg.* 2003;19 (2):73-78.

⁶¹ L'Heureux-Lebeau B1, Odobescu A, Harris PG, Guertin L, Danino AM. Chimaeric subscapular system free flap for complex oro-facial defects. *J Plast Reconstr Aesthet Surg.* 2013 Jul;66(7):900-5.

⁶² Brown J, Bekiroglu F, Shaw R. Indications for the scapular flap in reconstructions of the head and neck. *Br J Oral Maxillofac Surg.* 2010 Jul;48(5):331-7.

⁶³ Brown JS, Rogers SN, McNally DN, Boyle M. A modified classification for the maxillectomy defect. *Head Neck.* 2000 Jan;22(1):17-26.

⁶⁴ Pagedar NA, Gilbert RW, Chan H, Daly MJ, Irish JC, Siewerdsen JH. Maxillary reconstruction using the scapular tip free flap: a radiologic comparison of 3D morphology. *Head Neck.* 2012 Oct;34(10):1377-82.

⁶⁵ Brown JS, Lowe D, Kanatas A, Schache A. Mandibular reconstruction with vascularised bone flaps: a systematic review over 25 years. *Br J Oral Maxillofac Surg.* 2017 Feb;55(2):113-126.

⁶⁶ Schultes G, Gaggl A, K€archer H. Stability of dental implants in microvascular osseous transplants. *Plast Reconstr Surg* 2002;109:916–921.

⁶⁷ Maxwell GP, Manson PN, Hoopes JE. Experience with thirteen Latissimus dorsi myocutaneous free flaps. *Plast Reconstr Surg* , 1979; 64:641

⁶⁸ Hamdi M, Decorte T, Demuyneck M, et al. Shoulder function after harvesting a thoracodorsal artery perforator flap. *Plast Reconstr Surg* 2008;122:1111—7.

⁶⁹ Clark JR, Vesely M, Gilbert R. Scapular angle osteomyogenous flap in postmaxillectomy reconstruction: defect, reconstruction, shoulder function, and harvest technique. *Head Neck.* 2008;30(1):10-20.

⁷⁰ Ferrari S, Ferri A, Bianchi B, Varazzani A, Perlangeli G, Sesenna E. Donor site morbidity after scapular tip free flaps in head-and-neck reconstruction. *Microsurgery.* 2015 Sep;35(6):447-50.

⁷¹ Roll C, Prantl L, Ferer D, Nerlich M, Kinner B. Functional donor-site morbidity following (osteo-) fasciocutaneous parascapular flap transfer. *Ann Plast Surg* 2007; 59:410-414.

⁷² Howard MA, Davison SP, Piorkowski JR. Iliac crest donor defect: avoiding the limp. *Plast Reconstr Surg.* 2004 Feb;113(2):659-63.

⁷³ Brown JS, Thomas S, Chakrabati A, Lowe D, Rogers SN. Patient preference in placement of the donor-site scar in head and neck cancer reconstruction. *Plast Reconstr Surg.* 2008 Jul;122(1):20e-22e

Figure 1. Summary of critical aspects of thoracodorsal arterial anatomy relevant to the reconstructive surgeon.

Figure 2. The Latissimus Dorsi and tip of Scapula (LD-TSCAP) chimeric flap. Note the large volume of soft-tissue that is independently mobile from the bone.

Figure 3. Diagram to show where the greatest concentration of perforators >0.5 mm in diameter can be found (dashed yellow circle) when planning the thoracodorsal artery perforator flap (TDAP). The measurements are derived from Schaverien et al³⁷. Note that the TDAP flap is predominantly centred over the vertical branch of the thoracodorsal artery.

Figure 4. Intra-operative photograph demonstrating the thoracodorsal artery perforator (TDAP) flap being raised. Note that the large skin paddle and long length of pedicle that can be achieved. Also note the extensive muscular dissection required to raise this flap.


Figure 5. A chimeric flap incorporating the thoracodorsal artery perforator flap and composite parascapular flap. The point at which angular branch takes off from the thoracodorsal artery is labeled to demonstrate that the tip of scapula flap can also be incorporated should the ablative defect require two independent bone flaps (e.g. in combined maxillary and mandibular reconstruction).

Figure 6. Intra-operative series demonstrating the use of the chimeric Latissimus Dorsi and tip of Scapula (LD-TSCAP) flap in complex post radiotherapy mandibular reconstruction. A) Complex through-and-through defect involving skin, bone and oral mucosa. B) Contralateral neck used for anastomotic vessel access. The ruler demonstrates the length of pedicle required. C) The harvested LD-TSCAP flap with independently mobile soft and hard tissue components. D) The flap inset to the defect.

Figure 7. The Tip of Scapula (TSCAP) flap used for midface reconstruction. In this case the bone was positioned cranio-caudally (vertically) to reconstruct a Brown Class V Orbito-maxillary defect. The bone supports both the overlying soft-tissue as well as the orbital contents.

Figure

[Click here to download high resolution image](#)




Figure

[Click here to download high resolution image](#)



Figure

[Click here to download high resolution image](#)



Figure

[Click here to download high resolution image](#)



Figure

[Click here to download high resolution image](#)




Figure
[Click here to download high resolution image](#)



Figure

[Click here to download high resolution image](#)

