

# A multilinear algebra proof of the Cauchy-Binet formula and a multilinear version of Parseval's identity

TAKIS KONSTANTOPOULOS

1 May 2013

## Abstract

We give a short proof of the Cauchy-Binet determinantal formula using multilinear algebra by first generalizing it to an identity *not* involving determinants. By extending the formula to abstract Hilbert spaces we obtain, as a corollary, a generalization of the classical Parseval identity.

*Keywords and phrases.* CAUCHY-BINET THEOREM, DETERMINANT, MATRIX IDENTITIES, HILBERT SPACE, PARSEVAL'S IDENTITY, MULTILINEAR ALGEBRA, EXTERIOR PRODUCTS, PROJECTIONS, PYTHAGOREAN THEOREM, FOCK SPACE

*AMS 2000 subject classifications.* Primary 15A15,15A69; secondary 15A24,46C05

## 1 Introduction and overview

The classical Cauchy-Binet formula states that if  $A, B$  are two matrices over  $\mathbb{R}$  (or any field) of sizes  $n \times N$ ,  $N \times n$ , respectively, with  $n \leq N$ , then

$$\det(AB) = \sum_{\sigma} \det(A_{\sigma}) \det(B^{\sigma}) \quad (1)$$

where the sum is taken over all  $\sigma = (\sigma_1 < \sigma_2 < \dots < \sigma_n)$ , with  $\sigma_i \in \{1, \dots, N\}$ , and where  $A_{\sigma}$  (respectively  $B^{\sigma}$ ) is the  $n \times n$  submatrix of  $A$  (respectively submatrix of  $B$ ) obtained by deleting all columns (respectively all rows) except those with indices in  $\sigma$ .

There are many proofs of this formula, each telling its own story, explaining the formula from a different point of view. The most direct way of proving the formula is by writing down the determinant as a sum over permutations and performing algebraic manipulations. This is the approach taken in many linear algebra books; see, e.g., Marcus and Minc [8, Theorem 6.1, p. 128] and Gohberg *et al.* [7, Theorem A.2.1, p. 651]. A probabilistic interpretation and proof of the formula (which starts by using the formula for a determinant) is also available [4, 5]. On the other hand, there are many combinatorial proofs. Suffice, perhaps, to refer to

the one chosen to be included in the “Proofs from The Book” [2] by Aigner and Ziegler. This is a nice proof (after all, it is a proof from The Book) based on the beautiful Gessel-Vienot lemma which states that, in a finite weighted acyclic directed graph, the determinant of the path matrix between two sets of vertices of cardinality  $n$  each equals a sum over all possible vertex-disjoint path systems; see [2, Chap. 29, p. 196] and [1] for details. Another very simple proof appears in the recent book by Terence Tao [13, p. 298]) on random matrices. This proof is based on a relation between the characteristic polynomials of  $AB$  and  $BA$ .

On the other hand, it is well-known that the Cauchy-Binet formula is a generalization of the Pythagorean theorem. Indeed, let  $A$  be a  $n \times N$  real matrix,  $n \leq N$ , and take  $B = A^T$ , the transpose of  $A$ . Since  $B^\sigma = (A^T)^\sigma = (A_\sigma)^T$ , the formula gives

$$\det(AA^T) = \sum_{\sigma} \det(A_\sigma)^2,$$

which can be interpreted geometrically as follows: The parallelotope in  $\mathbb{R}^N$  generated by the  $n$  row vectors of  $A$  has  $n$ -dimensional Lebesgue measure  $\sqrt{\det(AA^T)}$ . Therefore the formula says that the square of the  $n$ -dimensional measure of an  $n$ -dimensional parallelotope, embedded in a higher-dimensional Euclidean space, equals the sum of the squares of the measures of its projections onto all possible  $n$ -dimensional coordinate hyperplanes. If  $n = 1$  this reduces to the Pythagorean theorem.

The goal of this short article is to give a proof of the Cauchy-Binet formula which is as simple as possible, from an algebraic-geometric viewpoint. If  $n = 1$ , the Cauchy-Binet formula is a triviality: it states that the inner product of two  $N$ -dimensional vectors equals the sum of the products of their components:

$$(a_1 \dots, a_N) \cdot (b_1, \dots, b_N)^T = \sum_{\sigma=1}^N a_\sigma b_\sigma.$$

There is no need to take determinants here, because both sides involve  $1 \times 1$  matrices, i.e., real numbers. What we show is that the general case, when  $n \geq 1$ , is the same, but on bigger vector spaces. In Section 2 we give an account of the ingredients we need, and, in Section 3, we state and prove the main formula (Theorem 1) without determinants and in a more general setup; a corollary of it is the classical Cauchy-Binet formula. Then, in Section 4, we see that the formula can be extended to a Hilbert space, giving a generalization of the classical Parseval identity. We conclude with a few bibliographic remarks.

## 2 The main ingredients

The main theorem, Theorem 1 below, requires two ingredients.

(i) The first is the notion of the determinant of a linear transformation  $F : X \longrightarrow X$  on a vector space  $X$  of dimension  $d$ . The dimension of the linear space  $\bigwedge^m X$  of alternating  $m$ -linear maps  $\omega : X^m \longrightarrow \mathbb{R}$  is  $\binom{d}{m}$ . For each  $m$ , the  $m$ -th level dual  $F^* : \bigwedge^m X \longrightarrow \bigwedge^m X$

of  $F$  is defined by

$$F^*\omega[x_1, \dots, x_m] := \omega[Fx_1, \dots, Fx_m]. \quad (2)$$

See, e.g., [12]. (Duals obey the standard composition rules:  $(GF)^* = F^*G^*$ .) Since  $\bigwedge^d X$  is 1-dimensional, the  $d$ -th level dual  $F^*$  is multiplication by a constant. This constant is, by definition, the determinant of  $F$ :

$$F^*\omega = (\det F) \cdot \omega, \quad \omega \in \bigwedge^d X. \quad (3)$$

(ii) The second ingredient is very simple too. Let  $X, Y, Z$  be vector spaces, and  $F : X \rightarrow Y, G : Y \rightarrow Z$  linear maps. Suppose  $Y$  is the direct sum of  $Y_1, \dots, Y_K$ . Let  $P_i : Y \rightarrow Y_i, 1 \leq i \leq K$ , be the projections corresponding to this direct sum (so  $\text{id}_Y = P_1 + \dots + P_K$  is a partition of the identity on  $Y$ ), and let  $E_i : Y_i \rightarrow Y$  be the natural embedding of  $Y_i$  into  $Y$ . Then, clearly,

$$GF = \sum_{i=1}^K (GE_i)(P_iF). \quad (4)$$

See Diagram 1.

### 3 An abstract version of the Cauchy-Binet formula

Let  $U, V, W$  be finite-dimensional vector spaces of arbitrary dimensions, and let  $B : U \rightarrow V, A : V \rightarrow W$  be two linear maps. Fix  $n \in \mathbb{N}$  and consider the  $n$ -th level duals  $B^* : \bigwedge^n V \rightarrow \bigwedge^n U, A^* : \bigwedge^n W \rightarrow \bigwedge^n V$ . Let  $N$  be the dimension of  $V$  and let  $f_1, \dots, f_N$  be a basis for  $V$ . See Diagram 2. Denote by  $\mathcal{S}_n(N)$  the set of subsets of  $\{1, \dots, N\}$  of size  $n$ . For each  $\sigma \in \mathcal{S}_n(N)$ , let  $V_\sigma$  be the subspace of  $V$  spanned by  $\{f_i, i \in \sigma\}$  and consider the direct sum

$$V = V_\sigma \oplus V_{\bar{\sigma}}, \quad (5)$$

where  $\bar{\sigma} := \{1, \dots, N\} \setminus \sigma$ , letting

$$P_\sigma : V \rightarrow V_\sigma$$

be the projection of  $V$  onto  $V_\sigma$  along  $V_{\bar{\sigma}}$ , and

$$E_\sigma : V_\sigma \rightarrow V$$

the natural embedding of  $V_\sigma$  into  $V$ .

**Theorem 1.**

$$(AB)^* = \sum_{\sigma \in \mathcal{S}_n(N)} (P_\sigma B)^* (AE_\sigma)^*, \quad (6)$$

*Proof.* The  $\binom{N}{n}$ -dimensional space  $\bigwedge^n V$  is the direct sum of the 1-dimensional spaces  $\bigwedge^n V_\sigma$ , where  $\sigma$  ranges in  $\mathcal{S}_n(N)$ :

$$\bigwedge^n V = \bigoplus_{\sigma \in \mathcal{S}_n(N)} \bigwedge^n V_\sigma. \quad (7)$$

Let  $\mathcal{P}_\sigma : \bigwedge^n V \longrightarrow \bigwedge^n V_\sigma$  be projections corresponding to this direct sum, and let  $\mathcal{E}_\sigma : \bigwedge^n V_\sigma \longrightarrow \bigwedge^n V$  be natural embedding. Using (4) (with  $A^*$ ,  $B^*$  in place of  $F$ ,  $G$ , respectively, and  $K = \binom{N}{n}$ ), we have

$$B^*A^* = \sum_{\sigma \in \mathcal{S}_n(N)} (B^*\mathcal{E}_\sigma)(\mathcal{P}_\sigma A^*).$$

See Diagram 2. Since (see Lemma 1 below)

$$\mathcal{E}_\sigma = P_\sigma^*, \quad \mathcal{P}_\sigma = E_\sigma^*,$$

the theorem follows from the composition rules of the duals.  $\square$

**Lemma 1.**

$$\mathcal{P}_\sigma = E_\sigma^*, \quad \mathcal{E}_\sigma = P_\sigma^*.$$

*Proof.* We identify  $\mathcal{S}_n(N)$  with the set of strictly increasing sequences of length  $n$  with values in  $\{1, \dots, N\}$ . Thus, if  $\sigma$  is a subset of  $\{1, \dots, N\}$  we let  $(\sigma_1, \dots, \sigma_n)$  be a listing of its elements in increasing order. To prove the first equality it suffices to show that

$$\mathcal{P}_\sigma \omega[v_1, \dots, v_n] = \omega[E_\sigma v_1, \dots, E_\sigma v_n],$$

for all  $\omega \in \Lambda_n(V)$  and all  $v_1, \dots, v_n \in V_\sigma$ . But then  $E_\sigma v_i = v_i$  and, since  $V_\sigma$  is spanned by  $f_{\sigma_1}, \dots, f_{\sigma_n}$ , it suffices to show that

$$\mathcal{P}_\sigma \omega[f_{\sigma_{\pi(1)}}, \dots, f_{\sigma_{\pi(n)}}] = \omega[f_{\sigma_{\pi(1)}}, \dots, f_{\sigma_{\pi(n)}}],$$

where  $\pi$  is a permutation of  $\{1, \dots, n\}$ . Since  $\omega = \sum_{\tau \in \mathcal{S}_n(N)} \mathcal{P}_\tau \omega$  [this is the partition of the identity on  $\bigwedge^n V$  corresponding to (7)] we may replace  $\omega$  by  $\mathcal{P}_\tau \omega$  in the last display:

$$\mathcal{P}_\sigma \mathcal{P}_\tau \omega[f_{\sigma_{\pi(1)}}, \dots, f_{\sigma_{\pi(n)}}] = \mathcal{P}_\tau \omega[f_{\sigma_{\pi(1)}}, \dots, f_{\sigma_{\pi(n)}}].$$

But then, if  $\tau = \sigma$  the two sides are obviously equal, and if  $\tau \neq \sigma$  the left-hand side equals zero and  $\mathcal{P}_\tau \omega[f_{\sigma_{\pi(1)}}, \dots, f_{\sigma_{\pi(n)}}] = 0$ .

To prove the second equality it suffices to show that

$$\mathcal{E}_\sigma \omega[v_1, \dots, v_n] = \omega[P_\sigma v_1, \dots, P_\sigma v_n],$$

for all  $\omega \in \Lambda_n(V_\sigma)$  and all  $v_1, \dots, v_n \in V$ . But then  $\mathcal{E}_\sigma \omega = \omega$ . Since  $v_i = P_\sigma v_i + P_{\overline{\sigma}} v_i$  [corresponding to (5)], we have

$$\mathcal{E}_\sigma \omega[v_1, \dots, v_n] = \omega[P_\sigma v_1 + P_{\overline{\sigma}} v_1, \dots, P_\sigma v_n + P_{\overline{\sigma}} v_n].$$

Using the multilinearity of  $\omega$  we split the latter into  $2^n$  terms, all of which are zero except the one involving only  $P_\sigma v_i$  as arguments.  $\square$

Consider now the case where  $W = U$ . Moreover, take the number  $n$  in Theorem 1 to be equal to their common dimension. Assume  $n \leq N = \dim V$  to avoid trivialities. Then the linear maps  $(AB)^*$ ,  $(P_\sigma B)^*$ , and  $(AE_\sigma)^*$ , appearing in formula (6), are maps between

1-dimensional spaces. Since the spaces  $V_\sigma$  and  $U$  have common dimension  $n$ , we can identify them by means of a linear bijection

$$\varphi_\sigma : V_\sigma \longrightarrow U.$$

Then

$$(P_\sigma B)^*(AE_\sigma)^* = (P_\sigma B)^*\varphi_\sigma^*(\varphi_\sigma^{-1})^*(AE_\sigma)^* = (\varphi_\sigma P_\sigma B)^*(AE_\sigma\varphi_\sigma^{-1})^*,$$

and so

$$(AB)^* = \sum_{\sigma \in \mathcal{S}_n(N)} (\varphi_\sigma P_\sigma B)^*(AE_\sigma\varphi_\sigma^{-1})^*. \quad (8)$$

Since all three linear maps  $AB$ ,  $\varphi_\sigma P_\sigma B$ ,  $AE_\sigma\varphi_\sigma^{-1}$  are linear maps on the same 1-dimensional vector space  $U$ , it follows, from the definition of the determinant, that

$$\det(AB) = \sum_{\sigma \in \mathcal{S}_n(N)} \det(\varphi_\sigma P_\sigma B) \det(AE_\sigma\varphi_\sigma^{-1}). \quad (9)$$

(The role of  $\varphi_\sigma$  is to force all maps be on the same space, so we can talk about determinants.) In the case where  $U = \mathbb{R}^n$ ,  $V = \mathbb{R}^N$ , this proves the classical Cauchy-Binet formula (1). If  $N = n$ , then we have shown that the determinant of the product is the product of the determinants.

Therefore (1) follows from (9). The latter is a restatement of (8). But (8) is a special case of (6) because in (6) we allow  $U, V, W$  to be different with dimensions that may be distinct from  $n$ .

## 4 Multilinear Parseval's identity

We are now going to replace the middle space  $V$  of the previous setup by a separable Hilbert space  $H$  over the complex numbers  $\mathbb{C}$ , having inner product  $\langle x, y \rangle$ . Let  $f_1, f_2, \dots$  be an orthonormal basis for  $H$ . Let  $\bigwedge^n H$  be the collection of all *continuous* alternating multilinear functionals  $\omega : H^n \longrightarrow \mathbb{C}$ . In particular,  $\bigwedge^1 H = H^*$  is the Hilbert space dual of  $H$ . By the Riesz-Fischer theorem,  $f_1, f_2, \dots$  forms a basis for  $\bigwedge^1 H$  in the sense that every  $\omega \in \bigwedge^1 H$  can be uniquely written as  $\omega[x] = \sum_{\sigma=1}^{\infty} a_\sigma \langle f_\sigma, x \rangle$ , for  $a_\sigma \in \mathbb{C}$  such that  $\sum_{\sigma} |a_\sigma|^2 < \infty$ . More generally,  $\bigwedge^n H$  is a separable Hilbert space with orthonormal (with respect to a suitably defined inner product) basis

$$f_{\sigma_1} \wedge \dots \wedge f_{\sigma_n}, \quad \sigma = (\sigma_1, \dots, \sigma_n) \in \mathcal{S}_n(\mathbb{N}),$$

where  $\mathcal{S}_n(\mathbb{N})$  is the collection of all  $n$ -tuples  $(\sigma_1, \dots, \sigma_n)$  of positive integers such that  $\sigma_1 < \dots < \sigma_n$ . Recall that the wedge product satisfies, by definition,

$$(f_1 \wedge f_2)[x, y] = f_1[x]f_2[y] - f_1[y]f_2[x],$$

and, more generally,  $f_{\sigma_1} \wedge \dots \wedge f_{\sigma_n}$  is obtained by antisymmetrization of the tensor product of  $f_{\sigma_1}, \dots, f_{\sigma_n}$ . Incidentally, the direct sum of  $\bigoplus_{n=0}^{\infty} \bigwedge^n H$  (where  $\bigwedge^0 H := \mathbb{C}$ ) is the so-called

alternating Fock (or fermionic) space [11]. Wedge products can be defined, by linearity, between any finite number of elements of this space.

If  $H_1, H_2$  are two Hilbert spaces and  $F : H_1 \longrightarrow H_2$  is a continuous linear function then  $F^* : \bigwedge^n H_2 \longrightarrow \bigwedge^n H_1$  is defined as before—see (2)—and is, moreover, continuous.

**Theorem 2.** *Let  $H$  be a separable Hilbert space over  $\mathbb{C}$  with orthonormal basis  $f_1, f_2, \dots$ , and let  $n$  be a positive integer. For each  $\sigma \in \mathcal{S}_n(\mathbb{N})$ , let  $H_\sigma$  be the subspace spanned by  $f_{\sigma_1}, \dots, f_{\sigma_n}$ . Let  $E_\sigma : H_\sigma \longrightarrow H$  be the natural embedding of  $H_\sigma$  into  $H$  and  $P_\sigma : H \longrightarrow H_\sigma$  the orthogonal projection of  $H$  onto  $H_\sigma$ . If  $U, W$  are finite-dimensional vector spaces over  $\mathbb{C}$  and  $B : U \longrightarrow H, A : H \longrightarrow W$  continuous linear maps, then*

$$(AB)^* = \sum_{\sigma \in \mathcal{S}_n(\mathbb{N})} (P_\sigma B)^* (AE_\sigma)^*.$$

*If  $W = U$  with common dimension  $n$ , and if  $\varphi_\sigma : H_\sigma \longrightarrow U$  is any linear bijection, then*

$$(AB)^* = \sum_{\sigma \in \mathcal{S}_n(\mathbb{N})} (\varphi_\sigma P_\sigma B)^* (AE_\sigma \varphi_\sigma^{-1})^*.$$

*In particular,*

$$\det(AB) = \sum_{\sigma \in \mathcal{S}_n(\mathbb{N})} \det(\varphi_\sigma P_\sigma B) \det(AE_\sigma \varphi_\sigma^{-1}).$$

The proof of this theorem is exactly as in the finite-dimensional case. Infinite sums have to be understood in the Hilbert space sense.

Consider now  $H = L^2[0, 1]$  with inner product  $\langle x, y \rangle = \int_0^1 x(t) \overline{y(t)} dt$  and the standard orthonormal basis  $e_k(t) = \exp(i2\pi kt)$ ,  $k \in \mathbb{Z}$ , and let  $U = W = \mathbb{C}^n$ , for a given positive integer  $n$ . A continuous linear map  $A : L^2[0, 1] \longrightarrow \mathbb{C}^n$  is necessarily (Riesz representation theorem) of the form

$$Ax = (\langle x, a_1 \rangle, \dots, \langle x, a_n \rangle) = \left( \int_0^1 \overline{a_1(t)} x(t) dt, \dots, \int_0^1 \overline{a_n(t)} x(t) dt \right), \quad x \in L^2[0, 1],$$

where  $a_1, \dots, a_n \in L^2[0, 1]$ . A linear map  $B : \mathbb{C}^n \longrightarrow L^2[0, 1]$  is of the form

$$(Bu)(t) = u_1 b_1(t) + \dots + u_n b_n(t), \quad u \in \mathbb{C}^n,$$

where  $b_1, \dots, b_n \in L^2[0, 1]$ . Hence the  $jk$ -entry of the matrix of  $AB : \mathbb{C}^n \longrightarrow \mathbb{C}^n$ , with respect to the standard basis on  $\mathbb{C}^n$ , is given by

$$(AB)_{jk} = \int_0^1 \overline{a_j(t)} b_k(t) dt.$$

Consider now  $\sigma \in \mathcal{S}_n(\mathbb{Z})$ , i.e.,  $\sigma = (\sigma_1, \dots, \sigma_n) \in \mathbb{Z}^n$  with  $\sigma_1 < \dots < \sigma_n$ . (There is no difficulty in replacing  $\mathbb{N}$  in the above theorem by  $\mathbb{Z}$ .) Then  $H_\sigma$  is the subspace of  $L^2[0, 1]$  spanned by  $e_{\sigma_1}, \dots, e_{\sigma_n}$ . So the orthogonal projection  $P_\sigma : H \longrightarrow H_\sigma$  is given by

$$P_\sigma x = \widehat{x}(\sigma_1) e_{\sigma_1} + \dots + \widehat{x}(\sigma_n) e_{\sigma_n},$$

where

$$\widehat{x}(k) := \int_0^1 x(t) \exp(-i2\pi kt) dt, \quad k \in \mathbb{Z},$$

are the Fourier coefficients of  $x$ . Letting  $\varphi_\sigma : H_\sigma \rightarrow \mathbb{C}^n$  be the linear bijection that takes  $e_{\sigma_r}$  into the  $r$ -th standard basis vector of  $\mathbb{C}^n$ , for  $r = 1, \dots, n$ , we see that the  $jk$ -entry of the matrix of  $\varphi_\sigma P_\sigma B$  is

$$(\varphi_\sigma P_\sigma B)_{jk} = \widehat{b}_k(\sigma_j).$$

Arguing analogously, the  $jk$ -entry of the matrix of  $AE_\sigma \varphi_\sigma^{-1}$  is

$$(AE_\sigma \varphi_\sigma^{-1})_{jk} = \overline{\widehat{a}_j(\sigma_k)}.$$

Hence the last formula of Theorem 2 gives

$$\begin{aligned} \det_{1 \leq j, k \leq n} \int_0^1 \overline{a_j(t)} b_k(t) dt &= \sum_{\sigma \in \mathcal{S}_n(\mathbb{Z})} \det_{1 \leq j, k \leq n} [\overline{\widehat{a}_j(\sigma_k)}] \det_{1 \leq j, k \leq n} [\widehat{b}_j(\sigma_k)] \\ &= \frac{1}{n!} \sum_{\sigma_1 \in \mathbb{Z}} \cdots \sum_{\sigma_n \in \mathbb{Z}} \det_{1 \leq j, k \leq n} [\overline{\widehat{a}_j(\sigma_k)}] \det_{1 \leq j, k \leq n} [\widehat{b}_j(\sigma_k)], \end{aligned}$$

where the second equality follows from the fact that applying the permutation of  $(\sigma_1, \dots, \sigma_n)$  to both matrices will change the sign of both determinants simultaneously and the fact that repeated indices result into zero determinants. For  $n = 1$ , this is the standard Parseval identity.

Of course, there is nothing special with the Lebesgue measure. We can obtain formulas for any other  $L^2$  space or other separable Hilbert spaces.

## 5 Remarks

My motivation for this article was due to my desire to understand some elements of random matrix theory [3] and determinantal point processes [6]. In particular, the derivation of the ubiquitous Tracy-Widom probability distribution [3] involves several applications of Cauchy-Binet type formulas. When I looked at it first, a standard computational proof was not too satisfactory. I discovered that there are many proofs, which can be roughly classified into combinatorial and algebraic ones. The version presented in this short article was inspired by the simple observation that the Cauchy-Binet formula is a version of Pythagorean theorem: it is a version of the Pythagorean theorem on  $\bigwedge^n \mathbb{R}^N$ , with  $n \leq N$  (which is of course isomorphic to  $\mathbb{R}^{\binom{N}{n}}$ ).

Several years ago, Zeilberger [14] “complained” that, to most contemporary mathematicians, matrices and linear transformations are practically interchangeable notions and that the mainstream ‘Bourbakian’ establishment, with its profound disdain for the concrete, goes as far as to frown at the mere mention of the word ‘matrix’. He then explains how “to [him], as well as to other ‘dissidents’ called ‘combinatorialists’, a matrix has nothing whatsoever to do with that intimidating abstract concept called ‘a linear transformation between linear

vector spaces' ” and, by thinking of matrices as putting weights on a graph, he develops a combinatorial way of interpreting and proving fundamental results such as the Cayley-Hamilton theorem. The Cauchy-Binet formula has found a nice proof, in the Zeilberger sense, as a corollary of the Gessel-Vienot lemma. We also mention Zeng's proof [15] which also uses Zeilberger's methods.

In a sense then, what we have done here is in exactly the opposite of Zeilberger's spirit, because the proof presented uses nothing else but the concept of a linear map between vector spaces (and lots of definitions). Each point of view has its own merits in that, for instance, it leads to different kind of extensions. (Extensions to infinite matrices are not easy when the combinatorial point of view is adopted.)

We finally remark that there are generalizations of the Cauchy-Binet formula for the case where the matrices contain elements of a noncommutative ring [9]. We do not know how to extend the ideas above to this case.

## Acknowledgments

I thank Svante Janson [10] for pointing out reference [9] to me and for his comments on this article, and Richard Ehrenborg [4] for kindly making his notes available to me.

## References

- [1] Martin AIGNER. Lattice paths and determinants. *Comput. discrete mathematics*, 1-12, *Lect. Notes Comp. Sci.* **2122**, Springer, Berlin, 2001.
- [2] Martin AIGNER and Günter ZIEGLER. *Proofs from the Book*. Springer, 4th ed., New York, 2009.
- [3] Greg ANDERSON, Alice GUIONNET and Ofer ZEITOUNI. *An Introduction to Random Matrices*. Cambridge Univ. press, 2009.
- [4] Richard EHRENBORG. Personal communication, 2013.
- [5] Richard EHRENBORG and Gian-Carlo ROTA. Topics on polynomials. Lecture Notes, MIT, 1991.
- [6] John Ben HOUGH, Manjunath KRISHNAPUR, Yuval PERES and Balint VIRAG. *Zeros of Gaussian Analytic Functions and Determinantal Point Processes*. American Math. Society, 2009.
- [7] Israel GOHBERG, Peter LANCASTER and Leiba RODMAN. *Invariant Subspaces of Matrices with Applications*. SIAM Classics in Applied Mathematics **51**, Wiley, New York, 2986.
- [8] Marvin MARCUS, Henryk MINC. *Introduction to Linear Algebra*. MacMillan, New York, 1965; Reprinted by Dover, New York, 1965.
- [9] Sergio CARACCILO, Alan SOKAL and Andrea SPORTIELLO. Noncommutative determinants, Cauchy-Binet formulae, and Capelli-type identities. I. Generalizations of the Capelli and Turnbull identities. *Electron. J. Combin.* **16**, no. 1, Research Paper 103, 43 pp., 2009.
- [10] Svante JANSON. Personal communication, 2012.
- [11] Michael REED and Barry SIMON. *Methods of Modern Mathematical Physics*, Vol. II. Academic Press, 1975.



- [12] Michael SPIVAK. *Calculus on Manifolds*. Addison-Wesley, 1965.
- [13] Terence TAO. *Topics in Random Matrix Theory*:  
<http://terrytao.wordpress.com/books/topics-in-random-matrix-theory/>
- [14] Doron ZEILBERGER. A combinatorial approach to matrix algebra. *Discr. Math.* **56**, 61-72, 1985.
- [15] Jiang ZENG. A bijective proof of Muir's identity and the Cauchy-Binet formula. *Linear Alg. Appl.* **184**, 79-82, 1993.

TAKIS KONSTANTOPOULOS  
 DEPARTMENT OF MATHEMATICS  
 UPPSALA UNIVERSITY  
 751 06 UPPSALA  
 SWEDEN  
 takis@math.uu.se  
 www.math.uu.se/~takis




Diagram 1

$$Y = \bigoplus_{i=1}^K Y_i$$

$$GF = \sum_{i=1}^K (GE_i)(P_iF)$$




Diagram 2

$$\wedge^n V = \bigoplus_{\sigma \in \mathcal{S}_n(N)} \wedge^n V_\sigma$$

$$B^*A^* = \sum_{\sigma \in \mathcal{S}_n(N)} (B^*\mathcal{E}_\sigma)(\mathcal{P}_\sigma A^*)$$