

CLASSIFICATION: BIOLOGICAL SCIENCES (Neuroscience)

Ultrafast glutamate sensors resolve synaptic short-term plasticity

Nordine Helassa^{a1*}, Céline D. Dürst^{b*}, Catherine Coates^a, Silke Kerruth^a, Urwa Arif^a, Christian Schulze^b, J. Simon Wiegert^b, Michael Geeves^c, Thomas G. Oertner^b and Katalin Török^{a2}

^aMolecular and Clinical Sciences Research Institute, St George's, University of London, London SW17 0RE, UK; ^bInstitute for Synaptic Physiology, Center for Molecular Neurobiology Hamburg, Hamburg 20251, Germany; ^cSchool of Biosciences, University of Kent, Canterbury CT2 7NZ, UK

*These authors contributed equally to this work.

¹Current address: Department of Cellular and Molecular Physiology, Institute of Translational Medicine, University of Liverpool, Crown Street, Liverpool L69 3BX, UK.

²To whom correspondence should be addressed: Dr Katalin Török, Molecular and Clinical Sciences Research Institute, St George's, University of London, London SW17 0RE, UK, Telephone: +442087255832; E-mail: ktorok@sgul.ac.uk

Keywords: glutamate, synaptic transmission, post-tetanic potentiation, hippocampus, two-photon imaging, short-term plasticity

ABSTRACT

Glutamatergic synapses display a rich repertoire of plasticity mechanisms on many different time scales, involving dynamic changes in the efficacy of transmitter release as well as changes in the number and function of postsynaptic glutamate receptors. The genetically encoded glutamate sensor iGluSnFR enables visualization of glutamate release from presynaptic terminals at frequencies up to ~10 Hz. However, to resolve glutamate dynamics during high frequency bursts, faster indicators are required. Here we report the development of fast (iGlu_f) and ultrafast (iGlu_u) variants with comparable brightness, but increased K_d for glutamate (137 μ M and 600 μ M, respectively). Compared to iGluSnFR, iGlu_u has a 6-fold faster dissociation rate *in vitro* and 5-fold faster kinetics in synapses. Fitting a three-state model to kinetic data, we identify the large conformational change after glutamate binding as the rate-limiting step. In rat hippocampal slice culture stimulated at 100 Hz, we find that iGlu_u is sufficiently fast to resolve individual glutamate release events, revealing that glutamate is rapidly cleared from the synaptic cleft. Depression of iGlu_u responses during 100 Hz trains correlates with depression of postsynaptic EPSPs, indicating that depression during high frequency stimulation is purely presynaptic in origin. At individual boutons, the recovery from depression could be predicted from the amount of glutamate released on the second pulse (paired pulse facilitation/depression), demonstrating differential frequency-dependent filtering of spike trains at Schaffer collateral boutons.

Significance Statement

Excitatory synapses convert presynaptic action potentials into chemical signals that are sensed by postsynaptic glutamate receptors. To eavesdrop on synaptic transmission, genetically encoded fluorescent sensors for glutamate have been developed. However, even the best available sensors lag behind the very fast glutamate dynamics in the synaptic cleft. Here we report the development of an ultrafast genetically encoded glutamate sensor, iGlu_u, which allowed us to image glutamate clearance and synaptic depression during 100 Hz spike trains. We found that only boutons showing paired-pulse facilitation were able to rapidly recover from depression. Thus, presynaptic boutons act as frequency-specific filters to transmit select features of the spike train to specific postsynaptic cells.

INTRODUCTION

The efficacy of synaptic transmission is not constant, but changes dynamically during high-frequency activity. In terms of information processing, different forms of short-term plasticity act as specific frequency filters: Facilitating synapses are most effective during high frequency bursts, while depressing synapses preferentially transmit isolated spikes preceded by silent periods (Markram et al., 1998). Mechanistically, a number of pre- and postsynaptic parameters change dynamically during high frequency activity, e.g. the number of readily releasable vesicles, presynaptic Ca^{2+} dynamics, and the properties of postsynaptic receptors, which may be altered by Ca^{2+} -activated enzymes (Regehr, 2012; Jenkins and Traynelis, 2012).

Electrophysiological analysis of short-term plasticity, by monitoring postsynaptic responses, is complicated by the fact that neurons are often connected by more than one synapse. In addition, it is not straightforward to distinguish between pre- and postsynaptic plasticity mechanisms. Directly measuring glutamate concentrations inside the synaptic cleft during high-frequency activity would allow isolating the dynamics of the vesicle release machinery from potential changes in glutamate receptor properties (e.g. desensitization,

phosphorylation, lateral diffusion). Early fluorescent glutamate sensors, constructed by chemical labelling of the fused glutamate binding lobes of ionotropic glutamate receptor GluA2 (termed S1S2) (Best and Török, 2005; Chen and Gouaux, 1997; Kuusinen et al., 1995) and later of the bacterial periplasmic glutamate/aspartate binding protein (GluBP) (de Lorimier et al., 2002; Hu et al., 2008), were not suitable for quantitative single-synapse experiments due to their low dynamic range. Genetically encoded FRET-based fluorescent glutamate sensors e.g. FLIPE, GluSnFR and SuperGluSnFR (**Fig. 1a**) have relatively low FRET efficiency, since glutamate binding causes only a small conformational change in GluBP (Hires et al., 2008; Okumoto et al., 2005; Tsien, 2005). A breakthrough in visualizing glutamate release in intact tissue was achieved with iGluSnFR, a single-fluorophore glutamate sensor (Marvin et al., 2013). Following the concept developed for the GCaMP family of genetically encoded Ca^{2+} sensors (Nakai et al., 2001), iGluSnFR was constructed from circularly permuted (cp) EGFP (Baird et al., 1999) inserted into the GluBP sequence, creating a large fragment iGlu_N (residues 1-253) at the N-terminus and a small fragment iGlu_C (residues 254-279) at the C-terminus (**Fig. 1a**). Upon glutamate binding GluBP is reconstituted from its two fragments, pulling the cpEGFP β -barrel together, resulting in a ~5-fold fluorescence increase. Extracellular expression was achieved by fusion with a PDGFR peptide segment (Marvin et al., 2011).

iGluSnFR has high glutamate affinity and a large dynamic range, but reacts relatively slowly. Its fluorescence response is reported to have a decay half-time ($t_{1/2}$) of 92 ms upon synaptic glutamate release (Marvin et al., 2013). Imaging iGluSnFR in cultured hippocampal neurons during 10 Hz stimulation shows summation, which, without deconvolution, might indicate that glutamate accumulates during stimulation (Taschenberger et al., 2016). Deconvolution of the data suggests that glutamate is cleared between release events (Taschenberger et al., 2016). iGluSnFR itself is too slow for accurate tracking of synaptic glutamate dynamics during high frequency transmission. Here we introduce two fast iGluSnFR variants, iGlu_f (for ‘fast’) and iGlu_u (for ‘ultrafast’) and identify the rate-limiting step leading to bright fluorescence upon glutamate binding. In organotypic slice cultures of rat hippocampus, iGlu_u directly reports discrete synaptic glutamate release events at 100 Hz. Combining high-speed two-photon imaging and electrophysiology, we show that short-term depression of Schaffer collateral AMPA responses is fully accounted for by the depression of glutamate release. Furthermore, we show a tight correlation between paired-pulse facilitation and rapid recovery from post-tetanic depression at individual boutons, suggesting that differential use of presynaptic resources (readily releasable vesicles) determines the filtering properties of CA3 pyramidal cell boutons.

RESULTS

Affinity variants of iGluSnFR by binding site mutations. Six variants of iGluSnFR were generated in which residues coordinating glutamate or in the vicinity of the binding site were mutated (Hires et al., 2008). The variants had a broad range of glutamate affinities with varied fluorescence dynamic ranges (**Fig. S1a** and **Table S1**). We selected the two variants with the fastest response kinetics, iGluSnFR E26D (termed iGlu_f) and iGluSnFR S73T (termed iGlu_u) for detailed characterization with regard to their biophysical properties as isolated proteins and as membrane-bound glutamate sensors on HEK293T cells and pyramidal neurons (**Fig. 1b**). Ligand selectivity, brightness and $\text{p}K_a$ for iGlu_f and iGlu_u were similar to those for iGluSnFR (**Fig. S1b-e** and **Table S1-S3**). *In vitro* measurements gave a dissociation constant (K_d) for glutamate of 33 μM for iGluSnFR, a similar value to that previously reported (Marvin et al., 2013), while iGlu_f and iGlu_u had increased K_d values of 137 μM and 600 μM , respectively (**Fig. 1c**, **Table S4**). When expressed on the membrane of HEK293T cells, K_d values for glutamate were reduced to $3.1 \pm 0.3 \mu\text{M}$ for iGluSnFR, 26 ± 2

μM for iGlu_f and $53 \pm 4 \mu\text{M}$ for iGlu_u (measured at 37°C, **Fig. 1d,e**). A similar reduction of the K_d in the cellular environment compared to that in solution was reported for iGluSnFR (Marvin et al., 2013). The *in situ* fluorescence dynamic range $(F_{+\text{Glu}} - F_{-\text{Glu}})/F_{-\text{Glu}}$ or $\Delta F/F_0$ was 1.0 ± 0.1 for both iGluSnFR and iGlu_f, but 1.7-fold larger for iGlu_u.

Kinetic measurements of iGluSnFR variants in vitro and in situ. Based on their large K_d values, we expected iGlu_f and iGlu_u to have faster glutamate release kinetics than iGluSnFR. Fluorescence measurements in a stopped-flow instrument indeed revealed faster *off*-rates for the new variants: using the non-fluorescent high-affinity GluBP 600n (Okumoto et al., 2005) in excess (0.67 mM) to trap released glutamate, k_{off} values of 110 s^{-1} ($\tau_{\text{off}} = 9 \text{ ms}$), 283 s^{-1} ($\tau_{\text{off}} = 4 \text{ ms}$) and 468 s^{-1} ($\tau_{\text{off}} = 2 \text{ ms}$) were obtained for iGluSnFR, iGlu_f and iGlu_u, respectively, at 20°C (**Fig. 1f-h** and **Table S4**). To compare *in vitro* response kinetics to physiological measurements, the temperature dependencies of the *off*-rates of iGluSnFR and the fast variants were determined. Linear Arrhenius plots were obtained between 4°C and 34°C (**Fig. S1f,g**). For the fast variants, values exceeding the temporal precision of our stopped-flow device were linearly extrapolated (dotted line in **Fig. S1f,g**). At 34°C, decay rates were $233 \pm 3 \text{ s}^{-1}$ for iGluSnFR ($\tau_{\text{off}} = 4.3 \text{ ms}$), $478 \pm 5 \text{ s}^{-1}$ for iGlu_f ($\tau_{\text{off}} = 2.1 \text{ ms}$) and $1481 \pm 74 \text{ s}^{-1}$ for iGlu_u ($\tau_{\text{off}} = 0.68 \text{ ms}$). Thus, we were able to improve iGluSnFR kinetics by a factor of 6.3.

To image glutamate dynamics in the synaptic cleft, we expressed the newly generated iGluSnFR variants in CA3 pyramidal cells in organotypic slice culture of rat hippocampus (**Fig. 1i, j**). Fluorescence was monitored at single Schaffer collateral terminals in CA1 by spiral line scanning (**Fig. 1k**) while action potentials were triggered by brief (2 ms) depolarizing current injections into the soma of the transfected CA3 neuron. Glutamate release was detected as sharp increase in green fluorescence (**Fig. 1l**). The iGluSnFR response started $4.5 \pm 1.6 \text{ ms}$ (mean \pm SD) after the peak of the somatic action potential, consistent with a short propagation delay between CA3 and CA1. Consistent with the stochastic nature of glutamate release, individual boutons showed different release probabilities (median $p_r = 0.56$, range 0.05 – 1.0). For kinetic analysis, boutons with high release probability and good signal-to-noise ratio were selected. The measured fluorescence decay time constants (τ_{off}) were $13.8 \pm 3.8 \text{ ms}$ for iGluSnFR, $5.2 \pm 2.0 \text{ ms}$ for iGlu_f, and $2.6 \pm 1.0 \text{ ms}$ for iGlu_u (**Fig. 1m-o**). Thus, compared to iGluSnFR, detected by iGlu_u synaptic responses were revealed to be faster by a factor of 5.3.

Synaptic glutamate dynamics during high frequency stimulation. With decay kinetics of 1-2 milliseconds, iGlu_f and iGlu_u were promising tools for direct tracking of synaptic glutamate during high frequency stimulation. The response of iGluSnFR, iGlu_f and iGlu_u to paired-pulse stimulation (**Fig. 2** and **Fig. S2**) and to trains of 10 Action potentials (APs) at 50, 67 and 100 Hz (**Fig. S3**) was tested. While the responses of iGluSnFR and iGlu_f suggested build-up of glutamate during high frequency stimulation, iGlu_u responses revealed that even at 100 Hz stimulation, glutamate was completely cleared from the synaptic cleft between action potentials (**Fig. 2f, Fig. S2i**). Interestingly, the amplitude of synaptic fluorescence signals ($\Delta F/F_0$) were similar for all three indicators, suggesting that the *on*-rate, not the overall affinity, determined the number of glutamate-bound indicator molecules in the synaptic cleft.

Excitatory postsynaptic potentials (EPSPs) in CA1 become strongly depressed during high-frequency stimulation (Kim et al., 2012). We were interested whether EPSP depression during 100 Hz stimulation could be fully accounted for by depression of glutamate release from presynaptic boutons. In paired recordings from connected CA3-CA1 pyramidal cells, we triggered APs in the CA3 cell by brief current injections while monitoring postsynaptic potentials (EPSPs) in the CA1 cell. The protocol consisted of a short high frequency burst (10 APs at 100 Hz) followed by a single AP 500 ms after the burst to probe recovery of

synaptic function (Tsodyks and Markram, 1997). We repeated the protocol up to 100 times at 0.1 Hz and averaged the recorded traces (**Fig. 3a**). The decay time constant of the recovery response was used to extract the amplitude of individual responses during the 100 Hz train by deconvolution (**Fig. 3b,d**). As expected, connected CA3-CA1 pyramidal cell pairs showed strong depression during the high frequency train. The response to the recovery test pulse (#11) was not significantly different from the first EPSP in the train, indicating full recovery of synaptic function. To investigate depression and recovery of glutamate release, we evaluated $iGlu_u$ signals during identical stimulation (**Fig. 3c,e**). Due to the extremely fast kinetics of the indicator, deconvolution of the fluorescence time course was not necessary: We read the peak amplitudes during the 100 Hz train directly from the averaged fluorescence time course (average of 10 individual trials). Glutamate release decreased during the train with a time course that matched EPSP depression (**Fig. 3c**). This result points to a purely presynaptic origin of depression, which is consistent with AMPA receptors rapidly recovering from desensitization after each release event ($\tau_{\text{recovery}} = 5$ ms (Crowley et al., 2007)). However, glutamate release 500 ms after the tetanus was significantly depressed (two-tailed student's test, p -value: 0.0016) while AMPA receptor currents were not. This discrepancy suggests that the response of AMPA receptors to cleft glutamate was in fact potentiated 500 ms after the high frequency train, perfectly compensating for the reduced output of Schaffer collateral boutons.

Paired-pulse facilitation correlates with rapid recovery from depression. The rapid kinetics of $iGlu_u$ allowed us to analyze frequency filtering at individual boutons. On the second AP, boutons showed a wide range of facilitated (3 out of 13 boutons) or depressed responses (10 out of 13 boutons, **Fig. 3e**). The response to the tenth AP was strongly depressed in all boutons (18% of response amplitude to first AP), with no correlation between the second and the tenth response ($R^2 = 0.005$, **Fig. 3f**). Interestingly, a highly significant correlation was observed between the response to the second AP and the recovery response 500 ms after the high frequency train ($R^2 = 0.76$, **Fig. 3g**): Synapses that showed pronounced paired-pulse facilitation were also able to recover rapidly from depression, both of which is indicative of a low utilization of presynaptic resources (Tsodyks and Markram, 1997). Such boutons are optimized for the transmission of high-frequency activity (spike bursts). In contrast, boutons that showed paired-pulse depression were still depressed 500 ms after the high-frequency train. These boutons acting as low-pass filters: They preferentially transmit isolated APs preceded by a silent period.

Response kinetics of $iGlu_{\text{SnFR}}$ and variants $iGlu_f$ and $iGlu_u$ are based on the rate of structural change. Finally, we investigated the response mechanism of $iGlu_{\text{SnFR}}$ and its fast variants using fluorescence stopped-flow with millisecond time resolution. In association kinetic experiments (20°C), the fluorescence response rates (k_{obs}) showed hyperbolic glutamate concentration dependence, approaching saturating rates of 643 s^{-1} and 1240 s^{-1} for $iGlu_{\text{SnFR}}$ and $iGlu_f$, respectively (**Fig. 4a-d**). For $iGlu_u$, in contrast, k_{obs} was found to be concentration-independent at 604 s^{-1} (**Fig. 4e,g**). k_{off} values of 110 s^{-1} , 283 s^{-1} and 468 s^{-1} were obtained for $iGlu_{\text{SnFR}}$, $iGlu_f$ and $iGlu_u$, respectively (**Table S4**). We considered two different reaction pathways to explain our kinetic data (**Fig. 4g**). $iGlu_{\text{SnFR}}$ is represented as a complex of the large fragment of the GluBP domain (GluBP 1-253, $iGlu_l$), N-terminally flanking cpEGFP and of the C-terminally fused small GluBP fragment (GluBP 254-279, $iGlu_s$). The term $iGlu_l \sim iGlu_s$, indicates that the large GluBP fragment $iGlu_l$ and the small fragment $iGlu_s$ are within one molecule, albeit separated by the interjecting cpEGFP. In **Scheme 1**, the binding of glutamate to $iGlu_l$ in $iGlu_l \sim iGlu_s$ is the primary step (no change in fluorescence). Glutamate binding is followed by a conformational change induced by the reattachment of $iGlu_s$ to Glu-bound $iGlu_l$, resulting in the highly fluorescent $\text{Glu} \cdot iGlu_c^*$ complex (rate limiting step). According to **Scheme 1**, the hyperbolic dependence of the observed rate k_{obs} on the glutamate concentration $[\text{Glu}]$ has the intercept of the y-axis at k_2

(see **SI Kinetic Theory, eq. 7**). At low [Glu], the initial linear slope gives $k_{+2}K_1$. At high [Glu], k_{obs} tends to $k_{+2}+k_{-2}$. Although k_{obs} for iGlu_u appears essentially concentration independent, its kinetics is consistent with **Scheme 1**, with $k_{+2}+k_{-2}$ having a similar value to k_{-2} (**Table S5**).

In the alternative pathway (**Scheme 2**), the reattachment of iGlu_s to iGlu_l occurs without prior binding of glutamate. Therefore, iGlu_l~iGlu_s with the GluBP fragments separated and complete GluBP domain (iGlu_c*) are in equilibrium. The conformational change that represents the reattachment of the two GluBP fragments is expected to generate a fluorescent state of cpEGFP. However, the equilibrium is likely to be strongly shifted to the separated, non-fluorescent state (iGlu_l~iGlu_s). Assuming that this equilibrium is fast and glutamate binding stabilizes the fluorescent state, at low [Glu], a linear dependence of k_{obs} on [Glu] is predicted with a slope of $K_3k_{+4}/(1+K_3)$ and an intercept of the y-axis at k_{-2} (see **SI Kinetic Theory, eq. 15**). Although at low [Glu], mono-exponential fluorescence changes are expected, as [Glu] increases, the concentration of iGlu_c* cannot be assumed to be at steady-state and slow isomerisation will limit k_{obs} , in a similar pattern to that for Scheme 1. Thus, at high [Glu], even if iGlu_c* and Glu.iGlu_c* have equal relative fluorescence intensities, biphasic fluorescence changes would be expected for the association reactions. As all the reactions studied here for the three variants had a single exponential appearance, we can exclude **Scheme 2** as a possible reaction pathway. In conclusion, **Scheme 1** provides an excellent fit to our measurements (**Table S5**), pointing to ‘Venus fly-trap’ closure by glutamate binding as a required first step for the conformational change that increases iGluSnFR fluorescence.

DISCUSSION

The development of iGluSnFR was a breakthrough in fluorescent glutamate sensors towards investigating neurotransmission in living organisms (Xie et al., 2016). Here we describe how to overcome one of the key limitations of iGluSnFR, its slow response kinetics, and use the new ultrafast variant iGlu_u to investigate synaptic transmission and frequency filtering at individual Schaffer collateral boutons.

For all tested variants, synaptic *off*-kinetics were slower by a factor of 2.5 - 3.8 compared to temperature-matched *in vitro* measurements on isolated protein. This is consistent with the much higher affinities of HEK293T cell-expressed glutamate sensors compared to soluble protein. These systematic differences, also noted in the original characterization of iGluSnFR (Marvin et al., 2013), may be attributed to the tethering of the molecule to a membrane anchor, slowing down conformational changes compared to free-floating sensor molecules. Nevertheless, the relative differences in affinity and kinetics of the new versions compared to iGluSnFR were preserved *in vitro* and *in situ*. The *on*- and *off*-rates of iGlu_u are greater (2- and 5-6 fold, respectively) compared to iGluSnFR. Interestingly, iGlu_u was a faster reporter in the hippocampal slice than iGlu_f, even though the latter has a faster limiting *on*-rate. iGlu_u may be put at an advantage over iGlu_f by its concentration-independent response kinetics. It must be noted that the kinetics of iGluSnFR-type indicators are ultimately limited by the structural change that reconstitutes the fluorescent complex, similar to calcium-sensing GCaMPs. The constraints of the mechanism with regard to the onset of fluorescence suggest that it cannot be engineered to resolve sub-millisecond glutamate dynamics. To achieve microsecond response times, it might be necessary to develop hybrid glutamate indicators using synthetic dyes.

Synaptic iGlu_u imaging revealed complete clearance of glutamate between release events even at 100 Hz stimulation frequency. The time course of synaptic glutamate transients has previously been estimated from the decay of NMDA receptor responses in primary cell culture: Kinetic analysis of the displacement of a competitive NMDA receptor antagonist suggest rapid glutamate clearance with $\tau = 1.2$ ms (Clements et al., 1992). Our

optical measurements in hippocampal tissue are consistent with these earlier estimates, but, due to the intrinsic limits of the iGluSnFR mechanism, we might still underestimate the true kinetics of free glutamate in the synaptic cleft. What we can say with confidence is that accumulation of glutamate in the synaptic cleft does not contribute to short-term plasticity at Schaffer collateral synapses.

Glutamate release showed strong depression during 100 Hz firing, in line with the expected depletion of release-ready vesicles. As we controlled the generation of every action potential by somatic current injections, we can exclude decreased afferent excitability as a source of depression in these experiments (Kim et al., 2012). AMPA receptor currents during 100 Hz firing did not show more run-down than iGlu_u responses, suggesting that AMPA receptor desensitization did not play a major role in the decrease of synaptic efficacy during the train. Paradoxically, AMPA responses were fully recovered 500 ms after the train while glutamate release was still significantly depressed. This finding points to an unexpected increase in sensitivity of the postsynaptic compartment to glutamate. It has been shown previously that post-tetanic potentiation in hippocampal slice culture can be blocked by antagonists of protein kinase C, which was thought to be a presynaptic effect (Brager et al., 2003). However, active PKC is able to phosphorylate GluA1-Ser831, thereby enhancing AMPA receptor conductance (Jenkins and Traynelis, 2012). Thus, it is possible that elevated Ca²⁺ levels in the spine during our high frequency protocol activated PKC and enhanced AMPA receptor currents, compensating for the reduced glutamate release 500 ms after the tetanus.

The surprisingly tight correlation between paired-pulse facilitation and rapid recovery from depression at individual boutons provides direct evidence that differential use of presynaptic resources determines the neural code between pyramidal cells (Tsodyks and Markram, 1997; Markram et al., 1998). Using Schaffer collateral synapses as an example, we show that iGlu_u is a useful tool for a mechanistic analysis of high frequency synaptic transmission, interrogating presynaptic function independently of postsynaptic transmitter receptors.

METHODS

We provide a detailed description of the methods, data analysis and kinetic modeling in the on-line Supplemental Information.

Materials. pCMV(MinDis).iGluSnFR and pRSET FLIPE-600n plasmids were a gift from Loren Looger (Addgene Plasmid #41732) and Wolf Frommer (Addgene plasmid # 13537), respectively. Site-directed mutagenesis was carried out following the QuikChange II XL protocol (Agilent Technologies).

Fluorescence spectroscopies. Glutamate association and dissociation kinetic experiments of iGluSnFR proteins were carried out on a Hi-Tech Scientific SF-61DX2 stopped-flow system equipped with a temperature manifold (Walklate and Geeves, 2015). Fluorescence spectra and equilibrium glutamate titrations were recorded on a Fluorolog3 (Horiba Scientific).

In situ glutamate titration. HEK293T cells were cultured on 24-well glass bottom plates in DMEM containing non-essential amino-acids (Life Technologies), 10% heat inactivated FBS (Life Technologies) and penicillin/streptomycin (100 U/ml, 100 mg/ml, respectively), at 37°C in an atmosphere of 5% CO₂. Cells were allowed 24 h to adhere before transfection with Lipofectamine 2000 (Invitrogen). Cells were examined at 37°C (OKO lab incubation chamber) with a 3i Marianas spinning-disk confocal microscope equipped with a Zeiss

AxioObserver Z1, a 40x/NA1.3 oil immersion objective and a 3i Laserstack as excitation light source (488 nm).

Synaptic measurements. Organotypic hippocampal slices (400 μm) were prepared from male Wistar rats at postnatal day 5 as described (Gee et al., 2017). iGluSnFR and variant plasmids were electroporated into 2-3 CA3 pyramidal cells at 40 ng/ μl (iGluSnFR) or 50 ng/ μl (iGlu_f, iGlu_u) together with tdimer2 (20 ng/ μl), a cytoplasmic red fluorescent protein (Wiegert et al., 2017). 2 - 4 days after electroporation (at DIV 14-30), slice cultures were placed in the recording chamber of a two-photon microscope and superfused with artificial cerebrospinal fluid (ACSF) containing (in mM) 25 NaHCO₃, 1.25 NaH₂PO₄, 127 NaCl, 25 D-glucose, 2.5 KCl, 2 CaCl₂, 1 MgCl₂. Whole-cell recordings from a transfected CA3 pyramidal cell were made with a Multiclamp 700B amplifier (Molecular Devices). Red and green fluorescence was detected through the objective (LUMPLFLN 60XW, 60x, NA 1.0, Olympus) and through the oil immersion condenser (NA 1.4, Olympus) using 2 pairs of photomultiplier tubes (H7422P-40SEL, Hamamatsu).

Acknowledgements. We thank Dr Zoltan Ujfalusi (University of Kent) for assistance with stopped-flow experiments, Iris Ohmert for the preparation of organotypic cultures and Dr David Trentham for comments on the manuscript. Use of The Institute of Translational Medicine, University of Liverpool Imaging Facility is gratefully acknowledged. This project was supported by the Wellcome Trust (Grant 094385/Z/10/Z to K.T.) and BBSRC (Grant BB/M02556X/1 to K.T.); the German Research Foundation (SPP 1665, SFB 936, FOR 2419 to T.G.O; SPP 1926, FOR 2419 to J.S.W) and the European Research Council (ERC-2016-StG 714762 to J.S.W.).

Reference List

Baird GS, Zacharias DA, Tsien RY (1999) Circular permutation and receptor insertion within green fluorescent proteins. *Proc Natl Acad Sci U S A* 96:11241-11246.

Best SL, Török K (2005) Development of a fluorescent glutamate binding protein. *Biophysical Journal* 88:46a.

Brager DH, Cai X, Thompson SM (2003) Activity-dependent activation of presynaptic protein kinase C mediates post-tetanic potentiation. *Nat Neurosci* 6:551-552.

Chen GQ, Gouaux E (1997) Overexpression of a glutamate receptor (GluR2) ligand binding domain in *Escherichia coli*: application of a novel protein folding screen. *Proc Natl Acad Sci U S A* 94:13431-13436.

Clements JD, Lester RA, Tong G, Jahr CE, Westbrook GL (1992) The time course of glutamate in the synaptic cleft. *Science* 258:1498-1501.

Crowley JJ, Carter AG, Regehr WG (2007) Fast vesicle replenishment and rapid recovery from desensitization at a single synaptic release site. *J Neurosci* 27:5448-5460.

de Lorimier RM, Smith JJ, Dwyer MA, Looger LL, Sali KM, Paavola CD, Rizk SS, Sadigov S, Conrad DW, Loew L, Hellinga HW (2002) Construction of a fluorescent biosensor family. *Protein Sci* 11:2655-2675.

Gee CE, Ohmert I, Wiegert JS, Oertner TG (2017) Preparation of Slice Cultures from Rodent Hippocampus. *Cold Spring Harb Protoc* 2017: Feb 1;2017(2):pdb.prot094888. doi: 10.1101/pdb.prot094888.db.

Hires SA, Zhu Y, Tsien RY (2008) Optical measurement of synaptic glutamate spillover and reuptake by linker optimized glutamate-sensitive fluorescent reporters. *Proc Natl Acad Sci U S A* 105:4411-4416.

Hu Y, Fan CP, Fu G, Zhu D, Jin Q, Wang DC (2008) Crystal structure of a glutamate/aspartate binding protein complexed with a glutamate molecule: structural basis of ligand specificity at atomic resolution. *J Mol Biol* 382:99-111.

Jenkins MA, Traynelis SF (2012) PKC phosphorylates GluA1-Ser831 to enhance AMPA receptor conductance. *Channels (Austin)* 6:60-64.

Kim E, Owen B, Holmes WR, Grover LM (2012) Decreased afferent excitability contributes to synaptic depression during high-frequency stimulation in hippocampal area CA1. *J Neurophysiol* 108:1965-1976.

Kuusinen A, Arvola M, Keinänen K (1995) Molecular dissection of the agonist binding site of an AMPA receptor. *EMBO J* 14:6327-6332.

Markram H, Wang Y, Tsodyks M (1998) Differential signaling via the same axon of neocortical pyramidal neurons. *Proc Natl Acad Sci U S A* 95:5323-5328.

Marvin JS, Borghuis BG, Tian L, Cichon J, Harnett MT, Akerboom J, Gordus A, Renninger SL, Chen TW, Bargmann CI, Orger MB, Schreiter ER, Demb JB, Gan WB, Hires SA, Looger LL (2013) An optimized fluorescent probe for visualizing glutamate neurotransmission. *Nat Methods* 10:162-170.

Marvin JS, Schreiter ER, Echevarria IM, Looger LL (2011) A genetically encoded, high-signal-to-noise maltose sensor. *Proteins* 79:3025-3036.

Nakai J, Ohkura M, Imoto K (2001) A high signal-to-noise Ca²⁺ probe composed of a single green fluorescent protein. *Nat Biotechnol* 19:137-141.

Okumoto S, Looger LL, Micheva KD, Reimer RJ, Smith SJ, Frommer WB (2005) Detection of glutamate release from neurons by genetically encoded surface-displayed FRET nanosensors. *Proc Natl Acad Sci U S A* 102:8740-8745.

Regehr WG (2012) Short-term presynaptic plasticity. *Cold Spring Harb Perspect Biol* 4:a005702.

Taschenberger H, Woehler A, Neher E (2016) Superpriming of synaptic vesicles as a common basis for intersynapse variability and modulation of synaptic strength. *Proc Natl Acad Sci U S A* 113:E4548-E4557.

Tsien RY (2005) Building and breeding molecules to spy on cells and tumors. *FEBS Lett* 579:927-932.

Tsodyks MV, Markram H (1997) The neural code between neocortical pyramidal neurons depends on neurotransmitter release probability. *Proc Natl Acad Sci U S A* 94:719-723.

Walklate J, Geeves MA (2015) Temperature manifold for a stopped-flow machine to allow measurements from -10 to +40 degrees C. *Anal Biochem* 476:11-16.

Wiegert JS, Gee CE, Oertner TG (2017) Single-Cell Electroporation of Neurons. *Cold Spring Harb Protoc* 2017: Feb 1;2017(2):pdb.prot094904. doi: 10.1101/pdb.prot094904.

Xie Y, Chan AW, McGirr A, Xue S, Xiao D, Zeng H, Murphy TH (2016) Resolution of High-Frequency Mesoscale Intracortical Maps Using the Genetically Encoded Glutamate Sensor iGluSnFR. *J Neurosci* 36:1261-1272.

LEGENDS

FIGURE 1. Genetically encoded glutamate indicators (GEGI). (a) Domain structure and design of FRET- and single fluorophore-based GEGI; key: (GluBP) (blue), cpEGFP (green), IgG kappa secretion tag (pink), hemagglutinin (HA) tag (purple), myc tag (grey) and a PDGFR transmembrane domain (brown); iGluSnFR lacks the hemagglutinin tag, GluBP 1-253 and 254-279 fragments are in light and dark blue, respectively; $\Delta 8$ aa and $\Delta 5$ aa specify deletions at the N- and C-terminus of GluBP introduced in GluSnFR. (b) Design of selected iGluSnFR variants. Crystal structure of GluBP (PDB 2VHA, adapted from Hu *et al.* (Hu *et al.*, 2008)). Selected mutated residues around the glutamate site are shown as red and green backbone as specified. Bound glutamate is represented in orange space filling display. (c) Equilibrium glutamate binding titrations at 20°C for iGluSnFR (●), iGluSnFR E26D (iGlu_f) (▼) and iGluSnFR S73T (iGlu_u) (■) *in vitro*; (d) Glutamate titrations *in situ*. iGluSnFR, iGlu_f and iGlu_u were expressed in HEK293T cells and titrated with glutamate. Data derived from iGluSnFR (n = 19), iGlu_f (n = 41) and iGlu_u (n = 33). (e) Representative images of HEK293T cells prior to glutamate addition and at saturating (1, 3 and 10 mM, respectively) glutamate. The scale bar represents 10 μ m. Glutamate dissociation kinetics of (f) iGluSnFR, (g) iGlu_f and (h) iGlu_u determined by stopped-flow fluorimetry. Experimental data (dotted lines) are overlaid by curves fitted to single exponentials (solid lines). Fluorescence changes are normalised to F_{\max} of 1. Imaging glutamate release from single presynaptic terminals. (i) Schematic representation of organotypic hippocampal slice culture with transfected and patch-clamped CA3 pyramidal cell; (j) Imaging axonal projections in CA1 (two-photon stack, maximum intensity projection); (k) Individual bouton with spiral scan path for 500 Hz sampling; (l) Unfolded scan lines (64 lines, 2 ms/line), single trial. The scan line intersected the fusion site of the vesicle in two positions; Fluorescence time course ($\Delta F/F_0$) upon glutamate release stimulated by paired pulse stimulation (48 ms ISI) by iGlu_u. Decay time (τ_{off}) measurements with bleach correction (solid lines) for individual experiments by single exponential fit for (m) iGluSnFR (n = 13 boutons, 500 Hz sampling rate) and variants (n) iGlu_f (n = 7 boutons, 1 kHz sampling rate) and (o) iGlu_u (n = 7 boutons, 1 kHz sampling rate).

FIGURE 2. Imaging glutamate release from single presynaptic terminals. Spiral line scans at 500 Hz were used to cover the entire surface of individual boutons, intersecting the release

site multiple times. Responses of (a,d) iGluSnFR, (b,e) iGlu_f and (c,f) iGlu_u-expressing boutons stimulated by 2 somatic action potentials at 48 ms (a-c) and 10 ms inter-stimulus interval (d-f). Upper traces: single trial responses. Lower traces: averages of 3-6 responses.

FIGURE 3. Depression and recovery of synaptic transmission during 100 Hz trains. (a) Example of patch-clamp recording from a connected pair of CA3-CA1 pyramidal cells. Black trace: Induced action potentials (APs) in CA1 pyramidal cell, 100 Hz train and single AP. Gray trace: EPSPs in CA1 pyramidal cell (average of 50 sweeps). The single AP response (right) was used to extract EPSP amplitudes from the burst response (dotted line). Green trace: Single-bouton iGlu_u response to identical stimulation (average of 10 sweeps). (b) EPSPs (deconvolved amplitudes) show strong depression during the 100 Hz train, full recovery 500 ms later (n = 5 CA3-CA1 pairs), two-tailed student's test comparing EPSP #1 and EPSP #11. (c) Glutamate release showed strong depression during the 100 Hz train, partial recovery 500 ms later (n = 13 boutons, 8 cells), two-tailed student's test comparing response #1 and response #11 (p<0.01). (d) Individual paired recordings show consistent depression (response 10) and recovery (response #11). (e) Individual Schaffer collateral boutons show large variability in 2nd response and in recovery response (#11) (f) iGlu_u responses to second AP (paired-pulse facilitation/depression) were not correlated with total depression (response #10). (g) iGlu_u responses to second AP were highly correlated with recovery after 500 ms (response #11).

FIGURE 4. Kinetics of glutamate binding by iGluSnFR variants (20°C). (a, c, e) Glutamate association kinetics of iGluSnFR, iGlu_f and iGlu_u, respectively. Stopped-flow records of iGluSnFR, iGlu_f and iGlu_u reacting with the indicated concentrations of glutamate. Experimental data (dotted lines) are overlaid with curves fitted to single exponentials (solid lines); (b, d, f) Plot of observed association rates, $k_{obs(on)}$ of iGluSnFR, iGlu_f and iGlu_u as a function of glutamate concentration; (g) Cartoon diagram depicting the putative molecular transitions of iGluSnFR and its fast variants to the fluorescent state. Key: cpEGFP (green), GluBP 1-253 (iGlu_l) (light blue) and 254-279 (iGlu_s) (dark blue) fragments, glutamate (orange).

Figure 1




Figure 2




Figure 3




Figure 4

