

From socio-economic disadvantage to obesity: The mediating role of psychological distress
and emotional eating.

Jade Spinosa^{1,2}, Paul Christiansen¹, Joanne M. Dickson³, Valentina Lorenzetti⁴, & Charlotte
A. Hardman^{1*}

¹ Department of Psychological Sciences, University of Liverpool, UK

² Clinical Psychology Department, St Helens Hospital, UK

³ Psychology Department, Edith Cowan University, Australia

⁴ Australian Catholic University, Melbourne, Australia

Keywords: Socio-economic Status, Obesity, Emotional Eating

Running Title: Psychological pathway between lower SES and obesity.

* **Correspondence to:** Dr. Charlotte Hardman, Department of Psychological Sciences, 2.08, Eleanor Rathbone Building, Bedford Street South, University of Liverpool, Liverpool, L69 7ZA, UK. Phone: +44 (0)151 7941480 Email: Charlotte.Hardman@liverpool.ac.uk

Word Count: 3896

Conflict of Interest: Charlotte Hardman and Paul Christiansen receive research funding from the American Beverage Association. Jade Spinosa, Valentina Lorenzetti and Joanne Dickson declare no conflicts of interest.

STUDY IMPORTANCE QUESTIONS

* What is already known about this subject?

- In developed countries, lower socio-economic status is robustly associated with obesity.
- Socio-economic disparities are widely attributed to differences in dietary intake.
- Socio-economic disadvantage is associated with psychological distress.

* What does this study add?

- Socio-economic status is associated with obesity via psychological distress and subsequent emotional eating.
- Higher levels of resilience did not provide a protective effect.
- New insight into psychological mediators of socio-economic disparities in obesity.

Abstract

Objective: Lower socio-economic status is robustly associated with obesity, however the underpinning psychological mechanisms remain unclear. The current study sought to determine whether the relationship between lower socio-economic status and obesity is explained by psychological distress and subsequent emotional eating as a coping strategy. It also examined whether psychological resilience plays a protective role in this pathway.

Method: Participants ($N=150$), from a range of socio-economic backgrounds, completed questionnaire measures of psychological distress, emotional eating, and resilience. They reported their income and education level as an indicator of socio-economic status, and their height and weight in order to calculate body mass index (BMI). **Results:** There was a significant indirect effect of socio-economic status on BMI via psychological distress and emotional eating; specifically, lower socio-economic status was associated with higher distress, higher distress was associated with higher emotional eating, and higher emotional eating was associated with higher BMI, $b(SE) = -.02 (.01)$, 95% CI = $-.04, -.01$. However, resilience was not a significant moderator of this association. **Conclusions:** Psychological distress and subsequent emotional eating represent a serial pathway that links lower socio-economic status with obesity. Targeting these maladaptive coping behaviours may be one strategy to reduce obesity in low-income populations.

Introduction

The increasing prevalence of obesity in many countries worldwide has been labelled as an “epidemic”. Globally, the number of individuals with overweight and obesity increased by 27.5% for adults and 47.1% for children between 1980 and 2013 (1). In the UK, around two-thirds of adults are overweight or obese, and obesity prevalence almost doubled between 1993 and 2015 (2). Costs to the National Health Service associated with obesity or being overweight are projected to reach £10 billion by 2050, with wider economic costs (such as days of employment missed) expected to reach £50 billion per year (3). Current weight management strategies primarily focus upon two areas; improving the quality of dietary intake and reducing sedentary lifestyles. However, their success has been limited, especially with regard to longer-term maintenance of weight loss (4).

The causes of obesity are complex and vary between individuals. However, one factor that has been reliably associated with obesity is socio-economic status (SES). In developed countries, obesity disproportionately affects individuals from lower social classes and this is particularly the case for women (5-9). Recent research (10) has shown how socio-economic disparities in child and adolescent body weight have reversed over time; in the 1940s through to the 1970s, low SES was associated with *lower* weight, however in 2001, low SES was associated with *higher* weight. The reason for this socio-economic disparity is not well understood but it is often attributed to the greater availability of low-cost, calorie-dense foods in more deprived areas relative to more affluent neighbourhoods (11). However, there is limited evidence for an association between local food environments and obesity (12) indicating that other factors also play a role.

To date, there has been relatively little consideration of the underlying psychological mechanisms which may explain why individuals from lower socio-economic groups are vulnerable to developing obesity. In view of this, Hemmingsson (13) recently proposed a

theoretical model which emphasises the role of psychological, emotional and social factors. According to this model, socio-economic disadvantage causes psychological and emotional distress. This distress is transferred from parents to children thus creating a disharmonious family environment. Ultimately, the adult and/or child experiences psychological and emotional overload which leads to maladaptive coping strategies, such as eating energy-dense foods to alleviate negative emotions and stress. These maladaptive eating behaviours, coupled with stress-induced disturbances to metabolic signals, are thought to promote weight gain and obesity over time. The negative social, psychological, emotional and behavioural consequences of obesity exacerbate psychological distress and maladaptive eating behaviours, thus creating a cyclic mechanism which perpetuates the difficulties.

Whilst Hemmingsson's model (13) is yet to be empirically tested in its entirety, there is considerable support for some of the proposed relationships. For example, numerous studies have shown a link between socio-economic disadvantage (e.g., income inequality and lower social status) and psychological distress, such as higher rates of depression and lower mental wellbeing in lower socio-economic groups relative to more affluent groups (14-17). Poverty and poor mental health are inter-related such that poverty can be both a cause and consequence of mental health problems (18). With regard to obesity, in an experimental study, participants who were experimentally induced to feel poor consumed significantly more calories from snack foods compared to participants who were induced to feel wealthy (19). Notably, this study also found that higher social anxiety was directly associated with increased consumption, and this was particularly true for participants who had a strong need to belong (19). There is also a body of evidence linking emotional or stress-induced eating with higher BMI and consumption of energy-dense sweet and/or fatty foods in adults (20-22) and in children and adolescents (for a meta-analysis see (23)). Furthermore, the tendency to eat palatable foods as a coping strategy predicted increases in BMI over 2 years in young

adults (24). So while there is general support for direct relationships between these key variables, to the authors' knowledge, no studies have directly examined whether psychological distress and emotional eating mediate the association between SES and BMI.

Hemmingsson's model (13) also proposed that there are protective factors which can act as buffers thus preventing the link between socio-economic disadvantage and psychological distress. Resilience refers to an individual's capacity to cope with stressors and to withstand the potential depressive consequences of such stressors (25). In previous studies, lower levels of resilience were associated with higher incidences of depression (26, 27). Low resilience has also been found to independently predict higher BMI and waist circumference (28). This suggests that individuals who are high in resilience may cope better with socio-economic disadvantage and thus be protected from increased psychological distress and subsequent maladaptive eating.

The current study aimed to elucidate the associations between SES, psychological distress, emotional eating and BMI. It was predicted that lower SES would be indirectly associated with (higher) BMI via psychological distress and emotional eating (i.e., whereby lower SES is associated with higher psychological distress; higher psychological distress is associated with higher emotional eating; higher emotional eating is associated with higher BMI). It was also predicted that resilience would moderate the association between lower SES and psychological distress such that this association would be most pronounced in individuals with lower relative to higher levels of resilience.

Method

Participants

Participants were recruited using online advertisements (via internal university websites and externally using social media) and through an Urban Community and

Neighbourhood Centre (UCAN) situated in the town of Bolton in the North West of England. UCANs provide support and advice to local residents within an identified geographical area of socio-economic deprivation and Bolton is one of the most deprived local authorities in England (29). Inclusion criteria for the study were being aged between 18 and 65 years with a good level of English language skills. 194 participants were recruited in total and commenced the study. Complete data were obtained for 150 of these participants. Ethical approval for the study was obtained through the University of Liverpool's Research Ethics Committee.

All participants were provided with written information outlining the nature and purpose of the study. Written informed consent was obtained prior to study commencement. As compensation for their time, participants were given the option to be entered into a prize draw upon completion of the study.

Measures

Demographic information: Each participant was asked to provide their age (in years), gender and ethnicity.

Socio-economic status (SES): Consistent with previous approaches (30, 31), participants reported their employment status (employed full-time, employed part-time, unemployed looking for work, unemployed not looking for work, retired, student, unable to work due to health or disability, housewife/husband, voluntary employment), their total household income (9 – point scale: 1 = < £5 200, 2 = £5 200 - £10 399, 3 = 10 400 - £15 599, 4 = £15 600 - £20 799, 5 = £20 800 - £25 999, 6 = £26 000 - £36 399, 7 = £36 400 - £51 999, 8 = £52 000 - £77 999, 9 = >£78 000) and their level of education (8-point scale: 1 = none, 2 = GCSE grade D or below, 3 = GCSE grade C or above, 4 = A-level or equivalent, 5 = university degree or equivalent, 6 = postgraduate qualification or equivalent, 7 = Masters or equivalent, 8 = PHD or equivalent).

Psychological Distress: The 21-item Depression, Anxiety and Stress Scale (DASS) (32) was used to measure three related states of psychological distress: depression, anxiety and stress. Participants responded to each item (e.g., “I found it hard to wind down”) using a 4-point scale (0 = never, 1 = sometimes, 2 = often, 3 = almost always). Cronbach’s α values for the current study were as follows: depression $\alpha = .92$, anxiety $\alpha = .83$, stress $\alpha = .86$, total distress scale $\alpha = .94$.

Emotional Eating: The 13-item emotional eating subscale from the Dutch Eating Behaviour Questionnaire (DEBQ) was used (33). Participants responded to each item (e.g. “Do you have a desire to eat when you are emotionally upset?”) on a 5-point scale (1 = never, 2 = seldom, 3 = sometimes, 4 = often, 5 = very often). The total subscale score was calculated as the mean of responses to the items. Cronbach’s α for the current data was $\alpha = .95$.

Body Mass Index (BMI): Participants reported their current weight (in kg or in stones and pounds) and height (in cm or in feet and inches). Data were converted to metric units, where necessary, to calculate BMI using the formula $\text{weight}(\text{kg})/\text{height}(\text{m})^2$. Previous research indicates that self-reported and objectively measured weight data are highly correlated (34, 35).

Resilience: The 6-item Brief Resilience Scale (BRS) was used (36). Participants responded to each item (e.g. “I tend to bounce back quickly after hard times”) on a 5-point scale (1 = strongly disagree, 2 = disagree, 3 = neutral, 4 = agree, 5 = strongly agree). The total scale score was calculated as the mean of responses to the items with higher scores indicating higher levels of resilience. Cronbach’s α for the current data was $\alpha = .86$.

Life Events: The Life Events Scale (LES) measures the occurrence of stressful life events (37). Participants are asked to indicate how many times in the past 5 years they have experienced 14 events (e.g. death of a child/partner/relative/friend, end of intense relationship, serious or long-lasting financial problems, serious or longstanding work

problem). The frequencies for each event are totalled to provide the total scale score with higher scores indicating a greater frequency of stressful life events.

Procedure

The questionnaires were hosted online using Qualtrics. Participants recruited using online advertisements could access the questionnaires via a web link. Participants at the UCAN centre were recruited using paper-based advertisements posted within the centre which provided them with the web link to the online survey. Alternatively, participants at the UCAN centre were given the opportunity to meet with the researcher and were then given access to the online survey at a computer in the centre or, depending on their personal preference, they completed paper-based copies of the questionnaires ($N=20$ opted to complete paper-based copies, 13% of the total sample). We took a flexible approach to recruitment to ensure that participants without internet access were not precluded from taking part.

Upon commencing the study, participants first viewed the Participant Information Sheet and provided their consent to participate in the subsequent consent form. They then completed the demographic (including height and weight) and SES information followed by the four questionnaires in the following order: DASS, BRS, DEBQ and LES. Upon completion of the study participants were provided with a debrief information sheet.

Statistical Analysis

According to guidance on sample size for mediation analyses (38), a minimum of 71 participants are needed to detect the hypothesised associations between the key variables (80% power with medium effect sizes). Data were checked for outliers alongside tests of multi-collinearity, homogeneity, and homoscedasticity to ensure assumptions for further analysis were met. Due to the variation in measurement scales used, all variables were log transformed to standardise the data prior to running further analyses. Data sets for four participants had missing data for a single item, within either the DEBQ or DASS. Missing

data points were handled using valid mean substitution (VMS). VMS uses the average of participants' other responses to generate a value for the missing data. It has been shown to be a valid method, when the measure in question employs multiple items to gauge a single construct and where participants have answered all remaining questions related to that construct (39). A composite score was generated for SES using the two measures with numerical scales; total household income (9-point scale; higher scores indicate higher income) and level of education (8-point scale; higher scores indicate higher levels of education), which have been used as indicators of SES in previous studies (30, 31). The DASS provides sub-scores for depression, anxiety and stress whilst also providing a total score as a more general dimension of psychological distress. Initial correlations indicated a high level of association across the three sub-scores (r values $\geq .68$), therefore, the DASS total score was used in the analysis.

The primary hypothesis predicted that lower SES would be indirectly associated with (higher) BMI via psychological distress and emotional eating. This was tested by a serial multiple mediation analysis in PROCESS (40); the independent variable (IV) was SES, the dependent variable (DV) was BMI, and the serial mediators were psychological distress (first mediator) and emotional eating (second mediator) (Figure 1). PROCESS compares the magnitude of the direct effect (IV-DV; controlling for the mediators) with the total effect of the IV on the DV including the indirect pathway via the mediators. It produces bias-corrected bootstrapped confidence intervals (CIs) for indirect effects via individual mediators and for the serial effect of the two mediators in the serial mediation model. A significant indirect effect is inferred by upper and lower CIs that do not include zero.

Exploratory analyses were conducted to determine whether lower SES was associated with higher psychological distress due to greater frequency of negative life events (measured using the Life Events Scale). This was tested by a serial multiple mediation analysis; the

independent variable was SES, the dependent variable was BMI, and the serial mediators were negative life events (first mediator), psychological distress (second mediator) and emotional eating (third mediator).

It was also predicted that resilience would moderate the association between lower SES and psychological distress. This was tested using a moderated mediation analysis in PROCESS in which the indirect effect of SES on emotional eating via psychological distress was examined at three levels of the moderator (resilience scores; $-1 SD$, mean, $+1 SD$).

All models controlled for age and gender as covariates.

Results

Descriptive characteristics of the study participants are shown in Table 1 ($N = 150$). Boxplots illustrating the spread of the data for highest education level and yearly household income can be found in the supplementary material (Figure S1). The majority of the sample were female (83%) and white (93%). With regard to employment status, 52% of the sample were employed full time, 16% were employed part-time, 16% were students, 9% were unemployed and/or looking for work, 3% were unable to work due to health or disability, 3% reported being a housewife/husband, <1% were retired, and <1% were in voluntary employment. The mean BMI of the sample was 26.3 kg/m^2 (scores >25 indicative of being overweight). 4% of participants were underweight, 44% were of healthy weight, 32% were overweight, and 20% had obesity. The correlations between the variables are shown in Table 2.

Effect of SES on BMI via Psychological Distress and Emotional Eating (Figure 1)

In the serial multiple mediation model, there was no significant total effect of SES on BMI, $b(SE) = -.01 (.04)$, $p = .79$. However, as predicted, there was a significant indirect effect of SES on BMI via psychological distress and emotional eating; $b(SE) = -.02 (.01)$, 95% CI =

-.04, -.01. That is, lower SES predicted higher psychological distress which, in turn, predicted higher emotional eating which, in turn, predicted higher BMI. There was also a significant simple indirect effect of SES on BMI via emotional eating; $b(SE) = .06 (.02)$, 95% CI = .03, .10. However, contrary to prediction this pathway indicated that higher SES predicted higher emotional eating. The simple indirect effect of SES on BMI via psychological distress was not significant $b(SE) = .01 (.01)$, 95% CI = -.01, .04. Taken together, the effect of SES on BMI in the model accounting for the all mediators explained 15% of the variance ($R^2 = .15$, $p = .0003$).

Insert Figure 1 here

The exploratory analysis with the inclusion of negative life events as an additional serial mediator found no evidence of a significant indirect effect via the three mediator serial pathway (i.e., SES → negative life events → distress → emotional eating → BMI; $b(SE) = -.001 (.001)$, 95% CI = -.01, .00). Please see the supplementary material (Figure S2) for additional detail of this analysis.

Resilience as a Moderator of the Indirect Effect of SES on Emotional Eating via Psychological Distress

Resilience was found to be an independent predictor of psychological distress, whereby higher resilience was associated with lower psychological distress, $b(SE) = -1.21 (.20)$, $p < .001$. However, the significant indirect effect of SES on emotional eating via psychological distress remained evident at all three levels of the moderator (Low, Medium and High resilience) (Table 3). The total index of moderated mediation indicated that no significant moderation occurred $b(SE) = -.122 (.18)$, 95% CI = -.51, .19.

Discussion

The current study aimed to elucidate the associations between SES, psychological distress, emotional eating and BMI. As predicted, there was a significant indirect effect of SES on BMI via psychological distress and emotional eating; namely, lower SES was predictive of higher psychological distress, higher psychological distress predicted higher emotional eating which, in turn, predicted higher BMI. This finding directly supports components of the theoretical model proposed by Hemmingsson (13) which emphasises the key role of psychological distress and maladaptive coping strategies in explaining the association between socio-economic disadvantage and obesity.

In the current study, the simple indirect effect of SES on BMI via psychological distress was not significant. This indicates that psychological distress did not significantly mediate the (cross-sectional) relationship between SES and BMI; the pathway required the addition of emotional eating as a coping strategy for distress. This finding suggests that it is not distress *per se* but people's coping strategies for dealing with distress that may be critical in explaining the link between socio-economic disadvantage and body weight. Consistent with this, a recent study (41) found that although lower SES was associated with both greater psychosocial stress and weight gain over a 9-year period, perceived stress did not mediate the higher weight gain associated with lower SES.

Higher resilience was an independent predictor of lower psychological distress, in line with previous research (26, 27). However, contrary to our hypothesis, resilience did not moderate the relationship between SES and psychological distress. This indicates that being high on trait resilience alone was not sufficient to protect those with lower SES from experiencing greater levels of psychological distress. Hemmingsson's theoretical model emphasises a process whereby multiple factors create a cumulative protective effect (i.e. resilience, social support, self-esteem, functional coping). It is possible therefore that

resilience alone in the current study, without other additional protective mechanisms, is not sufficient to moderate the relationship between SES and psychological distress. Future studies in this area should seek to measure a range of potential protective factors.

The current study also revealed an unexpected finding, whereby *higher* SES was predictive of *higher* emotional eating in the simple indirect pathway, independent of psychological distress. Previous research has found that male participants with degree level qualifications (indicative of higher SES) had significant levels of stress-related eating (22). The positive direct association between SES and emotional eating contrasts with the previously-discussed negative indirect association between SES and emotional eating via psychological distress; this phenomenon of opposing directions of direct and indirect effects in a mediation analysis is known as a suppression effect. This finding suggests that, for individuals in higher social-economic positions, emotional eating is also prevalent, however, critically this is not in response to significant psychological distress. In addition, the DEBQ emotional eating measure used in the current study assesses the tendency to eat in response to a variety of emotions, some of which imply coping (e.g. in response to low mood) but others which do not (e.g., boredom). It is, therefore, possible that participants of higher SES may be eating in response to other emotions, not directly related to coping with distress.

The exploratory analyses failed to find a significant association between lower SES and greater frequency of negative life events. This suggests that it is not an increased likelihood of negative life events *per se* which makes individuals from lower SES more vulnerable to experiencing greater psychological distress. Other studies have suggested that individuals who experience socio-economic disadvantage may have more limited access to resources (e.g., material, interpersonal and intrapersonal) (43). It is possible that this, rather than negative life events themselves, may make them more vulnerable to experiencing psychological distress and subsequent maladaptive coping behaviours. Other psychological

experiences, for example, feeling lower in social rank or feeling deprived, may also underpin some of the vulnerability posed by socio-economic disadvantage, and should be explored further.

We had a relatively small sample size and a limited number of participants with obesity (20%) based on BMI. However, the average BMI was in the overweight range and is in line with the national average for BMI in the UK (27 women, 27.4 men (44)). BMI was self-reported within the current study and, whilst previous research has indicated that self-reported and objectively measured weight data are highly correlated (34, 35), it is possible that discrepancies may have occurred. Future research would benefit by using objective measures of body weight. The sample recruited was a predominantly female, white population and further research is needed to explore the role of gender and ethnicity. Yet our findings are relevant as previous studies indicate that socio-economic disparities in obesity are most pronounced in women (5). In addition, the propensity to use increased consumption of food as a coping strategy may be more prevalent in women than men (21). Examination of economic circumstances and population weight in 67 countries found that whilst in more economically developed nations lower SES was associated with higher BMI, in less developed countries the opposite was found (i.e., higher SES associated with higher BMI) (45). Future research would benefit from the comparison of studies conducted in economically developed and less developed countries, to consider the generalisability of the current findings. Further, the data from the current study are cross-sectional. Whilst the results provide evidence for association it is not possible to make causal inferences about the relationships reported. Notably, Hemmingsson's model predicts that the consequences of obesity exacerbate psychological distress and maladaptive eating behaviours, thus creating a vicious circle of negative affect and weight gain. Longitudinal studies which measure a range

of socio-economic, psychological, behavioural, social and environmental factors are needed to test the model in full to determine the temporal sequence of the variables of interest.

The high prevalence of obesity in many countries worldwide is a major concern and the development of effective intervention and preventive approaches is at the forefront of health agendas. The present study suggests an important role for psychological and emotional factors in eating behaviour and body weight regulation particularly for those of lower SES. Therefore weight management initiatives should encompass psychological factors alongside existing strategies such as the promotion of healthy eating messages and exercise promotion. Initiatives and interventions that target psychological distress and teach people to develop more positive coping strategies (e.g. problem-solving, positive help-seeking, relaxation techniques etc.) may be particularly effective. This is consistent with recent recommendations that tailored approaches are needed that meet the needs of the local population and consider the impact of wider socio-economic and community factors on obesity prevalence (46).

The present study shows that the relationship between SES and obesity may be partly explained by psychological distress and subsequent emotional eating as a coping strategy. Resilience was not found to be a protective factor in this relationship. Overall, these findings suggest that psychological interventions may play an important role in public health and weight management strategies particularly in lower SES populations.

Acknowledgements: The Authors would like to acknowledge the University of Liverpool Clinical Psychology Doctorate Programme.

Conflict of Interest: Charlotte Hardman and Paul Christiansen receive research funding from the American Beverage Association. Jade Spinosa, Valentina Lorenzetti and Joanne Dickson declare no conflicts of interest.

References

1. Ng M, Fleming T, Robinson M, Thomson B, Graetz N, Margono C, et al. Global, regional and national prevalence of overweight and obesity in children and adults during 1980 – 2013: a systematic analysis for the Global Burden of Disease Study 2013. *The Lancet*, 2014; 384 (9945): 766-781.
2. National Health Service Statistics on Obesity, Physical Activity and Diet [Web page]. <https://digital.nhs.uk/data-and-information/publications/statistical/statistics-on-obesity-physical-activity-and-diet/statistics-on-obesity-physical-activity-and-diet-england-2017>. Accessed September 6, 2018.
3. Government Office for Science. Foresight report: Tackling obesity: Future choices 2007. [Web page]. <https://www.gov.uk/government/collections/tackling-obesities-future-choices>. Accessed September 6, 2018.
4. Johansson K, Neovius M, Hemmingsson E. Effects of anti-obesity drugs, diet, and exercise on weight-loss maintenance after a very-low-calorie diet or low-calorie diet: a systematic review and meta-analysis of randomized controlled trials. *Am J Clin Nutr* 2014;99(1):14-23.
5. Parsons TJ, Power C, Logan S, Summerbell CD. Childhood predictors of adult obesity: a systematic review. *Int J Obes* 1999;23.
6. Sobal J, Stunkard AJ. Socioeconomic status and obesity: a review of the literature. *Psychol Bull* 1989;105(2):260.
7. Wardle J, Brodersen NH, Cole TJ, Jarvis MJ, Boniface DR. Development of adiposity in adolescence: five year longitudinal study of an ethnically and socioeconomically diverse sample of young people in Britain. *BMJ* 2006;332(7550):1130.
8. Wardle J, Waller J, Jarvis MJ. Sex differences in the association of socioeconomic status with obesity. *Am J Public Health Res* 2002;92(8):1299-304.

9. Stamatakis E, Primatesta P, Chinn S, Rona R, Falaschetti E. Overweight and obesity trends from 1974 to 2003 in English children: what is the role of socioeconomic factors? *Arch Dis Child* 2005;90(10):999-1004.
10. Bann D, Johnson W, Li L, Kuh D, Hardy R. Socioeconomic inequalities in childhood and adolescent body-mass index, weight, and height from 1953 to 2015: an analysis of four longitudinal, observational, British birth cohort studies. *The Lancet Public Health* 2018;3(4):194-203.
11. Larson NI, Story MT, Nelson MC. Neighborhood environments: disparities in access to healthy foods in the U.S. *Am J Prev Med* 2009;36(1):74-81.
12. Cobb LK, Appel LJ, Franco M, Jones-Smith JC, Nur A, Anderson CAM. The relationship of the local food environment with obesity: a systematic review of methods, study quality, and results. *Obes* 2015;23(7):1331-44.
13. Hemmingsson E. A new model of the role of psychological and emotional distress in promoting obesity: conceptual review with implications for treatment and prevention. *Obes Rev* 2014;15(9):769-79.
14. Lorant V, Croux C, Weich S, Deliège D, Mackenbach J, Anseau M. Depression and socio-economic risk factors: 7-year longitudinal population study. *Br J Psychiatry* 2007;190(4):293-8.
15. Power M, Uphoff E, Kelly B, Pickett KE. Food insecurity and mental health: an analysis of routine primary care data of pregnant women in the Born in Bradford cohort. *J Epidemiol Community Health* 2016; 71(4): 324-328.
16. Timms DWG. Social mobility and mental health in a Swedish cohort. *Soc Psychiatry Psychiatr Epidemiol* 1996;31(1):38-48.
17. Pickett KE, Wilkinson RG. Inequality: an underacknowledged source of mental illness and distress. *Br J Psychiatry* 2010;197(6):426-8.

18. Elliott I. Poverty and Mental Health: A review to inform the Joseph Rowntree Foundation's Anti-Poverty Strategy. *London: Mental Health Foundation* 2016.
19. Bratanova B, Loughnan S, Klein O, Claassen A, Wood R. Poverty, inequality, and increased consumption of high calorie food: Experimental evidence for a causal link. *Appetite* 2016;100:162-71.
20. Gibson EL. The psychobiology of comfort eating: implications for neuropharmacological interventions. *Behav Pharmacol* 2012;23(5-6):442-60.
21. Jääskeläinen A, Nevanperä N, Remes J, Rahkonen F, Järvelin M-R, Laitinen J. Stress-related eating, obesity and associated behavioural traits in adolescents: a prospective population-based cohort study. *BMC public health* 2014;14(1):321.
22. Laitinen J, Ek E, Sovio U. Stress-related eating and drinking behavior and body mass index and predictors of this behavior. *J Prev Med* 2002;34(1):29-39.
23. Hill DC, Moss RH, Sykes-Muskett B, Conner M, O'Connor DB. Stress and eating behaviors in children and adolescents: Systematic review and meta-analysis. *Appetite* 2018;123:14-22.
24. Boggiano MM, Wenger LE, Turan B, Tatum MM, Morgan PR, Sylvester MD. Eating tasty food to cope. Longitudinal association with BMI. *Appetite*. 2015;87:365-70.
25. Luthar SS, Cicchetti D. The construct of resilience: Implications for interventions and social policies. *Dev Psychopathol* 2000;12(4):857-85.
26. Fredrickson BL, Tugade MM, Waugh CE, Larkin GR. What good are positive emotions in crisis? A prospective study of resilience and emotions following the terrorist attacks on the United States on September 11th, 2001. *J Pers Soc Psychol* 2003;84(2):365.
27. Sharpley CF, Bitsika V, Jesulola E, Fitzpatrick K, Agnew LL. The association between aspects of psychological resilience and subtypes of depression: implications for focussed clinical treatment models. *Int J Psychiatry Clin Pract* 2016;20(3):151-6.

28. Stewart-Knox B, Duffy ME, Bunting B, Parr H, de Almeida MDV, Gibney M. Associations between obesity (BMI and waist circumference) and socio-demographic factors, physical activity, dietary habits, life events, resilience, mood, perceived stress and hopelessness in healthy older Europeans. *BMC Public Health* 2012;12(1):424.
29. Bolton Council [Web page].
<http://www.bolton.gov.uk/website/pages/Deprivationstatistics.aspx>. Accessed May 2, 2018
30. Clark HR, Goyder E, Bissell P, Blank L, Walters SJ, Peters J. A pilot survey of socio-economic differences in child-feeding behaviours among parents of primary-school children. *Public Health Nutr* 2008;11(10):1030-6.
31. Shavers VL. Measurement of socioeconomic status in health disparities research. *J Natl Med Assoc* 2007;99(9):1013-23.
32. Henry JD, Crawford JR. The short- form version of the Depression Anxiety Stress Scales (DASS- 21): Construct validity and normative data in a large non- clinical sample. *Br J Health Psychol* 2005;44(2):227-39.
33. Van Strien T, Frijters JER, Bergers GPA, Defares PB. The Dutch Eating Behavior Questionnaire for assessment of restrained emotional and external eating behavior. *The Int J Eat Disord* 1986;5:295-316.
34. Ng SP, Korda R, Clements M, Latz I, Bauman A, Bambrick H, et al. Validity of self-reported height and weight and derived body mass index in middle- aged and elderly individuals in Australia. *Aust N Z J Public Health* 2011;35(6):557-63.
35. Pursey K, Burrows TL, Stanwell P, Collins CE. How accurate is web-based self-reported height, weight, and body mass index in young adults? *J Med Internet Res* 2014;16(1):4.
36. Smith BW, Dalen J, Wiggins K, Tooley E, Christopher P, Bernard J. The brief resilience scale: assessing the ability to bounce back. *Int J Behav Med* 2008;15(3):194-200.

37. Mooy JM, De Vries H, Grootenhuis PA, Bouter LM, Heine RJ. Major stressful life events in relation to prevalence of undetected type 2 diabetes: the Hoorn Study. *Diabetes Care* 2000;23(2):197-201.
38. Fritz MS, Mackinnon DP. Required sample size to detect the mediated effect. *Psychol Sci* 2007;18(3):233-9.
39. Dodeen HM. Effectiveness of Valid Mean Substitution in Treating Missing Data in Attitude Assessment. *Assess Eval High Educ* 2003;28(5):505-13.
40. Hayes AF. PROCESS: A versatile computational tool for observed variable mediation, moderation, and conditional process modeling [White paper]. Retrieved from <http://www.afhayes.com/public/process2012.pdf>. 2012.
41. Ball K, Schoenaker DAJM, Mishra GD. Does psychosocial stress explain socioeconomic inequities in 9- year weight gain among young women? *Obes* 2017;25(6):1109-14.
42. Krause MR, Serlin RC, Ward SE, Rony RYZ, Ezenwa MO, Naab F. Testing Mediation in Nursing Research: Beyond Baron and Kenny. *Nurs Res* 2010;59(4):288-94.
43. Matthews K, A., Gallo L, C. Psychological Perspectives on Pathways Linking Socioeconomic Status and Physical Health. *Annu Rev Psychol* 2011;62(1):501-30.
44. Di Cesare M, Betham J, Stevens GA, and Markey O. Trends in adult body-mass index in 200 countries from 1975 to 2014: a pooled analysis of 1698 population-based measurement studies with 19.2 million participants. *The Lancet*, 2016; 387 (10026): 1377-1396.
45. Pampel FC, Denney JT, Krueger PM. Obesity, SES, and economic development: A test of the reversal hypothesis. *Soc Sci Med* 2012;74(7):1073-81.

46. Local Government. Tackling obesity. Local government's new public health role. 2013 [Web page] <https://www.local.gov.uk/sites/default/files/documents/tackling-obesity-local-go-1cf.pdf>. Accessed July 2, 2018.

Figure Legend

Figure 1. Serial multiple mediation analysis with socio-economic status as the independent variable (IV), BMI as the dependent variable (DV), and psychological distress and emotional eating as the first and second mediators, respectively. Values are unstandardized regression coefficients (SEs in parentheses) and associated p-values. Bracketed association = direct effect (controlling for indirect effects). Solid lines indicate significant pathways and dashed lines indicate non-significant pathways.




Figure 1. Serial multiple mediation analysis with socio-economic status as the independent variable (IV), BMI as the dependent variable (DV), and psychological distress and emotional eating as the first and second mediators, respectively. Values are unstandardized regression coefficients (SEs in parentheses) and associated p-values. Bracketed association = direct effect (controlling for indirect effects). Solid lines indicate significant pathways and dashed lines indicate non-significant pathways.

Table 1. *Sample descriptives and questionnaire scores (N = 150)*

	Mean	Standard deviation	Range
Age in years	35.35	10.90	18 - 65
BMI kg/m ²	26.31	6.00	16.3 - 45.2
DASS	15.06	11.42	0 - 56
LES	6.14	6.39	0 - 38
DEBQ	2.58	1.02	1 - 5
Resilience	3.25	0.80	1 - 5
Highest education level ^a	5.15	1.87	1 - 8
Yearly household income ^b	5.67	2.46	1 - 9

Note. BMI = Body Mass Index, DASS = Depression Anxiety Stress Scale, LES = Life Events Scale, DEBQ = Dutch Eating Behaviour Questionnaire (Emotional Eating subscale).

^a 8-point scale: 1 = none, 2 = GCSE grade D or below, 3 = GCSE grade C or above, 4 = A-level or equivalent, 5 = University degree or equivalent, 6 = Postgraduate qualification or equivalent, 7 = Masters or equivalent, 8 = PHD or equivalent.

^b 9-point scale: 1 = < £5 200, 2 = £5 200 - £10 399, 3 = 10 400 - £15 599, 4 = £15 600 - £20 799, 5 = £20 800 - £25 999, 6 = £26 000 - £36 399, 7 = £36 400 - £51 999, 8 = £52 000 - £77 999, 9 = >£78 000.

Table 2. Pearson's correlation coefficients (r) between SES, questionnaire measures, and BMI.

	1	2	3	4	5	6
1. SES	-					
2. DASS	-0.34**	-				
3. DEBQ	0.18*	0.26**	-			
4. Resilience	0.14	-0.49**	-0.13	-		
5. BMI	-0.06	0.04	0.33**	0.05	-	
6. LES	-0.16	0.21*	0.07	-0.16	0.20*	-

Note. * $p < .05$ ** $p < .01$. SES = Socio-economic status composite score, DASS = Depression Anxiety Stress Scale, DEBQ = Dutch Eating Behaviour Questionnaire (Emotional Eating sub-scale), BMI = Body Mass Index, LES = Life Events Scale. Higher scores on the DASS indicate higher emotional distress, higher scores on the DEBQ indicate higher levels of emotional eating, higher scores on the Resilience scale indicate higher levels of resilience, higher scores on the LES indicate a greater number of stressful life events.

Table 3. Moderated mediation by resilience of indirect effect of SES on emotional eating via psychological distress

	Effect (<i>SE</i>)	95% CI
Low resilience	-.08 (.04)	[-.18, -.03]
Medium resilience	-.10 (.04)	[-.19, -.04]
High resilience	-.11 (.04)	[-.22, -.04]

Note. Resilience scores; Low = -1 *SD*, Medium = mean, High = +1 *SD*.




Figure S1. Boxplots to illustrate the data for highest education level (left panel) and yearly household income (right panel). Highest education level used an 8-point scale (1 = none, 2 = GCSE grade D or below, 3 = GCSE grade C or above, 4 = A-level or equivalent, 5 = University degree or equivalent, 6 = Postgraduate qualification or equivalent, 7 = Masters or equivalent, 8 = PHD or equivalent). Yearly current household income used a 9-point scale (1 = <£5 200, 2 = £5 200 - £10 399, 3 = 10 400 - £15 599, 4 = £15 600 - £20 799, 5 = £20 800 - £25 999, 6 = £26 000 - £36 399, 7 = £36 400 - £51 999, 8 = £52 000 - £77 999, 9 = >£78 000).

Additional data analysis: Mediation with the addition of Life events as a first mediator.




Figure S2. Serial multiple mediation analysis with socio-economic status as the independent variable (IV), BMI as the dependent variable (DV), and life events, psychological distress and emotional eating as the first, second and third mediators, respectively. Values are unstandardized regression coefficients (SEs in parentheses) and associated p-values. Bracketed association = direct effect (controlling for indirect effects).