

Psychoradiological patterns of small-world properties and a systematic review of connectome studies of patients with 6 major psychiatric disorders

Xueling Suo^{a,1} PhD, Du Lei^{b,1} PhD, Lei Li^{a,1} MD, Wenbin Li^a MM, Jing Dai^c MD, Song Wang^c PhD, Manxi He^c MD, Hongyan Zhu^d MD, PhD, Graham J Kemp^e DSc, Qiyong Gong^{a, f, *} MD, PhD.

Author Affiliations

^aHuaxi MR Research Center (HMRRC), Department of Radiology, West China Hospital of Sichuan University, Chengdu, Sichuan, 610041 China

^bDepartment of Psychosis Studies, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, SE5 8AF United Kingdom

^cDepartment of Psychoradiology, Chengdu Mental Health Center, Chengdu, Sichuan, 610041 China

^dLaboratory of Stem Cell Biology, State Key Laboratory of Biotherapy, West China Hospital of Sichuan University, Chengdu, Sichuan, 610041 China

^eLiverpool Magnetic Resonance Imaging Centre (LiMRIC) and Institute of Ageing and Chronic Disease, University of Liverpool, Liverpool, L69 3GE United Kingdom

^fDepartment of Psychology, School of Public Administration, Sichuan University, Chengdu, Sichuan, 610041 China

¹ Xueling Suo, Du Lei and Lei Li contributed to this work equally.

***Correspondence to:** Prof. Qiyong Gong, Huaxi MR Research Center (HMRRC), Department of Radiology, West China Hospital, Sichuan University, No. 37 Guo Xue Xiang, Chengdu, Sichuan 610041, China. E-mail: qiyonggong@hmrrc.org.cn; Tel: 086-028 81812593; Fax: 086-028 85423503

Word count for the text: 6315

Word count for the abstract: 277

Abstract

Background: Brain connectome research based on graph theoretical analysis shows that “small-world” topological properties play an important role in the structural and functional alterations observed in patients with psychiatric disorders. However, the reported global topological alterations in small-world properties are controversial, are not consistently conceptualized according to agreed-upon criteria, and are not critically examined for consistent alterations in patients with each major psychiatric disorder. Here, we describe the main patterns of altered small-world properties and then systematically review the evidence for these alterations in the structural and functional connectomes in patients with 6 different major psychiatric disorders.

Methods: Based on a comprehensive PubMed search we selected 40 studies of schizophrenia, 33 studies of depression, 5 studies of attention-deficit/hyperactivity disorder, 5 studies of bipolar disorder, 7 studies of obsessive compulsive disorder and 5 studies of posttraumatic stress disorder using non-invasive neuroimaging data and graph theory approaches.

Results: The following 4 patterns of altered small-world properties are defined from the perspectives of segregation and integration: ‘regularization’, ‘randomization’, ‘stronger small-worldization’ and ‘weaker small-worldization.’ Although more differences than similarities are noted in patients with these disorders, one prominent trend is the structural regularization versus functional randomization in patients with schizophrenia.

Limitations: Differences in demographic and clinical characteristics, preprocessing steps, and analytical methods can produce contradictory results, increasing the difficulty of integrating results across different studies.

Conclusions: Four patterns of altered small-world properties are proposed. The analysis of altered small-world properties may provide novel insights into the pathophysiological mechanisms underlying psychiatric disorders from a connectomic perspective. In future connectome studies, the global network measures of both segregation and integration should be calculated to fully evaluate altered small-world properties in patients with a particular disease.

1. Introduction

1.1. The need for a systematic approach to examine connectome abnormalities

Recent advances in neuroscience methodology have underpinned a large increase in human brain research studies. In particular, by taking advantage of non-invasive imaging approaches, the developing field of psychoradiology (<https://radiopaedia.org/articles/psychoradiology>) promises not only insights into abnormal brain function and underlying neuropsychopathologies, but also potential clinical utility in patients with mental illness, from diagnosis to planning and monitoring therapeutic interventions.¹ Psychoradiological studies have revealed abnormal brain networks in patients with several major psychiatric disorders.²⁻⁵

However, reported neuroimaging studies to date have sometimes neglected the interactions among multiple brain regions. Widespread and subtle pathology across the brain may be only imperfectly detected using a traditional regional or voxel-based analysis.⁶ In contrast, a large-scale network analysis enables the assessment of overall connectivity patterns among all regions of the brain. This approach has been stimulated by pioneering studies modeling the brain as a complex network in the so-called ‘connectome’.⁷ In these terms, the human brain network exhibits topological properties known as ‘small-worldness’, an optimal balance between the segregation and integration of information processing procedures.⁸ We suggest that a clear definition of the main patterns of altered small-world properties might yield new insights into how these processes are perturbed in patients with pathological brain conditions, with potential benefits to diagnosis and therapy.

Graph theory provides a rigorous mathematical framework to analyze the topology

of complex networks.⁹ Graph theoretical approaches have been widely applied to connectome studies to investigate the topological organization of large-scale functional and structural brain networks in healthy and disease states.¹⁰⁻¹³ Brain disorders clearly cause significant alterations in global network topology;¹⁴ however, connectome studies have often reported inconsistent quantitative accounts of these. Occasionally, similar alterations have been described in terms of different patterns of change. For example, two functional connectome studies in patients with schizophrenia reported a decreased clustering coefficient (C_p) and an increased path length (L_p) compared to healthy controls (HC).^{15, 16} However, one study interpreted this result as a tendency toward more random organization,¹⁵ whereas the other study described the result as what we will call here ‘weaker small-worldization’, i.e., a change toward less-marked small-world properties.¹⁶ Conversely, different topological alterations may be interpreted as showing the same pattern. For example, two functional studies of patients with schizophrenia reported a tendency toward a more random organization^{15, 17}; however, one study observed an increased L_p and decreased C_p ,¹⁵ whereas the other study noted a decreased L_p (marginally group difference) and decreased C_p .¹⁷ Definitions of unified criteria for characterizing small-world properties in terms of global topological parameters are urgently needed.

Furthermore, functional and structural networks may show discrepancies in alterations in small-world properties in patients with the same disorder. For example, in patients with schizophrenia, increased C_p and L_p have been reported in the structural connectome,¹⁸ but decreased C_p and L_p in the resting-state functional connectome.¹⁵ Meanwhile, other studies on the structural and functional connectomes have reported consistent alterations in small-world properties, e.g., decreased global efficiency (E_{glob}) and a maintained local efficiency (E_{loc})^{19, 20} and increased C_p and

L_p .^{18, 21} Clearly, further studies are required to explore whether there are potential common trends in functional and structural networks.

Several reviews have recently summarized findings for both the functional and structural connectomes in patients with psychiatric disorders.^{2, 22-25} However, to date, no study has reviewed the clinically altered small-world properties of the human brain connectome from the perspectives of information segregation and integration. In this review, we explore the main patterns of altered small-world properties using a graph theoretical analysis, and then summarize recent findings from studies of patients with several major psychiatric disorders.

First, we introduce the construction of the brain connectome, connectome measures, and network types (for additional details, please refer to the study by Liao et al.²⁵).

1.2. Rationale for brain connectome construction

In graph theory, a brain network is defined as a graph $G(N, K)$, with N and K representing the number of nodes and edges, respectively. Modern brain mapping techniques include structural magnetic resonance imaging (MRI), functional MRI (fMRI), electroencephalography (EEG) and magnetic-encephalography (MEG); we did not consider the latter two techniques, in order to reduce the confounding effects of heterogeneous image acquisition techniques.

Brain nodes are often defined based on anatomical landmarks, functional activation,²⁶⁻²⁸ and regional connectivity profiles.²⁹ Anatomical landmarks include the automated anatomical labeling (AAL) atlas, Harvard-Oxford atlas and Desikan-Killiany atlas; the AAL atlas has also been used,³⁰ but different investigators have used different numbers of regions. Some researchers have used software tools such as FreeSurfer to parcellate the cortex into a given number, x , of nodes. The majority of

templates have constructed brain networks with < 1000 nodes.³¹ Independent component analysis (ICA) is the most popular of the many data-driven analyses for delineating resting-state brain functional networks. The number of components identified using the ICA approach is typically less than a hundred for a whole-brain analysis.²⁶ There is no gold standard for node parcellation. Researchers typically choose network nodes based on anatomic templates for convenience. The use of an anatomical atlas is known to introduce potential bias because of its inhomogeneity, and a reasonable approach used in most neuroimaging studies is to employ a single atlas for analysis.³² A recent functional study investigating the impact of parcellation choice on group differences found that patient-specific global network properties are robustly observed using different parcellation schemes, whereas graph metrics characterizing impairments of individual nodes vary considerably.³³ Overall, the correct definition of nodes in connectome studies is still an open question that needs further technical investigation.

Once the nodes have been determined, the edges are defined. The edges of the brain functional connectome are constructed by calculating the Pearson, partial, or wavelet correlation for every possible pair of regional time series among the nodes. The Pearson correlation measures the interdependence between two time series. However, correlation may result from indirect relationships with common sources. To avoid this problem, in the partial correlation approach the brain activity in one region is correlated with that in another region after the activity in all other regions is regressed out; this attempts to remove the effects of indirect paths³⁴ and provides a better estimate of the true macroscopic functional connectome.³⁵ However, this strategy has the opposite problem of losing real connections, which is likely to underestimate topological measures.³⁶ The best recent evidence is that Pearson correlation is more

valid and reliable.³⁷

After construction, the edges of the brain structural connectome can be measured in two ways. The first, using structural data directly, is by computing the chosen (partial or Pearson) correlation coefficients of the morphological features of the brain (cortical thickness or gray matter volume) among nodes. A more advanced approach is by reconstructing diffusion MRI-traced white matter (WM) using either deterministic or probabilistic tractography. The tracking procedure used in deterministic tractography terminates when it reaches regions with fiber crossings,³⁸ and this tends to reduce sensitivity. Probabilistic tractography overcomes this shortcoming, and offers a natural approach to modeling uncertainty by generating multiple curves originating from a seed point,³⁹ providing a more comprehensive picture of connectivity. The probabilistic algorithm has better test–retest reliability and is more robust than deterministic tractography in areas of high uncertainty (notably fiber crossings).^{40, 41} However, the lower computational cost of deterministic tractography makes it more often used in clinical applications. High-field imaging and the application of advanced diffusion reconstruction algorithms offer the potential to characterize more complex fiber patterns when edges are defined.

Once the nodes and edges are defined, a matrix representation is generated where each entry records the edge weight of corresponding node pairs denoted by the row and column index of the matrix. The reconstructed network is classified as the directed or undirected type, depending on whether the edges display directionality. Depending on whether the edges are assigned with different strengths, a binary network is obtained by thresholding the weights of the edges, where the edges are either 1 (connected) or 0 (unconnected); otherwise, the network is a weighted type.

1.3. Brain connectome measures

Once the brain connectome is constructed, the connectome measurement is calculated. In graph theory, a complex network is characterized by two types of measures, integration and segregation, which represent crucial information processing patterns of the brain and ensure efficient global communication and functional specialization.⁴²

1.3.1. Integration measures: L_p , λ and E_{glob}

Specifically, topological integration refers to the efficiency of global information communication or the ability to integrate distributed information in the network, which is measured by the parameters L_p , λ and E_{glob} . The characteristic path length L_p is calculated by averaging the minimum number of connections that link any paired nodes in the network.⁴³ The normalized characteristic path length λ is the normalized L_p , which is calculated for the mean L_p of 100 matched random networks that preserve the same number of nodes and edges as the real network. The global efficiency E_{glob} measures how efficiently information is exchanged at the global level.⁴⁴

1.3.2. Segregation measures: C_p , γ and E_{loc}

In contrast, segregation refers to the ability of densely interconnected groups of brain regions to perform specialized processing procedures and is measured by the parameters C_p , γ and E_{loc} . The network clustering coefficient C_p is calculated by averaging the C_p over all nodes in the network: C_p is equivalent to the fraction of the node's neighbors that are also neighbors of each other.⁴³ The normalized clustering coefficient γ is the normalized C_p , which is calculated for the mean C_p of 100 matched random networks that preserve the same number of nodes and edges as the real network. Local efficiency, E_{loc} , measures how efficiently information is exchanged at

the local level.⁴⁴ The level of clustering measured by E_{loc} expresses the level of local connectedness of a network, with high levels of clustering interpreted as high levels of local organization of the network.²²

1.3.3. Small-world index σ

The small-world index σ is γ / λ , which is > 1 when the small-world fulfills the conditions of $\gamma > 1$ and $\lambda \approx 1$.⁴³ Graph measures are discussed in more detail elsewhere.⁴²

1.4. Brain network types: small-world, regular and random networks

Based on the perspectives of segregation and integration, networks are divided into three main types: small-world networks (high C_p and short L_p), regular networks (high C_p and long L_p) and random networks (low C_p and short L_p). A regular network is characterized by a high C_p (the probability that neighboring nodes are also interconnected with other neighboring nodes) and a long L_p (the average distance from one node to any other node in the network, expressed as the number of links that must be traveled). A random network is characterized by a low C_p and a short L_p . A small-world network has a higher C_p than but a similar L_p to a random network, and has both high global and local information transformation capacities and maintains an optimized balance between segregation and integration. Quantitatively, the small-world index σ should be > 1 for small-world networks, as measured by both $\gamma > 1$ and $\lambda \approx 1$.

Although connectome studies have confirmed that the brain networks of patients and HCs all exhibit small-world topology ($\sigma > 1$ with $\gamma > 1$ and $\lambda \approx 1$), significant group differences in global network measures have been reported, implying different types of alterations in small-world properties in patients with different disorders. However,

to date, the conceptualization of these altered small-world properties has varied, with no agreed definitive criteria.

1.5. The aims of this review

This review is designed to address the following questions: (1) Can unified criteria be defined for the altered small-world properties? (2) What are the consistently altered small-world properties in patients with specific psychiatric disorders? (3) Are common alterations observed between functional and structural networks? Specifically, we first propose 4 defined patterns of altered small-world properties, and then systematically review published connectome studies to identify the most consistently altered small-world properties in patients with one of 6 specific psychiatric disorders [schizophrenia, depression, ADHD (attention-deficit/hyperactivity disorder), BD (bipolar disorder), OCD (obsessive compulsive disorder), and PTSD (posttraumatic stress disorder)]. Structural and functional studies are treated separately to assess their consistency and evaluate any discrepancies.

2. Methods

2.1. Study selection

An online search of the PubMed database was conducted using the following search terms: [(schizophrenia) or (depression) or (attention-deficit/hyperactivity disorder or ADHD) or (bipolar disorder or BD) or (obsessive compulsive disorder or OCD) or (posttraumatic stress disorder or PTSD)] and (graph analysis or graph theory or small world or connectome). Nine hundred forty-three studies published before December 1, 2016 were screened. The exclusion criteria were 1) studies that did not examine global properties (C_p , γ , E_{loc} , L_p , λ , E_{glob} and σ); 2) studies that did not compare patients with one of the aforementioned psychiatric disorders with HCs; and 3)

studies using EEG or MEG.

In addition to this online search, we manually cross-referenced the studies with the bibliographies of recent reviews in the field to ensure that no studies of significance were omitted from the review,^{2, 23, 45} after which 1 additional paper was included.⁴⁶ Ninety-five papers reporting global properties were finally included in the current review, including 40 studies of schizophrenia, 33 studies of depression, 5 studies of ADHD, 5 studies of BD, 7 studies of OCD and 5 studies of PTSD.

2.2. Patterns of altered small-world properties

Based on the perspectives of segregation (C_p , γ and E_{loc}) and integration (L_p , λ and E_{glob}), we propose 4 patterns of altered small-world properties (**Table 1 and Figure 1**): regularization, in which the network transforms from a small-world network to a relatively regular network; randomization, in which the network transforms from a small-world network to a relatively random network; stronger small-worldization, in which the network transforms from a small-world network to a relatively stronger small-world network; and weaker small-worldization, in which the network transforms from a small-world network to a relatively weaker marked small-world network.

Regularization was defined where at least one altered measurement of segregation (increased C_p , γ or E_{loc}) and/or at least one altered measurement of integration (decreased E_{glob} or increased L_p or λ) were present. Randomization was defined where at least one altered measurement of segregation (decreased C_p , γ or E_{loc}) and/or at least one altered measurement of integration (increased E_{glob} or decreased L_p or λ) were present. Stronger small-worldization was defined where at least one altered measurement of segregation (increased C_p , γ or E_{loc}) and at least one altered

measurement of integration (increased E_{glob} or decreased L_p or λ) were present. Weaker small-worldization was defined where at least one altered measurement of segregation (decreased C_p , γ or E_{loc}) and at least one altered measurement of integration (decreased E_{glob} or increased L_p or λ) were present.

3. Results

3.1. Methodological review of connectome studies

Ninety-five papers reporting on the global properties of functional and structural connectomes were included in our systematic review (**Tables 2-4**). Fifty-three papers used AAL, 7 used the Desikan-Killiany atlas, 6 used the Harvard-Oxford atlas, 1 used the Destrieux atlas, 1 used a spatially unbiased infratentorial template, 1 used a voxelwise approach, and 11 used FreeSurfer to parcellate the cortex into diverse numbers of nodes. The brain nodes in 14 papers were defined based on functional activation, 11 of which performed an ICA. The brain nodes in 1 paper were defined based on the regional connectivity profiles.

Twenty-nine, 11, and 6 papers used the Pearson, partial, and wavelet correlations of the regional time series, respectively, to construct the edges of the functional connectome. One paper estimated functional connectivity in the brain using the higher-order statistical dependence. One paper used the IMAGES algorithm to detect functional connections. In 1 paper, functional networks were built using the correlations among ICA time courses of brain nodes. For the edges of structural connectome of morphological features in the brain, 3 and 2 papers, respectively, computed the Pearson and partial correlations of cortical thickness; 3 papers calculated the Pearson correlations of gray matter volume. For the structural connectome reconstructing diffusion MRI data, 30 and 7 papers, respectively,

performed deterministic and probabilistic tractography. One paper used the generalized q-sampling imaging (GQI) to construct the edges.⁴⁷

3.2. Global topological alterations in patients with schizophrenia

Forty-seven datasets from 22 functional and 18 structural connectome studies were included, of which the majority (26) used fMRI, followed by 20 studies using diffusion MRI, and 1 study using structural MRI (**Table 2**). Of the functional studies, 2 recruited subgroups of patients, 1 recruited patients with familial and sporadic schizophrenia and analyzed the data with and without global signal regression (GSR),⁴⁸ and the other study recruited patients with and without auditory verbal hallucinations (AVH).¹⁶ One study examined global properties during a task and at rest,¹⁵ another study investigated the networks of the left and right hemispheres.⁴⁹ One study explored both the structural and functional connectomes.¹⁶

3.2.1. Functional connectome studies in patients with schizophrenia

Fourteen studies showed a shift toward randomization,^{15, 17, 48-57} including 7 reports of decreased segregation, and 7 reports of decreased segregation and increased integration. Specifically, in the study by Ma et al. (2012), patients with schizophrenia showed decreased segregation (decreased C_p) and increased integration (decreased L_p) at rest.¹⁵ Yu et al. (2011) reported decreased segregation (decreased γ) in the right hemisphere task-related network.⁴⁹ Zhu et al. (2016) observed decreased segregation (decreased γ and E_{loc}) with GSR, and decreased segregation (decreased γ and E_{loc}) and increased integration (decreased λ) without GSR, in patients with familial schizophrenia, as well as decreased segregation (decreased E_{loc}) in patients with sporadic schizophrenia without GSR.⁴⁸

Seven datasets from 6 studies showed a tendency toward weaker small-worldization

with decreased segregation and integration.^{15, 16, 49, 58-60} Specifically, patients with schizophrenia showed decreased segregation (decreased C_p) and decreased integration (increased L_p) during the auditory oddball (AOD) task,¹⁵ and decreased segregation (decreased γ) and decreased integration (decreased E_{glob} and increased L_p) in the left hemisphere task-related network.⁴⁹ Patients with schizophrenia with and without AVH showed decreased segregation (decreased C_p and E_{loc}) and decreased integration (decreased E_{glob} and increased L_p).¹⁶

Four studies showed a tendency toward regularization,^{20, 21, 61, 62} including 3 reports of both increased segregation and decreased integration, and 1 report of decreased integration. One study showed a tendency toward stronger small-worldization, with increased segregation (increased C_p) and increased integration (increased E_{glob}).⁶³ Two studies, however, did not report any significant difference in global properties.^{48, 64}

3.2.2. Structural connectome studies in patients with schizophrenia

Twelve studies showed a tendency toward regularization,^{16, 18, 19, 65-73} including 11 reports of decreased integration and 1 report of both decreased integration and increased segregation. Five studies showed a tendency toward weaker small-worldization, with decreased segregation and integration.^{71, 74-77} Specifically, of the 2 datasets in the study by van den Heuvel et al. (2013), one dataset showed a tendency toward regularization with decreased integration (decreased E_{glob}), and the other independent dataset showed a tendency toward weaker small-worldization with decreased segregation (decreased C_p) and decreased integration (decreased E_{glob}) in patients with schizophrenia.⁷¹ One study showed a tendency toward randomization with increased integration.⁷⁸ Three studies, however, did not report any significant difference in global properties.^{16, 79, 80}

3.2.3. A summary of altered small-world properties in patients with schizophrenia

Eight hundred and eighteen patients with schizophrenia and 791 HCs, along with 28 datasets from 22 functional studies, as well as 945 patients with schizophrenia and 912 HCs, along with 21 datasets from 19 structural studies, were included. In structural MRI studies, the largest proportion (12 of 21 datasets) of altered small-world properties included regularization in 675 of 945 patients, but in fMRI studies, randomization was observed more frequently (14 of 28 datasets, 390 of 818 patients). Most patients with schizophrenia were currently taking medications; only three datasets comprised drug-naïve and medication-washout patients, and in five datasets medication status was not available. For the datasets including currently medicated patients with schizophrenia, the most prominent trend was still structural regularization (10 of 16 datasets) rather than functional randomization (11 of 25 datasets).

3.3. Global topological alterations in patients with depression

Thirty-four datasets from 16 functional and 17 structural connectome studies were included, and the majority (17) used fMRI, followed by 12 DTI studies and 5 structural MRI studies (**Table 3**). One of the fMRI studies recruited patients with late-life depression (LLD) with and without comorbid amnesic mild cognitive impairment (aMCI).⁸¹

3.3.1. Functional connectome studies in patients with depression

Four studies showed a tendency toward randomization, including 2 reports of increased integration,^{13, 82} and 2 reports of decreased segregation.^{81, 83} Specifically, one dataset from the study by Li et al. (2015) showed decreased segregation

(decreased E_{loc}) in patients with LLD.⁸¹ Five studies showed a tendency toward regularization, including 3 reports of decreased integration characterized by increased L_p ^{84, 85} or decreased E_{glob} and increased L_p ,⁸⁶ 1 report of increased segregation (increased E_{loc}),⁸⁷ and 1 report of increased segregation (increased E_{loc}) and decreased integration (decreased E_{glob}).⁸⁸

Three studies showed a tendency toward weaker small-worldization with decreased segregation and integration, including decreased C_p with decreased E_{glob} and increased L_p ,⁸⁹ and decreased E_{loc} with decreased E_{glob} and increased L_p .^{81, 90} Four studies, however, reported no significant differences.^{46, 91-93} One study showed a decreased σ .⁹⁴

3.3.2. Structural connectome studies in patients with depression

Four studies showed a tendency toward regularization, including 1 report of increased segregation (higher transitivity used as an alternative to C_p),⁹⁵ 1 report of decreased integration (decreased E_{glob} and increased L_p),⁹⁶ 2 reports of both increased segregation and decreased integration characterized by an increased γ with a decreased E_{glob} and increased λ ,⁹⁷ and increased C_p and E_{loc} with increased L_p .⁴⁷

Two studies showed a shift toward randomization with decreased segregation characterized by a decreased γ and E_{loc} ,⁹⁸ as well as a decreased C_p .⁹⁹ Two studies showed a tendency toward weaker small-worldization with decreased segregation and decreased integration: one study showed a decreased γ and increased λ ,¹⁰⁰ and the other study showed a decreased C_p and E_{loc} along with a decreased E_{glob} .¹⁰¹ One study showed a tendency toward stronger small-worldization with increased segregation (increased E_{loc}) and increased integration (increased E_{glob} and decreased L_p and λ).¹⁰² Eight studies did not report significant differences.¹⁰³⁻¹¹⁰

3.3.3. A summary of altered small-world properties in patients with depression

In summary, global topology was not significantly altered in 12 reports, followed by 9 studies showing regularization, 6 showing randomization, 5 showing weaker small-worldization, and 1 showing stronger small-worldization. In one study, the global network alterations, which depend on one single measurement, σ (see Discussion), were difficult to classify. In sixteen datasets, patients with depression were currently taking medicine, 14 datasets comprised drug-naïve and medication-washout patients, and medication status was not available in four datasets. No prominent trends in altered small-world properties were observed either in the medicated patients or in the medication-free patients.

3.4. Global topological alterations in patients with ADHD

Only 5 studies, 3 using fMRI and 2 using DTI, were included (**Table 4**).

Of the fMRI studies, one study showed a shift toward regularization by increased segregation (increased E_{loc}) in patients with ADHD.¹¹¹ One additional fMRI study reported decreased segregation (decreased E_{loc}) in the visual attention network of children with ADHD during a task, suggesting a shift toward randomization.¹¹² Another fMRI study reported no significant between-group differences in C_p , γ , L_p , or λ .¹¹³

The WM networks in boys with ADHD exhibited decreased integration (decreased E_{glob} and increased L_p and λ),¹¹⁴ showing a shift toward regularization. A WM study in adults with ADHD reported no significant between-group differences in C_p , γ , λ , or E_{glob} .¹¹⁵

3.5. Global topological alterations in patients with BD

Only 5 studies, 3 using fMRI and 2 DTI, were included (**Table 4**).

In the fMRI studies, patients with BD type I showed decreased integration (increased L_p) during a task, indicating a shift toward regularization.⁸⁴ Two resting-state fMRI studies included patients with BD types I and II, one suggesting a shift toward regularization characterized by increased segregation (increased C_p) and decreased integration (decreased E_{glob}),¹¹⁶ and the other showing a lack of significant between-group differences in C_p , L_p , E_{loc} or E_{glob} .⁹²

Two WM network studies included patients with BD type I; one study showed a shift toward regularization characterized by decreased integration (decreased E_{glob}),¹¹⁷ and the other study suggested a shift toward weaker small-worldization characterized by decreased segregation (decreased C_p) and decreased integration (decreased E_{glob} and increased L_p).¹¹⁸

3.6. Global topological alterations in patients with OCD

Seven studies, including 4 fMRI and 3 structural MRI studies, were included (**Table 4**).

Of the 4 fMRI studies, 1 study found increased segregation (increased γ) of the top-down control network in patients with OCD, suggesting a shift toward regularization.¹² Another 2 fMRI studies showed a tendency toward randomization in patients with OCD, one study (focusing on treatment) reported decreased segregation (decreased γ and E_{loc}) at baseline,¹¹⁹ and the other study reported decreased segregation (decreased γ) and increased integration (decreased λ) in the orbitofrontal-striato-thalamic circuit.¹²⁰ Decreased σ has been reported in children with OCD.¹²¹

Of the 2 WM network studies, one study reported increased segregation (increased γ) and decreased integration (decreased E_{glob} and increased L_p and λ), indicating a

tendency toward regularization,¹²² while the other study reported no significant differences in λ or γ .¹²³ One cortical thickness study showed no significant between-group differences in E_{glob} or E_{loc} .¹²⁴

3.7. Global topological alterations in patients PTSD

Only 5 studies, including 2 fMRI and 3 structural MRI studies, were included (**Table 4**).

Of the 2 fMRI studies, one study reported increased segregation (increased C_p and E_{loc}) and increased integration (increased E_{glob} and decreased L_p) in adults with PTSD, showing a shift toward stronger small-worldization,¹¹ whereas the other study reported increased segregation (increased C_p and E_{loc}) and decreased integration (increased λ) in children with PTSD, suggesting a shift toward regularization.¹²⁵

Of the 2 WM studies, one study reported increased integration (decreased L_p and λ) in adults with PTSD, indicating a shift toward randomization,¹²⁶ whereas the other study reported a shift toward weaker small-worldization in children with PTSD characterized by decreased segregation (decreased E_{loc}) and decreased integration (decreased E_{glob} and increased L_p).¹²⁷ In an additional structural network study of gray matter, decreased σ was reported in patients with PTSD.¹²⁸

4. Discussion

Our systematic review highlights several important findings in brain connectome research in patients with psychiatric disorders.

4.1. Summary of the approaches and findings

Based on the perspectives of information segregation and integration, we explicitly propose 4 patterns of altered small-world properties: regularization, randomization, stronger small-worldization and weaker small-worldization. According to these

criteria, the most prominent trend in the psychiatric studies reviewed here is the regularization of the structural connectome with randomization of the functional connectome in patients with schizophrenia (**Figure 2**). However, no consistent alterations were observed in patients with depression, ADHD, BD, OCD or PTSD.

4.2. Methodology

Brain connectome research based on graph theoretical analysis has proven that ‘small-world’ topological properties are important in structural and functional alterations in the brains of patients with psychiatric disorders. However, the reported alterations of small-world topological properties in previous connectome studies are controversial and often contradictory (see the Introduction), which may be partially due to the lack of unified criteria for conceptualizing these alterations. Here, we have attempted to address this issue from the perspectives of segregation and integration to classify altered small-world properties into 4 patterns: regularization, randomization, stronger small-worldization and weaker small-worldization. In future studies we recommend comprehensive calculation of the global network measures of segregation (C_p , γ , and E_{loc}) and integration (L_p , λ , and E_{glob}). Additionally, the small-world index σ should be calculated to define the small-world network: σ is calculated as γ / λ , which is > 1 for a small-world organization that only fulfills the conditions of $\gamma > 1$ and $\lambda \approx 1$. An important technical point is that reporting $\sigma > 1$ is not sufficient to prove small-worldness:^{94, 121, 128} A regular network with a large γ (e.g., $\gamma = 3$) and a long λ (e.g., $\lambda = 2$) is possible, which together yield $\sigma > 1$.¹²⁹ In other words, regular networks may have $\sigma > 1$, and therefore, both $\gamma > 1$ and $\lambda \approx 1$ are required to prove small-worldness.

4.3. Findings from patients with major psychiatric disorders

We systematically reviewed all current reports examining altered small-world properties in both the structural and functional connectomes of patients with one of 6 major psychiatric disorders. The prominent trend is the structural regularization and functional randomization in patients with schizophrenia. In the structural connectome, large proportions of the reported connectomes were constructed using diffusion MRI, and most node and edge definitions used the AAL 90 template and deterministic tractography to construct weighted networks. In contrast, in the functional connectome, obviously consistent definitions for nodes and edges were not used. In addition, neural activity is shaped, but not defined, by the underlying anatomy,¹⁶ as relatively fixed structural organizations produce diverse functional network patterns.¹³⁰ Nevertheless, structural connections constrain and shape the diverse patterns of functional connections, while the functional connection patterns reflect the architecture of structural connections.¹³¹

Patients with depression did not display consistent alterations in small-world properties. A substantial proportion of alterations preserved global organization, rather than the falling into one of the 4 patterns defined here. Where there were significant between-group differences in the global network measures, most studies on the structural connectome in patients with depression seems to indicate a trend toward regularization. Our attempt to identify consistent altered small-world properties might be limited by the heterogeneity of the patient samples (e.g., age of the patients). An exploratory subgroup analysis of patients with MDD revealed that the structural connectome in patients with LLD^{95-97, 99, 104, 105} seemed to show a shift toward regularization.⁹⁵⁻⁹⁷

To date, there is too little brain connectome research in patients with ADHD, BD, OCD and PTSD for us to infer consistent alterations in small-world properties. In

patients with BD type I, the brain networks appear to show a trend toward regularization. Future research is needed to confirm this speculation.

Several potential factors likely contributed to the variable findings. The first is the preprocessing procedure. Global signal regression is controversial, possibly leading to false-positive results.^{132, 133} Only a few studies stated that they did not regress out the global signal, and in most studies we could not ascertain any information. Another potential influence is the choice of the correlation metric. Most functional connectome studies used Pearson's correlation coefficient to define network edges. According to a reported test-retest analysis, the reliability of global topological properties is modulated by the correlation metrics and the global signal, with the highest reliability observed for Pearson's correlation-based brain networks without global signal removal.³⁷ The second confounding factor is the brain parcellation schemes. Lord et al. showed that different parcellation schemes for global network properties are comparably consistent.³³ However, they only presented a comparison between two atlases (AAL and Dosenbach) for the graph theoretical analysis, which might generalize to many other commonly used atlases. Third, we cannot exclude the potential confounding effects of medications, although the trend toward structural regularization rather than functional randomization was still prominent in the datasets comprising medicated patients with schizophrenia. Several studies showed appearance of significant structural and functional alterations in the brain following the administration of anti-psychotics.^{3, 134, 135} Fourth, variability may result from the use of different diagnostic categories for patients with the same disorder. A recent study showed a greater stability for some patients with a first psychosis diagnosis, as assigned using ICD-10 criteria rather than DSM-IV criteria.¹³⁶ Fifth, subtype heterogeneity may be a potential confounding factor. For example, two

schizophrenia studies recruited subgroups of patients: one study recruited patients with familial and sporadic schizophrenia, and the other study recruited patients with and without auditory verbal hallucinations. Two DTI studies recruited patients with the most common paranoid subtype of schizophrenia: one study recruited a population considered a homogeneous genetic subtype of schizophrenia, namely, the 22q11.2 deletion syndrome (22q11.2DS). A detailed description of the subtypes analyzed in the remaining studies was not reported. We performed an exploratory analysis of subtypes of LLD and BD type I, which was provided by the original reports. Despite the significant variability among the different studies, two prominent trends of structural regularization rather than functional randomization emerged in patients with schizophrenia. Future studies will be facilitated by strategies overcoming these issues of methodological boundaries and patient heterogeneity.

Different alterations in the global property measures belonging to the same perspective were identified. For example, in the structural connectome, decreased γ and increased E_{loc} , both of which belong to the segregation perspective, were found in patients with depression.¹⁰² The analysis of γ and E_{loc} revealed seemingly opposite results; γ decreased, whereas E_{loc} increased in this group of patients. Initially, the results appeared to be contradictory. Notably, γ is normalized relative to C_p of the 100 matched random networks that preserve the number of nodes, edges, and degree distribution of the real network. The ambiguity was likely the result of the choice of a null model for normalization, which was similar to the finding of increased γ and decreased C_p in patients with focal epilepsy.¹³⁷ The E_{loc} of each node was similar to its C_p . In this circumstance, we believe that the alterations in small-worldness should be evaluated using non-normalized global measurements.

Inconsistent alterations in several original reports were reported compared to the

current criteria. A lower σ in the functional networks in the brains of patients with schizophrenia displays a tendency toward randomization during an AOD task.¹⁵ However, according to the current criteria, decreased segregation (decreased C_p) and decreased integration (increased L_p) suggest a shift toward weaker small-worldization. In another AOD study, decreased segregation (decreased γ) in both the left and right hemisphere task-related networks in patients schizophrenia were reported to be similar to randomization.⁴⁹ However, according to the current criteria, decreased segregation (decreased γ) and decreased integration (decreased E_{glob} and increased L_p) indicated a trend toward weaker small-worldization in the left hemisphere task-related networks. In the structural connectome of children with PTSD, decreased integration (decreased E_{glob} and increased L_p) suggested a shift toward regularization.¹²⁷ However, according to the current criteria, the decreased integration (decreased E_{glob} and increased L_p) and decreased segregation (decreased E_{loc}) showed a shift toward weaker small-worldization. Those inconsistencies might be attributable to the previous unclear definition of the altered small-world properties.

4.4. Limitations of the present study

This study has several limitations. First, we described the patterns of altered small-world properties from the perspectives of segregation and integration. However, 10 reports only investigated 1 perspective, including 9 reports from 8 studies involving only integration^{65, 66, 84, 107, 108, 116, 117, 121} and 1 study involving only segregation.⁵⁵ Based on the reported findings from those 10 studies, our hypothesis may be not accurate. For example, increased segregation alone without information on integration will appear as regularization, but stronger small-worldization may actually occur. Second, 3 studies^{94, 121, 128} only reported changes in σ , which is not sufficient to determine small-worldness alterations. Third, the differences in preprocessing steps,

and analytical methods (e.g., parcellation and correlation calculations)^{37, 138-140} can produce inconsistent results, and thus the integration of results across different studies is difficult. Fourth, although 3 studies reported altered small-world properties in patients with schizophrenia and LLD,^{17, 95, 110} these differences did not reach statistical significance. Fifth, combining evidence from patients with different disease subtypes will increase the variance of the results. However, detailed subtype information was not available for most studies included. Sixth, most studies reported p values for between-group comparisons and did not show the values of the global properties. A better strategy is to report the magnitude of the effects and differences, because the p value may be a biased estimate. We were consequently unable to perform analyses of effect sizes and publication bias.

4.5. Conclusions and future work

In conclusion, we propose 4 patterns of altered small-world properties, namely, regularization, randomization, stronger small-worldization and weaker small-worldization, and use these patterns to summarize the altered small-world properties in patients with one of several major psychiatric disorders. Although more differences than similarities are observed in patients with these disorders, one prominent trend is the prevalent report of regularization of the structural connectome with randomization of the functional connectome in patients with schizophrenia. An analysis of altered small-world properties may provide novel insights into the pathophysiological mechanisms underlying psychiatric disorders from a connectomic perspective. However, several important challenges for future research remain.

First, as described above, mixed results for the brain networks have been reported. The heterogeneity of the patient samples may be a major reason for these discrepancies. Researchers will need to select more homogeneous samples by more

detailed consideration of demographic variables. The accumulation of validated evidence from connectome studies will reveal biological markers that indicate specific phenotypes of psychiatric disorders. Second, the combination of multimodal imaging modalities in future studies will provide additional integrative information to map the patterns of the brain connectome. Third, most of these connectome changes were detected based on cross-sectional data and thus may be influenced by inter-subject variability and unbalanced cohort distributions. Investigations of longitudinal networks are needed. Fourth, caution should be exercised in choosing the most reliable methods for analyzing structural and functional connectomes. For resting-state fMRI studies, higher test-retest network reliabilities are obtained using functionally rather than structurally defined nodes, and Pearson rather than partial correlations.^{37, 140} Finally, the emergence of novel connectome analytical approaches (e.g., dynamic connectivity) and imaging protocols (e.g., multiband fMRI) should dramatically increase our knowledge of network dysfunction in patients with brain disorders. Additional imaging research on the connectome in patients with the major psychiatric disorders is urgently needed.

Conflicts of interest: All of the authors report no biomedical financial interests or potential conflicts of interest.

Acknowledgements: This study was supported by the National Natural Science Foundation (Grant Nos. 81501452, 81621003, 81761128023, 81220108013, 81227002 and 81030027), the Program for Changjiang Scholars and Innovative Research Team in University (PCSIRT, grant IRT16R52) of China, the Changjiang Scholar Professorship Award (Award No. T2014190) of China, and the CMB Distinguished a Professorship Award (Award No. F510000/ G16916411) administered by the Institute of International Education. D.L. supported by Newton International Fellowship from the Royal Society.

References

1. Lui S, Zhou XJ, Sweeney JA, et al. Psychoradiology: The Frontier of Neuroimaging in Psychiatry. *Radiology* 2016;281(2):357-72.
2. Gong Q, He Y. Depression, neuroimaging and connectomics: a selective overview. *Biol Psychiatry* 2015;77(3):223-35.
3. Gong Q, Lui S, Sweeney JA. A Selective Review of Cerebral Abnormalities in Patients With First-Episode Schizophrenia Before and After Treatment. *Am J Psychiatry* 2016;173(3):232-43.
4. Sarpal DK, Lencz T, Malhotra AK. In Support of Neuroimaging Biomarkers of Treatment Response in First-Episode Schizophrenia. *Am J Psychiatry* 2016;173(7):732-3.
5. Gong Q. Response to Sarpal et al.: Importance of Neuroimaging Biomarkers for Treatment Development and Clinical Practice. *Am J Psychiatry* 2016;173(7):733-4.
6. Li F, Lui S, Yao L, et al. Altered White Matter Connectivity Within and Between Networks in Antipsychotic-Naive First-Episode Schizophrenia. *Schizophr Bull* 2017.
7. Sporns O, Tononi G, Kotter R. The human connectome: A structural description of the human brain. *PLoS Comput Biol* 2005;1(4):e42.
8. Sporns O. Network attributes for segregation and integration in the human brain. *Curr Opin Neurobiol* 2013;23(2):162-71.
9. Fornito A, Zalesky A, Breakspear M. Graph analysis of the human connectome: promise, progress, and pitfalls. *Neuroimage* 2013;80:426-44.
10. He Y, Chen ZJ, Evans AC. Small-world anatomical networks in the human brain revealed by cortical thickness from MRI. *Cereb Cortex* 2007;17(10):2407-19.
11. Lei D, Li K, Li L, et al. Disrupted Functional Brain Connectome in Patients with Posttraumatic Stress Disorder. *Radiology* 2015;276(3):818-27.

12. Zhang T, Wang J, Yang Y, et al. Abnormal small-world architecture of top-down control networks in obsessive-compulsive disorder. *J Psychiatry Neurosci* 2011;36(1):23-31.
13. Zhang J, Wang J, Wu Q, et al. Disrupted brain connectivity networks in drug-naive, first-episode major depressive disorder. *Biol Psychiatry* 2011;70(4):334-42.
14. Aerts H, Fias W, Caeyenberghs K, et al. Brain networks under attack: robustness properties and the impact of lesions. *Brain* 2016;139(Pt 12):3063-83.
15. Ma S, Calhoun VD, Eichele T, et al. Modulations of functional connectivity in the healthy and schizophrenia groups during task and rest. *Neuroimage* 2012;62(3):1694-704.
16. Zhu J, Wang C, Liu F, et al. Alterations of Functional and Structural Networks in Schizophrenia Patients with Auditory Verbal Hallucinations. *Front Hum Neurosci* 2016;10:114.
17. He H, Sui J, Yu Q, et al. Altered small-world brain networks in schizophrenia patients during working memory performance. *PLoS One* 2012;7(6):e38195.
18. Zhang Y, Lin L, Lin CP, et al. Abnormal topological organization of structural brain networks in schizophrenia. *Schizophr Res* 2012;141(2-3):109-18.
19. Wang Q, Su TP, Zhou Y, et al. Anatomical insights into disrupted small-world networks in schizophrenia. *Neuroimage* 2012;59(2):1085-93.
20. Su TW, Hsu TW, Lin YC, et al. Schizophrenia symptoms and brain network efficiency: A resting-state fMRI study. *Psychiatry Res* 2015;234(2):208-18.
21. Yu Q, Sui J, Rachakonda S, et al. Altered topological properties of functional network connectivity in schizophrenia during resting state: a small-world brain network study. *PLoS One* 2011;6(9):e25423.
22. Filippi M, van den Heuvel MP, Fornito A, et al. Assessment of system

dysfunction in the brain through MRI-based connectomics. *Lancet Neurol* 2013;12(12):1189-99.

23. Fornito A, Zalesky A, Pantelis C, et al. Schizophrenia, neuroimaging and connectomics. *Neuroimage* 2012;62(4):2296-314.

24. Mears D, Pollard HB. Network science and the human brain: Using graph theory to understand the brain and one of its hubs, the amygdala, in health and disease. *J Neurosci Res* 2016;94(6):590-605.

25. Liao X, Vasilakos AV, He Y. Small-world human brain networks: Perspectives and challenges. *Neurosci Biobehav Rev* 2017;77:286-300.

26. Shen X, Tokoglu F, Papademetris X, et al. Groupwise whole-brain parcellation from resting-state fMRI data for network node identification. *Neuroimage* 2013;82:403-15.

27. Power JD, Cohen AL, Nelson SM, et al. Functional network organization of the human brain. *Neuron* 2011;72(4):665-78.

28. Dosenbach NU, Nardos B, Cohen AL, et al. Prediction of individual brain maturity using fMRI. *Science* 2010;329(5997):1358-61.

29. Craddock RC, James GA, Holtzheimer PE, 3rd, et al. A whole brain fMRI atlas generated via spatially constrained spectral clustering. *Hum Brain Mapp* 2012;33(8):1914-28.

30. Tzourio-Mazoyer N, Landeau B, Papathanassiou D, et al. Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *Neuroimage* 2002;15(1):273-89.

31. Xia M, He Y. Functional connectomics from a "big data" perspective. *Neuroimage* 2017.

32. Park B, Ko JH, Lee JD, et al. Evaluation of node-inhomogeneity effects on the

functional brain network properties using an anatomy-constrained hierarchical brain parcellation. *PLoS One* 2013;8(9):e74935.

33. Lord A, Ehrlich S, Borchardt V, et al. Brain parcellation choice affects disease-related topology differences increasingly from global to local network levels. *Psychiatry Res* 2016;249:12-9.

34. Smith SM, Miller KL, Salimi-Khorshidi G, et al. Network modelling methods for FMRI. *Neuroimage* 2011;54(2):875-91.

35. Smith SM, Vidaurre D, Beckmann CF, et al. Functional connectomics from resting-state fMRI. *Trends Cogn Sci* 2013;17(12):666-82.

36. Zalesky A, Fornito A, Bullmore E. On the use of correlation as a measure of network connectivity. *Neuroimage* 2012;60(4):2096-106.

37. Liang X, Wang J, Yan C, et al. Effects of different correlation metrics and preprocessing factors on small-world brain functional networks: a resting-state functional MRI study. *PLoS One* 2012;7(3):e32766.

38. Mori S, van Zijl PC. Fiber tracking: principles and strategies - a technical review. *NMR Biomed* 2002;15(7-8):468-80.

39. Ratnarajah N, Simmons A, Davydov O, et al. A novel approach for improved tractography and quantitative analysis of probabilistic fibre tracking curves. *Med Image Anal* 2012;16(1):227-38.

40. Behrens TE, Woolrich MW, Jenkinson M, et al. Characterization and propagation of uncertainty in diffusion-weighted MR imaging. *Magn Reson Med* 2003;50(5):1077-88.

41. Buchanan CR, Pernet CR, Gorgolewski KJ, et al. Test-retest reliability of structural brain networks from diffusion MRI. *Neuroimage* 2014;86:231-43.

42. Rubinov M, Sporns O. Complex network measures of brain connectivity: uses

and interpretations. *Neuroimage* 2010;52(3):1059-69.

43. Watts DJ, Strogatz SH. Collective dynamics of 'small-world' networks. *Nature* 1998;393(6684):440-2.

44. Latora V, Marchiori M. Efficient behavior of small-world networks. *Phys Rev Lett* 2001;87(19):198701.

45. Cao M, Shu N, Cao Q, et al. Imaging functional and structural brain connectomics in attention-deficit/hyperactivity disorder. *Mol Neurobiol* 2014;50(3):1111-23.

46. Lord A, Horn D, Breakspear M, et al. Changes in community structure of resting state functional connectivity in unipolar depression. *PLoS One* 2012;7(8):e41282.

47. Chen VC, Shen CY, Liang SH, et al. Assessment of abnormal brain structures and networks in major depressive disorder using morphometric and connectome analyses. *J Affect Disord* 2016;205:103-11.

48. Zhu J, Zhuo C, Liu F, et al. Distinct disruptions of resting-state functional brain networks in familial and sporadic schizophrenia. *Sci Rep* 2016;6:23577.

49. Yu Q, Sui J, Rachakonda S, et al. Altered small-world brain networks in temporal lobe in patients with schizophrenia performing an auditory oddball task. *Front Syst Neurosci* 2011;5:7.

50. Alexander-Bloch AF, Gogtay N, Meunier D, et al. Disrupted modularity and local connectivity of brain functional networks in childhood-onset schizophrenia. *Front Syst Neurosci* 2010;4:147.

51. Liu F, Zhuo C, Yu C. Altered Cerebral Blood Flow Covariance Network in Schizophrenia. *Front Neurosci* 2016;10:308.

52. Lo CY, Su TW, Huang CC, et al. Randomization and resilience of brain functional networks as systems-level endophenotypes of schizophrenia. *Proc Natl*

Acad Sci U S A 2015;112(29):9123-8.

53. Lynall ME, Bassett DS, Kerwin R, et al. Functional connectivity and brain networks in schizophrenia. *J Neurosci* 2010;30(28):9477-87.

54. Wang L, Metzak PD, Honer WG, et al. Impaired efficiency of functional networks underlying episodic memory-for-context in schizophrenia. *J Neurosci* 2010;30(39):13171-9.

55. Yu Q, Plis SM, Erhardt EB, et al. Modular Organization of Functional Network Connectivity in Healthy Controls and Patients with Schizophrenia during the Resting State. *Front Syst Neurosci* 2011;5:103.

56. Alexander-Bloch AF, Vertes PE, Stidd R, et al. The anatomical distance of functional connections predicts brain network topology in health and schizophrenia. *Cereb Cortex* 2013;23(1):127-38.

57. Tomasi D, Volkow ND. Mapping small-world properties through development in the human brain: disruption in schizophrenia. *PLoS One* 2014;9(4):e96176.

58. Liu Y, Liang M, Zhou Y, et al. Disrupted small-world networks in schizophrenia. *Brain* 2008;131(Pt 4):945-61.

59. Sheffield JM, Repovs G, Harms MP, et al. Fronto-parietal and cingulo-opercular network integrity and cognition in health and schizophrenia. *Neuropsychologia* 2015;73:82-93.

60. Yu Q, Erhardt EB, Sui J, et al. Assessing dynamic brain graphs of time-varying connectivity in fMRI data: application to healthy controls and patients with schizophrenia. *Neuroimage* 2015;107:345-55.

61. Hadley JA, Kraguljac NV, White DM, et al. Change in brain network topology as a function of treatment response in schizophrenia: a longitudinal resting-state fMRI study using graph theory. *NPJ Schizophr* 2016;2:16014.

62. Li X, Xia S, Bertisch HC, et al. Unique topology of language processing brain network: a systems-level biomarker of schizophrenia. *Schizophr Res* 2012;141(2-3):128-36.
63. Yu Q, Sui J, Kiehl KA, et al. State-related functional integration and functional segregation brain networks in schizophrenia. *Schizophr Res* 2013;150(2-3):450-8.
64. Fornito A, Yoon J, Zalesky A, et al. General and specific functional connectivity disturbances in first-episode schizophrenia during cognitive control performance. *Biol Psychiatry* 2011;70(1):64-72.
65. Chen J, Yao Z, Qin J, et al. Abnormal inter- and intra-hemispheric integration in male paranoid schizophrenia: a graph-theoretical analysis. *Shanghai Arch Psychiatry* 2015;27(3):158-66.
66. Griffa A, Baumann PS, Ferrari C, et al. Characterizing the connectome in schizophrenia with diffusion spectrum imaging. *Hum Brain Mapp* 2015;36(1):354-66.
67. Kim DJ, Kent JS, Bolbecker AR, et al. Disrupted modular architecture of cerebellum in schizophrenia: a graph theoretic analysis. *Schizophr Bull* 2014;40(6):1216-26.
68. Ottet MC, Schaer M, Debbane M, et al. Graph theory reveals disconnected hubs in 22q11DS and altered nodal efficiency in patients with hallucinations. *Front Hum Neurosci* 2013;7:402.
69. Sun Y, Chen Y, Lee R, et al. Disruption of brain anatomical networks in schizophrenia: A longitudinal, diffusion tensor imaging based study. *Schizophr Res* 2016;171(1-3):149-57.
70. Sun Y, Chen Y, Collinson SL, et al. Reduced Hemispheric Asymmetry of Brain Anatomical Networks Is Linked to Schizophrenia: A Connectome Study. *Cereb Cortex* 2017;27(1):602-15.

71. van den Heuvel MP, Sporns O, Collin G, et al. Abnormal rich club organization and functional brain dynamics in schizophrenia. *JAMA Psychiatry* 2013;70(8):783-92.
72. Yeo RA, Ryman SG, van den Heuvel MP, et al. Graph Metrics of Structural Brain Networks in Individuals with Schizophrenia and Healthy Controls: Group Differences, Relationships with Intelligence, and Genetics. *J Int Neuropsychol Soc* 2016;22(2):240-9.
73. Zalesky A, Fornito A, Seal ML, et al. Disrupted axonal fiber connectivity in schizophrenia. *Biol Psychiatry* 2011;69(1):80-9.
74. Collin G, Kahn RS, de Reus MA, et al. Impaired rich club connectivity in unaffected siblings of schizophrenia patients. *Schizophr Bull* 2014;40(2):438-48.
75. Sun Y, Lee R, Shen K, et al. Structural connectivity analysis reveals topological aberrations in patients with schizophrenia. *Conf Proc IEEE Eng Med Biol Soc* 2013;2013:1386-9.
76. Yan H, Tian L, Wang Q, et al. Compromised small-world efficiency of structural brain networks in schizophrenic patients and their unaffected parents. *Neurosci Bull* 2015;31(3):275-87.
77. Zhang R, Wei Q, Kang Z, et al. Disrupted brain anatomical connectivity in medication-naive patients with first-episode schizophrenia. *Brain Struct Funct* 2015;220(2):1145-59.
78. Cabral J, Fernandes HM, Van Hartevelt TJ, et al. Structural connectivity in schizophrenia and its impact on the dynamics of spontaneous functional networks. *Chaos* 2013;23(4):046111.
79. Hu M, Zong X, Zheng J, et al. Risperidone-induced topological alterations of anatomical brain network in first-episode drug-naive schizophrenia patients: a

- longitudinal diffusion tensor imaging study. *Psychol Med* 2016;46(12):2549-60.
80. van den Heuvel MP, Mandl RC, Stam CJ, et al. Aberrant frontal and temporal complex network structure in schizophrenia: a graph theoretical analysis. *J Neurosci* 2010;30(47):15915-26.
81. Li W, Douglas Ward B, Liu X, et al. Disrupted small world topology and modular organisation of functional networks in late-life depression with and without amnesic mild cognitive impairment. *J Neurol Neurosurg Psychiatry* 2015;86(10):1097-105.
82. Guo H, Cheng C, Cao X, et al. Resting-state functional connectivity abnormalities in first-onset unmedicated depression. *Neural Regen Res* 2014;9(2):153-63.
83. Borchardt V, Lord AR, Li M, et al. Preprocessing strategy influences graph-based exploration of altered functional networks in major depression. *Hum Brain Mapp* 2016;37(4):1422-42.
84. Manelis A, Almeida JR, Stiffler R, et al. Anticipation-related brain connectivity in bipolar and unipolar depression: a graph theory approach. *Brain* 2016;139(Pt 9):2554-66.
85. Borchardt V, Krause AL, Starck T, et al. Graph theory reveals hyper-functionality in visual cortices of Seasonal Affective Disorder patients. *World J Biol Psychiatry* 2015;16(2):123-34.
86. Meng C, Brandl F, Tahmasian M, et al. Aberrant topology of striatum's connectivity is associated with the number of episodes in depression. *Brain* 2014;137(Pt 2):598-609.
87. Ye M, Yang T, Qing P, et al. Changes of Functional Brain Networks in Major Depressive Disorder: A Graph Theoretical Analysis of Resting-State fMRI. *PLoS One* 2015;10(9):e0133775.

88. Park CH, Wang SM, Lee HK, et al. Affective state-dependent changes in the brain functional network in major depressive disorder. *Soc Cogn Affect Neurosci* 2014;9(9):1404-12.
89. Luo Q, Deng Z, Qin J, et al. Frequency dependant topological alterations of intrinsic functional connectome in major depressive disorder. *Sci Rep* 2015;5:9710.
90. Wang Z, Yuan Y, Bai F, et al. Altered topological patterns of brain networks in remitted late-onset depression: a resting-state fMRI study. *J Clin Psychiatry* 2016;77(1):123-30.
91. Bohr IJ, Kenny E, Blamire A, et al. Resting-state functional connectivity in late-life depression: higher global connectivity and more long distance connections. *Front Psychiatry* 2012;3:116.
92. He H, Yu Q, Du Y, et al. Resting-state functional network connectivity in prefrontal regions differs between unmedicated patients with bipolar and major depressive disorders. *J Affect Disord* 2016;190:483-93.
93. Peng D, Shi F, Shen T, et al. Altered brain network modules induce helplessness in major depressive disorder. *J Affect Disord* 2014;168:21-9.
94. Jin C, Gao C, Chen C, et al. A preliminary study of the dysregulation of the resting networks in first-episode medication-naive adolescent depression. *Neurosci Lett* 2011;503(2):105-9.
95. Mak E, Colloby SJ, Thomas A, et al. The segregated connectome of late-life depression: a combined cortical thickness and structural covariance analysis. *Neurobiol Aging* 2016;48:212-21.
96. Bai F, Shu N, Yuan Y, et al. Topologically convergent and divergent structural connectivity patterns between patients with remitted geriatric depression and amnesic mild cognitive impairment. *J Neurosci* 2012;32(12):4307-18.

97. Ajilore O, Lamar M, Leow A, et al. Graph theory analysis of cortical-subcortical networks in late-life depression. *Am J Geriatr Psychiatry* 2014;22(2):195-206.
98. Wang T, Wang K, Qu H, et al. Disorganized cortical thickness covariance network in major depressive disorder implicated by aberrant hubs in large-scale networks. *Sci Rep* 2016;6:27964.
99. Li X, Steffens DC, Potter GG, et al. Decreased between-hemisphere connectivity strength and network efficiency in geriatric depression. *Hum Brain Mapp* 2017;38(1):53-67.
100. Singh MK, Kesler SR, Hadi Hosseini SM, et al. Anomalous gray matter structural networks in major depressive disorder. *Biol Psychiatry* 2013;74(10):777-85.
101. Chen JH, Yao ZJ, Qin JL, et al. Aberrant Global and Regional Topological Organization of the Fractional Anisotropy-weighted Brain Structural Networks in Major Depressive Disorder. *Chin Med J (Engl)* 2016;129(6):679-89.
102. Long Z, Duan X, Wang Y, et al. Disrupted structural connectivity network in treatment-naive depression. *Prog Neuropsychopharmacol Biol Psychiatry* 2015;56:18-26.
103. Ajilore O, Lamar M, Kumar A. Association of brain network efficiency with aging, depression, and cognition. *Am J Geriatr Psychiatry* 2014;22(2):102-10.
104. Charlton RA, Leow A, GadElkarim J, et al. Brain connectivity in late-life depression and aging revealed by network analysis. *Am J Geriatr Psychiatry* 2015;23(6):642-50.
105. Lim HK, Jung WS, Aizenstein HJ. Aberrant topographical organization in gray matter structural network in late life depression: a graph theoretical analysis. *Int Psychogeriatr* 2013;25(12):1929-40.
106. Qin J, Wei M, Liu H, et al. Abnormal brain anatomical topological

organization of the cognitive-emotional and the frontoparietal circuitry in major depressive disorder. *Magn Reson Med* 2014;72(5):1397-407.

107. Sacchet MD, Prasad G, Foland-Ross LC, et al. Elucidating Brain Connectivity Networks in Major Depressive Disorder Using Classification-Based Scoring. *Proc IEEE Int Symp Biomed Imaging* 2014;2014:246-9.

108. Sacchet MD, Prasad G, Foland-Ross LC, et al. Support vector machine classification of major depressive disorder using diffusion-weighted neuroimaging and graph theory. *Front Psychiatry* 2015;6:21.

109. Tymofiyeva O, Connolly CG, Ho TC, et al. DTI-based connectome analysis of adolescents with major depressive disorder reveals hypoconnectivity of the right caudate. *J Affect Disord* 2017;207:18-25.

110. Korgaonkar MS, Fornito A, Williams LM, et al. Abnormal structural networks characterize major depressive disorder: a connectome analysis. *Biol Psychiatry* 2014;76(7):567-74.

111. Wang L, Zhu C, He Y, et al. Altered small-world brain functional networks in children with attention-deficit/hyperactivity disorder. *Hum Brain Mapp* 2009;30(2):638-49.

112. Xia S, Foxe JJ, Sroubek AE, et al. Topological organization of the "small-world" visual attention network in children with attention deficit/hyperactivity disorder (ADHD). *Front Hum Neurosci* 2014;8:162.

113. Cocchi L, Bramati IE, Zalesky A, et al. Altered functional brain connectivity in a non-clinical sample of young adults with attention-deficit/hyperactivity disorder. *J Neurosci* 2012;32(49):17753-61.

114. Cao Q, Shu N, An L, et al. Probabilistic diffusion tractography and graph theory analysis reveal abnormal white matter structural connectivity networks in drug-naive

- boys with attention deficit/hyperactivity disorder. *J Neurosci* 2013;33(26):10676-87.
115. Sidlauskaite J, Caeyenberghs K, Sonuga-Barke E, et al. Whole-brain structural topology in adult attention-deficit/hyperactivity disorder: Preserved global - disturbed local network organization. *Neuroimage Clin* 2015;9:506-12.
116. Spielberg JM, Beall EB, Hulvershorn LA, et al. Resting State Brain Network Disturbances Related to Hypomania and Depression in Medication-Free Bipolar Disorder. *Neuropsychopharmacology* 2016;41(13):3016-24.
117. Collin G, van den Heuvel MP, Abramovic L, et al. Brain network analysis reveals affected connectome structure in bipolar I disorder. *Hum Brain Mapp* 2016;37(1):122-34.
118. Leow A, Ajilore O, Zhan L, et al. Impaired inter-hemispheric integration in bipolar disorder revealed with brain network analyses. *Biol Psychiatry* 2013;73(2):183-93.
119. Shin DJ, Jung WH, He Y, et al. The effects of pharmacological treatment on functional brain connectome in obsessive-compulsive disorder. *Biol Psychiatry* 2014;75(8):606-14.
120. Jung WH, Yucel M, Yun JY, et al. Altered functional network architecture in orbitofronto-striato-thalamic circuit of unmedicated patients with obsessive-compulsive disorder. *Hum Brain Mapp* 2017;38(1):109-19.
121. Armstrong CC, Moody TD, Feusner JD, et al. Graph-theoretical analysis of resting-state fMRI in pediatric obsessive-compulsive disorder. *J Affect Disord* 2016;193:175-84.
122. Zhong ZX, Zhao TD, Luo J, et al. Abnormal topological organization in white matter structural networks revealed by diffusion tensor tractography in unmedicated patients with obsessive-compulsive disorder. *Progress in Neuro-*

Psychopharmacology & Biological Psychiatry 2014;51:39-50.

123. Reess TJ, Rus OG, Schmidt R, et al. Connectomics-based structural network alterations in obsessive-compulsive disorder. *Translational Psychiatry* 2016;6.

124. Kim SG, Jung WH, Kim SN, et al. Disparity between dorsal and ventral networks in patients with obsessive-compulsive disorder:evidence revealed by graph theoretical analysis based on cortical thickness from MRI. *Frontiers in Human Neuroscience* 2013;7.

125. Suo X, Lei D, Li K, et al. Disrupted brain network topology in pediatric posttraumatic stress disorder: A resting-state fMRI study. *Hum Brain Mapp* 2015;36(9):3677-86.

126. Long Z, Duan X, Xie B, et al. Altered brain structural connectivity in post-traumatic stress disorder: a diffusion tensor imaging tractography study. *J Affect Disord* 2013;150(3):798-806.

127. Suo X, Lei D, Chen F, et al. Anatomic Insights into Disrupted Small-World Networks in Pediatric Posttraumatic Stress Disorder. *Radiology* 2017;282(3):826-34.

128. Mueller SG, Ng P, Neylan T, et al. Evidence for disrupted gray matter structural connectivity in posttraumatic stress disorder. *Psychiatry Res* 2015;234(2):194-201.

129. Bullmore ET, Bassett DS. Brain graphs: graphical models of the human brain connectome. *Annu Rev Clin Psychol* 2011;7:113-40.

130. Park HJ, Friston K. Structural and functional brain networks: from connections to cognition. *Science* 2013;342(6158):1238411.

131. Wang Z, Dai Z, Gong G, et al. Understanding structural-functional relationships in the human brain: a large-scale network perspective. *Neuroscientist* 2015;21(3):290-305.

132. Fox MD, Zhang D, Snyder AZ, et al. The global signal and observed anticorrelated resting state brain networks. *J Neurophysiol* 2009;101(6):3270-83.
133. Murphy K, Birn RM, Handwerker DA, et al. The impact of global signal regression on resting state correlations: are anti-correlated networks introduced? *Neuroimage* 2009;44(3):893-905.
134. Navari S, Dazzan P. Do antipsychotic drugs affect brain structure? A systematic and critical review of MRI findings. *Psychol Med* 2009;39(11):1763-77.
135. Fusar-Poli P, Smieskova R, Kempton MJ, et al. Progressive brain changes in schizophrenia related to antipsychotic treatment? A meta-analysis of longitudinal MRI studies. *Neurosci Biobehav Rev* 2013;37(8):1680-91.
136. Salvatore P, Baldessarini RJ, Tohen M, et al. McLean-Harvard International First-Episode Project: two-year stability of ICD-10 diagnoses in 500 first-episode psychotic disorder patients. *J Clin Psychiatry* 2011;72(2):183-93.
137. Vytvarova E, Marecek R, Fousek J, et al. Large-scale cortico-subcortical functional networks in focal epilepsies: The role of the basal ganglia. *Neuroimage Clin* 2017;14:28-36.
138. Wang J, Wang L, Zang Y, et al. Parcellation-dependent small-world brain functional networks: a resting-state fMRI study. *Hum Brain Mapp* 2009;30(5):1511-23.
139. Liao XH, Xia MR, Xu T, et al. Functional brain hubs and their test-retest reliability: a multiband resting-state functional MRI study. *Neuroimage* 2013;83:969-82.
140. Wang JH, Zuo XN, Gohel S, et al. Graph theoretical analysis of functional brain networks: test-retest evaluation on short- and long-term resting-state functional MRI data. *PLoS One* 2011;6(7):e21976.

Figure legends

Figure 1. Representation of four main patterns of altered small-world properties in the brain connectome, ‘regularization’, ‘randomization’, ‘stronger small-worldization’ and ‘weaker small-worldization’, based on the perspectives of segregation and integration. Regularization is defined as increased segregation and/or decreased integration. Randomization is defined as decreased segregation and/or increased integration. Stronger small-worldization is defined as increased segregation and increased integration. Weaker small-worldization is defined as decreased segregation and decreased integration.

Figure 2. Graph showing the number of reports describing either the ‘randomization’, ‘regularization’, ‘weaker small-worldization’, ‘stronger small-worldization’ or ‘no difference’ in functional and structural brain connectomes between patient groups compared with healthy controls. ADHD, attention-deficit/hyperactivity disorder; BD, bipolar disorder; OCD, obsessive compulsive disorder; PTSD, posttraumatic stress disorder; HC, healthy controls.