

High macrophage activities are associated with advanced periductal fibrosis in chronic Opisthorchis viverrini infection

Journal:	Parasite Immunology
Manuscript ID	Draft
Manuscript Type:	Original Paper
Date Submitted by the Author:	n/a
Complete List of Authors:	Salao, Kanin; Khon Kaen University Faculty of Medicine, Microbiology Watakulsin, Krongkarn; Khon Kaen University Faculty of Medicine, Microbiology Mairiang, Eimorn; Khon Kaen University Faculty of Medicine, Radiology Suttiprapa, Sutas; Khon Kaen University Faculty of Medicine, Academic Affairs Tangkawattana, Sirikachorn; Khon Kaen University Faculty of Veterinary Medicine, Pathobiology Edwards, Steven; University of Liverpool School of Life Sciences, Integrative Biology Sripa, Banchob; Faculty of Medicine,Khon Kaen University, Department of Pathology
Key Words:	Macrophage < Cell, Peripheral blood mononuclear cells < Cell, Human < Host species, Innate immunity < Immunological terms, Opisthorchis viverrini < Parasite, Flow cytometry < Tools and techniques
	·

SCHOLARONE[™] Manuscripts

1 2		
3 4 5	1	High macrophage activities are associated with advanced periductal fibrosis in chronic
6 7	2	Opisthorchis viverrini infection
8 9 10	3	Running title: Innate immunity in opisthorchiasis
11 12	4	
13 14 15	5	Kanin Salao ^{1,7} , Krongkarn Watakulsin ^{1,7} , Eimorn Mairiang ² , Sutas Suttiprapa ^{3, 7} , Sirikachorn
15 16 17	6	Tangkawattan ^{4, 7} , Steven W. Edwards ⁵ , Banchob Sripa ^{6, 7*}
18 19	7	
20 21	8	¹ Department of Microbiology, Faculty of Medicine, Khon Kaen University, Khon Kaen 40002,
22 23	9	THAILAND, Email: <u>kaninsa@kku.ac.th</u>
24	10	¹ Department of Microbiology, Faculty of Medicine, Khon Kaen University, Khon Kaen 40002,
25 26	11	THAILAND, Email: <u>Watakarn@kkumail.com</u>
27 28	12	² Department of Radiology, Faculty of Medicine, Khon Kaen University, Khon Kaen 40002,
29	13	THAILAND, Email: <u>eimmai@kku.ac.th</u>
30 31	14	³ Tropical Medicine Graduate Program (International Program), Academic Affairs,, Faculty of
32 33	15	Medicine, Khon Kaen University, Khon Kaen 40002, THAILAND, Email: sutasu@kku.ac.th
34	16	⁴ Department of Pathobiology, Faculty of Veterinary Medicine, Khon Kaen University, Khon
36	17	Kaen 40002, THAILAND, Email: sirikach@kku.ac.th
37 38	18	⁵ Institute of Integrative Biology, Faculty of Health and Life Sciences, University of Liverpool,
39 40	19	Liverpool L69 7ZB, UK, Email: <u>S.W.Edwards@liverpool.ac.uk</u>
41	20	⁶ Department of Pathology, Faculty of Medicine, Khon Kaen University, Khon Kaen 40002,
42 43	21	THAILAND, Email: <u>banchob@kku.ac.th</u>
44 45	22	⁷ WHO Collaborating Centre for Research and Control of Opisthorchiasis (Southeast Asian Liver
46 47	23	Fluke Disease), Tropical Disease Research Center, Faculty of Medicine, Khon Kaen University,
47 48	24	Khon Kaen 40002, THAILAND
49 50	25	
51 52	26	* Corresponding author: Dr.Banchob Sripa, Tropical Disease Research Center (TDRC),
53 54 55 56 57 58 59	27	Department of Pathology, Faculty of Medicine, Khon Kaen University, Khon Kaen 40002, 1

1		
2		
3		
4	28	Thailand. Tel: 66-43-363113, Fax: 66-43-204359, E-mail: <u>banchob@kku.ac.th</u>
5	20	
7	29	
, 8		
9		
10		
11		
12		
13		
14		
15		
16		
17 10		
10		
20		
21		
22		
23		
24		
25		
26		
27		
28		
29		
30		
32		
33		
34		
35		
36		
37		
38		
39		
40 41		
41		
43		
44		
45		
46		
47		
48		
49		
50 51		
51 52		
53		
54		
55		
56		
57		
58		
59		
60		

Abstract

Parasite Immunology

1	
2	
2	
ر ۸	
4 7	
5	
6	
7	
8	
9	
10	
11	
12	
12	
13	
14	
15	
16	
17	
18	
19	
20	
21	
22	
22	
23	
24	
25	
26	
27	
28	
29	
30	
31	
32	
22	
JJ ⊃∧	
24	
35	
36	
37	
38	
39	
40	
41	
42	
43	
44	
<u>4</u> 5	
7J 76	
40	
47	
48	
49	
50	
51	
52	
53	
54	
55	
56	
50	
57	
20	
79	

31	Liver fluke infection caused by Opisthorchis viverrini induces several hepatobiliary conditions
32	including advanced periductal fibrosis (APF) and cholangiocarcinoma (CCA), but >25% of the
33	infected population develops APF and 1% develop CCA. The innate immune response is the first
34	line of defense, and macrophages are critical regulators of fibrosis. We hypothesized that
35	macrophages from infected individuals have different capacities to either promote or suppress
36	periductal fibrosis. We compared phagocytic activities of macrophages of healthy individuals and
37	O. viverrini-infected individuals ±APF, and found that macrophages from infected individuals
38	with APF ingested significantly higher numbers of beads compared with healthy controls and O.
39	viverrini-infected individuals without APF. To further investigate proteolytic activity, we
40	monitored real-time phagosomal proteolysis of beads conjugated to DQ BODIPY-BSA using
41	live-cell imaging. We show that macrophages from O. viverrini-infected individuals with APF
42	also have elevated phagosomal proteolysis activity, which is consistent with their increased
43	phagocytic activity. Additionally, stimulated ROS production by blood monocytes was higher in
44	individuals with APF compared with healthy controls and infected individuals without APF.
45	These results suggest that during O. viverrini infection, macrophages with high phagocytic and
46	proteolytic activities together with elevated ROS production, are the phenotypes that can promote
47	tissue damage, which results in periductal fibrosis.
48	

Keywords: Opisthorchis viverrini; Innate immunity; Peripheral blood mononuclear cells, 49 50 Macrophages; Human; Flow cytometry

3

51

1 2		
3 4 5	52	
6 7	53	1. INTRODUCTION
8 9 10	54	Opisthorchis viverrini infection is endemic in the Lower Mekong regions of Southeast Asia,
10 11 12	55	including Thailand, with approximately 8 million people infected, especially in the northeast of
13 14	56	Thailand where consumption of raw freshwater fishes harboring the infective stage of the parasite
15 16 17	57	is common (1). Infection can cause several disease manifestations of the bile duct including
18 19	58	cholangitis, cholelithiasis, advanced periductal fibrosis (APF) and the most severe complication,
20 21 22	59	cholangiocarcinoma (CCA) (2-6). APF provides the basis for malignant transformation to CCA.
22 23 24	60	Fibrosis occurs when tissues are damaged and normal wound healing responses persist or become
25 26	61	dysregulated (7), usually in response to repetitive tissue injury (8) such as chronic O. viverrini
27 28 20	62	infection (5). Repeated tissue damage results from a combination of factors induced by O.
30 31	63	viverrini such as 1) physical damage induced by feeding, 2) release of products such as reactive
32 33	64	oxygen species from innate immune cells, 3) and O. viverrini-induced host inflammation through
34 35 36	65	both innate and adaptive immunity (9). The latter mechanism, which involves host immune
37 38	66	responses, is well supported by several findings from our group and others, which have
39 40	67	confirmed that chronic O. viverrini infection promotes adaptive proinflammatory responses (10,
41 42	68	11), i.e. IL-6 production (11). However, to date, far less is known about <i>O. viverrini</i> effects on
43 44 45	69	innate immunity.
46 47	70	Macrophages provide defense in innate immunity and these innate immune cells are
48 49	71	among the earliest infiltrating cells after acute O. viverrini infection (9). Apart from host defense,
50 51 52	72	macrophages also play a pivotal role in tissue homeostasis and wound repair. In chronic O.
53 54	73	viverrini infection that usually leads to persistent inflammation, macrophages produce large
55 56		4
57 58		
60		

Page 5 of 27

Parasite Immunology

amounts of mediators that are potentially toxic to the parasites but these may also cause bystander damage to surrounding host tissues. To compensate for this injury, a small subset of macrophages induce tissue repair, which results in the formation of fibrosis or fibrogenesis (12). We hypothesized that different rates of development of APF may be due to the consequence of different levels of innate immune responses that vary among individuals. As macrophages are part of the innate immune system and critical regulators of fibrosis (13), we sought to determine if O. viverrini-infected individuals with APF possess active innate immune responses (e.g. phagocytic, proteolytic activities and ROS production), and hence have a higher risk of developing APF compared with individuals without APF. Unraveling this association should take us a step closer toward identifying a high-risk group and possibly providing data for early and effective diagnosis or primary prevention of APF and CCA.

2. MATERIALS AND METHODS

2.1 Sample recruitment

The current study is an analysis of baseline data collected from a community-based, case-control study of risk factors associated with the development of APF using high resolution ultrasonography, as previously described (14), from ten villages including Bankae, Banbor, Nongtu, Nonmakum, Saadsomsri, Jikngam, Songplui, Nhonkho, Huahad, and Somhong in Kalasin province (Thailand). Overall, 332 individuals aged between 20 and 60 years old were recruited into this study. The control group consisted of 102 O. viverrini-infected individuals (60 males, 42 females) without APF (APF-), while the case group consisted of 104 O. viverrini-infected individuals (58 males and 46 females) with APF (APF +). A group of healthy donors

consisted of 126 O. viverrini-negative participants (68 males and 58 females) (Ov-). Written informed consents were obtained from all participants. This study complied with the standard good clinical practice (GCP) guideline and was approved by the Ethics Committee of Khon Kaen University, Khon Kaen, Thailand, reference number HE591185 and HE480528. 2.2 Sample size calculation A statistical power analysis was performed for sample size estimation based on data from a published study (14). The effect size in this study was 0.5, which is considered medium by Cohen's (1988) criteria. With an alpha = 0.05 and power = 0.80, the projected sample size needed

for this effect size (G*Power 3.1 analysis) is approximately n= 51 per group for the simplest
between-group comparison. Thus, our sample size of >100 per group was more than adequate for
the main objective of this study.

2.3 Ultrasonography

A detailed description of the ultrasonography methods used in this study can be found in previous publications (14, 15). Using a mobile, high-resolution ultrasound (US) machine (GE model LOGIQ Book XP, GE healthcare, WI), hepatobiliary abnormalities including portal vein radical echoes, echoes in liver parenchyma, indistinct gallbladder wall, gallbladder size, sludge and suspected CCA were graded and recorded. Individuals were classified as having "non-advanced periductal fibrosis" or "APF-" if the US grade was 0 or 1, and "advanced periductal fibrosis" or "APF+" if the US grade was 2 or 3. Individuals with alcoholic liver disease, which is seen as fatty liver by US examination, were excluded. Individuals with marked hepatic fibrosis not

1 2			
3 4 5	118	related to O. viverrini infection (e.g., cirrhosis from HBV or HCV) were also excluded from this	5
6 7	119	study.	
8 9 10	120		
11 12	121	2.4 Blood collection	
13 14 15	122	A total of 20 mL of venous blood collected from the participants by venipuncture, 8 mL of which	h
15 16 17	123	was collected in sodium heparin sprayed coated tubes (cat# 367871, Becton Dickinson, NJ) and	
18 19	124	used for monocyte isolation.	
20 21 22	125		
22 23 24	126	2.5 Monocyte isolation for subsequent macrophage generation	
25 26	127	The monocyte-derived macrophages (MDMs) used in this study were from peripheral blood	
27 28 20	128	monocytes isolated using a double gradient centrifugation method (16). Briefly, 8 mL of blood	
29 30 31	129	was overlaid on top of 6 mL of Ficoll solution (1.077 g/ml)(cat# 17-1440-02, GE Healthcare Life	è
32 33	130	Sciences, Singapore) in a 15 mL Falcon tube (cat# 352097, Becton Dickinson, NJ) before	
34 35 26	131	centrifuging at $400x g$ without braking for 45 min at room temperature. The white band of	
36 37 38	132	peripheral blood mononuclear cells (PBMCs) at the interface was collected and washed 3 times	
39 40	133	with PBS and resuspended in 9 mL of RPMI-1640 medium (Cat #11875-093, Thermo Fisher	
41 42	134	Scientific, CA). For the second density gradient, 9 mL of the RPMI-resuspended PBMCs were	
43 44 45	135	placed on top of 6 mL of Percoll solution (1.131 g/mL) (cat# 28-9038-33, GE Healthcare Life	
46 47	136	Sciences, Singapore) in a 15-mL Falcon tube and centrifuged at 550x g without braking for 45	
48 49	137	min at room temperature. The band of monocytes at the interface between the two phases was	
50 51 52	138	collected and washed 3 times with PBS. Monocytes were allowed to differentiate into	
53 54	139	macrophages by culturing in RPMI medium supplemented with 10% autologous serum on a 35-	
55 56			7
57 58 59			

Page 8 of 27

1	
' 2	
2	
3	
4	
5	
6	
7	
/	
8	
9	
10	
11	
12	
12	
13	
14	
15	
16	
17	
10	
IÖ	
19	
20	
21	
22	
22	
23	
24	
25	
26	
27	
28	
20	
29	
30	
31	
32	
33	
34	
35	
20	
30	
37	
38	
39	
40	
10	
41	
42	
43	
44	
45	
46	
17	
47	
48	
49	
50	
51	
52	
52	
53	
54	
55	
56	
57	
50	
20	
59	

60

140	mm petri dish (cat# CLS430165, Corning Inc., NY) for 7 days. The MDMs were then cultured in
141	complete medium (RPMI medium containing 100 µg/mL streptomycin (Cat #15140-122, Thermo
142	Fisher Scientific, CA), 2 mM L-glutamine (Cat #25030-081, Thermo Fisher Scientific, CA), 50
143	μ M 2-mecaptoethanol, and 10% heat-inactivated fetal calf serum (Cat #14190-250, Thermo
144	Fisher Scientific, CA).
145	
146	2.7 Preparation of Alexa Fluor 594 dye and bodipy-BSA conjugated silica beads
147	The preparation of silica beads was adapted from the protocol described by Yates and Russell
148	(17). Briefly, 3.0-µm carboxylate-modified silica particles (Cat #PSi-3.0COOH, Kisker Products
149	for Biotechnologies, Germany) were conjugated with Alexa Fluor 594 Succinimidyl Esters
150	(mixed isomers) (Alexa594-SE, Cat #A20004, Thermo Fisher Scientific, CA) or DQ green
151	BODIPY bovine serum albumin (DQ-BODIPY BSA, Cat #D-12050, Thermo Fisher Scientific,
152	CA). For opsonization, 0.5 mL of silica beads (14 mg/mL) was mixed with 0.5 mL purified goat-
153	IgG (5 μ g/mL) (Cat # I9140, Sigma-Aldrich, MO) and incubated at 37°C in 5% CO ₂ for 30 min.
154	The opsonized silica beads were then washed, and resuspended in PBS at a concentration of 1
155	mg/mL.
156	
157	2.8 Synchronized phagocytosis
158	MDMs that adhered to a 35-mm petri dish were washed 3 times with PBS and resuspended in
159	complete medium and then placed on ice for 10 min. Five μL (2x10 ⁶ particles/mL) silica beads
160	conjugated with red Alexa Fluor 594 fluorescent dye were added to the petri dish containing
161	adhered MDMs and mixed gently. The MDMs were then incubated on ice for a further 10 min to
	8

1 2		
3 4 5	162	allow the beads to attach to the cell surface. Synchronised phagocytosis was then triggered by
6 7	163	warming the dish to 37 °C.
8 9 10	164	
11 12	165	2.9 Rate of Phagocytosis
13 14	166	After 60 min of synchronized phagocytosis of the beads by the MDMs at 37° C in a 5% CO ₂
15 16 17	167	incubator, phagocytosis was stopped by 3 washes with PBS before adding 2 mL of 4%
18 19	168	paraformaldehyde (Cat # P6148, Sigma-Aldrich, MO) and incubated at room temperature for 10
20 21	169	min for fixation. The fixed MDMs were then washed 3 times with PBS. Randomized images of a
22 23 24	170	total of 100 images were captured using scanning modes for red Alexa Fluor 594 fluorescent dye
24 25 26	171	(excitation 490 nm, emission 594 nm) using BioStation IM/IM-Q software version 2.21 available
27 28	172	from BioStation IM-Q Time Lapse Imaging System (Nikon, Japan). Images were acquired using
29 30 21	173	a x40 objective with randomized automated acquisition following the manufacturer's
32 33	174	instructions. The phagocytosed beads were manually counted by an investigator blinded to the
34 35	175	group assignment.
36 37	176	
38 39 40	177	2.10 Rate of Proteolysis
41 42	178	MDMs that adhered to the petri dish after undergoing synchronized phagocytosis were placed
43 44	179	into a 37 °C chamber containing 5% CO_2 on BioStation IM. The fluorescence intensities of green
45 46 47 48 49	180	reporter DQ BODIPY dye (excitation 490 nm, emission 525 nm) were acquired in 60 sec
	181	intervals for 60 min. Approximately 100 cells were analyzed per sample. The fluorescence
50 51	182	intensity was measured using an ImageJ (64-bit) software (NIH, imagej.nih.gov/ij/download/)
52 53 54		
54 55 56		9
57		

2 3		
4 5	183	and plotted against time using Prism 6 (GraphPad Software Inc., CA) for ratiometric data
6 7	184	analysis of intraphagosomal proteolysis.
8 9 10	185	
10 11 12	186	2.11 Measurement of ROS production
13 14	187	50 μ l of whole blood was incubated ± GM-CSF (5 ng/mL) in a total volume of 100 μ l in
15 16	188	phosphate-buffered saline. After 30 min incubation at 37°C, samples were treated with 5 μ M
17 18 19	189	dihydrorhodamine 123 followed by addition of either $1\mu M$ fMet-Leu-Phe or $0.1 \mu g/mL$ PMA for
20 21	190	a further 5- or 15 min, respectively at 37°C. Flow cytometry was used to measure the production
22 23	191	of ROS.
24 25 26	192	
27 28	193	2.12 Statistical analysis
29 30	194	All data were expressed as the mean \pm SEM. Statistical comparisons were performed using the <i>t</i>
31 32 22	195	test between groups. When serial measurements were taken over time, groups were compared
33 34 35	196	using two-way repeated-measures ANOVA with Bonferroni correction for multiple comparisons.
36 37	197	A <i>P value</i> < 0.05 was considered statistically significant. All data were analyzed using GraphPad
38 39	198	Prism 6.0 statistical software.
40 41 42	199	
43 44	200	3. RESULTS
45 46	201	3.1 MDMs from <i>O. viverrini</i> -infected individuals with APF had high phagocytic activity.
47 48 49	202	MDMs from O. viverrini-infected individuals with APF phagocytosed more beads per cell than
50 51	203	MDMs from healthy controls (Figures 1, A-D: 3.50 ± 0.09 vs 1.58 ± 0.87 ; n=102/group with
52 53	204	>100 cells analyzed per sample, p=0.01, unpaired t test) and O. viverrini-infected individuals
54 55		10
56 57 58		
59		

Page 11 of 27

3		
4 5	205	without APF (Figures 1, A-D: 3.50 ± 0.09 vs 1.48 ± 0.05 ; n=102/group with >100 cells analyzed
6 7	206	per patient, p<0.001, unpaired t test). We then calculated the percentage of macrophages that
8 9 10	207	phagocytosed the beads and found that the percentage of phagocytic cells was increased in
11 12	208	individuals with OV infection and increased further in APF+ individuals (Figure 1E). These data
13 14	209	provided the first indication of a possible association between macrophages with high phagocytic
15 16 17	210	activity and the presence of APF in chronic O. viverrini infection.
17 18 19	211	
20 21	212	3.2 MDMs from O. viverrini-infected individuals with APF had elevated phagosomal
22 23	213	proteolysis.
24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42	214	We then measured the rate of intraphagosomal proteolysis of individual ingested beads, as the
	215	rate of increased fluorescence of individual particles (increased fluorescence per ingested bead)
	216	by fluorescence imaging microscopy. The rate of proteolysis of BSA protein after phagocytosis
	217	(measured by BODIPY fluorescence intensity) by MDMs from O. viverrini-infected individuals
	218	with APF (Figure 2C) was greater than that from non-APF individuals (Figure 2B) and healthy
	219	controls (Figure 2A) at 30 min and 60 min after phagocytosis. The time course of BSA
	220	proteolysis clearly indicated that MDMs from patients with O. viverrini-induced APF proteolysed
	221	BSA faster than MDMs from the healthy controls (Figure 2D, n=102/group with 10-15 individual
43 44	222	beads measured for each sample, p<0.001, two-way repeated-measures ANOVA) and O.
45 46	223	viverrini-infected individuals without APF (Figure 2D, p<0.001, two-way repeated-measures
47 48 49	224	ANOVA).
50 51 52 53 54 55 56 57 58 59	225	11
59 60		

3.3 ROS production is elevated in monocytes from O. viverrini-infected individuals with APF.

Because macrophages used in previous two experiments are derived from peripheral blood monocytes, we then measured the production of ROS from circulating monocytes in these three cohorts. We found that there were no differences between groups if the cells in unprimed cells regardless of the stimulants. For example, there were no differences in ROS production in GM-CSF primed or unprimed cells when stimulated by the protein kinase C activator, PMA (data not shown). However, significant differences in ROS production were observed in response to the peptide, fMet-Leu-Phe which activates receptor-mediated stimulation of ROS, particularly after exposure to priming agents. In the absence of the priming agent, GM-CSF, there were no differences in ROS production between the three cohorts and fMet-Leu-Phe did not activate ROS in the absence of GM-CSF (Figure 3A-C). However, monocytes from O. viverrini-infected individuals with APF produced significantly more ROS than healthy control monocytes and monocytes from O. viverrini-infected individuals without APF (Figure 3D). These data suggest that circulating monocytes from APF+ individual are capable of generated higher levels of ROS after priming and stimulation.

4. DISCUSSION

Liver fluke infection has been widely known for more than two decades as a major cause of several hepatobiliary diseases including APF (4). The immunopathogenesis of APF is well described and it is partly induced by host adaptive immune responses through interaction of several immune cells (9) such as T cells (18), B cells (18), cytokines (11) and reactive oxygen

Page 13 of 27

Parasite Immunology

species (19). However, far less is known about the role of host innate immune responses on APF generation. Our studies demonstrate for the first time that MDMs from O. viverrini-infected individuals who have APF have high phagocytosis (Figures 1) and high proteolysis activity (Figure 2) when compared with both non-O. viverrini-infected controls and O. viverrini-infected individuals without APF. Furthermore, circulating monocytes from infected individuals with APF have elevated ability to generate ROS, compared to infected individuals without APF or healthy controls. This suggests that the presence of APF may associate with the pro-inflammatory phenotype of MDMs, consistent with several findings from our group and others (11, 14). Phagocytosis is one of several ways for uptake of antigen by macrophages (20). This process involves the vesicular internalization of solid insoluble particles and can subsequently activate NF-kB signal transduction to produce pro-inflammatory cytokines and antimicrobial peptides (21). These pro-inflammatory products are toxic to invading pathogens, but could also damage host tissues when deregulated, especially in chronic infection/inflammation. Thus, enhanced phagocytosis found in MDMs from APF+ individuals could result in an excessive production of pro-inflammatory mediators such as reactive oxygen species. Indeed, we measured enhanced ROS production by monocytes in whole blood and found that this was significantly elevated in infected individuals with APF. Reactive oxygen species could damage host tissues surrounding the bile ducts, which in turn, undergo repair processes, which result in the development of fibrosis (22). In support of this hypothesis, a previous study showed that a subset of MDMs promotes fibrogenesis after chronic O. viverrini-induced tissue damage (23). Another possible mechanism by which pro-inflammatory MDMs could lead to more advanced fibrosis is by elevated proteolytic activity. Proteolysis activity is tightly regulated by a

number of protease enzymes (24) and phagosomal pH (25) because effective proteolysis determines the fate of antigens, influencing antigen processing and presentation, which determines whether an efficient adaptive immune response is generated. As such, the elevated proteolysis that observed in this study in MDMs of APF+ individuals could induce stronger adaptive immune responses, which in turn, could promote robust inflammation, resulting in more severe host tissue damage and repair. Chronic infection leads to inflammation and the production of inflammatory mediators that are toxic to invading pathogens (25). In the case of helminth infection, the helminthes could modulate host immune responses and switch immune responses from proinflammatory (Th1) to anti-inflammatory (Th2) responses (26). The key innate immune cells that play a significant role in this switching are MDMs, which can switch from M1- to M2-like phenotypes that can induce Th1 and Th2, respectively. Once recruited to chronic inflamed sites in response to local stimuli, M1 macrophages drive Th1 responses for pathogen clearance, whereas M2 macrophages drive Th2 responses for tissue remodeling and fibrosis (12). Therefore, in future studies, identification of MDM subsets from these populations of *O. viverrini* infected individuals will provide a better understanding of the mechanisms underlying the development of APF. In conclusion, we measured and compared the phagocytic and proteolytic activities of MDMs between non-O. viverrini-infected control subjects, and O. viverrini-infected individuals with and without APF. We also showed elevated ROS production in those with APF. We propose that in the case of chronic O. viverrini infection, tissue resident macrophages with high phagocytic and proteolytic activities are the phenotypes that could damage bile duct epithelial and surrounding tissues, which results in the development of APF. Excessive production of

Page 15 of 27

Parasite Immunology

1 2		
3 4	292	proinflammatory mediators from these macrophages might have effects on the parasites, but
5 6 7	293	could also lead to excessive injury of surrounding tissue and hence result in extensive periductal
, 8 9	294	fibrosis.
10 11	295	
12 13	296	
14 15		
16 17	297	FUNDING
18 19	298	This work was supported by the Thailand Research Fund (TRF) grant number RTA 5680006, the
20 21	299	Office of Higher Education Commission of Thailand - the Newton Fund Institutional Links -
22 23 24	300	British Council, and partially from the National Institute of Allergy and Infectious Diseases
24 25 26	301	(NIAID), NIH, grant number P50AI098639. BS is a TRF Senior Research Scholar.
27 28	302	
29 30	303	ACKNOWLEDGEMENTS
31 32	304	We would like to thank John T. Cathey for editing the manuscript via the Publication Clinic
33 34 35	305	KKU, Thailand. The content is solely the responsibility of the authors and does not necessarily
36 37	306	represent the official views of the TRF, NIAID, OHEC-Newton Fund, or the NIH or the funders.
38 39	307	AUTHOR CONTRIBUTION
40 41	308	Conceived and design experiments: KS BS. Performed the experiments: KS EM BS. Analysis
42 43	309	and/or data interpretation: KS ST SS BS. Contributed reagents/materials/analysis/technical
44 45	310	expertise tools: EM SS ST SWE BS. Wrote the paper: KS SS ST SWE BS.
46 47	311	
48 49	312	CONFLICT OF INTERESTS STATEMENT
50 51 52 53	313	The authors declare no conflict of interests.
54 55		15
56 57		
58		
59 60		

1 2						
3 4 5	314	REFERENCES				
6 7	315	1.	Sripa B, Bethony JM, Sithithaworn P et al. Opisthorchiasis and Opisthorchis-			
8 9 10	316		associated cholangiocarcinoma in Thailand and Laos. Acta Trop 2011; 120 Suppl 1:			
11 12	317		S158-168.			
13 14 15	318	2.	Elkins DB, Mairiang E, Sithithaworn P et al. Cross-sectional patterns of hepatobiliar	у		
15 16 17	319		abnormalities and possible precursor conditions of cholangiocarcinoma associated			
18 19	320		with Opisthorchis viverrini infection in humans. Am J Trop Med Hyg 1996; 55: 295-	-		
20 21 22	321		301.			
23 24	322	3.	Mairiang E, Chaiyakum J, Chamadol N et al. Ultrasound screening for Opisthorchis			
25 26	323		viverrini-associated cholangiocarcinomas: experience in an endemic area. Asian Pa	IC		
27 28 29	324		<i>J Cancer Prev</i> 2006; 7: 431-433.			
30 31	325	4.	Mairiang E, Elkins DB, Mairiang P et al. Relationship between intensity of			
32 33	326		Opisthorchis viverrini infection and hepatobiliary disease detected by			
34 35 36	327		ultrasonography. <i>J Gastroenterol Hepatol</i> 1992; 7: 17-21.			
37 38	328	5.	Mairiang E and Mairiang P. Clinical manifestation of opisthorchiasis and treatment.			
39 40	329		<i>Acta Trop</i> 2003; 88: 221-227.			
41 42 43	330	6.	Sripa B, Kaewkes S, Sithithaworn P et al. Liver fluke induces cholangiocarcinoma.			
44 45	331		<i>PLoS Med</i> 2007; 4: e201.			
46 47 48	332	7.	Wynn TA and Ramalingam TR. Mechanisms of fibrosis: therapeutic translation for			
40 49 50	333		fibrotic disease. <i>Nat Med</i> 2012; 18: 1028-1040.			
51 52	334	8.	Borthwick LA, Wynn TA and Fisher AJ. Cytokine mediated tissue fibrosis. <i>Biochim</i>			
53 54 55	335		<i>Biophys Acta</i> 2013; 1832: 1049-1060.	16		
55 56 57				10		
58 59						
60						

Page 17 of 27

1

Parasite Immunology

2 3			
4 5 6 7 8 9 10 11 12	336	9.	Sripa B, Brindley PJ, Mulvenna J et al. The tumorigenic liver fluke Opisthorchis
	337		<i>viverrini</i> multiple pathways to cancer. <i>Trends Parasitol</i> 2012; 28: 395-407.
	338	10.	Pinlaor S, Sripa B, Sithithaworn P and Yongvanit P. Hepatobiliary changes, antibody
	339		response, and alteration of liver enzymes in hamsters re-infected with Opisthorchis
13 14 15	340		viverrini. Exp Parasitol 2004; 108: 32-39.
15 16 17	341	11.	Sripa B, Thinkhamrop B, Mairiang E et al. Elevated plasma IL-6 associates with
 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 	342		increased risk of advanced fibrosis and cholangiocarcinoma in individuals infected
	343		by <i>Opisthorchis viverrini. PLoS Negl Trop Dis</i> 2012; 6: e1654.
	344	12.	Murray PJ and Wynn TA. Protective and pathogenic functions of macrophage
	345		subsets. <i>Nat Rev Immunol</i> 2011; 11: 723-737.
	346	13.	Weidenbusch M and Anders HJ. Tissue microenvironments define and get reinforced
	347		by macrophage phenotypes in homeostasis or during inflammation, repair and
	348		fibrosis. <i>J Innate Immun</i> 2012; 4: 463-477.
	349	14.	Mairiang E, Laha T, Bethony JM et al. Ultrasonography assessment of hepatobiliary
	350		abnormalities in 3359 subjects with Opisthorchis viverrini infection in endemic
	351		areas of Thailand. <i>Parasitol Int</i> 2012; 61: 208-211.
41 42 43	352	15.	Sripa B, Mairiang E, Thinkhamrop B et al. Advanced periductal fibrosis from
44 45	353		infection with the carcinogenic human liver fluke Opisthorchis viverrini correlates
46 47	354		with elevated levels of interleukin-6. <i>Hepatology</i> 2009; 50: 1273-1281.
48 49 50	355	16.	Menck K, Behme D, Pantke M et al. Isolation of human monocytes by double gradient
50 51 52	356		centrifugation and their differentiation to macrophages in teflon-coated cell culture
53 54	357		bags. <i>J Vis Exp</i> 2014: e51554.
55 56 57			17
58 59			
60			

Page 18 of 27

1 2 3			
3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	358	17.	Yates RM and Russell DG. Phagosome maturation proceeds independently of
	359		stimulation of toll-like receptors 2 and 4. <i>Immunity</i> 2005; 23: 409-417.
	360	18.	Wongratanacheewin S, Good MF, Sithithaworn P and Haswell-Elkins MR. Molecular
	361		analysis of T and B cell repertoires in mice immunized with Opisthorchis viverrini
	362		antigens. Int J Parasitol 1991; 21: 719-721.
	363	19.	Saichua P, Yakovleva A, Kamamia C et al. Levels of 8-oxodG predict hepatobiliary
	364		pathology in Opisthorchis viverrini endemic settings in Thailand. PLoS Negl Trop Di
	365		2015; 9: e0003949.
	366	20.	Stuart LM and Ezekowitz RA. Phagocytosis: elegant complexity. <i>Immunity</i> 2005; 22:
25 26	367		539-550.
27 28 29	368	21.	Kawai T and Akira S. The role of pattern-recognition receptors in innate immunity:
30 31	369		update on Toll-like receptors. <i>Nat Immunol</i> 2010; 11: 373-384.
32 33	370	22.	Duffield JS, Lupher M, Thannickal VJ and Wynn TA. Host responses in tissue repair
34 35 36	371		and fibrosis. <i>Annu Rev Pathol</i> 2013; 8: 241-276.
37 38	372	23.	Bility MT and Sripa B. Chronic Opisthorchis viverrini infection and associated
39 40	373		hepatobiliary disease is associated with iron loaded M2-like macrophages. Korean J
41 42 43	374		<i>Parasitol</i> 2014; 52: 695-699.
44 45	375	24.	Reiser J, Adair B and Reinheckel T. Specialized roles for cysteine cathepsins in health
46 47	376		and disease. <i>J Clin Invest</i> 2010; 120: 3421-3431.
40 49 50	377	25.	Watts C. The endosome-lysosome pathway and information generation in the
51 52	378		immune system. Biochim Biophys Acta 2012; 1824: 14-21.
53 54			1
55 56 57			
58 59			
60			

1 2 2				
5 4 5	379	26.	Salgame P, Yap GS and Gause WC. Effect of helminth-induced immunity on infection	ons
6 7 8	380		with microbial pathogens. <i>Nat Immunol</i> 2013; 14: 1118-1126.	
9 10				
11 12 13	382			
14 15				
16 17 18				
19 20				
21 22 23				
24 25				
26 27 28				
28 29 30				
31 32				
33 34 35				
36 37				
38 39 40				
41 42				
43 44 45				
46 47				
48 49 50				
50 51 52				
53 54				10
55 56 57				19
58 59				
60				

383 FIGURE LEGENDS

7 384

FIGURE 1 MDMs from individuals with *O. viverrini*-induced APF phagocytosed more beads.

Numbers of phagocytosed opsonized silica beads covalently coupled to Alexa Fluor 594 were manually counted after synchronized phagocytosis for 60 min with MDMs from healthy individuals (A), O viverrini-infected individual with no APF (B) and O viverrini-infected individual with APF (C). Average number of phagocytosed beads per cell was calculated by manual counts (D). The % of phagocytic cells was also calculated (E). Data represents mean ± SEM analyzed using the *t* test. * indicates a p value of <0.05, and mean values are of at least 20 individual cells (chosen blinded) for each of the 102 samples in each cohort.

30 394

395 FIGURE 2 Proteolysis of MDMs from individuals with *O. viverrini-*induced with and

without APF and healthy controls. Live MDMs that had undergone synchronized phagocytosis of silica beads covalently coupled with DQ-BODIPY BSA were continuously monitored for 60 min using an inverted live cell BioStation IM microscope. Representative images are shown in Figures A-C. Upper panels are bright-field images while lower panels are fluorescence images. The time course of proteolytic activity within the phagosome, measured as gain of fluorescence,

401 is shown in Figure **D**. Data are mean \pm SEM analyzed using two-way repeated-measures

402 ANOVA of 10-15 individual beads measured in each of the 102 samples in each cohort. *

403 indicates a p value of <0.05

Parasite Immunology

1 2 3						
3 4 5	405					
6 7 8 9 10 11 23 14 15 16 7 8 9 10 11 23 24 25 26 27 8 9 30 32 33 45 36 37 8 9 40 41 23 44 56 7 8 9 0 12 23 24 25 26 27 8 9 30 31 23 34 56 7 8 9 40 41 45 46 7 8 9 20 21 22 24 25 26 7 8 9 30 31 23 34 56 7 8 9 40 41 45 46 7 8 9 20 21 22 23 24 52 67 8 9 30 31 23 34 53 67 7 8 9 40 41 42 44 45 67 7 8 9 20 21 22 32 45 26 7 8 9 30 31 23 34 5 36 7 8 9 40 41 42 44 45 67 8 9 40 41 42 44 45 67 8 9 40 41 42 44 45 67 8 9 40 41 42 44 45 67 8 9 40 41 42 44 45 67 8 9 9 0 12 23 24 25 26 27 28 9 30 32 33 45 36 7 8 9 40 41 42 44 45 67 55 55 54 55 55 54 55 55 55 55 55 55 55	406					
	407	Figure 3 ROS production of monocytes in whole blood				
	408	Whole blood from healthy controls (OV-) or OV-infected individuals without (OV+/APF-) or				
	409	with advanced periductal fibrosis (OV+/APF+) were incubated in the absence (unprimed) or				
	410	presence of 5 ng/mL GM-CSF for 30 min. DHR 123 (at 5 μ M) was then added and then primed				
	411	or unprimed cells were incubated with or without fMet-Leu-Phe for 5 min before analysis by				
	412	flow cytometry and gating of monocyte responses. Figure 3A-C show mean fluorescent				
	413	intensities of DHR123 from OV-, OV+/APF-, OV+/APF+ respectively. Figure 3D, () shows				
	414	fMet-IPhe responses in unprimed cells, () GM-CSF responses in the absence of fMet-				
	415	Leu-Phe and () ROS production (
	416	GM-CSF primed responses. Data represents mean \pm SEM analyzed using the <i>t</i> test. * indicates a				
	417 418	p value of <0.05.				
		1				
55 56 57 58 59		21				



142x185mm (300 x 300 DPI)

calculated (E). Data represents mean \pm SEM analyzed using the t test. * indicates a p value of <0.05, and

mean values are of at least 20 individual cells (chosen blinded) for each of the 102 samples in each cohort.



FIGURE 2 Proteolysis of MDMs from individuals with O. viverrini-induced with and without APF and healthy controls. Live MDMs that had undergone synchronized phagocytosis of silica beads covalently coupled with DQ-BODIPY BSA were continuously monitored for 60 min using an inverted live cell BioStation IM microscope. Representative images are shown in Figures A-C. Upper panels are bright-field images while lower panels are fluorescence images. The time course of proteolytic activity within the phagosome, measured as gain of fluorescence, is shown in Figure D. Data are mean ± SEM analyzed using two-way repeated-measures ANOVA of 10-15 individual beads measured in each of the 102 samples in each cohort. * indicates a p value of <0.05

168x113mm (300 x 300 DPI)





Whole blood from healthy controls (OV-) or OV-infected individuals without (OV+/APF-) or with advanced periductal fibrosis (OV+/APF+) were incubated in the absence (unprimed) or presence of 5 ng/mL GM-CSF for 30 min. DHR 123 (at 5 μM) was then added and then primed or unprimed cells were incubated with or without fMet-Leu-Phe for 5 min before analysis by flow cytometry and gating of monocyte responses. Figure 3A-C show mean fluorescent intensities of DHR123 from OV-, OV+/APF-, OV+/APF+ respectively. Figure 3D, () shows fMet-Leu-Phe responses in unprimed cells, () GM-CSF responses in the absence of fMet-Leu-Phe and () ROS production by fMet-Leu-Phe in GM-CSF cells. N = 51 per group for GM-CSF primed responses. Data represents mean ± SEM analyzed using the t test. * indicates a p value of <0.05.

225x186mm (300 x 300 DPI)

Reviewer: 1

Comments to the Author

Review on "Association between macrophages phagocytosis activities and advanced periductal fibrosis in chronic opisthorchiasis"

The presented manuscript analyses in vitro generated macrophages (MDM) from patients infected with O.viverrini exhibiting (APF+) or free from (APF-) signs of advanced periductal fibrosis. The authors report an enhanced uptake of opsonised silica beads by MDM from APF+ patients. The study seems for the most part technically sound and is well described. However the conclusions drawn seem rather speculative and are only loosely based on the presented data. Overall the extent of the presented findings is very limited and would need further experimental work to corroborate the authors claims.

<u>Response:</u> Thank you for your comments! We agree that the work presented in the previous submission was limited as only phagocytotic and proteolytic assays were included. We have therefore performed more experiments to support our claims and conclusions. We have now measured and compared ROS among these three groups. The results can be found from lines 230-245 and Figure 3.

Individual comments:

The conclusions drawn by the authors seem speculative and one-sided. Phagocytosis per se is not pro-inflammatory but can be anti-inflammatory and prevent chronic inflammation (e.g. phagocytosis of apoptotic cells during cystic fibrosis). Thus, the authors would need to include further data on cytokine production or other inflammatory mediators to clarify whether the enhanced rate of phagocytosis promotes or protects from APF.

Furthermore, the authors repeatedly imply that the altered phagocytosis rate observed in macrophages from APF+ patients may be causative for the formation of fibrosis. However, this is not supported by the presented data. Indeed, since none of the APF- patients (some of whom will develop APF eventually) show altered macrophage phagocytosis it would seem this is a consequence rather than a cause of fibrosis.

<u>Response:</u> In order to address this, we have now performed ROS assays. We found that ROS production by blood monocytes was higher in individuals with APF compared with healthy controls and infected individuals without APF. ROS is a key inflammatory mediator that could cause tissue damage if overexpressed. We believe that elevated ROS production can therefore contribute to tissue damage and promote persistent inflammation. Also, we have re-examined the phagocytosis data and have measured the % of monocytes that underwent phagocytosis (new Figure 1E). This new figure shows that in infected individuals without APF, a greater % of the population of macrophages were capable of phagocytosis, compared to non-infected controls. Therefore, infection *per se* leads to a great phagocytic activity of these macrophages.

The data presented in figure 3 is inconclusive and does not reflect on the rate of proteolysis in MDM. If I understand the experimental procedure correctly macrophages were incubated with BSA-Bodipy coupled silica beads and the release of Bodipy measured over time. However, the authors have already shown that MDM from APF+ patients take up more beads in a similar assay. Thus, the increased fluorescent signal might not reflect an altered rate of proteolysis but rather a higher uptake of substrate by the APF+ MDM. Therefore this data cannot be interpreted beyond what has already been shown in figure 1&2 and should be removed. Alternatively the authors would need to provide evidence of substrate normalisation.

<u>Response:</u> This is a very good point raised by the referee but unfortunately, we did not adequately explain the experiment or how the data were obtained. In fact, we actually measured the increase in fluorescence of individual ingested beads. We tracked the release of bodiopy from individual engulfed beads over the first 60 min of phagocytosis, measuring the fluorescence of 10-15 individual beads per patient sample (n= 102 patient samples). We apologise for not making the nature of this measurement clear in our first submission. We added a clearer explanation in the manuscript in line 218-220.

The flowcytometry data presented in figure 2 and the microscopy pictures shown in figure 1 seem to contradict each other. Whereas the microscopy suggests virtually all macrophages do phagocytose opsonised silica beads only about 1% of cells do so according to the flow cytometry data. Is this due to variation in donors and the specific datasets chosen? If so it might be more prudent to chose more representative images / flow-plots.

Also, please include the flow cytometry data as % positive cells rather than or in addition to the MFI of the whole population.

<u>Response:</u> We agree with these comments. We believe that the phagocytic capacity of the cultured macrophages was grossly perturbed when we scraped them from the culture plates prior to flow cytometry, which explains their very low activity. Therefore, we have removed this figure from the revised manuscript as we believe that the data are not representative of the true phagocytic activity of these cells. The images of phagocytosis shown in Figure 1A-C are single images but we have quantified the number of ingested particles and (new Figure 1E) the number of cultured macrophages (expressed as a % of the total number), that could undergo phagocytosis. We analysed at least 100 cells/patient and believe that these are a truer representation of the phagocytic capacity of these cells.

Since the authors use autologous serum to generate MDMs in vitro they should include data on the differentiation and activation status (i.e. MHC-expression, costimulatory molecules etc.) of the macrophages from the three groups. Does serum from APF+ patients induce altered differentiation / pre-activation of MDM?

<u>Response:</u> These are extremely helpful comments, but we now cannot do this retrospectively. We will include such measurements in a new study.

Thank you for your invaluable comments.

Reviewer: 2

Comments to the Author

The manuscript entitled "Association between macrophages phagocytosis activities and advanced periductal fibrosis in chronic opisthorchiasis" addresses a major gap in the knowledge on the pathogenesis of chronic opisthorchiasis, namely the role innate immune cells such as macrophages. The authors demonstrate that elevated macrophage phagocytic and proteolytic activities are associated with chronic OV-induced liver fibrosis. This study will establish the basis for more mechanistic studies on the role of macrophages in OV-induced liver disease.

Response: Thank you for your comments!