

Sequential infection can decrease virulence in a fish-bacterium-fluke interaction: implications for aquaculture disease management

Anssi Karvonen¹, Andy Fenton² & Lotta-Riina Sundberg^{1,3}

¹ University of Jyväskylä, Department of Biological and Environmental Science, P.O. Box 35, 40014 University of Jyväskylä, Finland

² University of Liverpool, Institute of Integrative Biology, Liverpool, UK.

³ University of Jyväskylä, Nanoscience Center, P.O. Box 35, 40014 University of Jyväskylä, Finland

Correspondence: Anssi Karvonen (anssi.t.karvonen@jyu.fi)

Running head: Sequential infection and virulence

Acknowledgements

The authors would like to thank Jenni Marjakangas for assistance in the laboratory, and Elina Laanto and Heidi Kunttu for the bacterial isolates. Kayla King gave valuable comments on an earlier draft. This work was supported by the Finnish Centre of Excellence Program of the Academy of Finland; the CoE in Biological Interactions 2012-2017 (#252411), and by the Academy of Finland grants #263864, #266879, #292763, #310632, and #314939.

1 **Abstract**

2

3 Hosts are typically infected with multiple strains or genotypes of one or several parasite
4 species. These infections can take place simultaneously, but also at different times, i.e.
5 sequentially, when one of the parasites establishes first. Sequential parasite dynamics are
6 common in nature, but also in intensive farming units such as aquaculture. However,
7 knowledge of effects of previous exposures on virulence of current infections in intensive
8 farming is very limited. This is critical as consecutive epidemics and infection history of a
9 host could underlie failures in management practises and medical intervention of diseases.
10 Here, we explored effects of timing of multiple infection on virulence in two common
11 aquaculture parasites, the bacterium *Flavobacterium columnare* and the fluke *Diplostomum*
12 *pseudospathaceum*. We exposed fish hosts first to flukes and then to bacteria in two separate
13 experiments, altering timing between the infections from few hours to several weeks. We
14 found that both short-term and long-term difference in timing of the two infections resulted in
15 significant, genotype-specific decrease in bacterial virulence. Second, we developed a
16 mathematical model, parameterized from our experimental results, to predict the implications
17 of sequential infections for epidemiological progression of the disease, and levels of fish
18 population suppression, in an aquaculture setting. Predictions of the model showed that
19 sequential exposure of hosts can decrease the population-level impact of the bacterial
20 epidemic, primarily through the increased recovery rate of sequentially infected hosts,
21 thereby substantially protecting the population from the detrimental impact of infection.
22 However, these effects depended on bacterial strain – fluke genotype combinations,
23 suggesting the genetic composition of the parasite populations can greatly influence the
24 degree of host suppression. Overall, these results suggest that host infection history can have
25 significant consequences for the impact of infection at host population level, potentially

26 shaping parasite epidemiology, disease dynamics and evolution of virulence in farming
27 environments.

28

29 Keywords: Dynamic infection, Epidemiology, Multiple infection, Sequential infection,
30 Spatiotemporal variation

31

32

33 **1. Introduction**

34

35 Hosts are commonly infected with multiple parasite species or strains/genotypes of one
36 species at the same time (Graham, 2008; Read & Taylor, 2001; Salgame, Yap, & Gause,
37 2013; Telfer et al., 2010). Such infections can result in direct (interference competition) or
38 indirect (resource or host immune-mediated competition) interactions between parasites and
39 have significant implications for key parasite traits such as virulence, harm to the host (Bell,
40 Roode, Sim, & Read, 2006; Ben-Ami, Regoes, & Ebert, 2008; Davies, Fairbrother, &
41 Webster, 2002; de Roode, Helinski, Anwar, & Read, 2005). Recent studies have emphasised
42 the importance of multiple infections also between phylogenetically distant parasites (Ben-
43 Ami, Rigaud, & Ebert, 2011; Clay, Dhir, Rudolf, & Duffy, 2019; Doublet, Natsopoulou,
44 Zschiesche, & Paxton, 2015; Duncan, Agnew, Noel, & Michalakis, 2015; Fellous & Koella,
45 2009; Lohr, Yin, & Wolinska, 2010; Vojvodic, Boomsma, Eilenberg, & Jensen, 2012),
46 suggesting that interactions can take place at the scale of the entire parasite community of one
47 host. It is common that infections from different parasites do not occur only simultaneously,
48 but also sequentially at different times as the infection risk in nature varies over time (e.g.,
49 seasons (Faltýnková, Valtonen, & Karvonen, 2008; Karvonen, Seppälä, & Valtonen, 2004))
50 and space (e.g., spatial aggregation of infected hosts (Byers, Blakeslee, Linder, Cooper, &

51 Maguire, 2008; Jokela & Lively, 1995; King, Delph, Jokela, & Lively, 2009)). The timing
52 between different infections again can vary from a few hours to several weeks, or even years.
53 Consequently, each individual host can have a different infection history and immunological
54 status, thus making the landscape of disease outcomes complex and unpredictable. Empirical
55 examples of sequential infections of multiple parasites in plants (Hood, 2003; Laine, 2011;
56 Marchetto & Power, 2018), invertebrates (Ben-Ami, Mouton, & Ebert, 2008; Ben-Ami et al.,
57 2011; Gower & Webster, 2005; Lohr et al., 2010; Natsopoulou, McMahon, Doublet, Bryden,
58 & Paxton, 2015) and vertebrates (Graham, 2008; Hoverman, Hoye, & Johnson, 2013;
59 Klemme, Louhi, & Karvonen, 2016), suggest an effect of sequential infection of hosts on
60 parasite fitness-related traits such as infection success and virulence.

61

62 Infections from multiple parasites are common also in intensive production environments,
63 where high densities of susceptible hosts favour the spread of virulent pathogens (Kennedy et
64 al., 2016; Pulkkinen et al., 2010). Infections can cause significant economic loss by impairing
65 quality, condition and growth of crops and farmed animals. For example, in aquaculture,
66 parasitic infections are considered one of the most important threats for development of the
67 industry. Similar to natural conditions, parasitic epidemics in aquaculture are typically
68 consecutive with different parasites infecting their hosts in varying timescales. Disease
69 epidemics typically sweep through aquaculture units at different times in response to
70 variation in pathogen ecology and host susceptibility (e.g. cohorts of varying age) (Karvonen,
71 Rintamäki, Jokela, & Valtonen, 2010; Rintamäki-Kinnunen & Valtonen, 1997). This creates
72 favourable conditions for development of cumulative infection history of hosts that can affect
73 virulence in subsequent disease outbreaks. Earlier studies in fish have shown that a prior
74 parasite exposure can influence the outcome of simultaneous re-exposure of the host by
75 multiple parasite genotypes (Klemme et al., 2016), alter associations between parasite species

76 (Karvonen, Seppälä, & Valtonen, 2009), and influence parasite community composition of
77 the host (Benesh & Kalbe, 2016). However, knowledge of the implications of sequential
78 exposure of hosts to multiple parasites in farming environments is still very limited.
79 Consequently, most infections occurring in intensive farming units are commonly treated
80 instantaneously with very little consideration of previous or existing other infections, which,
81 among aquaculture fish, can range from viruses and bacteria (Mohanty & Sahoo, 2007;
82 Pulkkinen et al., 2010; Skall, Olesen, & Møllgaard, 2005; Tøbbach, Decostere, Hermans,
83 Haesebrouck, & Chiers, 2007) to protozoans and metazoans (Hakalahti & Valtonen, 2003;
84 Karvonen, Savolainen, Seppälä, & Valtonen, 2006; Rintamäki-Kinnunen & Valtonen, 1997).
85 Regardless, infection history of a host population could influence ongoing epidemics and
86 potentially underlie failures in management practises and medical intervention of diseases.

87

88 Here, we studied effects of host sequential exposure on parasite virulence in an interaction
89 between two widely distributed aquaculture parasites, the bacterium *Flavobacterium*
90 *columnare* and the fluke *Diplostomum pseudospathaceum*. Bacterium *F. columnare*, the
91 causative agent of the columnaris disease, is an opportunistic pathogen and currently
92 considered as one of the most severe disease threats in fish farming (Declercq, Haesebrouck,
93 Van den Broeck, Bossier, & Decostere, 2013). The disease can cause considerable losses if
94 not treated with antibiotics (Declercq, Haesebrouck, et al., 2013; Pulkkinen et al., 2010;
95 Wagner, Wise, Khoo, & Terhune, 2002), which in many cases has resulted in emergence of
96 antibiotic-resistant bacterial strains (Declercq, Boyen, et al., 2013). The trematode *D.*
97 *pseudospathaceum* causes local, but significant aquaculture problems by blinding fish
98 (Karvonen, 2012). Unlike *F. columnare*, infections of *D. pseudospathaceum* are not
99 transmitted directly between fish, but the life cycle includes three hosts (snail, fish and fish-
100 eating bird) and fish become infected when in contact with the parasite larvae (cercariae)

101 released from infected snails. Infections of *F. columnare* and *D. pseudospathaceum* can co-
102 occur in aquaculture fish (Karvonen et al., 2006; Sundberg et al., 2016). They also interact in
103 genotype-specific manner when infecting the host at the same time, which can result in
104 higher morbidity of fish, i.e. virulence, and higher infection success of the fluke (Louhi,
105 Sundberg, Jokela, & Karvonen, 2015).

106

107 We first exposed rainbow trout (family Salmonidae) hosts to both parasites in two
108 experiments manipulating the timing between the infections from few hours to several weeks
109 and monitoring the disease-related morbidity of the fish. Based on our previous results on
110 simultaneous infections of the two parasites (Louhi et al., 2015), we expected that both the
111 short-term and the long-term sequence between the infections would result in lower bacterial
112 virulence, possibly depending on the strain-genotype combinations of the parasites. Similarly,
113 we expected sequential infection to change infection success of the fluke. Second, we
114 developed a compartmental mathematical model capturing the disease dynamics in a host
115 population to explore how sequential exposure of hosts to the two parasites could influence
116 disease-related mortality, and total host population size, of farmed fish. Overall, our results
117 suggest that host infection history can potentially shape parasite virulence over a long time
118 period, which may have implications for evolution of virulence as well as for disease
119 prevention strategies in intensive farming systems.

120

121 **2. Material and methods**

122

123 *(a) Bacterial cultures*

124

125 Three *Flavobacterium columnare* strains (1-3, Supporting information, Table S1) differing in
126 their virulence were used (Kunttu, Sundberg, Pulkkinen, & Valtonen, 2012; Laanto,
127 Bamford, Laakso, & Sundberg, 2012). The strains had originally been isolated from fish
128 farms or from environment in 2008-2010 by standard culture methods using Shieh medium
129 (Song, Fryer, & Rohovec, 1988) and Shieh medium supplemented with tobramycin
130 (Decostere, Haesebrouck, & Devriese, 1997). Different sampling locations, sampling times,
131 sources of isolation (fish vs. environment), ARISA groups (Table S1), differences in
132 CRISPR-Cas sequences (Laanto, Hoikkala, Ravantti, & Sundberg, 2017) and the different
133 pathogenicity of the isolates (Kunttu et al., 2012; Laanto et al., 2012) ensured that the strains
134 differed in genetic and/or ecological characteristics. Cultures were stored at -80°C with 10%
135 glycerol and 10% fetal calf serum. Prior to the exposures, bacterial strains were grown
136 overnight in 2 ml of Shieh medium, then enriched in 1:10 in fresh medium and incubated at
137 25°C with 150 rpm agitation for 22 h. The optical density (OD, A570) of the culture was
138 measured with spectrophotometer, and the corresponding colony forming units (CFU) were
139 calculated using a previously determined relationship between OD and CFU (unpublished).

140

141 *(b) Sampling and genotyping of flukes*

142

143 *Lymnaea stagnalis* snails ($n = 42$), intermediate hosts for *D. pseudopathaceum*, shedding
144 clonal fluke cercariae were collected from Lake Vuojärvi (62° 24' 54" N, 25° 56' 14" E),
145 Finland. Fifteen cercariae were collected from each snail and stored individually in
146 Eppendorf tubes in 15 µl of lake water and frozen in -20°C for subsequent microsatellite
147 analysis to identify snails that were infected with one fluke genotype (Louhi, Karvonen,
148 Rellstab, & Jokela, 2010; Reusch, Rauch, & Kalbe, 2004) (Table S2). Parasite DNA was
149 extracted according to Criscione and Blouin (2004). Snails infected with one genotype were

150 stored individually in 1 l of water at 6°C and fed *ad libitum* with lettuce until the beginning of
151 the experiment. Note that all parasite genotypes are produced sexually in the avian definitive
152 host, which is why all infections in the snails are unique and individual genotypes persist in a
153 host population only for one complete round of the parasite life cycle.

154

155 (c) *Experimental exposure 1*

156

157 Naïve, uninfected juvenile rainbow trout (*Oncorhynchus mykiss*; age 2.5 months, average
158 length \pm s.e = 38.23 \pm 0.2 mm) were obtained from a hatchery in Central Finland. Fish were
159 maintained in aerated ground water with continuous water flow (17°C) for four weeks before
160 the experiments and fed daily with commercial fish food pellets. Prior to the exposures, the
161 water temperature was raised slowly to 25°C (2°C every second day) to allow fish
162 acclimation to experimental conditions. Three freshly grown strains of *F. columnare* (1-3, see
163 Table S1) and clonally produced cercaria larvae of *D. pseudospathaceum* from three
164 single-genotype infected snails (A-C; see Table S2) were used in the fish exposures. Three
165 hours prior to the exposures, the snails were placed individually in 2.5 dl of water (20°C), and
166 allowed to produce cercariae. Cercarial density from each snail (fluke genotype) was
167 estimated by counting ten times 1 ml subsamples from each container.

168

169 A pairwise infection design was then applied to test virulence and intensity of infection
170 across the combinations (Table S3). In the experiment, 20 rainbow trout were exposed
171 individually to single bacterial strains (5×10^3 colony forming units ml⁻¹; 3 \times 20 fish), single
172 fluke genotypes (50 cercariae/fish; 3 \times 20 fish), or co-exposed to both bacteria and flukes in
173 nine different combinations (9 \times 20 fish), totalling 300 fish. To explore the effect of short-
174 term sequential infection on virulence, the co-exposure matrix (9 \times 20 fish) was replicated

175 together with the simultaneous infections so that each of the nine co-exposure combinations
176 received the fluke first and the bacterium 4 h later. Bacterial single infections (3×20 fish)
177 were also repeated at this time with freshly grown strains to control for possible changes in
178 bacterial virulence. A negative control group of 30 fish receiving pure culture medium and/or
179 water instead of bacteria or flukes, respectively, was also established. Overall, the setup
180 totalled 570 fish (Table S3). The infection doses corresponded to those in natural conditions.
181 For example, fish infected with *F. columnare* can emit bacterial concentrations that are orders
182 of magnitude higher than those used here (Kunttu, Valtonen, Jokinen, & Suomalainen, 2009)
183 and one infected snail can release thousands of *D. pseudospathaceum* cercariae per day
184 (Karvonen, Kirsi, Hudson, & Valtonen, 2004; Karvonen, Rellstab, Louhi, & Jokela, 2012).
185 All fish were haphazardly assigned to the different treatment groups (single exposure to *F.*
186 *columnare*, single exposure to *D. pseudospathaceum*, exposure to both parasites) in the
187 simultaneous and sequential exposures.

188

189 The exposures and the subsequent monitoring took place in small containers with 500 ml
190 pre-aerated ground water (25°C). The fish were checked every hour for disease symptoms
191 and morbidity. Morbid fish that had lost their natural swimming buoyancy and did not
192 respond to external stimuli were considered dead and were euthanized using MS-222
193 anaesthetic every hour. This gave an accurate estimate of time of death (see Louhi et al.
194 (2015)). The fish were immediately sampled for presence of *F. columnare* on the skin and
195 gills (by culture on Shieh containing tobramycin (Decostere et al., 1997)), and dissected for
196 *D. pseudospathaceum* in the eye lenses. The establishment of *D. pseudospathaceum* in the
197 eye lenses takes place within few hours from exposure (Louhi et al., 2015). The dissection
198 protocol was used to determine the exact shape of the time-establishment relationship used in
199 estimation of differences in fluke abundance among the treatment groups (see below). The

200 experiment was terminated at 28 h post-exposure when 87.5% of the fish exposed to the
201 bacterium had died. All surviving fish were subsequently euthanized (MS-222) and examined
202 for bacterial and fluke infection as described above. Bacterial cultures confirming *F.*
203 *columnare* infection were incubated at 22°C for two days and checked for presence of
204 bacterial colonies.

205

206 Data on fish survival were analysed using Cox regression with sequential infection, *F.*
207 *columnare* strain and *D. pseudospathaceum* genotype as fixed covariates, and fish length as a
208 continuous covariate. Since the bacterial virulence changed slightly during the 4 h interval
209 between the infections (see results), fish groups exposed only to the bacterium at 0 h and 4 h
210 were used as reference categories in the analysis. Thus, the effect of sequential infection on
211 virulence would be seen as a significant three-way interaction between the fixed factors. In
212 addition, we applied analysis of covariance (ANCOVA) to data on fluke numbers in fish eyes
213 to identify factors that affected infection intensity in exposures together with the bacterial
214 strains. To correct for variation in fluke exposure and establishment time between fish
215 individuals showing different time of survival, we used the residuals of the non-linear
216 asymptotic regression predicting infection intensity as function of survival time as the
217 response variable (see Louhi et al. (2015)).

218

219 *(d) Experimental exposure 2*

220

221 To explore the longer-term effect of sequential exposure, 960 rainbow trout from the same lot
222 as in Experiment 1 were divided into 6 tanks, each with 160 fish and 70 l of water (16°C).
223 Three of the groups were exposed to a total of 480 *D. pseudospathaceum* cercariae per tank, 3
224 cercariae per fish, while the other three groups served as unexposed controls. A low-dose

225 exposure was used to keep the number of parasites establishing in eye lenses low so that the
226 parasite would not influence subsequent fish growth (Karvonen, 2012). Parasite cercariae
227 were produced from two *L. stagnalis* snails as described above and pooled for the exposure
228 (different genotypes to those used in the re-exposure below (D-F), or in Experiment 1 (A-C)).
229 During the exposure, the incoming water was turned off and was turned back on after 1 h. As
230 the cercarial infective lifespan is less than 30 h (Karvonen, Paukku, Valtonen, & Hudson,
231 2003), parasites that failed to locate a fish, if any, were eventually lost from the tanks. Fish
232 were then maintained for five weeks and fed daily with fish pellets. Possible acquired host
233 responses against *D. pseudospathaceum* are cross-reactive across parasite genotypes
234 (Rellstab, Karvonen, Louhi, & Jokela, 2013), which minimized genotype-specific responses,
235 if any, between the first infection and the re-exposure (see below). Water temperature was
236 then raised slowly to 25°C as described above. Fish with and without the previous fluke
237 infection were exposed either to single *Flavobacterium* (strains 1-3, Table S1; 2 × 3 × 20
238 fish), single *Diplostomum* (genotypes D-F, note different genotypes to Experiment 1 because
239 of mortality among the snails carrying genotypes A-C, Table S2; 2 × 3 × 20 fish), or to
240 pairwise combinations of the two (9 different combinations; 2 × 9 × 20 fish) (Table S4).
241 Doses of the bacteria (5×10^3 colony forming units ml⁻¹) and flukes (50 cercariae/fish) at re-
242 exposure were the same as in the first experiment. Each treatment had 20 replicate fish taken
243 randomly from groups of previously unexposed and exposed fish (the three replicate tanks
244 were pooled). A negative control group of 30 fish was also established. The entire setup
245 totalled 630 fish (Table S4). Again, fish were maintained individually and followed for
246 disease symptoms until 28 h post-exposure as described above. All fish that died or survived
247 the experiment were sampled for bacterial presence of the skin and gills, and dissected for the
248 number of flukes. Flukes originating from the first (3 cercariae per fish) and the second (50
249 cercariae per fish) exposure were separated according to their size.

250

251 Data were analysed using Cox regression with initial fluke infection, *F. columnare* strain and
252 *D. pseudospathaceum* genotype as fixed covariates, and fish length as a continuous covariate.
253 Analysis of covariance (ANCOVA) was applied to residual fluke numbers as described
254 above. All analyses were conducted in SPSS 24 statistical package. The experiments were
255 approved by Finnish Regional State Administrative Agency (license number
256 ESAVI/1375/04.10.03/2012) and they conformed to the animal care legislation of Finland.

257

258 *(e) Modelling the population-level effects of fluke infection on the impact of a bacterial*
259 *epidemic*

260

261 To explore the population-level consequences of the individual-level effects seen in
262 Experiments 1 and 2, we developed a mathematical model similar to previously-published
263 models of priority effects in multiple infections (Clay, Cortez, Duffy, & Rudolf, 2019; Clay,
264 Dhir, et al., 2019), parameterized with the data from our experiments, to predict the effects of
265 prior or subsequent fluke infections on the impact of a bacterial epidemic within a single
266 season (70 days) under aquaculture conditions. The model tracked changes in the proportion
267 of hosts in the population that were either (1) infected with just the bacteria ('*B*'), (2) infected
268 with just the fluke ('*F*'), (3) recovered from the bacteria infection (and assumed to be
269 immune to bacterial reinfection during the same season; '*R*'), (4) sequentially infected with
270 the fluke first, then the bacteria ('*C_{FC}*'), (5) sequentially infected with the bacteria first, then
271 the fluke ('*C_{BC}*'), (6) previously infected with both parasites, but had recovered from their
272 bacterial infection (but retained their fluke infection, and were immune to subsequent
273 bacterial infection; '*F_R*'), or (7) uninfected by either parasite ('*U*').

274

275 Transitions between the various classes depended on the transmission and recovery rates of
 276 the bacteria and fluke. As stated above, hosts that recover from bacterial infection were
 277 assumed to be resistant against subsequent bacterial reinfection; bacteria-only infected hosts
 278 (B) were assumed to recover to resistant hosts (R) at rate σ_B , whereas sequentially infected
 279 hosts C_{FC} and C_{BC} recover to fluke-infected resistant hosts (F_R) at rates σ_{FC} and σ_{BC} ,
 280 respectively. Hosts were assumed not to recover from fluke infections. Hosts susceptible to
 281 bacterial infections (i.e., all non-resistant hosts) were assumed to acquire bacterial infections
 282 at a rate dependent on the total abundance of all bacterial-infected hosts ($B_T = B + C_{FC} + C_{BC}$),
 283 and *per capita* rate β_B , resulting in the following transitions: $U \rightarrow B$ and $F \rightarrow C_{FC}$. Note, we
 284 assume these *per capita* transmission rates are the same regardless of fluke infection status,
 285 so prior or on-going fluke infection is assumed not to influence bacterial infectivity or
 286 shedding rates. Because the fluke life-cycle involves multiple life-stages in different host
 287 species, spanning long durations, we modelled fluke transmission as a constant force of
 288 infection parameter (Γ ; i.e., ignoring dynamic feedback between current infections and
 289 subsequent transmission rates), reflecting the population of cercariae present in the water
 290 throughout the season, resulting in the following transitions: $U \rightarrow F$, $B \rightarrow C_{BC}$ and $R \rightarrow F_R$.
 291 Hosts were assumed to die at infection-specific mortality rates μ_i , where i represents the
 292 infection class. Due to relative short duration of our simulations, we assumed no background
 293 mortality of uninfected fish or fluke-only infected fish. Similarly, we assumed no increases in
 294 host population size (e.g., through host reproduction, immigration, or input from external
 295 sources) corresponding to aquaculture conditions. Overall then this leads to the following set
 296 of equations describing the changes in abundance of each host class:

$$\frac{dU}{dt} = -U(\beta_B B_T + \Gamma)$$

$$\frac{dB}{dt} = \beta_B B_T U - B(\mu_B + \Gamma + \sigma_B)$$

$$\frac{dF}{dt} = U\Gamma - F\beta_F B_T$$

$$\frac{dR}{dt} = B\sigma_B - R\Gamma$$

$$\frac{dC_{FC}}{dt} = F\beta_F B_T - C_{FC}(\mu_{FC} + \sigma_{FC})$$

$$\frac{dC_{BC}}{dt} = B\Gamma - C_{BC}(\mu_{BC} + \sigma_{BC})$$

$$\frac{dF_R}{dt} = C_{FC}\sigma_{FC} + C_{BC}\sigma_{BC} + R\Gamma$$

297

298 We parameterized the model separately for each bacteria strain – fluke genotype combination
 299 using data from either Experiment 1 or Experiment 2, resulting in 18 parameter sets (3
 300 bacterial strains \times 3 fluke genotypes \times 2 experiments). For each combination the mortality
 301 rate of the appropriate infection classes (e.g., bacterial-only infected hosts, sequential fluke –
 302 bacterial infection hosts) were given by the inverse of the observed mean host survival times
 303 for those experimental categories; sequential bacteria – fluke infection hosts were assumed to
 304 die at the rate given by the simultaneous infection experiments. The bacterial recovery rates
 305 for infection class i were calculated based on the observed proportion of fish surviving each
 306 experimental exposure ($p_{surv,i}$); assuming constant recovery and mortality rates of host class i
 307 are σ_i and μ_i , respectively, the expected proportion surviving is given by $p_{surv,i} = \frac{\sigma_i}{\sigma_i + \mu_i}$.
 308 Since the $p_{surv,i}$ are known for each infection class i , and the μ_i can be estimated as described
 309 above, this equation can be rearranged to calculate the recovery rate $\sigma_i = \frac{p_{surv,i} \cdot \mu_i}{(1 - p_{surv,i})}$ that
 310 results in the observed proportion surviving from that class. The infection parameters in the
 311 system are unknown, so we chose an arbitrary value of bacterial transmission rate, β_B
 312 (although we varied this value by two orders of magnitude around this baseline value and
 313 found no qualitative effect on our results), and varied the cercarial force of infection (Γ) to

314 explore a range of scenarios of increasing fluke transmission pressure. All simulations were
315 assumed to start with 100 fish, of which 10 were infected with the bacteria, to seed the
316 epidemic. For each combination of bacterial strain \times fluke genotype \times experimental
317 parameters, we ran the model for a duration of 70 days, and assessed the effect of varying
318 cercarial infection pressure (Γ) on the end-of-season (day 70) total host abundance, compared
319 to the scenarios when either (1) there was no bacterial epidemic, or (2) there was a bacterial
320 epidemic, but no fluke infection. All models were run in R 3.5.1.

321

322 **3. Results**

323

324 *Experiment 1*

325

326 Virulence of the flavobacterial strains differed significantly in single infections so that the
327 strains 2 and 3 were more virulent compared to strain 1 [Cox regression: Wald = 70.48, $p <$
328 0.001 (strain)]. The virulence of the strains also slightly changed during the 4 h interval
329 between the infections [Wald = 25.77, $p <$ 0.001 (strain \times sequence)] (Fig. 1). *Diplostomum*
330 infection alone did not cause mortality of fish. No mortality was observed either among the
331 30 unexposed control fish.

332

333 Sequential infection with the 4h interval between the administrations of the two parasites
334 significantly reduced the virulence of the secondary bacterial infection (Fig. 1, Table 1).
335 However, this was bacterial strain-specific and most evident in strain 3 (interaction between
336 sequential infection, flavobacterial strain and fluke genotype; Fig. 1, Table 1). There was
337 also a significant increase in the proportion of fish surviving the experiment with sequential
338 infection in strain 1 when co-exposed with fluke genotypes B [increase from 35% (95% CI =

339 15.4-59.2%) to 85% surviving (62.1-96.8%)] and C [35% (15.4-59.2%) to 90% (68.3-
340 98.8%); Fig. 1, Table S3].

341

342 The residual number of flukes was significantly different between the fluke genotypes, with
343 genotype explaining large part of the variation (Fig. S1, Table S5). Sequential infection did
344 not affect the residual parasite numbers overall, but had an effect depending on the bacterial
345 strain, fluke genotype and their interaction (Fig. S1, Table S5). This suggests that the
346 administration of the bacterium four hours later also affected the fluke numbers in strain-
347 genotype-specific manner.

348

349 *Experiment 2*

350

351 Similarly to the Experiment 1, virulence of the flavobacterial strains 2 and 3 was higher
352 compared to strain 1 [Cox regression: Wald = 39.61, $p < 0.001$ (strain)] and this pattern was
353 independent of the previous exposure to flukes [Score = 1.33, $p = 0.515$ (prior
354 infection \times strain)] (Fig. 2). Prior infection with flukes five weeks earlier resulted in an
355 average of 1.68 ± 0.09 parasites established in the eyes of fish, which did not influence their
356 growth compared to the uninfected fish (mean body length \pm SE: 40.2 ± 0.2 mm (uninfected),
357 40.5 ± 0.2 mm (infected); t-test: $t_{591} = 1.069$, $p = 0.286$). The infection caused a small, but
358 significant reduction in the virulence of bacterial infection (Fig. 2, Table 1). Again, the effect
359 depended on the bacterial strain and was evident with strains 2 and 3 when co-exposed with
360 the fluke (Fig. 2, Table 1). However, in single bacterial exposures, the effect of decreased
361 virulence with the prior exposure to flukes was consistent across all strains (Wald = 5.472, p
362 = 0.019; Fig. 2). Single fluke infection did not cause significant mortality (two fish out of 120
363 exposed only to flukes died during the experiment). No mortality was observed among the 30

364 unexposed control fish. The residual number of flukes was significantly different between the
365 fluke genotypes, indicating that genotypes differed in infection success. However, effects of
366 the prior fluke infection, bacterial strain, or their interactions on fluke numbers in the second
367 exposure were not significant (Table S5).

368

369 *Predicted population-level effects of fluke infection on the impact of a bacterial epidemic*

370

371 Our model showed that the presence of fluke infections can protect the host population from
372 the detrimental effect of a bacterial epidemic, most notably when using parameter values
373 from Experiment 1, which assumed short-term sequential infections (Fig. 3). However, the
374 magnitude of this protective effect varied considerably across bacterial strain and fluke
375 genotype combinations; bacterial strain 1 (Fig. 3, top row) appeared to be the most easily-
376 overcome strain, with increasing cercarial force of infection (Γ) leading to progressively
377 higher levels of end-of-season fish abundance. The magnitude of these effects varied with
378 fluke genotype, from around 40% at the highest levels of Γ examined for fluke genotype A,
379 up to ~90% protection for fluke genotype C. However, population-level protection from the
380 other bacterial strains was negligible, regardless of the fluke genotype. Where population-
381 level protection was observed, (e.g., for bacterial strain 1), this was driven primarily by the
382 increased recovery rate from bacterial infection of sequentially-infected fish, as switching this
383 component off (i.e., assuming recovery rates were the same regardless of the individual's
384 prior fluke infection history) resulted in the loss of population-level protection (Fig. 4).
385 Running the model using parameter values from Experiment 2, assuming longer-term
386 sequential infections, revealed low levels of population-level protection, and now most
387 commonly observed for fluke genotype D and for bacterial strain 3 (Fig. S2).

388

389 4. Discussion

390

391 Temporally variable infections of multiple parasites are common in the wild, potentially
392 altering the outcomes of virulence in natural host-parasite interactions. Sequential parasite
393 dynamics are common also in farming environments such as aquaculture, but the knowledge
394 of the effects of temporal parasite dynamics on disease virulence in farmed animals is very
395 limited. We investigated how temporal spacing between the infections of the pathogenic
396 bacterium *F. columnare* and the fluke *D. pseudospathaceum* influenced the virulence of
397 infection (morbidity) in aquaculture fish hosts. Both short (few hours) and long (several
398 weeks) temporal difference between the infections resulted in reduction in bacterial virulence,
399 while this effect depended on the genetic interactions among the parasite species. Similarly,
400 timing of the infections changed the success of the fluke genotypes, suggesting influence also
401 on parasite fitness. Overall, these results suggest that previous infections in different temporal
402 scales can shape success and virulence of parasites with very different mechanisms of
403 transmission and infection, and their subsequent impact on host population dynamics.

404

405 Previously, we have shown that the virulence of simultaneous infections of *F. columnare* and
406 *D. pseudospathaceum* in fish is determined by complex genotype-specific interactions (Louhi
407 et al., 2015). Our present results, suggesting both short and long-term influence of sequential
408 infections, add yet another dimension to these G×G interactions. Mechanistically, the lower
409 virulence in sequential compared to simultaneous infections could be related, for example, to
410 reduction in the rate of bacterial invasion to the host's body (Louhi et al., 2015), or to higher
411 efficiency of the host's immune system to cope with two infections (Karvonen et al., 2012;
412 Klemme et al., 2016). Similar host-related factors could also influence the differences in
413 fluke establishment, although detailed mechanisms underlying the changes in virulence and

414 infection success are currently unclear. Overall, our results add significantly to earlier studies
415 on sequential infections between different parasite taxa, majority of which have used single
416 genotypes/strains (Ben-Ami et al., 2011; Clay, Dhir, et al., 2019; Doublet et al., 2015;
417 Hoverman et al., 2013; Lohr et al., 2010; Marchetto & Power, 2018; Natsopoulou et al.,
418 2015), by emphasising the importance of variation in infection outcomes depending on the
419 specific G×G parasite combinations. Indeed, combining G×G interactions in multiple
420 parasites with host infection history makes estimation of virulence and virulence evolution
421 increasingly challenging (Karvonen, Jokela, & Laine, 2019). Nevertheless, such interactions
422 could have important applied implications for scenarios of parasite prevention in intensive
423 farming environments.

424

425 Our model on infection dynamics in an aquaculture fish population, parametrized from the
426 experimental data showed that a previous fluke infection, particularly few hours earlier, can
427 protect the host population from the bacterial epidemic. However, this effect depended on the
428 cercarial force of infection and, most importantly, on the bacterial-fluke genetic
429 combinations. Interestingly, the protective effect increased dramatically at low cercarial
430 forces of infection and was notably strong in some of the strain/genotype combinations, with
431 up to ~90% of the population protected from the disease. This suggest that when flukes are
432 present in the tank water, as a result of parasite input via incoming water or from infected
433 snails inside the farm (Karvonen et al., 2006; Stables & Chappell, 1986), even a low-level
434 fluke infection could potentially decrease morbidity and mortality associated with an
435 imminent bacterial exposure. This is consistent with earlier results of less-virulent parasites
436 providing host with at least some degree of protection against later arriving virulent
437 strains/species in plants (Adame-Alvarez, Mendiola-Soto, & Heil, 2014; Seifi, Nonomura,
438 Matsuda, Toyoda, & Bai, 2012; Tollenaere, Susi, & Laine, 2016) and invertebrates (Ben-

439 Ami, Mouton, et al., 2008; Clay, Cortez, et al., 2019; Wuerthner, Hua, & Hoverman, 2017).
440 Similarly, the model on sequential infections five weeks apart predicted recovery in some of
441 the parasite combinations, although the magnitude of this protective effect was clearly lower.
442 While we could not compare the two experiments directly because they used different fluke
443 genotypes, the experimental results and the model predictions suggest that the protective
444 effect of sequential fluke infection against the bacterial outbreak might decay with time. If
445 true, this could reflect operation of different short and long-term host-level mechanisms such
446 as reduction in parasite facilitation (Louhi et al., 2015) or activation of host immune system
447 (Klemme et al., 2016) (see above). These are promising leads for future studies on the
448 detailed mechanisms underlying the effects of sequential infections on disease epidemiology
449 in this system.

450

451 Our model assumes single strain-genotype combinations between the parasites whereas in
452 reality aquaculture fish would likely to be exposed to mixed genotypes of both flukes (Rauch,
453 Kalbe, & Reusch, 2005) and bacteria (Kunttu et al., 2012; Sundberg et al., 2016),
454 significantly increasing the number of possible genotype combinations and outcomes of the
455 disease. Thus, the model predictions on effects of sequential infection in specific strain-
456 genotype combinations can be considered as extremes, ranging from no effect to nearly total
457 recovery of the host population. Given the likelihood of multiple strain-genotype infections,
458 the reality is likely to lie somewhere in between these extremes, depending on the genetic
459 composition of the bacterial and fluke populations in the water. Further, the model
460 predictions were driven mainly by host recovery while our experimental data showed that
461 sequential infection between the parasites also prolonged the lifetime of the hosts. While this
462 effect was small in these experimental conditions, where the disease progression from
463 exposure to morbidity is very fast (most fish had died within 28h), it could be speculated that

464 the effect of increase in lifetime in aquaculture conditions, where the length of an untreated
465 epidemic is typically several days or even weeks (Räihä, Sundberg, Ashrafi, Hyvärinen, &
466 Karvonen, 2019), could be stronger. Importantly, these increases in infected host lifetime,
467 though beneficial to the individual host, could have counter-productive effects on the host
468 population as a whole. Similar to host ‘tolerance’ responses to infection (Ayres & Schneider,
469 2012; Medzhitov, Schneider, & Soares, 2012; Råberg, Graham, & Read, 2009), by keeping
470 infected hosts alive, this prolongs the window of opportunity for infection to other hosts in
471 the population, increasing overall infection prevalence, and resulting in a net decrease in host
472 population abundance (e.g. Vale, Fenton, and Brown (2014)), and, as most clearly seen here,
473 in the absence of any effect of fluke infection on host recovery from bacterial infection (Fig.
474 4). However, the detailed epidemiological consequences of the prolonged lifetime are
475 currently unknown and require further work. Finally, competition/interactions between the
476 bacterial strains (Kinnula, Mappes, & Sundberg, 2017; Sundberg et al., 2016), in interaction
477 with the flukes infecting the hosts simultaneously (Louhi et al., 2015) or sequentially (this
478 study), could also shape the evolution of virulence. For example, as the order of parasite
479 arrival to host can significantly alter the outcome of virulence (Alizon, de Roode, &
480 Michalakis, 2013), factors such as G×G interactions between parasites could be important in
481 determining which virulence genotypes are favoured by selection under each infection
482 scenario (Karvonen et al., 2019). However, while our results support G×G variation in
483 infection outcomes, a formal study of implications of sequential infections on evolution of
484 virulence is beyond the scope of the present model/study.

485

486 To conclude, flavobacteria are currently considered among the most important bacterial fish
487 pathogens worldwide. Interactions between increasingly virulent strains of *F. columnare*
488 (Kinnula et al., 2017; Pulkkinen et al., 2010; Sundberg et al., 2016) and those between the

489 bacterium and other co-occurring parasite species (Bandilla, Valtonen, Suomalainen, Aphalo,
490 & Hakalahti, 2006; Louhi et al., 2015; Xu, Shoemaker, & LaFrentz, 2014) can significantly
491 alter the disease outcomes and influence predictions on virulence of infection. Our results,
492 suggesting protective effect of sequential infections from a less-virulent parasite against a
493 highly virulent one (see also King et al. (2016)), provide an interesting viewpoint into how
494 epidemics could be altered by the host infection history. While the epidemiological effects of
495 consecutive parasite outbreaks on subsequent disease occurrence are generally poorly known,
496 results from our model suggest that changes in disease epidemiology with sequential
497 exposure can also have important implications for the need and success of disease
498 management practices and medication protocols. In a wider perspective, possible reduction in
499 use of medication could also constrain environmental discharge of medical residues and
500 pathogen evolution for antibiotic resistance (Cabello, 2006; Martinez, 2009).

501

502 **Data archiving**

503 All data used in this paper are deposited in Dryad Digital Repository upon acceptance

504

505 **Competing interests**

506 We have no competing interests

507

508

509

510

511

512

513

514

515

516

517

518

519

520 **References**

521

522 Adame-Alvarez, R. M., Mendiola-Soto, J., & Heil, M. (2014). Order of arrival shifts
523 endophyte-pathogen interactions in bean from resistance induction to disease
524 facilitation. *FEMS Microbiology Letters*, 355(2), 100-107. doi:10.1111/1574-
525 6968.12454

526 Alizon, S., de Roode, J. C., & Michalakis, Y. (2013). Multiple infections and the evolution of
527 virulence. *Ecology Letters*, 16(4), 556-567. doi:10.1111/ele.12076

528 Ayres, J. S., & Schneider, D. S. (2012). Tolerance of infections. In W. E. Paul (Ed.), *Annual*
529 *Review of Immunology* (Vol. 30, pp. 271-294).

530 Bandilla, M., Valtonen, E. T., Suomalainen, L. R., Aphalo, P. J., & Hakalahti, T. (2006). A
531 link between ectoparasite infection and susceptibility to bacterial disease in rainbow
532 trout. *International Journal for Parasitology*, 36(9), 987-991.
533 doi:10.1016/j.ijpara.2006.05.001

534 Bell, A. S., Roode, J. C., Sim, D., & Read, A. F. (2006). Within-host competition in
535 genetically diverse malaria infections: parasite virulence and competitive success.
536 *Evolution*, 60(7), 1358-1371.

- 537 Ben-Ami, F., Mouton, L., & Ebert, D. (2008). The effects of multiple infections on the
538 expression and evolution of virulence in a *Daphnia*-endoparasite system. *Evolution*,
539 62(7), 1700-1711. doi:10.1111/j.1558-5646.2008.00391.x
- 540 Ben-Ami, F., Regoes, R. R., & Ebert, D. (2008). A quantitative test of the relationship
541 between parasite dose and infection probability across different host-parasite
542 combinations. *Proceedings of the Royal Society B-Biological Sciences*, 275(1636),
543 853-859.
- 544 Ben-Ami, F., Rigaud, T., & Ebert, D. (2011). The expression of virulence during double
545 infections by different parasites with conflicting host exploitation and transmission
546 strategies. *Journal of Evolutionary Biology*, 24(6), 1307-1316. doi:10.1111/j.1420-
547 9101.2011.02264.x
- 548 Benesh, D. P., & Kalbe, M. (2016). Experimental parasite community ecology: intraspecific
549 variation in a large tapeworm affects community assembly. *Journal of Animal*
550 *Ecology*, 85(4), 1004-1013. doi:10.1111/1365-2656.12527
- 551 Byers, J. E., Blakeslee, A. M. H., Linder, E., Cooper, A. B., & Maguire, T. J. (2008).
552 Controls of spatial variation in the prevalence of trematode parasites infecting a
553 marine snail. *Ecology*, 89(2), 439-451. doi:10.1890/06-1036.1
- 554 Cabello, F. C. (2006). Heavy use of prophylactic antibiotics in aquaculture: a growing
555 problem for human and animal health and for the environment. *Environmental*
556 *Microbiology*, 8(7), 1137-1144. doi:10.1111/j.1462-2920.2006.01054.x
- 557 Clay, P. A., Cortez, M. H., Duffy, M. A., & Rudolf, V. H. W. (2019). Priority effects within
558 coinfecting hosts can drive unexpected population-scale patterns of parasite
559 prevalence. *Oikos*, 128(4), 571-583. doi:10.1111/oik.05937

- 560 Clay, P. A., Dhir, K., Rudolf, V. H. W., & Duffy, M. A. (2019). Within-host priority effects
561 systematically alter pathogen coexistence. *American Naturalist*, *193*(2), 187-199.
562 doi:10.1086/701126
- 563 Criscione, C. D., & Blouin, M. S. (2004). Life cycles shape parasite evolution: comparative
564 population genetics of salmon trematodes. *Evolution*, *58*(1), 198-202. doi:10.1554/03-
565 359
- 566 Davies, C. M., Fairbrother, E., & Webster, J. P. (2002). Mixed strain schistosome infections
567 of snails and the evolution of parasite virulence. *Parasitology*, *124*(01), 31-38.
- 568 de Roode, J. C., Helinski, M. E. H., Anwar, M. A., & Read, A. F. (2005). Dynamics of
569 multiple infection and within-host competition in genetically diverse malaria
570 infections. *The American Naturalist*, *166*(5), 531-542.
- 571 Declercq, A. M., Boyen, F., Van den Broeck, W., Bossier, P., Karsi, A., Haesebrouck, F., &
572 Decostere, A. (2013). Antimicrobial susceptibility pattern of *Flavobacterium*
573 *columnare* isolates collected worldwide from 17 fish species. *Journal of Fish*
574 *Diseases*, *36*(1), 45-55. doi:10.1111/j.1365-2761.2012.01410.x
- 575 Declercq, A. M., Haesebrouck, F., Van den Broeck, W., Bossier, P., & Decostere, A. (2013).
576 Columnaris disease in fish: a review with emphasis on bacterium-host interactions.
577 *Veterinary Research*, *44*. doi:10.1186/1297-9716-44-27
- 578 Decostere, A., Haesebrouck, F., & Devriese, L. A. (1997). Shieh medium supplemented with
579 tobramycin for selective isolation of *Flavobacterium columnare* (*Flexibacter*
580 *columnaris*) from diseased fish. *Journal of Clinical Microbiology*, *35*(1), 322-324.
- 581 Doublet, V., Natsopoulou, M. E., Zschiesche, L., & Paxton, R. J. (2015). Within-host
582 competition among the honey bees pathogens *Nosema ceranae* and Deformed wing
583 virus is asymmetric and to the disadvantage of the virus. *Journal of Invertebrate*
584 *Pathology*, *124*, 31-34. doi:10.1016/j.jip.2014.10.007

- 585 Duncan, A. B., Agnew, P., Noel, V., & Michalakis, Y. (2015). The consequences of co-
586 infections for parasite transmission in the mosquito *Aedes aegypti*. *Journal of Animal*
587 *Ecology*, *84*(2), 498-508. doi:10.1111/1365-2656.12302
- 588 Faltýnková, A., Valtonen, E. T., & Karvonen, A. (2008). Spatial and temporal structure of the
589 trematode component community in *Valvata macrostoma* (Gastropoda,
590 Prosobranchia). *Parasitology*, *135*(14), 1691-1699. doi:10.1017/s0031182008005027
- 591 Fellous, S., & Koella, J. C. (2009). Infectious dose affects the outcome of the within-host
592 competition between parasites. *American Naturalist*, *173*(6), E177-E184.
593 doi:10.1086/598490
- 594 Gower, C. M., & Webster, J. P. (2005). Intraspecific competition and the evolution of
595 virulence in a parasitic trematode. *Evolution*, *59*(3), 544-553.
- 596 Graham, A. L. (2008). Ecological rules governing helminth-microparasite coinfection.
597 *Proceedings of the National Academy of Sciences of the United States of America*,
598 *105*(2), 566-570. doi:10.1073/pnas.0707221105
- 599 Hakalahti, T., & Valtonen, E. T. (2003). Population structure and recruitment of the
600 ectoparasite *Argulus coregoni* Thorell (Crustacea : Branchiura) on a fish farm.
601 *Parasitology*, *127*, 79-85. doi:10.1017/s0031182003003196
- 602 Hood, M. E. (2003). Dynamics of multiple infection and within-host competition by the
603 anther-smut pathogen. *American Naturalist*, *162*(1), 122-133. doi:10.1086/375539
- 604 Hoverman, J. T., Hoyer, B. J., & Johnson, P. T. J. (2013). Does timing matter? How priority
605 effects influence the outcome of parasite interactions within hosts. *Oecologia*, *173*(4),
606 1471-1480. doi:10.1007/s00442-013-2692-x
- 607 Jokela, J., & Lively, C. M. (1995). Spatial variation in infection by digenetic trematodes in a
608 population of fresh-water snails (*Potamopyrgus antipodarum*). *Oecologia*, *103*(4),
609 509-517. doi:10.1007/bf00328690

- 610 Karvonen, A. (2012). *Diplostomum spathaceum* and Related Species. In P. T. K. Woo & K.
611 Buchmann (Eds.), *Fish Parasites: Pathobiology and Protection* (pp. 260-269).
612 Oxfordshire, UK: CAB International.
- 613 Karvonen, A., Jokela, J., & Laine, A.-L. (2019). Importance of sequence and timing in
614 parasite coinfections. *Trends in Parasitology*, 35(2), 109-118.
615 doi:10.1016/j.pt.2018.11.007
- 616 Karvonen, A., Kirsi, S., Hudson, P. J., & Valtonen, E. T. (2004). Patterns of cercarial
617 production from *Diplostomum spathaceum*: terminal investment or bet hedging?
618 *Parasitology*, 129, 87-92. doi:10.1017/s0031182004005281
- 619 Karvonen, A., Paukku, S., Valtonen, E. T., & Hudson, P. J. (2003). Transmission, infectivity
620 and survival of *Diplostomum spathaceum* cercariae. *Parasitology*, 127, 217-224.
621 doi:10.1017/s0031182003003561
- 622 Karvonen, A., Rellstab, C., Louhi, K.-R., & Jokela, J. (2012). Synchronous attack is
623 advantageous: mixed genotype infections lead to higher infection success in
624 trematode parasites. *Proceedings of the Royal Society B-Biological Sciences*,
625 279(1726), 171-176. doi:10.1098/rspb.2011.0879
- 626 Karvonen, A., Rintamäki, P., Jokela, J., & Valtonen, E. T. (2010). Increasing water
627 temperature and disease risks in aquatic systems: climate change increases the risk of
628 some, but not all, diseases. *International Journal for Parasitology*, 40(13), 1483-
629 1488. doi:10.1016/j.ijpara.2010.04.015
- 630 Karvonen, A., Savolainen, M., Seppälä, O., & Valtonen, E. T. (2006). Dynamics of
631 *Diplostomum spathaceum* infection in snail hosts at a fish farm. *Parasitology*
632 *Research*, 99(4), 341-345. doi:10.1007/s00436-006-0137-8

- 633 Karvonen, A., Seppälä, O., & Valtonen, E. T. (2004). Parasite resistance and avoidance
634 behaviour in preventing eye fluke infections in fish. *Parasitology*, *129*, 159-164.
635 doi:10.1017/s0031182004005505
- 636 Karvonen, A., Seppälä, O., & Valtonen, E. T. (2009). Host immunization shapes interspecific
637 associations in trematode parasites. *Journal of Animal Ecology*, *78*(5), 945-952.
638 doi:10.1111/j.1365-2656.2009.01562.x
- 639 Kennedy, D. A., Kurath, G., Brito, I. L., Purcell, M. K., Read, A. F., Winton, J. R., & Wargo,
640 A. R. (2016). Potential drivers of virulence evolution in aquaculture. *Evolutionary*
641 *Applications*, *9*(2), 344-354. doi:10.1111/eva.12342
- 642 King, K. C., Brockhurst, M. A., Vasieva, O., Paterson, S., Betts, A., Ford, S. A., . . . Hurst, G.
643 D. D. (2016). Rapid evolution of microbe-mediated protection against pathogens in a
644 worm host. *Isme Journal*, *10*(8), 1915-1924. doi:10.1038/ismej.2015.259
- 645 King, K. C., Delph, L. F., Jokela, J., & Lively, C. M. (2009). The geographic mosaic of sex
646 and the red queen. *Current Biology*, *19*(17), 1438-1441.
647 doi:10.1016/j.cub.2009.06.062
- 648 Kinnula, H., Mappes, J., & Sundberg, L.-R. (2017). Coinfection outcome in an opportunistic
649 pathogen depends on the inter-strain interactions. *BMC Evolutionary Biology*, *17*.
650 doi:10.1186/s12862-017-0922-2
- 651 Klemme, I., Louhi, K.-R., & Karvonen, A. (2016). Host infection history modifies co-
652 infection success of multiple parasite genotypes. *Journal of Animal Ecology*, *85*(2),
653 591-597. doi:10.1111/1365-2656.12472
- 654 Kunttu, H. M. T., Sundberg, L.-R., Pulkkinen, K., & Valtonen, E. T. (2012). Environment
655 may be the source of *Flavobacterium columnare* outbreaks at fish farms.
656 *Environmental Microbiology Reports*, *4*(4), 398-402. doi:10.1111/j.1758-
657 2229.2012.00342.x

- 658 Kunttu, H. M. T., Valtonen, E. T., Jokinen, E. I., & Suomalainen, L.-R. (2009). Saprophytism
659 of a fish pathogen as a transmission strategy. *Epidemics*, *1*(2), 96-100.
660 doi:10.1016/j.epidem.2009.04.003
- 661 Laanto, E., Bamford, J. K. H., Laakso, J., & Sundberg, L.-R. (2012). Phage-driven loss of
662 virulence in a fish pathogenic bacterium. *PLoS ONE*, *7*(12), e53157.
- 663 Laanto, E., Hoikkala, V., Ravantti, J., & Sundberg, L.-R. (2017). Long-term genomic
664 coevolution of host-parasite interaction in the natural environment. *Nature*
665 *Communications*, *8*. doi:10.1038/s41467-017-00158-7
- 666 Laine, A.-L. (2011). Context-dependent effects of induced resistance under co-infection in a
667 plant-pathogen interaction. *Evolutionary Applications*, *4*(5), 696-707.
668 doi:10.1111/j.1752-4571.2011.00194.x
- 669 Lohr, J. N., Yin, M., & Wolinska, J. (2010). Prior residency does not always pay off - co-
670 infections in *Daphnia*. *Parasitology*, *137*(10), 1493-1500.
671 doi:10.1017/s0031182010000296
- 672 Louhi, K.-R., Karvonen, A., Rellstab, C., & Jokela, J. (2010). Is the population genetic
673 structure of complex life cycle parasites determined by the geographic range of the
674 most motile host? *Infection, Genetics and Evolution*, *10*(8), 1271-1277.
675 doi:10.1016/j.meegid.2010.08.013
- 676 Louhi, K.-R., Sundberg, L.-R., Jokela, J., & Karvonen, A. (2015). Interactions among
677 bacterial strains and fluke genotypes shape virulence of co-infection. *Proceedings of*
678 *the Royal Society B: Biological Sciences*, *282*(1821), 20152097.
679 doi:10.1098/rspb.2015.2097
- 680 Marchetto, K. M., & Power, A. G. (2018). Coinfection timing drives host population
681 dynamics through changes in virulence. *American Naturalist*, *191*(2), 173-183.
682 doi:10.1086/695316

- 683 Martinez, J. L. (2009). Environmental pollution by antibiotics and by antibiotic resistance
684 determinants. *Environmental Pollution*, 157(11), 2893-2902.
685 doi:10.1016/j.envpol.2009.05.051
- 686 Medzhitov, R., Schneider, D. S., & Soares, M. P. (2012). Disease tolerance as a defense
687 strategy. *Science*, 335(6071), 936-941. doi:10.1126/science.1214935
- 688 Mohanty, B. R., & Sahoo, P. K. (2007). Edwardsiellosis in fish: a brief review. *Journal of*
689 *biosciences*, 32(3), 1331-1344. doi:10.1007/s12038-007-0143-8
- 690 Natsopoulou, M. E., McMahon, D. P., Doublet, V., Bryden, J., & Paxton, R. J. (2015).
691 Interspecific competition in honeybee intracellular gut parasites is asymmetric and
692 favours the spread of an emerging infectious disease. *Proceedings of the Royal*
693 *Society B-Biological Sciences*, 282(1798). doi:10.1098/rspb.2014.1896
- 694 Pulkkinen, K., Suomalainen, L. R., Read, A. F., Ebert, D., Rintamaki, P., & Valtonen, E. T.
695 (2010). Intensive fish farming and the evolution of pathogen virulence: the case of
696 columnaris disease in Finland. *Proceedings of the Royal Society B-Biological*
697 *Sciences*, 277(1681), 593-600. doi:10.1098/rspb.2009.1659
- 698 Rauch, G., Kalbe, M., & Reusch, T. B. H. (2005). How a complex life cycle can improve a
699 parasite's sex life. *Journal of Evolutionary Biology*, 18(4), 1069-1075.
700 doi:10.1111/j.1420-9101.2005.00895.x
- 701 Read, A. F., & Taylor, L. H. (2001). The ecology of genetically diverse infections. *Science*,
702 292(5519), 1099-1102. doi:10.1126/science.1059410
- 703 Rellstab, C., Karvonen, A., Louhi, K.-R., & Jokela, J. (2013). Genotype-specific vs. cross-
704 reactive host immunity against a macroparasite. *PLoS One*, 8(10).
705 doi:10.1371/journal.pone.0078427

- 706 Reusch, T. B. H., Rauch, G., & Kalbe, M. (2004). Polymorphic microsatellite loci for the
707 trematode *Diplostomum pseudospathaceum*. *Molecular Ecology Notes*, 4(4), 577-579.
708 doi:10.1111/j.1471-8286.2004.00740.x
- 709 Rintamäki-Kinnunen, P., & Valtonen, E. T. (1997). Epizootiology of protozoans in farmed
710 salmonids at northern latitudes. *International Journal for Parasitology*, 27(1), 89-99.
711 doi:10.1016/s0020-7519(96)00162-2
- 712 Råberg, L., Graham, A. L., & Read, A. F. (2009). Decomposing health: tolerance and
713 resistance to parasites in animals. *Philosophical Transactions of the Royal Society B-
714 Biological Sciences*, 364(1513), 37-49. doi:10.1098/rstb.2008.0184
- 715 Rähkä, V., Sundberg, L.-R., Ashrafi, R., Hyvärinen, P., & Karvonen, A. (2019). Rearing
716 background and exposure environment together explain higher survival of aquaculture
717 fish during a bacterial outbreak. *Journal of Applied Ecology*, 00, 1-10.
718 doi:10.1111/1365-2664.13393
- 719 Salgame, P., Yap, G. S., & Gause, W. C. (2013). Effect of helminth-induced immunity on
720 infections with microbial pathogens. *Nature Immunology*, 14(11), 1118-1126.
721 doi:10.1038/ni.2736
- 722 Seifi, A., Nonomura, T., Matsuda, Y., Toyoda, H., & Bai, Y. L. (2012). An avirulent tomato
723 powdery mildew isolate induces localized acquired resistance to a virulent isolate in a
724 spatiotemporal manner. *Molecular Plant-Microbe Interactions*, 25(3), 372-378.
725 doi:10.1094/mpmi-06-11-0171
- 726 Skall, H. F., Olesen, N. J., & Møllergaard, S. (2005). Viral haemorrhagic septicaemia virus in
727 marine fish and its implications for fish farming - a review. *Journal of Fish Diseases*,
728 28(9), 509-529. doi:10.1111/j.1365-2761.2005.00654.x
- 729 Song, Y. L., Fryer, J. L., & Rohovec, J. S. (1988). Comparison of six media for the
730 cultivation of *Flexibacter columnaris*. *Fish Pathology*, 23(2), 91-94.

- 731 Stables, J. N., & Chappell, L. H. (1986). The epidemiology of diplostomiasis in farmed
732 rainbow trout from north-east Scotland. *Parasitology*, *92*, 699-710.
- 733 Sundberg, L.-R., Ketola, T., Laanto, E., Kinnula, H., Bamford, J. K. H., Penttinen, R., &
734 Mappes, J. (2016). Intensive aquaculture selects for increased virulence and
735 interference competition in bacteria. *Proceedings of the Royal Society B-Biological*
736 *Sciences*, *283*(1826). doi:10.1098/rspb.2015.3069
- 737 Telfer, S., Lambin, X., Birtles, R., Beldomenico, P., Burthe, S., Paterson, S., & Begon, M.
738 (2010). Species interactions in a parasite community drive infection risk in a wildlife
739 population. *Science*, *330*(6001), 243-246. doi:10.1126/science.1190333
- 740 Tobback, E., Decostere, A., Hermans, K., Haesebrouck, F., & Chiers, K. (2007). *Yersinia*
741 *ruckeri* infections in salmonid fish. *Journal of Fish Diseases*, *30*(5), 257-268.
742 doi:10.1111/j.1365-2761.2007.00816.x
- 743 Tollenaere, C., Susi, H., & Laine, A.-L. (2016). Evolutionary and epidemiological
744 implications of multiple infection in plants. *Trends in Plant Science*, *21*(1), 80-90.
745 doi:10.1016/j.tplants.2015.10.014
- 746 Wagner, B. A., Wise, D. J., Khoo, L. H., & Terhune, J. S. (2002). The epidemiology of
747 bacterial diseases in food-size channel catfish. *Journal of Aquatic Animal Health*,
748 *14*(4), 263-272. doi:10.1577/1548-8667(2002)014<0263:teobdi>2.0.co;2
- 749 Vale, P. F., Fenton, A., & Brown, S. P. (2014). Limiting damage during infection: Lessons
750 from infection tolerance for novel therapeutics. *PLoS Biology*, *12*(1).
751 doi:10.1371/journal.pbio.1001769
- 752 Vojvodic, S., Boomsma, J. J., Eilenberg, J., & Jensen, A. B. (2012). Virulence of mixed
753 fungal infections in honey bee brood. *Frontiers in Zoology*, *9*. doi:10.1186/1742-
754 9994-9-5

755 Wuerthner, V. P., Hua, J., & Hoverman, J. T. (2017). The benefits of coinfection: trematodes
 756 alter disease outcomes associated with virus infection. *Journal of Animal Ecology*,
 757 86(4), 921-931. doi:10.1111/1365-2656.12665

758 Xu, D. H., Shoemaker, C. A., & LaFrentz, B. R. (2014). Enhanced susceptibility of hybrid
 759 tilapia to *Flavobacterium columnare* after parasitism by *Ichthyophthirius multifiliis*.
 760 *Aquaculture*, 430, 44-49. doi:10.1016/j.aquaculture.2014.03.041

761

762

763

764

765

766

767

768

769

770

771 Table 1. Results of stepwise Cox regression analyses on mortality of rainbow trout co-
 772 exposed to three strains of the bacterium *Flavobacterium columnare* and three genotypes of
 773 the fluke *Diplostomum pseudospathaceum* in all possible combinations in Experiments 1 and
 774 2. Infection type (simultaneous vs. sequential (Exp 1) or no prior infection vs. with prior
 775 infection (Exp 2)), bacterial strain (G_B ; 1-3) and fluke genotype (G_F ; A-C (Exp 1) or D-F
 776 (Exp 2)) were used as categorical covariates, and fish length as a continuous covariate.

Experiment	Source	Wald	df	p	Exp (B)	95% CI
1	Sequential infection (S)	69.74	1	<0.001	0.41	0.33-0.51
	G_B	284.58	2	<0.001		

	G _F	51.32	3	<0.001		
	S×G _F	27.31	3	<0.001		
	G _B ×G _F	35.26	6	<0.001		
	S×G _B ×G _F	26.80	6	<0.001		
2	Prior infection (I)	23.34	1	<0.001	0.63	0.52-0.76
	G _B	213.65	2	<0.001		
	G _F	8.94	3	0.030		
	I×G _B	7.56	2	0.023		
	I×G _F	14.25	3	0.003		
	Fish length	10.55	1	0.001		

777

778

779

780

781

782 **Figure captions**

783

784 Fig. 1. Mean survival times (\pm SE) of rainbow trout co-exposed simultaneously (Sim, open
785 boxes) or sequentially (Seq, grey boxes) to three strains of the bacterium *Flavobacterium*
786 *columnare* (1-3) and three genotypes of the fluke *Diplostomum pseudospathaceum* (A-C) in
787 all possible combinations in the first experiment. No *Diplostomum* indicates survival of fish
788 exposed only to *F. columnare*. Boxes show data for fish that died during the experiment.
789 Black dots indicate the percentage of fish surviving in each combination.

790

791 Fig. 2. Mean survival times (\pm SE) of rainbow trout previously unexposed (Unexp., open
792 boxes) or exposed to *Diplostomum pseudospathaceum* (Exposed, grey boxes) when re-
793 exposed to three strains of the bacterium *Flavobacterium columnare* (1-3) and three
794 genotypes of the fluke *D. pseudospathaceum* (D-F) in all possible combinations in the second
795 experiment. No *Diplostomum* indicates survival of fish exposed only to *F. columnare*. Boxes
796 show data for fish that died during the experiment. Black dots indicate the percentage of fish
797 surviving in each combination.

798

799 Fig. 3. Model predictions of the end-of-season (day 70) total host abundance as a function of
800 increasing cercarial force of infection (Γ), for each bacterial strain – fluke genotype
801 combination, parameterised using data from Experiment 1. Solid black line = host abundance
802 in the presence of both the bacteria and fluke ('B&F'), dashed blue line = host abundance in
803 the presence of just the bacteria ('B only'), dashed green line = host abundance in the absence
804 of both bacteria and fluke ('Neither'). Other parameters: initial number of hosts = 100, initial
805 number of bacterial-infected hosts = 10, $\beta_B = 0.001$.

806

807 Fig. 4. As in Fig. 3, but ignoring any effect of fluke infection on recovery from bacterial
808 infection (prior or subsequent fluke fish are assumed to recover from bacterial infection at the
809 same rate as bacterial-only infected fish; any effects of fluke infection on host survival time
810 are retained, as in Fig. 3).

811




Fig. 1




Fig. 2

