

Acyclic Gambling Games

Rida Laraki* and Jérôme Renault†

July 20, 2019

Abstract

We consider 2-player zero-sum stochastic games where each player controls his own state variable living in a compact metric space. The terminology comes from gambling problems where the state of a player represents its wealth in a casino. Under standard assumptions (e.g. continuous running payoff and non-expansive transitions), we consider for each discount factor the value v_λ of the λ -discounted stochastic game and investigate its limit when λ goes to 0. We show that under a new acyclicity condition, the limit exists and is characterized as the unique solution of a system of functional equations: the limit is the unique continuous excessive and depressive function such that each player, if his opponent does not move, can reach the zone when the current payoff is at least as good as the limit value, without degrading the limit value. The approach generalizes and provides a new viewpoint on the Mertens-Zamir system coming from the study of zero-sum repeated games with lack of information on both sides. A counterexample shows that under a slightly weaker notion of acyclicity, convergence of (v_λ) may fail.

Keywords: Zero-Sum Stochastic Games, Markov Decision Process, Asymptotic and Uniform Value, Gambling Theory, Mertens-Zamir System, Splitting Games

Classification: Primary: 91A15, 91A60 ; secondary: 60G40; 90C40

1 Introduction

The model of zero-sum stochastic games was introduced by Shapley [Shapley, 1953]. A state variable $\omega \in \Omega$ follows a controlled Markov chain with transitions $Q(\tilde{\omega}|i, j, \omega)$ controlled by the actions of two competing players ($i \in I$ for player 1 and $j \in J$ for player 2). Shapley assumed the action and state spaces (I , J and Ω) to be finite and proved the existence of the value v_λ of the λ -discounted game using a dynamic programming principle, and characterized v_λ as the unique fixed point of what has

*CNRS, Université Paris Dauphine-PSL, France and University of Liverpool, UK. Email: rida.laraki@dauphine.fri

†Toulouse School of Economics, University of Toulouse Capitole, Toulouse, France. Email: jerome.renault@tse-fr.eu. **Acknowledgment:** Support from the ANR-3IA Artificial and Natural Intelligence Toulouse Institute is gratefully

been called *the Shapley operator* [Rosenberg and Sorin, 2001] and [Sorin, 2002]. For a recent survey, see [Laraki and Sorin, 2014].

Bewley and Kohlberg [Bewley and Kohlberg, 1976] using algebraic tools, proved the existence of the asymptotic value $v = \lim_{\lambda \rightarrow 0} v_\lambda$. Actually, when action and state spaces are finite, the equations that define v_λ may be described by finitely many polynomial inequalities, implying that v_λ is semi-algebraic and so is converging. The extension of this result to infinite stochastic games is a central question in mathematical game theory (see [Mertens et al., 2015], [Sorin and Neyman, 2003] and [Sorin, 2002]).

Recently, several important conjectures (see [Mertens, 1986], [Mertens et al., 2015]) were proved to be false. Vigerál [Vigerál, 2013] and Ziliotto [Ziliotto, 2016a] provided examples where the family $\{v_\lambda\}$ diverges as λ approaches zero. In Vigerál, the state space Ω is finite and the action sets I and J are semi-algebraic. In Ziliotto, the set of actions is finite but the state space Ω is compact, and can be seen as the space of common beliefs on a finite state variable, controlled but not observed by the players.

On the other hand, there are many classes of stochastic games with general state space and action sets where $\{v_\lambda\}$ converges. Many have in common some *irreversibility* in the transitions. In recursive games [Everett, 1957, Sorin and Vigerál, 2013] the current payoff is zero until the game is absorbed. In absorbing games ([Kohlberg, 1974] [Mertens et al., 2009] and [Rosenberg and Sorin, 2001]) there is only one non-absorbing state. In repeated games with incomplete information (see [Aumann et al., 1995], [Mertens and Zamir, 1971] and [Rosenberg and Sorin, 2001]) once a player reveals some information, he cannot withdraw it. Similarly in splitting games ([Laraki, 2001b], [Laraki, 2001a] and [Sorin, 2002]) the state follows a martingale which eventually converges. This leads many to anticipate that irreversibility has to do with convergence. Interestingly, in all those classes of “irreversible” stochastic games, not only we have convergence but also a characterization of the asymptotic value.

Our paper provides a weak and a strong definition of irreversibility (we call acyclicity) and prove that they constitute the frontier between convergence and divergence of $\{v_\lambda\}$: strong acyclicity guarantees convergence while the closely related weak acyclicity does not. To do so, we restrict ourself to a new class which embeds any product stochastic game [Flesch et al., 2008] and naturally extends gambling houses from one to two players.

A classical gambling house problem has three ingredients : a metric *state space* S , a Borel-measurable *utility function* $u : S \rightarrow \mathbb{R}$, and a *gambling house* Φ , where Φ is a set value function that assigns to each $s \in S$ a set $\Phi(s)$ in $\Delta(S)$ (set of Borel probability distributions over S). At each stage t , given the state s_t , the decision maker gets the reward $u(s_t)$, chooses $p_t \in \Phi(s_t)$, and the state moves to s_{t+1} according to the probability distribution p_t . The gambling house is called *leavable* if for every $s \in S$, δ_s (the dirac mass at s) belongs to $\Phi(s)$. This model was introduced in [Dubins and Savage, 1965], and was studied extensively by several authors, for instance [Maitra and Sudderth, 1996]. It is well known that any MDP (Markov Decision Process) can be mapped to a gambling house (by encoding actions in the state space) and any positive MDP can be mapped to a leavable gambling house, see for instance [Dubins et al., 2002], [Maitra and Sudderth, 1996] and [Schal, 1989].

In a *gambling game*, each player controls his gambling house: $\Gamma : X \rightarrow \Delta(X)$ for Player 1 and $\Lambda : Y \rightarrow \Delta(Y)$ for Player 2, and the utility function is now $u : X \times Y \rightarrow \mathbb{R}$ with the convention that player 1 wants to maximize u whereas player 2 wants to minimize u . At each stage t , both players knowing the state $\omega_t = (x_t, y_t)$, simultaneously Player 1 chooses p_t in $\Gamma(x_t)$ and Player 2 chooses $q_t \in \Lambda(y_t)$, the stage payoff is $u(x_t, y_t)$ and a new state (x_{t+1}, y_{t+1}) is selected according to the probability distribution $p_t \otimes q_t$. As for MDP, any product stochastic game [Flesch et al., 2008] can be mapped to a gambling game.

For each $\lambda \in (0, 1]$, one can define the λ -discounted game where the stream of payoffs is evaluated according to $\sum_{t=1}^{\infty} \lambda(1-\lambda)^{t-1} u(x_t, y_t)$. In economics, λ is usually called the discount rate, whereas $1-\lambda = \frac{1}{1+r} = \delta$ is called the discount factor and r is the interest rate. Hence, λ small means the player is patient and long-term optimizing.

Under some classical regularity assumptions, the λ -discounted game will have a value v_λ and the family $\{v_\lambda\}$ will be equi-continuous. The central questions this paper is concerned with are: when does $\lim_{\lambda \rightarrow 0} v_\lambda$ exist and if so, how to characterize it from the data of the game. A characterization is important if one wants to compute the limit value. This is a difficult question that this paper will not address. Let us just mention that computing the convex envelope of a given function, a very special case of our characterization, is already an NP-hard problem and there are, to our knowledge, only few algorithms that approximate it (see for instance [Laraki and J.B-Lasserre, 2008]).

Our first main result shows that if at least one of the gambling houses Γ or Λ is strongly acyclic (definition 2.14), $\{v_\lambda\}$ uniformly converges to a function v as λ goes to 0. Moreover, we provide several characterizations of the asymptotic value v that extend the well known Mertens-Zamir system of functional equations [Mertens and Zamir, 1971]. The convergence of the discounted values to the solution of the Mertens-Zamir system of equations in the special case of the splitting games was first proved in [Laraki, 2001a] and [Laraki, 2001b].

Our second result proves that under a slightly weaker notion of acyclicity, $\{v_\lambda\}$ may diverge (even if both houses Γ and Λ are weakly acyclic and both state spaces X and Y are finite). The example has similarities with an example in Ziliotto [Ziliotto, 2016a] for stochastic games where both players control the same state variable. Our example is the first in the class of product stochastic games, and is somehow simpler than the recent counterexamples.

Finally –in the appendix– under an idempotent assumption combined with a bounded variation hypothesis on the transitions, we prove existence of the uniform value¹ and provide simple uniform optimal strategies, extending a recent result in [Oliu-Barton, 2018] for the *splitting game*.

¹Recall that when a uniform optimal strategy σ exists, this means that for every $\varepsilon > 0$, there is $\lambda_0 > 0$ such that σ is ε -optimal in all λ -discounted games as soon as $0 < \lambda < \lambda_0$, i.e. a player needs not to know the exact value of the discount factor to play well, but only that it is small enough. In the MDP literature this property is called Blackwell ε -optimality. This is a very desirable robustness property.

2 Gambling Games

2.1 Notations

Given a compact metric space S , we denote by $\mathcal{B}(S)$, resp. by $\mathcal{C}(S)$, the set of bounded measurable, resp. continuous, functions from S to the reals, and by $\Delta(S)$ the set of Borel probabilities over S . For s in S , we denote by $\delta_s \in \Delta(S)$ the Dirac measure on s , and whenever possible we assimilate s and δ_s . For v in $\mathcal{B}(S)$, we denote by \tilde{v} its affine extension to $\Delta(S)$: $\tilde{v}(p) = \mathbb{I}E_p(v)$ for all p in $\Delta(S)$, where $\mathbb{I}E_p(v) := \int_S v(s)dp(s)$ is the expectation of v with respect to p . When there is no confusion, $\tilde{v}(p)$ will also be denoted by $v(p)$. Finally, $\Delta(S)$ is endowed with the weak-* topology, a compatible distance being the Kantorovich-Rubinstein (also called the Wasserstein of order 1) metric.

Definition 2.1. For p, p' in $\Delta(S)$, $d_{KR}(p, p') = \sup_{v \in E_1} |\tilde{v}(p) - \tilde{v}(p')|$, where E_1 is the set of 1-Lipschitz functions on S .

Observe that when $p = \delta_x$ and $p' = \delta_{x'}$ are Dirac measures, then $d_{KR}(p, p') = d(x, x')$, the metric distance between x and x' .

2.2 Model and examples

A gambling game is a zero-sum stochastic game where each player controls his own state variable. We will always assume in this paper that the state spaces are non empty metric and compact, and denote by X and Y the respective set of states² controlled by Player 1 and by Player 2. The transitions of Player 1 are given by a continuous³ multifunction $\Gamma : X \rightrightarrows \Delta(X)$ with non empty convex⁴ compact values: if the state of Player 1 is at x , he can select his new state according to any probability in $\Gamma(x)$. Similarly, a continuous multifunction $\Lambda : Y \rightrightarrows \Delta(Y)$ with non empty convex compact values, gives the transitions of Player 2. The players independently control their own state, and only interact through payoffs: the running payoff of Player 1 is given by a continuous mapping $u : X \times Y \rightarrow \mathbb{R}$, and the payoff to Player 2 is given by $-u$.

Gambling games extend the model of gambling houses [Dubins and Savage, 1965], which correspond to the single player case when Y is a singleton and Player 2 plays no role. It is well known that, by an adequate augmentation of the state space in order to encompass actions, any MDP can be mapped into a gambling house (see [Dubins et al., 2002], [Maitra and Sudderth, 1996] and [Schal, 1989]).

A standard gambling house is the red-and-black casino where $X = [0, 1]$ is a fortune space. Suppose that at each fortune $x \geq 0$, the gambler can stake any amount s in her possession. The gambler loses the stake with probability $1 - w$ where $w \in (0, 1)$ is fixed

²Both metrics are denoted by d , and we will use the metric $d((x, y), (x', y')) = d(x, x') + d(y, y')$ on $X \times Y$.

³i.e. $\forall \varepsilon > 0, \exists \alpha > 0, \forall x, x' \in X$ with $d(x, x') \leq \alpha, \forall p \in \Gamma(x), \exists p' \in \Gamma(x')$ s.t. $d_{KR}(p, p') \leq \varepsilon$. Since Γ has compact values, it implies that the graph of Γ is compact.

⁴If originally the gambling game has non convex values, then allowing as usual players to randomize, would lead to transitions with convex values.

and given, and wins back the stake and an additional equal amount with probability w . The corresponding transition multifunction reads:

$$\Phi_w(x) = \{w\delta_{\min\{x+s,1\}} + (1-w)\delta_{x-s} : 0 \leq s \leq x\}.$$

More generally, a casino is a gambling house Γ on $X = [0, +\infty)$ in which “a rich gambler can do whatever a poor one can do” and a “poor gambler can, on a small scale, imitate a rich one.” Formally, for $x \in X$, let $\Theta(x) := \{\theta \in \Delta(\mathbf{R}) : \exists \gamma \in \Gamma(x) \text{ such that } \gamma = [\theta + x]\}$ where $[a\theta + b]$ for some reals a and b is the probability measure s.t. for all $u \in \mathcal{B}(X)$, $\int_{y \in X} u(y) d[a\theta + b](y) = \int_{y \in X} u(ay + b) d\theta(y)$. With those notations, the gambling house Γ is a casino if for all $x \geq 0$ and $0 \leq t \leq 1$, $[t\Theta(x)] \subset \Theta(tx) \subset \Theta(x)$. A fundamental result of Dubins and Savage (1965) is the classification of casinos into four types: trivial, subfair, fair, and superfair. Here we need only to distinguish superfair casinos from those that are not superfair. A casino is superfair if there is $\theta \in \Theta(x)$ for some $x > 0$ such that $\int_X x d\theta(x) > 0$. The red-and-black casino is superfair if and only if $w > \frac{1}{2}$. Observe that a Casino is superfair if the player has a strategy that wins in expectation against the Casino for at least one $x > 0$ (and so for every $x > 0$ given the definition of a casino).

Another class of gambling houses are splitting problems where $X = \Delta(K)$ is a simplex (K is a finite set) and $\Gamma(x)$ is the set of Borel probabilities σ on X centered at x (for example $\sigma = \sum_{i=1}^n \alpha_i \delta_{x_i}$ where $x = \sum_{i=1}^n \alpha_i x_i$). The idea of splitting was introduced by Aumann and Maschler [Aumann et al., 1995] in the context of repeated games with incomplete information on one side. This “gambling house” type of problems is now very popular in the persuasion and information design literature, see [Kamenica and Gentzkow, 2011].

The above two gambling houses naturally extend to gambling games. One can consider a casino game where each player i controls a red-and-black house with parameter w_i , and the running payoff depends on the current pair of fortunes. Another example is a splitting game introduced in [Laraki, 2001b], [Laraki, 2001a], and [Sorin, 2002] where $X = \Delta(K)$ and $Y = \Delta(L)$ are simplexes, $\Gamma(x)$ is the set of Borel probability measures on X that are centered at x and $\Lambda(y)$ is the set of Borel probability measures on Y that are centered at y .

2.3 Discounted Evaluations

Given a discount factor $\lambda \in (0, 1]$ and an initial state (x_1, y_1) in $X \times Y$, the game $G_\lambda(x_1, y_1)$ is played as follows: at any stage $t \geq 1$, the payoff to Player 1 is $u(x_t, y_t)$ and both players knowing (x_t, y_t) , simultaneously Player 1 chooses p_{t+1} in $\Gamma(x_t)$ and Player 2 chooses q_{t+1} in $\Lambda(y_t)$. Then, x_{t+1} and y_{t+1} are independently selected according to p_{t+1} and q_{t+1} , the new states x_{t+1} and y_{t+1} are publicly announced, and the play goes to stage $t + 1$. Under our assumptions of compact state spaces, continuous transitions with convex compact values and continuous running payoff, it is easy to describe the value of such dynamic game.

Definition 2.2. v_λ is the unique element of $\mathcal{C}(X \times Y)$ s.t. $\forall(x, y) \in X \times Y$,

$$\begin{aligned} v_\lambda(x, y) &= \max_{p \in \Gamma(x)} \min_{q \in \Lambda(y)} (\lambda u(x, y) + (1 - \lambda) \tilde{v}_\lambda(p, q)), \\ &= \min_{q \in \Lambda(y)} \max_{p \in \Gamma(x)} (\lambda u(x, y) + (1 - \lambda) \tilde{v}_\lambda(p, q)). \end{aligned}$$

This is the standard characterization of the value of a discounted game by means of the Shapley operator (dynamic programming principle for games). Existence and uniqueness of v_λ follow from standard fixed-point arguments (see for instance [Mertens et al., 2015], [Rosenberg and Sorin, 2001]). We denote by $v_\lambda(x, y)$ the value of the game $G_\lambda(x, y)$.

The goal of the paper is to study the convergence of $(v_\lambda)_\lambda$ when λ goes to 0, i.e. when players become more and more patient.

Remark 2.3. Cesaro Evaluations. It is also standard to define the value of the n -stage games by: $v_1 = u$, and for $n \geq 1$ and $(x, y) \in X \times Y$:

$$\begin{aligned} v_{n+1}(x, y) &= \frac{1}{n+1} \max_{p \in \Gamma(x)} \min_{q \in \Lambda(y)} (u(x, y) + n \tilde{v}_n(p, q)), \\ &= \frac{1}{n+1} \min_{q \in \Lambda(y)} \max_{p \in \Gamma(x)} (u(x, y) + n \tilde{v}_n(p, q)). \end{aligned}$$

It is known that the uniform convergence of $(v_n)_n$ when n goes to infinity, is equivalent to the uniform convergence of $(v_\lambda)_\lambda$ when λ goes to 0, and in case of convergence both limits are the same (Theorem 2.2 in [Ziliotto, 2016b] applies here).

2.4 Non-expansive transitions

Without further assumptions, convergence of (v_λ) may fail even in the simple case where Γ and Λ are single-valued (“0 player case”, players have no choice), so we will assume throughout the paper that the gambling game is non-expansive, i.e. have non-expansive transitions.

Definition 2.4. *The game has non-expansive transitions if:*

$$\forall x \in X, \forall x' \in X, \forall p \in \Gamma(x), \exists p' \in \Gamma(x'), \text{ s.t. } d_{KR}(p, p') \leq d(x, x'),$$

$$\text{and similarly : } \forall y \in Y, \forall y' \in Y, \forall q \in \Lambda(y), \exists q' \in \Lambda(y'), \text{ s.t. } d_{KR}(q, q') \leq d(y, y').$$

The gambling game has non-expansive transitions if the transitions, viewed as mappings from X to $2^{\Delta(X)}$, and from Y to $2^{\Delta(Y)}$, are 1-Lipschitz for the Hausdorff distance on compact subsets of $\Delta(X)$ and $\Delta(Y)$. Also, when the transitions are deterministic (Dirac measures), we recover the standard definition of non-expansiveness in that case. Note that, with an adequate choice of the metric, the transitions are always non-expansive when X and Y are finite⁵. Moreover splitting games are non-expansive

⁵If X is finite and $d(x, x') = 2$ for $x \neq x'$, then $d_{KR}(p, p') = \|p - p'\|_1$ for p, p' in $\Delta(X)$.

[Laraki, 2001b], and red-and-black casino houses with parameter w are non-expansive if and only if $w \leq \frac{1}{2}$. More generally, a casino is non-expansive if and only if it is not superfair [Laraki and Sudderth, 2004].

Let us mention also Markov chain repeated games with incomplete information where each player observes a private and exogenous Markov chain. These repeated games lead to gambling houses with transitions of the form: X is a simplex $\Delta(K)$, and $\Gamma(x) = \{pM, p \in \Delta(X) \text{ centered at } x\}$ with M a fixed stochastic matrix. Here again, transitions are non-expansive, see [Gensbittel and Renault, 2015].

Let us mention immediately an important consequence of the non-expansive assumption. The proof is in the Appendix.

Proposition 2.5. *Assume the gambling game is non-expansive. Then the family $\{v_\lambda\}_{\lambda \in (0,1]}$ is equicontinuous.*

As a consequence, point-wise and uniform convergence of $\{v_\lambda\}$ are equivalent, and since $X \times Y$ is compact, to prove convergence of $\{v_\lambda\}$ it is enough to prove uniqueness of a limit point.⁶

Remark 2.6. It is not difficult to see that without non-expansivity, $\{v_\lambda\}$ may not be equicontinuous and the convergence may not be uniform. For instance in red-and-black casino with a single player, if the parameter $w > \frac{1}{2}$ and $u(x) = x$, v_λ is continuous for every λ but $v = \lim_{\lambda \rightarrow 0} v_\lambda$ is not : $v(x) = 0$ for $x = 0$ and $v(x) = 1$ for $x > 0$.

There may be cases where non-expansiveness does not hold but there is still equicontinuity of v_λ . In that case, Proposition 3.2 would still apply.

2.5 Excessive, depressive and balanced functions

Definition 2.7. *Let v be in $\mathcal{B}(X \times Y)$.*

1) *v is balanced if $\forall (x, y) \in X \times Y$,*

$$v(x, y) = \max_{p \in \Gamma(x)} \min_{q \in \Lambda(y)} \tilde{v}(p, q) = \min_{q \in \Lambda(y)} \max_{p \in \Gamma(x)} \tilde{v}(p, q).$$

2) *v is excessive (with respect to X) if: $\forall (x, y) \in X \times Y$, $v(x, y) = \max_{p \in \Gamma(x)} \tilde{v}(p, y)$.*

3) *v is depressive (with respect to Y) if: $\forall (x, y) \in X \times Y$, $v(x, y) = \min_{q \in \Lambda(y)} \tilde{v}(x, q)$.*

Observe that any uniform limit v of $(v_\lambda)_{\lambda \in (0,1]}$ is necessarily continuous and balanced (by passing to the limit in definition 2.2).

In a splitting game, excessive means concave with respect to the first variable, and depressive means convex with respect to the second variable.

Definition 2.8.

The gambling game is leavable if: $\forall x \in X, \delta_x \in \Gamma(x)$ and $\forall y \in Y, \delta_y \in \Lambda(y)$.

⁶By convergence or limit point of $\{v_\lambda\}$, we always mean when λ approaches 0.

The gambling game is leavable if each player can remain in any given state. This is a standard assumption [Dubins and Savage, 1965]. This is the case in casinos and splitting games. In the persuasion literature and in repeated games with incomplete information, not moving means revealing no information.

Remark 2.9. *If the game is leavable, any excessive and depressive function in $\mathcal{B}(X)$ is balanced (the converse is not true as example 2.10 shows). Indeed, since the game is leavable, $\delta_y \in \Lambda(y)$ and so*

$$\min_{q \in \Gamma(y)} \max_{p \in \Gamma(x)} \tilde{v}(p, q) \leq \max_{p \in \Gamma(x)} \tilde{v}(p, y) = v(x, y)$$

the equality being due to excessivity. By symmetry,

$$\max_{p \in \Gamma(x)} \min_{q \in \Gamma(y)} \tilde{v}(p, q) \geq v(x, y)$$

since $\max\min \leq \min\max$, the value exists and (δ_x, δ_y) is a saddle point, implying in particular that the game is balanced.

Example 2.10. Consider a gambling game where players 1 and 2 move on the same finite grid of a circle containing 6 nodes in equidistant positions. Any player can move one step to the left, one step to the right, or not move (and choose randomly between these 3 options, so that transitions have convex values). This game is leavable. It is here possible for a player to go from any state to any other state in at most 3 stages (the game may be called *cyclic*), so any excessive and depressive function is necessarily constant. Suppose that Player 1's payoff is 1 if he is at most one step away from Player 2, and Player 1's payoff is 0 otherwise. If the players start a distance at most 1, Player 1 can guarantee that this property will hold forever by not moving or moving one step to the direction of Player 2 and so, in this case we have $v_\lambda = 1$ for every λ . On the other hand, if the players start at a distance at least 2, Player 2 can insure that this property will hold forever, by not moving or moving one step in the opposite direction of Player 1. For these initial states, $v_\lambda = 0$ for every λ . Here, $v = \lim v_\lambda$ is balanced, but neither excessive nor depressive.

Definition 2.11. *A gambling game is standard if both state spaces X and Y are compact metric, the running payoff u is continuous, and the transitions Γ and Λ have non empty convex compact values and are leavable and non-expansive.*

Throughout the paper, we consider only *standard* gambling games.

2.6 Acyclicity

We now come to the main new conditions of the paper.

Definition 2.12.

The gambling house Γ of player 1 is weakly acyclic if there exists φ in $\mathcal{B}(X)$ lower semi-continuous such that:

$$\forall x \in X, \operatorname{Argmax}_{p \in \Gamma(x)} \tilde{\varphi}(p) = \{\delta_x\}.$$

Similarly, the gambling house Λ of player 2 is weakly acyclic if there exists ψ in $\mathcal{B}(Y)$ upper semi-continuous such that:

$$\forall y \in Y, \operatorname{Argmin}_{q \in \Lambda(y)} \tilde{\psi}(q) = \{\delta_y\}.$$

The gambling game is weakly acyclic if both gambling houses are weakly acyclic.

(Notice that both definitions are equivalent by considering $\psi = -\varphi$).

Observe that a weakly acyclic gambling game is necessarily leavable. Weak acyclicity is, to our knowledge, a new condition in the gambling house literature. If the house Γ is weakly acyclic, the “potential” φ decreases in expectation along non stationary trajectories, hence the irreversibility of the process in the space of probabilities over X .

Example 2.13. When $w \leq \frac{1}{2}$, a red-and-black casino is weakly acyclic and similarly for any casino which is not superfair (take φ to be strictly increasing and strictly concave). A splitting game is weakly acyclic (take φ to be any strictly concave function).

We will now define strong acyclicity, our central condition. For this, we need to consider transitions for several stages. We first extend linearly the transitions to $\Delta(X)$ and $\Delta(Y)$ by defining $\tilde{\Gamma} : \Delta(X) \rightrightarrows \Delta(X)$ and $\tilde{\Lambda} : \Delta(Y) \rightrightarrows \Delta(Y)$. More precisely, the graph of $\tilde{\Gamma}$ is defined as the closure of the convex hull of the graph of Γ (viewed as the subset $\{(\delta_x, p), x \in X, p \in \Gamma(x)\}$ of $\Delta(X) \times \Delta(X)$), and similarly the graph of $\tilde{\Lambda}$ is defined as the closed convex hull of the graph of Λ . Because Dirac measures are extreme points of $\Delta(X)$ and $\Delta(Y)$, we have $\tilde{\Gamma}(\delta_x) = \Gamma(x)$ and $\tilde{\Lambda}(\delta_y) = \Lambda(y)$ for each x in X and y in Y . Be careful that in general, for p in $\Delta(X)$ and q in $\Delta(Y)$:

$$\tilde{v}_\lambda(p, q) = \int_X \int_Y v(x', y') dp(x') dq(y') \neq \max_{p' \in \tilde{\Gamma}(p)} \min_{q' \in \tilde{\Lambda}(q)} (\lambda u(p, q) + (1 - \lambda) \tilde{v}_\lambda(p', q')).$$

We now define inductively a sequence of transitions $(\tilde{\Gamma}^n)_n$ from $\Delta(X)$ to $\Delta(X)$, by $\tilde{\Gamma}^0(p) = \{p\}$ for every state p in $\Delta(X)$, and⁷ for each $n \geq 0$, $\tilde{\Gamma}^{n+1} = \tilde{\Gamma}^n \circ \tilde{\Gamma}$. $\tilde{\Gamma}^n(\delta_x)$ represents the set of probabilities over states that Player 1 can reach in n stages from the initial state x in X . Similarly we define $\tilde{\Lambda}^n$ for each n .

Definition 2.14.

1) The reachable set of Player 1 from state x in X is the closure of $\bigcup_{n \geq 0} \tilde{\Gamma}^n(\delta_x)$ in $\Delta(X)$, and is denoted $\Gamma^\infty(x)$. Similarly, the reachable set of Player 2 from state y in Y is the subset $\Lambda^\infty(y)$ of $\Delta(Y)$ defined as the closure of $\bigcup_{n \geq 0} \tilde{\Lambda}^n(\delta_y)$.

⁷The composition being defined by $G \circ H(p) = \{p'' \in \Delta(X), \exists p' \in H(p) \text{ s.t. } p'' \in G(p')\}$.

2) The gambling house Γ of player 1 is strongly acyclic (or simply, acyclic) if there exists φ in $\mathcal{B}(X)$ lower semi-continuous such that:

$$\forall x \in X, \text{Argmax}_{p \in \Gamma^\infty(x)} \tilde{\varphi}(p) = \{\delta_x\}.$$

Similarly, the gambling house Λ of player 2 is strongly acyclic (or simply, acyclic) if there exists ψ in $\mathcal{B}(Y)$ upper semi-continuous such that:

$$\forall y \in Y, \text{Argmin}_{q \in \Lambda^\infty(y)} \tilde{\psi}(q) = \{\delta_y\}.$$

The gambling game is strongly acyclic (or simply, acyclic) if both gambling houses are strongly acyclic.

Observe that the notion of acyclicity is payoff irrelevant (does not depend on the utility function u). Also, notice that since Γ is standard, for every $x \in X$, $(\tilde{\Gamma}^n(\delta_x))_n$ is increasing (for set inclusion), and $\Gamma^\infty(x) = \lim_n \tilde{\Gamma}^n(\delta_x)$. Since $\Gamma^\infty(x) \supset \Gamma(x)$, strong acyclicity of Γ implies weak acyclicity of Γ .

In splitting games, $\tilde{\Gamma}^n(\delta_x) = \Gamma(x)$ for every n and x in X , so weak and strong acyclicity coincide.⁸ We will give in section 4.2 a standard gambling house Γ on a finite set X that is weakly but not strongly acyclic (i.e. Γ^∞ is not weakly acyclic).

If $\Gamma(x)$ is the set of splittings at x with support at most K , then $\Gamma^2(x)$ is the set of splittings at x with support at most K^2 , ..., and $\Gamma^\infty(x)$ is the set of all splittings at x .

One can show that Γ is weakly acyclic with respect to a continuous ϕ if and only if $\tilde{\Gamma}^n(\delta_x)$ is weakly acyclic with respect to ϕ for every $n < +\infty$. This shows that the weak acyclicity property may be lost at $n = +\infty$ and that the difference between weak and strong acyclicity is formally sharp.

3 Main Results

Our main result is the following.

Theorem 3.1. *Consider a standard gambling game.*

1. *If at least one of the players has a strongly acyclic gambling house, (v_λ) uniformly converges to the unique function v in $\mathcal{C}(X \times Y)$ satisfying:*

- a) *v is excessive (in X), i.e. $\forall(x, y) \in X \times Y, v(x, y) = \max_{p \in \Gamma(x)} \tilde{v}(p, y)$,*
- b) *v is depressive (in Y), i.e. $\forall(x, y) \in X \times Y, v(x, y) = \min_{q \in \Gamma(y)} \tilde{v}(x, q)$,*
- c) *v satisfies P1, i.e. $\forall(x, y) \in X \times Y, \exists p \in \Gamma^\infty(x), v(x, y) = \tilde{v}(p, y) \leq u(p, y)$,*
- d) *v satisfies P2, i.e. $\forall(x, y) \in X \times Y, \exists q \in \Lambda^\infty(y), v(x, y) = \tilde{v}(x, q) \geq u(x, q)$.*

Moreover $v = \lim v_\lambda$ is the largest excessive-depressive continuous function satisfying P1, and is also the smallest excessive-depressive continuous function satisfying P2.

2. *Even if both gambling houses are weakly acyclic, (v_λ) may not converge.*

The conditions of the positive result 1. may be interpreted as follows:

⁸For other examples of strongly acyclic houses, see appendix (section 8.5).

- a) and b) : *It is always safe not to move.* For each player, not moving ensures not to degrade the limit value.
- c) and d) : *Each player can reach, if his opponent does not move, the zone when the current payoff is at least as good than the limit value, without degrading the limit value.*

These interpretations will lead later to the construction of simple uniformly optimal strategies under some additional assumptions, see section 8. The convergence and characterization result 1. of theorem 3.1 relies on the following three propositions (proved in the appendix). Recall that, thanks to proposition 2.5, to get convergence of the values it is enough to show uniqueness of a limit point of $(v_\lambda)_\lambda$.

Proposition 3.2. *Assume that one of the players has a weakly acyclic gambling house. If v in $\mathcal{C}(X \times Y)$ is balanced, then v is excessive and depressive.*

Without weak acyclicity on one side, a balanced function may not be excessive and depressive as example 2.10 shows.

Proposition 3.3. *Let v be a limit point of (v_λ) for the uniform convergence. Then v is balanced, and satisfies P1 and P2.*

This proposition provides some properties that all limit points of (v_λ) should satisfy. The next proposition shows that strong acyclicity on one side implies that at most one function will satisfy those properties.

Proposition 3.4. *Assume that one of the players has a strongly acyclic gambling house. Then, any balanced continuous function satisfying P1 is smaller than any balanced continuous function satisfying P2. Consequently, there is at most one balanced continuous function satisfying P1 and P2.*


On the other hand, if none of the players has a strongly acyclic gambling house, one can prove that there may be infinitely many balanced continuous functions satisfying P1 and P2. This will be the case in our counterexamples of section 4.4, where both gambling houses are weakly acyclic. Moreover, one of the counterexamples show that the family of discounted values may not converge as λ goes to zero. This shows that our results are tight and that strong acyclicity condition on one side has very strong consequences both on the convergence of (v_λ) and on the characterization of the limit.

4 Examples

4.1 A strongly acyclic gambling house

Let us first illustrate our characterization on a simple example. Consider the following Markov decision process with 3 states: $X = \{a, b, c\}$ from [Sorin, 2002]. States b and c

are absorbing with respective payoffs 1 and 0. Start at a , choose $\alpha \in I = [0, 1/2]$, and move to b with proba α and to c with proba α^2 .



Here formally Y is a singleton (there is only one player, so we can omit the variable y), $\Gamma(b) = \{\delta_b\}$, $\Gamma(c) = \{\delta_c\}$, and $\Gamma(a) = \text{conv}\{(1 - \alpha - \alpha^2)\delta_a + \alpha\delta_b + \alpha^2\delta_c, \alpha \in I\}$. The payoffs are $u(a) = u(c) = 0$, $u(b) = 1$.

Γ has compact convex values, the transitions are 1-Lipschitz, and the game is leavable. The gambling game is strongly acyclic: just consider φ such that $\varphi(a) = 1$, and $\varphi(b) = \varphi(c) = 0$.


Player 1 can go from state a to state b in infinitely many stages with arbitrarily high probability, by repeating a choice of $\alpha > 0$ small (so that α^2 is much smaller than α), and the limit value v clearly satisfies:

$$v(a) = v(b) = 1, v(c) = 0.$$

This is the unique function $w : X \rightarrow \mathbb{R}$ satisfying the conditions a), b), c), d) of Theorem 3.1: $P1$ and $P2$ implies $u \leq w \leq 1$, and because b and c are absorbing states, $w(b) = 1$ and $w(c) = 0$. Finally, w excessive gives $w(a) = 1$. Notice that $\delta_b \in \Gamma_\infty(a)$ but for each n , $\delta_b \notin \tilde{\Gamma}_n(a)$.

4.2 A weakly acyclic gambling house

Let us modify the gambling house of the previous section 4.1. We still have a unique player and a state space $X = \{a, b, c\}$. The only difference is that state b is no longer absorbing : in state b the player also has to choose some $\alpha \in I = [0, 1/2]$, and then moves to a with probability α , to c with probability α^2 and remains in b with probability $1 - \alpha - \alpha^2$.



States a and b are now symmetric. This gambling house is weakly acyclic, with $\varphi(a) = \varphi(b) = 1$, $\varphi(c) = 0$, but it is not strongly acyclic since $a \in \Gamma^\infty(b)$ and $b \in \Gamma^\infty(a)$. We will later use this gambling house to construct our counterexample of theorem 3.1, 2).

4.3 An example with countable state spaces

We present here a (strongly) acyclic gambling game with countable state spaces, and illustrate the proof of proposition 3.2, that under weak acyclicity any continuous balanced function is also excessive and depressive. Consider the state space:

$$X = \{1 - \frac{1}{n}, n \in \mathbb{N}^*\} \cup \{1\} = \{x_1, \dots, x_n, \dots, x_\infty\},$$

where $x_n = 1 - \frac{1}{n}$ if n is finite, and $x_\infty = 1$. We use $d(x, x') = |x - x'|$, so that X is countable and compact. The transition is given by:

$$\Gamma(x_n) = \{\alpha\delta_{x_n} + (1 - \alpha)\delta_{x_{n+1}}, \alpha \in [0, 1]\}, \text{ and } \Gamma(x_\infty) = \{\delta_{x_\infty}\}.$$

The intuition is clear: Player 1 can stay at his location, or move 1 to the right. The gambling house (Y, Λ) of Player 2 is a copy of the gambling house of Player 1. Transitions are non-expansive (since $|\frac{1}{n+1} - \frac{1}{n'+1}| \leq |\frac{1}{n} - \frac{1}{n'}|$), and the game is strongly acyclic. The payoff u is any continuous function $X \times Y \rightarrow \mathbb{R}$, so that theorem 3.1 applies.

Consider $v : X \times Y \rightarrow \mathbb{R}$, and for simplicity we use $w(n, m) = v(x_n, x_m)$. Here v excessive means that $w(n, m)$ is weakly decreasing in n , and v depressive means that $w(n, m)$ is weakly increasing in m . The meaning of v balanced is the following: for each n and m , $w(n, m)$ is the value of the matrix game (“local game” at (n, m)):

$$\begin{pmatrix} w(n+1, m) & w(n+1, m+1) \\ w(n, m) & w(n, m+1) \end{pmatrix}.$$

Clearly, if v is excessive and depressive it is balanced, but proposition 3.2 tells that *if v is continuous*, then the converse also holds: balancedness implies excessiveness and depressiveness. The idea of the proof of proposition 3.2 can be seen here as follows.

Suppose v is balanced, but not excessive. Then one can find n and m such that $w(n+1, m) > w(n, m)$. Because $w(n, m)$ is the value of the local game at (n, m) , we necessarily have $w(n+1, m+1) \leq w(n, m)$, and $w(n+1, m) > w(n+1, m+1)$. Consider now the “local game” at $(n+1, m)$. $w(n+1, m)$ is the value of the matrix:

$$\begin{pmatrix} w(n+2, m) & w(n+2, m+1) \\ w(n+1, m) & w(n+1, m+1) \end{pmatrix}.$$

Since $w(n+1, m) > w(n+1, m+1)$, we obtain $w(n+2, m+1) \geq w(n+1, m)$. We have obtained: $w(n+2, m+1) \geq w(n+1, m) > w(n, m) \geq w(n+1, m+1)$, so

$$w(n+2, m+1) - w(n+1, m+1) > w(n+1, m) - w(n, m).$$

Iterating the argument, we obtain that for each p ,

$$w(n + p + 1, m + p) - w(n + p, m + p) \geq w(n + 1, m) - w(n, m) > 0.$$


And this is a contradiction with w being continuous at infinity.

To conclude with this example, consider the simple case where the running payoff is given by $u(x, y) = |x - y|$. Player 1 wants to be far from Player 2, and Player 2 wants to be close to Player 1. If initially $n < m$, it is optimal for each player not to move, so $w(n, m) = |x_n - x_m|$. Suppose on the contrary that initially $n \geq m$, so that Player 1 is more to the right than Player 2. Then Player 2 has a simple optimal strategy which is to move to the right if the current positions satisfy $x > y$, and to stay at y if $x = y$. No matter how large is the initial difference $n - m$, Player 2 will succeed in being close to player, so that $w(n, m) = 0$ if $n \geq m$.

4.4 A weakly acyclic game without limit value

Here we prove the second part of theorem 3.1 by providing a counterexample to the convergence of (v_λ) in a weakly acyclic non-expansive gambling house.

The states and transitions for Player 1 are as in example 4.2:



The set of states of Player 1 is $X = \{a, b, c\}$. The difference with example 4.2 is that the set of possible choices for α_a and α_b may be smaller than $[0, 1/2]$. Here α_a and α_b now belong to some fixed compact set $I \subset [0, 1/2]$ such⁹ that 0 is in the closure of $I \setminus \{0\}$. Then $0 \in I$, the transitions are leavable and non-expansive. States a and b are symmetric, this gambling house is weakly acyclic, with $\varphi(a) = \varphi(b) = 1, \varphi(c) = 0$, but not strongly acyclic since $a \in \Gamma^\infty(b)$ and $b \in \Gamma^\infty(a)$.

The gambling house of Player 2 is a copy of the gambling house of Player 1, with state space $Y = \{a', b', c'\}$ and a compact set of choices $J \subset [0, 1/2]$ such that 0 is in the closure of $J \setminus \{0\}$. The unique difference between the gambling houses of the players is that I and J may be different. Payoffs are simple:

$$u(x, y) = 0 \text{ if } x = y, \quad u(x, y) = 1 \text{ if } x \neq y.$$

⁹Except in part 2 of theorem 4.2 where we will consider the case $I = \{0, 1/4\}$.

The u function can be written as follows

	a'	b'	c'
a	0	1	1
b	1	0	1
c	1	1	0

, with a clear interpretation : Player 1 and Player 2 both move on a space with 3 points, Player 2 wants to be at the same location as Player 1, and Player 1 wants the opposite.

Here the gambling game is weakly acyclic but not strongly acyclic, and the following lemma shows that the uniqueness property of proposition 3.4 fails.

Lemma 4.1. *Let $v : X \times Y \rightarrow \mathbb{R}$. Then A) and B) are equivalent:*

- A) v is excessive, depressive, and satisfies P1 and P2,
- B) There exists $x \in [0, 1]$ such that v can be written:

	a'	b'	c'
a	x	x	1
b	x	x	1
c	0	0	0

Proof: Assume v satisfies A). Since u takes values in $[0, 1]$, so does v . Because c and c' are absorbing, we have $v(c, c') = u(c, c') = 0$.

Consider now $v(c, a')$. v being depressive, for any fixed $\beta^* > 0$ in J we have:

$$v(c, a') \leq \beta^* v(c, b') + \beta^{*2} v(c, c') + (1 - \beta^* - \beta^{*2}) v(c, a').$$

and we obtain $v(c, a') \leq \frac{1}{1+\beta^*} v(c, b')$. But symmetrically we also have $v(c, b') \leq \frac{1}{1+\beta^*} v(c, a')$, and we get $v(c, a') = v(c, b') = 0$.

Consider now $v(a, c')$. By P2, we obtain that $v(a, c') \geq u(a, c') = 1$, and $v(a, c') = 1$. Similarly, $v(b, c') = 1$.

Consider now $v(a, a')$. v being excessive, for any $\alpha > 0$ in I we have:

$$v(a, a') \geq \alpha v(b, a') + \alpha^2 v(c, a') + (1 - \alpha - \alpha^2) v(a, a').$$

Hence $v(a, a') \geq \frac{1}{1+\alpha} v(b, a')$, and by assumption on I we obtain $v(a, a') \geq v(b, a')$. By symmetry of the transitions between a and b , $v(a, a') = v(b, a')$. Similarly, $v(a, b') = v(b, b')$.

It only remains to prove that $v(a, a') = v(a, b')$. v being depressive, for any $\beta > 0$ in J , $v(a, a') \leq \beta v(a, b') + \beta^2 v(a, c') + (1 - \beta - \beta^2) v(a, a')$. By assumption on J , we get $v(a, a') \leq v(a, b')$. By symmetry of the transitions between b and b' , $v(a, b') = v(a, a')$, and v satisfies B).

One can easily check that B) implies A), and the proof of lemma 4.1 is complete.

The second part of theorem 3.1 is a direct consequence of point 3 of the following theorem.

Theorem 4.2.

1) If $I = J = [0, 1/4]$, the limit value exists and is:

	a'	b'	c'
a	1/2	1/2	1
b	1/2	1/2	1
c	0	0	0

2) If $J = [0, 1/4]$ and $I = \{0, 1/4\}$, the limit value exists and is:

	a'	b'	c'
a	0	0	1
b	0	0	1
c	0	0	0

3) If $J = [0, 1/4]$ and $I = \{\frac{1}{4^n}, n \in \mathbb{N}^*\} \cup \{0\}$, then v_λ diverges.

For the proof see appendix (section 7.3).

5 Extensions

5.1 Gambling houses (or Markov Decision Processes)

We assume here that there is a unique player, i.e. that Y is a singleton. Then non-expansiveness is enough to guarantee the uniform convergence of $(v_\lambda)_\lambda$ (as well as the uniform value, see [Renault, 2011]) and the limit v can be characterized as follows [Renault and Venel, 2016]: for all x in X ,

$$v(x) = \inf\{w(x), w : \Delta(X) \rightarrow \mathbb{R} \text{ affine } C^0 \text{ s.t.} \\ (1) \forall x' \in X, w(x') \geq \sup_{p \in \Gamma(x')} w(p), \text{ and } (2) \forall r \in R, w(r) \geq u(r)\},$$

where $R = \{p \in \Delta(X), (p, p) \in \text{Graph } \tilde{\Gamma}\}$ is interpreted as the set of *invariant measures* for the gambling house (which is not necessarily leavable here). If we moreover assume that the gambling house is leavable, then $R = \Delta(X)$ and we recover the fundamental theorem of gambling ([Dubins and Savage, 1965], [Maitra and Sudderth, 1996]), namely, (v_λ) uniformly converges to:

$$v = \min\{w \in \mathcal{C}(X), w \text{ excessive}, w \geq u\} = \min\{w \in \mathcal{B}(X), w \text{ excessive}, w \geq u\}.$$

It is also easy to see that $v(x) = \sup_{p \in \Gamma^\infty(x)} u(p)$ for each x .

Our approach will lead to other characterizations. We don't assume any acyclicity condition in the following theorem.

Theorem 5.1. *Consider a one player (leavable and non-expansive) gambling house. Then (v_λ) uniformly converges to the unique function v in $\mathcal{C}(X)$ satisfying: v is excessive, $P1 : \forall x \in X, \exists p \in \Gamma^\infty(x), v(x) = \tilde{v}(p) \leq \tilde{u}(p)$, and $P2 : v \geq u$.*

Proof: From proposition 3.3 any accumulation point of (v_λ) is excessive and satisfies $P1$ and $P2$. Thus, we just need to show uniqueness, which is a direct consequence of the following lemma.

Lemma 5.2. *If $v_1 \in \mathcal{B}(X)$ satisfies $P1$ and $v_2 \in \mathcal{C}(X)$ is excessive and satisfies $P2$, then $v_1 \leq v_2$.*

Proof: Take any $x \in X$. Then there is $p \in \Gamma^\infty(x)$ such that $v_1(x) = v_1(p) \leq u(p)$. Because v_2 is excessive and continuous, by lemma 7.3 $v_2(x) \geq v_2(p)$. Since $v_2 \geq u$, we have $v_2(p) \geq u(p)$. Consequently, one has $v_2(x) \geq u(p)$. Thus, $v_1(x) - v_2(x) \leq 0$, as desired.

Using the gambling fundamental theorem, we obtain new viewpoints on the characterization of the limit value in leavable gambling houses.

Corollary 5.3. *Consider a one player standard gambling house. Then the asymptotic value exists and is:*

- (1) *the smallest excessive function v in $\mathcal{B}(X)$ satisfying $P2$;*
- (2) *the largest excessive function v in $\mathcal{B}(X)$ satisfying $P1$;*
- (3) *the unique excessive function v in $\mathcal{B}(X)$ satisfying $P1$ and $P2$.*

Moreover, v is continuous.

5.2 Other characterizations and link with the Mertens Zamir system

Recall that we consider standard gambling games.

Definition 5.4. *Consider g in $\mathcal{C}(X \times Y)$. The excessive envelope of g , $Exc_\Gamma(g)$, is the smallest excessive (w.r.t. X) function greater or equal to g and the depressive envelope of g , $Dep_\Lambda(g)$ is the largest depressive (w.r.t. Y) function smaller or equal to g .*

The existence of $Exc_\Gamma(g)$ (and similarly $Dep_\Lambda(g)$) is standard: for every (x, y) , we have $Exc_\Gamma(g)(x, y) = \inf\{f(x, y) \text{ s.t. } f(x', y) \geq g(x', y) \text{ for all } x', \text{ and } f \text{ is bounded measurable and } \Gamma\text{-excessive}\}$. Clearly, $Exc_\Gamma(g) \geq g$. Moreover, for every $f \in \mathcal{B}(X, Y)$ which is Γ -excessive and satisfies $f(x', y) \geq g(x', y)$ for all x' , every $(x, y) \in X$, and every $\sigma \in \Gamma(x)$, we have $\int Exc_\Gamma(g)(x', y) d\sigma(x') \leq \int f(x', y) d\sigma(x') \leq f(x, y)$. Taking the infimum w.r.t. f , one obtains that $Exc_\Gamma(g)$ is Γ -excessive and so, the smallest excessive (w.r.t. X) function greater or equal to g .

Exc_Γ is usually called the *réduite* operator and $Dep_\Lambda(g) = -Exc_\Gamma(-g)$. In splitting games, $Exc_\Gamma = Cav_X$ is the concavification operator on X and $Dep_\Lambda(g) = Vex_Y$ is the convexification operator on Y . We introduce the following definition by analogy with the Mertens-Zamir characterization.

Definition 5.5. A function v in $\mathcal{C}(X \times Y)$ satisfies the *MZ-characterization* if:

$$\begin{aligned} \text{MZ1: } & \forall(x, y) \in X \times Y, v(x, y) = \text{Exc}_\Gamma \min(u, v)(x, y), \\ \text{and MZ2: } & \forall(x, y) \in X \times Y, v(x, y) = \text{Dep}_\Lambda \max(u, v)(x, y). \end{aligned}$$

We now introduce other properties, by analogy with the one established for splitting games and repeated games with incomplete information, see for instance [Laraki, 2001b], [Laraki, 2001a], and [Rosenberg and Sorin, 2001].

Definition 5.6. Let v be in $\mathcal{C}(X \times Y)$.

1) For each (x, y) in $X \times Y$,

$$\begin{aligned} x \text{ is extreme for } v(\cdot, y) & \text{ if } \arg \max_{p \in \Gamma^\infty(x)} \tilde{v}(p, y) = \{\delta_x\}. \\ y \text{ is extreme for } v(x, \cdot) & \text{ if } \arg \min_{q \in \Lambda^\infty(y)} \tilde{v}(x, q) = \{\delta_y\}. \end{aligned}$$

2) v satisfies the *E-characterization* if:

E1: for all $(x, y) \in X \times Y$, if x is extreme for $v(\cdot, y)$ then $v(x, y) \leq u(x, y)$,
and *E2*: for all $(x, y) \in X \times Y$, if y is extreme for $v(x, \cdot)$ then $v(x, y) \geq u(x, y)$.

Proposition 5.7. Consider a standard gambling game and let v in $\mathcal{C}(X \times Y)$ be excessive and depressive. Then:

$$\begin{aligned} v \text{ satisfies MZ1} & \implies v \text{ satisfies P1} \implies v \text{ satisfies E1}, \\ \text{and } v \text{ satisfies MZ2} & \implies v \text{ satisfies P2} \implies v \text{ satisfies E2}. \end{aligned}$$

Proof. Let v be a continuous excessive function that satisfies *MZ1*. Fix y and define for each x , $f(x) = \min(v(x, y), u(x, y))$. Then, for every x , $v(x, y) = \text{Exc}_\Gamma(f)(x)$. We consider the gambling house for Player 1 where the state of Player 2 is fixed to y and the payoff is given by f . From corollary 5.3, there is $p \in \Gamma^\infty(x)$ such that $v(x, y) = v(p, y) \leq f(p)$. Since $f(p) \leq u(p, y)$, v satisfies *P1*.

Now, let v be an excessive continuous function that satisfies *P1*. Take any x and y and suppose that x is extreme for $v(\cdot, y)$. By *P1*, there is $p^* \in \Gamma^\infty(x)$ such that $v(x, y) = v(p^*, y) \leq u(p^*, y)$. Because v is excessive and continuous, by lemma 7.3 we have $p^* \in \arg \max_{p \in \Gamma^\infty} \tilde{v}(p, y)$. Because x is extreme for $v(\cdot, y)$, $p^* = \delta_x$ and so, $v(x, y) \leq u(x, y)$. Consequently, *E1* is satisfied.

By symmetry, *MZ2* \implies *P2* \implies *E2*.

Remark 5.8. It is easy to find examples where *E1* is satisfied but *MZ1* is not. For instance, assume that Y is a singleton, and that $\Gamma(x) = \Delta(X)$ for each x in X . Consider the constant, hence excessive, functions $u = 0$ and $v = 1$. v has no extreme points hence trivially satisfies *E1*, but $\text{Exc}_\Gamma \min(u, v) = u$ and so, v does not satisfy *MZ1*.

Proposition 5.9. *Consider a standard gambling game and let v be an excessive-depressive function in $\mathcal{C}(X \times Y)$. Then:*

(Γ strongly acyclic) and (v satisfies $E1$) \implies (v satisfies $MZ1$),

and;

(Λ strongly acyclic) and (v satisfies $E2$) \implies (v satisfies $MZ2$).

Consequently, if the gambling game is strongly acyclic, characterizations MZ , P and E are equivalents.

Proof. Let v be excessive-depressive that satisfies $E1$. Fix $y \in Y$. We want to show that $v(x, y) = g(x, y)$ where $g = \text{Exc}_\Gamma(f)$ and $f = \min(u, v)$. g is continuous by corollary 5.3. Since v is excessive and $v \geq f$, we have $v \geq g$. Let $Z = \arg \max_{x \in X} v(x, y) - g(x, y)$ and let $x_0 = \arg \min_{x \in Z} \varphi(x)$, where φ comes from the definition of acyclicity. It is enough to prove that $v(x_0, y) \leq g(x_0, y)$.

Suppose not. We have $g(x_0, y) \geq f(x_0, y)$, so (1) $g(x_0, y) \geq u(x_0, y)$. Now, let $p_0 \in \Gamma^\infty(x_0)$ such that $v(x_0, y) = v(p_0, y)$. Because g is excessive and continuous, $g(p_0, y) \leq g(x_0, y)$ (lemma 7.3). Consequently, $v(p_0, y) - g(p_0, y) \geq v(x_0, y) - g(x_0, y)$. Consequently, p_0 is supported on Z . Thus, $\varphi(x_0, y) \leq \varphi(p_0, y)$, and by strong acyclicity $p_0 = \delta_{x_0}$. Thus x_0 is an extreme point of $v(\cdot, y)$. By $E1$, we have (2) $v(x_0, y) \leq u(x_0, y)$. By (1) and (2), $v(x_0, y) - g(x_0, y) \leq 0$. A contradiction.

6 Conclusion, open problems and future directions

We introduce the class of gambling games. It is a sub-class of stochastic games which includes MDP problems, splitting games and product stochastic games. We define a strong notion of acyclicity under which we prove existence of the asymptotic value v and we establish several characterizations of v which are linked to the Mertens-Zamir system of functional equations (re-formulated in our more general set-up). We also prove that our condition is tight: a slight weakening of acyclicity implies non-existence of the asymptotic value. Our example is the first in the class of product stochastic games and is probably the simplest known counterexample of convergence for finite state spaces and compact action set (the first counterexample in this class was established in [Vigeral, 2013]). Many questions merit to be investigated in a future research:

- In standard gambling games, is it possible to characterize the asymptotic value in models where we know it exists (for example when X and Y are finite, transition function is polynomial and Γ and Λ are definable [Bolte et al., 2015] in an o-minimal structure)? We know that the asymptotic value is balanced and satisfy P1 and P2, but we may have infinitely many functions satisfying those properties.
- Is there an asymptotic value if one house is strongly acyclic and the other not necessarily leavable? As seen, even when both houses are weakly acyclic, we may have divergence: strong acyclicity of one of the two houses is necessary.

- Recently splitting games have been extended to continuous time and linked to differential games with incomplete information [Gensbittel and Rainer, 2017]. How our model and results extend to continuous time? The one player game has been investigated in [Buckdahn et al., 2015] and [Li et al., 2016].
- It would be interesting to study the non-zero sum analogue of this model. Actually, a static version of non-zero-sum splitting games, with a discontinuous payoff function, has been recently explored by [Koessler et al., 2018] and one natural extension is the dynamic model. Observe that for each discount factor λ , under some regularity assumptions, one can prove existence of subgame perfect equilibrium payoffs E_λ , and establish a standard recursive structure. An interesting question is : does E_λ converges as λ goes to zero -if all players have a strongly acyclic gambling house- and if so, how to characterize this limit.

References

- [Aliprantis and Border, 2006] Aliprantis, C. D. and Border, K. (2006). *Infinite Dimensional Analysis: a Hitchhiker's Guide*. Springer.
- [Aumann et al., 1995] Aumann, R., Maschler, M., and Stearns, R. (1995). *Repeated Games with Incomplete Information*. MIT Press.
- [Bewley and Kohlberg, 1976] Bewley, T. and Kohlberg, E. (1976). The asymptotic theory of stochastic games. *Mathematics of Operations Research*, 1:197–208.
- [Bolte et al., 2015] Bolte, J., Gaubert, S., and Vigerel, G. (2015). Definable zero-sum stochastic games. *Mathematics of Operations Research*, 40:171–191.
- [Buckdahn et al., 2015] Buckdahn, R., Quincampoix, M., and Renault, J. (2015). On representation formulas for long run averaging optimal control problem. *Journal of Differential Equations*, 259:5554–5581.
- [Dubins et al., 2002] Dubins, L., Maitra, A. P., and Sudderth, W. D. (2002). Invariant gambling problems and markov decision processes. In Feinberg, E. A. and Shwartz, A., editors, *Handbook of Markov Decision Processes*. Kluwer.
- [Dubins and Savage, 1965] Dubins, L. and Savage, L. J. (1965). *Inequalities for Stochastic Processes*. McGraw-Hill.
- [Everett, 1957] Everett, H. (1957). Recursive games. In Tucker, A. and Wolfe, P., editors, *Contributions to the Theory of Games III*. Princeton University Press.
- [Flesch et al., 2008] Flesch, J., Schoenmakers, G., and Vrieze, K. (2008). Stochastic games on a product state space. *Mathematics of Operations Research*, 33(2):403–420.

- [Gensbittel and Rainer, 2017] Gensbittel, F. and Rainer, C. (2017). A probabilistic representation for the value of zero-sum differential games with incomplete information on both sides. *SIAM Journal on Control Optimization*, 55(2):693–723.
- [Gensbittel and Renault, 2015] Gensbittel, F. and Renault, J. (2015). The value of markov chain games with lack of information on both sides. *Mathematics of Operations Research*, 40(4):820–841.
- [Kamenica and Gentzkow, 2011] Kamenica, E. and Gentzkow, M. (2011). Bayesian persuasion. *The American Economic Review*, 101(6):2590–2615.
- [Koessler et al., 2018] Koessler, F., Laclau, L., and Tomala, T. (2018). Interactive information design. *Preprint*.
- [Kohlberg, 1974] Kohlberg, E. (1974). Repeated games with absorbing states. *Annals of Statistics*, 2:724–738.
- [Laraki, 2001a] Laraki, R. (2001a). The splitting game and applications. *International Journal of Game Theory*, 30:359–376.
- [Laraki, 2001b] Laraki, R. (2001b). Variational inequalities, systems of functional equations and incomplete information repeated games. *SIAM Journal on Control Optimization*, 40:516–524.
- [Laraki and J.B-Lasserre, 2008] Laraki, R. and J.B-Lasserre (2008). Computing uniform convex approximations for convex envelopes and convex hull. *Journal of Convex Analysis*, 3:635–654.
- [Laraki and Sorin, 2014] Laraki, R. and Sorin, S. (2014). Advances in zero-sum dynamic games. In Young, H. P. and Zamir, S., editors, *Handbook of Game Theory IV*, pages 27–93. North Holland.
- [Laraki and Sudderth, 2004] Laraki, R. and Sudderth, W. D. (2004). The preservation of continuity and lipschitz continuity by optimal rewards operators. *Mathematics of Operations Research*, 29:672–685.
- [Li et al., 2016] Li, X., Quincampoix, M., and Renault, J. (2016). Limit value for optimal control with general means. *Discrete and Continuous Dynamical System A*, 36(4):2113–2132.
- [Maitra and Sudderth, 1996] Maitra, A. and Sudderth, W. D. (1996). *Discrete Gambling and Stochastic Games*. Springer Verlag.
- [Mertens, 1986] Mertens, J.-F. (1986). Repeated games. In *Proceedings of the International Congress of Mathematicians*, pages 1528–1577. American Mathematical Society.

- [Mertens et al., 2009] Mertens, J.-F., Neyman, A., and Rosenberg, D. (2009). Absorbing games with compact action spaces. *Mathematics of Operations Research*, 34:257–262.
- [Mertens et al., 2015] Mertens, J.-F., Sorin, S., and Zamir, S. (2015). *Repeated Games*. Cambridge University Press.
- [Mertens and Zamir, 1971] Mertens, J.-F. and Zamir, S. (1971). The value of two-person zero-sum repeated games with lack of information on both sides. *International Journal of Game Theory*, 1:39–64.
- [Oliu-Barton, 2018] Oliu-Barton, M. (2018). The splitting game: Uniform value and optimal strategies. *Dynamic Games and Applications*, 8:157–179.
- [Renault, 2011] Renault, J. (2011). Uniform value in dynamic programming. *Journal of the European Mathematical Society*, 13(2):309–330.
- [Renault and Venel, 2016] Renault, J. and Venel, X. (2016). Long-term values in markov decision processes and repeated games, and a new distance for probability spaces. *Mathematics of Operations Research*, 42(2):249–276.
- [Rosenberg and Sorin, 2001] Rosenberg, D. and Sorin, S. (2001). An operator approach to zero-sum repeated games. *Israel Journal of Mathematics*, 121:221–246.
- [Schal, 1989] Schal, M. (1989). On stochastic dynamic programming: A bridge between markov decision processes and gambling. In *Markov Processes and Control Theory*, number 54 in Mathematical Research. Akademie-Verlag, Berlin, Germany.
- [Shapley, 1953] Shapley, L. S. (1953). Stochastic games. *Proceedings of the National Academy of Sciences of the U.S.A.*, 39:1095–110.
- [Sorin, 2002] Sorin, S. (2002). *A First Course on Zero-Sum Repeated Games*. Springer.
- [Sorin and Neyman, 2003] Sorin, S. and Neyman, A., editors (2003). *Stochastic Games and Applications*. NATO Science Series C 570. Kluwer Academic Publishers.
- [Sorin and Vigerál, 2013] Sorin, S. and Vigerál, G. (2013). Existence of the limit value of two person zero-sum discounted repeated games via comparison theorems. *JOTA*, 157:564–576.
- [Vigerál, 2013] Vigerál, G. (2013). A zero-sum stochastic game with compact action sets and no asymptotic value. *Dynamic Games and Applications*, 3(2):172–186.
- [Ziliotto, 2016a] Ziliotto, B. (2016a). Annals of probability. *Zero-Sum Repeated Games: Counterexamples to the Existence of the Asymptotic Value and the Conjecture $\max \min = \lim v(n)$* , 44(2):1107–1133.
- [Ziliotto, 2016b] Ziliotto, B. (2016b). A tauberian theorem for nonexpansive operators and applications to zero-sum stochastic games. *Mathematics of Operations Research*, 41(4):1522–1534.

7 Appendix A: missing proofs

7.1 Proof of proposition 2.5

u being uniformly continuous over the compact set $X \times Y$, we consider a concave modulus of continuity $\omega : \mathbb{R}_+ \rightarrow \mathbb{R}_+$:

$$|u(x, y) - u(x', y')| \leq \omega(d(x, x') + d(y, y')), \quad \forall x, x' \in X, \forall y, y' \in Y.$$

ω is non decreasing, concave and $\lim_0 \omega = 0$. Denote by \mathcal{C} the set of functions v in $\mathcal{C}(X \times Y)$ satisfying: $|v(x, y) - v(x', y')| \leq \omega(d(x, x') + d(y, y')), \quad \forall x, x' \in X, \forall y, y' \in Y$. We start with a lemma.

Lemma 7.1. *For v in \mathcal{C} , p, p' in $\Delta(X)$, q, q' in $\Delta(Y)$,*

$$|\tilde{v}(p, q) - \tilde{v}(p', q')| \leq \omega(d_{KR}(p, p') + d_{KR}(q, q')).$$

Proof of lemma 7.1: By the Kantorovich duality theorem, there exists μ in $\Delta(X \times X)$ with first marginal p and second marginal p' satisfying: $d_{KR}(p, p') = \int_{X \times X} d(x, x') d\mu(x, x')$. Similarly there exists ν in $\Delta(Y \times Y)$ with first marginal q and second marginal q' satisfying: $d_{KR}(q, q') = \int_{Y \times Y} d(y, y') d\nu(y, y')$. We have for all x, x', y, y' :

$$v(x, y) \geq v(x', y') - \omega(d(x, x') + d(y, y')).$$

We integrate the above inequality with respect to the probability $\mu \otimes \nu$, and obtain using the concavity of ω :

$$\begin{aligned} \tilde{v}(p, q) &\geq \tilde{v}(p', q') - \int_{X^2 \times Y^2} \omega(d(x, x') + d(y, y')) d\mu(x, x') d\nu(y, y'), \\ &\geq \tilde{v}(p', q') - \omega \left(\int_{X^2 \times Y^2} d(x, x') + d(y, y') d\mu(x, x') d\nu(y, y') \right) \\ &= \tilde{v}(p', q') - \omega(d_{KR}(p, p') + d_{KR}(q, q')). \end{aligned}$$

We now return to the proof of proposition 2.5. Fix λ in $(0, 1]$. Given v in \mathcal{C} , define $\Phi(v) : X \times Y \rightarrow \mathbb{R}$ by: $\Phi(v)(x, y) = \sup_{p \in \Gamma(x)} \inf_{q \in \Lambda(y)} \lambda u(x, y) + (1 - \lambda) \tilde{v}(p, q)$.

Consider the zero-sum game with strategy spaces $\Gamma(x)$ and $\Lambda(y)$ and payoff function $(p, q) \mapsto \lambda u(x, y) + (1 - \lambda) \tilde{v}(p, q)$. The strategy spaces are convex compact and the payoff function is continuous and affine in each variable, hence by Sion's theorem we have:

$$\Phi(v)(x, y) = \max_{p \in \Gamma(x)} \min_{q \in \Lambda(y)} \lambda u(x, y) + (1 - \lambda) \tilde{v}(p, q) = \min_{q \in \Lambda(y)} \max_{p \in \Gamma(x)} \lambda u(x, y) + (1 - \lambda) \tilde{v}(p, q).$$

Consider (x, y) and (x', y') in $X \times Y$, and let p in $\Gamma(x)$ be an optimal strategy of Player 1 in the zero-sum game corresponding to (x, y) . The gambling game has

non-expansive transitions, so there exists $p' \in \Gamma(x')$ such that $d_{KR}(p, p') \leq d(x, x')$. Consider any q' in $\Lambda(y')$, there exists q in $\Gamma(y)$ with $d_{KR}(q, q') \leq d(y, y')$. Now, using lemma 7.1 we write:

$$\begin{aligned} & \lambda u(x', y') + (1 - \lambda) \tilde{v}(p', q') \\ & \geq \lambda(u(x, y) - \omega(d(x, x') + d(y, y'))) + (1 - \lambda)(\tilde{v}(p, q) - \omega(d_{KR}(p, p') + d_{KR}(q, q'))), \\ & \geq \Phi(v)(x, y) - \omega(d(x, x') + d(y, y')). \end{aligned}$$

We obtain $\Phi(v)(x', y') \geq \Phi(v)(x, y) - \omega(d(x, x') + d(y, y'))$, and $\Phi(v)$ belongs to \mathcal{C} .

The rest of the proof is very standard. \mathcal{C} is a complete metric space for $\|v - w\| = \sup_{(x,y) \in X \times Y} |v(x, y) - w(x, y)|$, and we have $\|\Phi(v) - \Phi(w)\| \leq (1 - \lambda)\|v - w\|$, so Φ is $(1 - \lambda)$ -contracting. Hence Φ has a unique fixed point which is v_λ . Each v_λ is in \mathcal{C} , and we obtain that the family $(v_\lambda)_{\lambda \in (0,1]}$ is equicontinuous, ending the proof of proposition 2.5.

7.2 Proof of Theorem 3.1, part 1.

We prove here propositions 3.2, 3.3 and 3.4.

7.2.1 Proof of proposition 3.2

By symmetry, suppose that Γ is weakly acyclic and Λ Leavable. Let us prove that any balanced continuous function v is excessive-depressive. Let us first prove that v is excessive.

Fix any (x_0, y_0) in $X \times Y$, and $p_1 \in \Gamma(x_0)$. A direct consequence of balancedness is the existence of q_1 in $\Lambda(y_0)$ such that $v(p_1, q_1) \leq v(x_0, y_0)$. Now, p_1 is in $\Delta(X)$ and q_1 is in $\Delta(Y)$. One has to be careful that there may not exist $p_2 \in \tilde{\Gamma}(p_1)$ such that for all $q_2 \in \tilde{\Lambda}(q_1)$, $v(p_2, q_2) \geq v(p_1, q_1)$. This is because, \tilde{v} being affine in each variable, $v(p_1, q_1)$ can be interpreted as the value of the auxiliary game where first x and y are chosen according to $p_1 \otimes q_1$ and *observed by the players*, then players respectively choose $p \in \Gamma(x)$ and $q \in \Lambda(y)$ and finally Player 1's payoff is $v(p, q)$. And to play well in this game Player 1 has to know the realization of q_1 before choosing p . However since y_0 is a Dirac measure, balancedness implies that there exists $p_2 \in \tilde{\Gamma}(p_1)$ such that $v(p_2, q_1) \geq v(p_1, y_0)$. We have obtained the following lemma.

Lemma 7.2. *Given (x_0, y_0) in $X \times Y$, and $p_1 \in \Gamma(x_0)$, there exists q_1 in $\Lambda(y_0)$ and $p_2 \in \tilde{\Gamma}(p_1)$ such that: $v(p_1, q_1) \leq v(x_0, y_0)$ and $v(p_2, q_1) \geq v(p_1, y_0)$.*

We now prove the proposition. Define, for x in X ,

$$h(x) = \text{Max}\{v(p, y) - v(x, y), y \in Y, p \in \Gamma(x)\}.$$

h is continuous. We put $Z = \text{Argmax}_{x \in X} h(x)$, and consider $x_0 \in \text{Argmin}_{x \in Z} \varphi(x)$, where φ comes from the definition of Γ weakly acyclic.

$x_0 \in Z$, so v is excessive if and only if $h_0 =_{def} h(x_0) \leq 0$. By definition of $h(x_0)$, there exists $p_1 \in \Gamma(x_0)$ and y_0 in Y such that $v(p_1, y_0) - v(x_0, y_0) = h_0$.

By lemma 7.2, there exists q_1 in $\Lambda(y_0)$ and $p_2 \in \tilde{\Gamma}(p_1)$ such that $v(p_1, q_1) \leq v(x_0, y_0)$ and $v(p_2, q_1) \geq v(p_1, y_0)$. Consequently,

$$v(p_2, q_1) - v(p_1, q_1) \geq v(p_1, y_0) - v(x_0, y_0) = h_0.$$

One can now find y_1 in Y such that $v(p_2, y_1) - v(p_1, y_1) \geq h_0$, and since $p_2 \in \tilde{\Gamma}(p_1)$ it implies that $\text{Supp}(p_1) \subset Z$. But $p_1 \in \Gamma(x_0)$, so by definition of x_0 and weak acyclicity, we obtain that $p_1 = \delta_{x_0}$. So $h_0 = 0$, and v is excessive.

Let us now prove that when v is balanced and excessive then it is depressive. For every (x, y) , and every $p \in \Gamma(x)$, we have $v(x, y) \geq v(p, y)$. Thus, for every (x, y) , $p \in \Gamma(x)$ and every $q \in \Lambda(y)$, $v(x, q) \geq v(p, q)$ and consequently, $\min_{q \in \Lambda(y)} v(x, q) \geq \min_{q \in \Lambda(y)} v(p, q)$. Taking the maximum in $p \in \Gamma(x)$ and using that v is balanced implies that $\min_{q \in \Lambda(y)} v(x, q) \geq v(x, y)$. Since Λ is leavable, we have equality and so v is depressive with respect to Y .

7.2.2 Proof of proposition 3.3

Let $(\lambda_n)_n$ be a vanishing sequence of discount factors such that $\|v_{\lambda_n} - v\| \rightarrow_{n \rightarrow \infty} 0$. We show first that v is balanced. From the Shapley equation, for every (x, y) in $X \times Y$ one has:

$$\begin{aligned} v_{\lambda_n}(x, y) &= \max_{p \in \Gamma(x)} \min_{q \in \Lambda(y)} (\lambda_n u(x, y) + (1 - \lambda_n) \tilde{v}_{\lambda_n}(p, q)), \\ &= \lambda_n u(x, y) + (1 - \lambda_n) \max_{p \in \Gamma(x)} \min_{q \in \Lambda(y)} (\tilde{v}_{\lambda_n}(p, q)). \end{aligned}$$

where max and min above commute (the value exists). Going to the limit and using the fact that the operator $(f \mapsto \max_{p \in \Gamma(x)} \min_{q \in \Lambda(y)} (\tilde{f}(p, q)))$ on $\mathcal{C}(X, Y)$ is non-expansive and so continuous for the uniform topology, imply that v is balanced.

Fix (x, y) in $X \times Y$, by symmetry it is enough to show that there exists $p \in \Gamma^\infty(x)$ such that $v(x, y) \leq v(p, y) \leq u(p, y)$. If $v(x, y) \leq u(x, y)$, it is enough to consider $p = \delta_x$, so we assume $v(x, y) > u(x, y)$. For n large enough, $v_{\lambda_n}(x, y) > u(x, y) + \lambda_n$.

Fix n . We define inductively a sequence $(p_t^n)_{t=0, \dots, T_n}$ in $\Delta(X)$, with $T_n \geq 1$, by:

- 1) $p_0^n = \delta_x$,
- 2) for each $t \geq 0$ such that $v_{\lambda_n}(p_t^n, y) > u(p_t^n, y) + \lambda_n$, we define p_{t+1}^n in $\tilde{\Gamma}(p_t^n)$ by:

$$p_{t+1}^n \in \text{Argmax}_{p \in \tilde{\Gamma}(p_t^n)} (\lambda_n u(p_t^n, y) + (1 - \lambda_n) v_{\lambda_n}(p, y)).$$

We have $\max_{p \in \tilde{\Gamma}(p_t^n)} (\lambda_n u(p_t^n, y) + (1 - \lambda_n) v_{\lambda_n}(p, y)) \geq v_{\lambda_n}(p_t^n, y)$, so:

$$\lambda_n u(p_t^n, y) + (1 - \lambda_n) v_{\lambda_n}(p_{t+1}^n, y) \geq v_{\lambda_n}(p_t^n, y). \quad (1)$$

Since $u(p_t^n, y) < v_{\lambda_n}(p_t^n, y) - \lambda_n$, we obtain:

$$v_{\lambda_n}(p_{t+1}^n, y) \geq v_{\lambda_n}(p_t^n, y) + \frac{\lambda_n^2}{1 - \lambda_n} > v_{\lambda_n}(p_t^n, y). \quad (2)$$

Since $\frac{\lambda_n^2}{1 - \lambda_n} > 0$ and v_{λ_n} is bounded, there exists a first integer $t = T_n$ where $v_{\lambda_n}(p_{T_n}^n, y) \leq u(p_{T_n}^n, y) + \lambda_n$, and we stop here the definition of the sequence $(p_t^n)_{t=0, \dots, T_n}$. Inequalities (1) and (2) above give:

$$v_{\lambda_n}(x, y) \leq v_{\lambda_n}(p_{T_n-1}^n, y) \leq \lambda_n u(p_{T_n-1}^n, y) + (1 - \lambda_n) v_{\lambda_n}(p_{T_n}^n, y). \quad (3)$$

Define now $p^n = p_{T_n}^n$ for each n . $p^n \in \tilde{\Gamma}^{T_n}(x) \subset \Gamma^\infty(x)$ for each n , and we consider a limit point $p^* \in \Gamma^\infty(x)$ of $(p^n)_n$. Because v_{λ_n} is an equicontinuous family converging to v , we obtain the convergence (along a subsequence) of $v_{\lambda_n}(p_{T_n}^n, y)$ to $v(p^*, y)$. Passing to the limit in the inequality defining T_n then gives:

$$v(p^*, y) \leq u(p^*, y).$$

Finally, passing to the limit in (3) shows: $v(x, y) \leq v(p^*, y)$.

7.2.3 Proof of proposition 3.4

We start with a lemma.

Lemma 7.3. *Assume v in $\mathcal{C}(X \times Y)$ is excessive. Then for all (x_0, y_0) in $X \times Y$ and all $p \in \Gamma^\infty(x_0)$, we have: $v(p, y_0) \leq v(x_0, y_0)$.*

Proof: $p = \lim_n p_n$, with $p_n \in \tilde{\Gamma}^n(x_0)$ for each n . It is enough to prove that $v(p_n, y_0) \leq v(x_0, y_0)$ for each n , and we do the proof by induction on n . The case $n = 1$ is clear by definition of v excessive. Since $p_{n+1} \in \tilde{\Gamma}(p_n)$, it is enough to prove that for p' in $\Delta(X)$ and $p'' \in \tilde{\Gamma}(p')$, we have $v(p'', y_0) \leq v(p', y_0)$. By definition of $\tilde{\Gamma}$, (p', p'') is in the closure of $\text{conv}(\text{Graph}\Gamma)$.

y_0 is fixed, and the function $h : p \rightarrow v(p, y_0)$ is affine continuous on $\Delta(X)$. The set

$$D =_{\text{def}} \{(p', p'') \in \Delta(X) \times \Delta(X), h(p'') \leq h(p')\}.$$

is convex and compact, and we want to show that $\text{Graph}(\tilde{\Gamma}) = \overline{\text{conv}}(\text{Graph}\Gamma) \subset D$. It is enough to prove that $\text{Graph}(\Gamma) \subset D$, and this is implied by the fact that v is excessive. This concludes the proof of lemma 7.3.

We now prove the proposition. Assume one of the gambling houses is strongly acyclic, and let v_1 and v_2 satisfying the conditions of proposition 3.4 (are continuous, balanced, v_1 satisfies $P1$ and v_2 satisfies $P2$). We will show that $v_1 \leq v_2$.

By symmetry, suppose that Γ is strongly acyclic. From Proposition 3.2, v_1 and v_2 are excessive (in X) and depressive (in Y). $v_1 - v_2$ being continuous on $X \times Y$, define the compact set:

$$Z = \text{Argmax}_{(x,y) \in X \times Y} v_1(x, y) - v_2(x, y).$$

Consider now φ u.s.c. given by the strong acyclicity condition of Γ . The set Z being compact, there exists (x_0, y_0) minimizing $\varphi(x)$ for (x, y) in Z .

By v_2 satisfying $P2$, there exists q in $\Lambda^\infty(y_0)$ such that $v_2(x_0, y_0) = v_2(x_0, q) \geq u(x_0, q)$. Thus, there is $y'_0 \in \text{Supp}(q)$ such that $v_2(x_0, y'_0) \geq u(x_0, y'_0)$.

Because v_1 is depressive, by lemma 7.3 we have $v_1(x_0, q) \geq v_1(x_0, y_0)$ and we obtain $v_1(x_0, q) - v_2(x_0, q) \geq v_1(x_0, y_0) - v_2(x_0, y_0)$. Since (x_0, y_0) is in Z , $\{x_0\} \times \text{Supp}(q) \subset Z$. Thus, $(x_0, y'_0) \in Z$. Obviously, (x_0, y'_0) also minimizes $\varphi(x)$ for (x, y) in Z (the minimum value remains unchanged: $\varphi(x_0)$).

By v_1 satisfying $P1$, there exists p in $\Gamma^\infty(x_0)$ such that: $v_1(x_0, y'_0) = v_1(p, y'_0) \leq u(p, y'_0)$. Because v_2 is excessive, by lemma 7.3 we have $v_2(p, y'_0) \leq v_2(x_0, y'_0)$ and we obtain $v_1(p, y'_0) - v_2(p, y'_0) \geq v_1(x_0, y'_0) - v_2(x_0, y'_0)$. Since (x_0, y'_0) is in Z , $\text{Supp}(p) \times \{y'_0\} \subset Z$. By definition of (x_0, y'_0) , this implies: $\varphi(p) \geq \varphi(x_0)$. The definition of φ now gives that p is the Dirac measure on x_0 . We obtain that $v_1(x_0, y'_0) = v_1(p, y'_0) \leq u(x_0, y'_0)$. So $\text{Max}_{(x,y) \in X \times Y} v_1(x, y) - v_2(x, y) = v_1(x_0, y'_0) - v_2(x_0, y'_0) \leq u(x_0, y'_0) - u(x_0, y'_0) = 0$, and thus $v_1 \leq v_2$.

7.3 Proof of theorem 4.2

We start with considerations valid for the 3 cases of the theorem. We fix $J = [0, 1/4]$ in all the proof, and only assume for the moment that I is a compact subset of $[0, 1/4]$ containing 0 and $1/4$.

Consider $\lambda \in (0, 1)$. It is clear that $v_\lambda(c, c') = 0$, and $v_\lambda(a, c') = v_\lambda(b, c') = 1$. By symmetry of the payoffs and transitions, we have $v_\lambda(a, a') = v_\lambda(b, b')$, $v_\lambda(b, a') = v_\lambda(a, b')$ and $v_\lambda(c, c') = v_\lambda(c, b')$, so we can write v_λ as:

	a'	b'	c'
a	x_λ	y_λ	1
b	y_λ	x_λ	1
c	z_λ	z_λ	0

with x_λ, y_λ and z_λ in $(0, 1)$.

z_λ is indeed easy to compute. If the game is at (c, a') , Player 1 can not move, and Player 2 wants to reach c' as fast as possible, so he will choose $\beta = 1/4$ and we have (see definition 2.12): $z_\lambda = \lambda 1 + (1 - \lambda)(\frac{1}{16} 0 + \frac{15}{16} z_\lambda)$, so that:

$$z_\lambda = \frac{16\lambda}{1 + 15\lambda} \leq 16\lambda. \quad (4)$$

Proposition 7.4. *Assume $J = [0, 1/4]$, $\min I = 0$ and $\max I = 1/4$. Then for λ small enough,*

$$z_\lambda < x_\lambda < y_\lambda, \quad (5)$$

$$\lambda x_\lambda = (1 - \lambda) \max_{\alpha \in I} (\alpha(y_\lambda - x_\lambda) + \alpha^2(z_\lambda - x_\lambda)), \quad (6)$$

$$\lambda y_\lambda = \lambda + (1 - \lambda) \min_{\beta \in J} (\beta(x_\lambda - y_\lambda) + \beta^2(1 - y_\lambda)). \quad (7)$$

(6) expresses the fact that at (a, a') or (b, b') , it is optimal for Player 2 to play the pure strategy $\beta = 0$ (stay at the same location and wait until Player 1 has moved), and Player 1 can play a pure strategy α there. Similarly, (7) expresses the fact that at (a, b') or (b, a') , it is optimal for Player 1 not to move. In spite of these simple intuitions, the proof of the proposition is rather technical, and is proved separately below.

Taking for granted proposition 7.4, we now proceed to the proof of theorem 4.2. It is simple to study the maximization problem of Player 2 given by (7), which is simply minimizing a concave polynomial on the interval $J = [0, 1/4]$.

If $y_\lambda - x_\lambda \leq \frac{1}{2}(1 - y_\lambda)$, the minimum in (7) is achieved for $\beta = \frac{y_\lambda - x_\lambda}{2(1 - y_\lambda)}$, and otherwise it is achieved for $\beta = \frac{1}{4}$. Hence for λ small enough:

$$\text{If } y_\lambda - x_\lambda \leq \frac{1}{2}(1 - y_\lambda), \quad 4\lambda(1 - y_\lambda)^2 = (1 - \lambda)(y_\lambda - x_\lambda)^2. \quad (8)$$

$$\text{If } y_\lambda - x_\lambda \geq \frac{1}{2}(1 - y_\lambda), \quad (1 - y_\lambda)(1 + 15\lambda) = 4(1 - \lambda)(y_\lambda - x_\lambda). \quad (9)$$

Notice that the two (in)equalities of (9) imply $1 + 15\lambda \geq 2(1 - \lambda)$, which is not possible for λ small. Consequently, for small discount factors (8) holds, and we have proved, for λ small enough, the main equality of the proof:

$$4\lambda(1 - y_\lambda)^2 = (1 - \lambda)(y_\lambda - x_\lambda)^2. \quad (10)$$

This clearly implies:

$$y_\lambda - x_\lambda \xrightarrow{\lambda \rightarrow 0} 0.$$

For each $\lambda > 0$, denote by $\alpha_\lambda \in I$ a maximizer in the expression (6), so that

$$\lambda x_\lambda = (1 - \lambda)\alpha_\lambda(y_\lambda - x_\lambda) + (1 - \lambda)\alpha_\lambda^2(z_\lambda - x_\lambda). \quad (11)$$

And since $x_\lambda > 0$, $\alpha_\lambda > 0$.

The following lemma implies part 2) of theorem 4.2.

Lemma 7.5. *If 0 is an isolated point in I, then y_λ and x_λ converge to 0.*

Proof: In this case there exists $\alpha^* > 0$ such that $\alpha_\lambda \geq \alpha^*$ for all λ . Passing to the limit in (11) gives the result.

We will now prove parts 1) and 3) of the theorem. The fact that $I \subset J$ gives an advantage to Player 2, which can be quantified as follows.

Lemma 7.6. *Assume that y_{λ_n} and x_{λ_n} converge to v in $[0, 1]$. Then $v \leq 1/2$, and $y_{\lambda_n} - x_{\lambda_n} \sim 2\sqrt{\lambda_n}(1 - v)$.*

Proof: $(1 - \lambda)(y_\lambda - x_\lambda) = 2\sqrt{\lambda}\sqrt{1 - \lambda}(1 - y_\lambda)$, so

$$x_\lambda(\lambda + \alpha_\lambda^2) = \lambda x_\lambda \alpha_\lambda^2 + (1 - \lambda)\alpha_\lambda^2 z_\lambda + 2\alpha_\lambda \sqrt{\lambda}\sqrt{1 - \lambda}(1 - y_\lambda) \geq 2\alpha_\lambda \sqrt{\lambda} x_\lambda,$$

since $\lambda + \alpha_\lambda^2 - 2\alpha_\lambda \sqrt{\lambda} \geq 0$. Dividing by $\alpha_\lambda \sqrt{\lambda}$ and passing to the limit gives $2(1 - v) \geq 2v$, so $v \leq 1/2$, and the lemma is proved.

Consider again the concave optimization problem of Player 1 given by equation (6), and denote by $\alpha^*(\lambda) = \frac{y_\lambda - x_\lambda}{2(x_\lambda - z_\lambda)} > 0$ the argmax of the unconstrained problem if Player 1 could choose any $\alpha \geq 0$. If y_λ and x_λ converge to $v > 0$, then $\alpha^*(\lambda) \sim \sqrt{\lambda} \frac{1 - v}{v}$, and Player 1 would like to play in the λ -discounted game at (a, a') some α close to $\sqrt{\lambda} \frac{1 - v}{v}$.

Lemma 7.7. *Let λ_n be a vanishing sequence of discount factors such that $\sqrt{\lambda_n} \in I$ for each n . Then y_{λ_n} and x_{λ_n} converge to $1/2$.*

Proof: By considering a converging subsequence we can assume that y_{λ_n} and x_{λ_n} converge to some v in $[0, 1]$. By the previous lemma, $v \leq 1/2$, and we have to show that $v \geq 1/2$. We have for each λ in the subsequence, since Player 1 can choose to play $\alpha = \sqrt{\lambda}$:

$$\lambda x_\lambda \geq (1 - \lambda)\sqrt{\lambda}(y_\lambda - x_\lambda) + (1 - \lambda)\lambda(z_\lambda - x_\lambda),$$

so

$$x_\lambda(2 - \lambda) \geq (1 - \lambda)z_\lambda + (1 - \lambda)\frac{y_\lambda - x_\lambda}{\sqrt{\lambda}}.$$

By passing to the limit, we get $2v \geq 2(1 - v)$, and $v \geq 1/2$.

Lemma 7.8. *Let λ_n be a vanishing sequence of discount factors such that for each n , the open interval $(\frac{1}{2}\sqrt{\lambda_n}, 2\sqrt{\lambda_n})$ does not intersect I . Then $\limsup_n y_{\lambda_n} \leq 4/9$.*

Proof: Suppose that (up to a subsequence) x_{λ_n} and y_{λ_n} converges to some $v \geq 4/9$. It is enough to show that $v = 4/9$. We know that $v \leq 1/2$ by lemma 7.6, and since $\alpha^*(\lambda) \sim \sqrt{\lambda} \frac{1 - v}{v}$ we have $\frac{1}{2}\sqrt{\lambda} \leq \alpha^*(\lambda) \leq 2\sqrt{\lambda}$ for λ small in the sequence. By assumption $(\frac{1}{2}\sqrt{\lambda}, 2\sqrt{\lambda})$ contains no point in I and the objective function of Player 1 is increasing from 0 to $\alpha^*(\lambda)$ and decreasing after $\alpha^*(\lambda)$. There are 2 possible cases:

If $\alpha_\lambda \leq \frac{1}{2}\sqrt{\lambda}$ we have:

$$\lambda x_\lambda \leq \frac{1}{2}(1 - \lambda)\sqrt{\lambda}(y_\lambda - x_\lambda) + \frac{1}{4}(1 - \lambda)\lambda(z_\lambda - x_\lambda).$$

Dividing by λ and passing to the limit gives: $v \leq 1 - v - \frac{1}{4}v$, i.e. $v \leq \frac{4}{9}$.

Otherwise, $\alpha_\lambda > 2\sqrt{\lambda}$ and we have:

$$\lambda x_\lambda \leq 2(1 - \lambda)\sqrt{\lambda}(y_\lambda - x_\lambda) + 4(1 - \lambda)\lambda(z_\lambda - x_\lambda).$$

Again, dividing by λ and passing to the limit gives: $v \leq 4(1 - v) - 4v$, i.e. $v \leq \frac{4}{9}$.

Finally, lemma 7.7 proves part 1) of theorem 4.2, whereas lemmas 7.7 and 7.8 together imply part 3), concluding the proof of theorem 4.2.

Remarks 7.9.

- In case 3) of theorem 4.2, it is possible to show, using lemma 7.8, that $\liminf x_\lambda = \liminf y_\lambda = 4/9$, and $\limsup x_\lambda = \limsup y_\lambda = 1/2$.

- It is not difficult to adapt lemma 7.8 to show the divergence of (v_λ) as soon as $J = [0, 1/4]$ and I satisfies:

- a) there exists a sequence (λ_n) converging to 0 such that $\sqrt{\lambda_n} \in I$ for each n , and

- b) there exist $\eta > 0$ and a sequence (λ_n) converging to 0 such that for each n , I does not intersect the interval $[\sqrt{\lambda_n}(1 - \eta), \sqrt{\lambda_n}(1 + \eta)]$.

- It is important for the counterexample that $I = \{\frac{1}{2^{2^n}}, n \in \mathbb{N}^*\} \cup \{0\}$ is not semi-algebraic. Indeed, it has been showed that if we assume X and Y finite, and the transitions Γ , Λ and the payoff u to be definable in some o-minimal structure, then $(v_\lambda)_\lambda$ converges [Bolte et al., 2015].

7.4 Proof of proposition 7.4

We proceed in 4 steps.

1. The game at (b, a') : It is intuitively clear that $y_\lambda \geq x_\lambda$ since Player 1 is better off when the players have different locations. We now formalize this idea. Consider the game at (b, a') . The current payoff is 1, and Player 1 has the option not to move, so we obtain by definition 2.12:

$$y_\lambda \geq \lambda + (1 - \lambda) \min_{\beta \in J} (\beta x_\lambda + (1 - \beta - \beta^2)y_\lambda + \beta^2 1).$$

This implies

$$\lambda y_\lambda \geq \lambda + (1 - \lambda) \min_{\beta \in J} (\beta(x_\lambda - y_\lambda) + \beta^2(1 - y_\lambda)),$$

and since $y_\lambda \leq 1$, we obtain: $x_\lambda \leq y_\lambda$. Now, $\min_{\beta \in J} (\beta(x_\lambda - y_\lambda) + \beta^2(1 - y_\lambda)) \geq \min_{\beta \in J} \beta(x_\lambda - y_\lambda) + \min_{\beta \in J} \beta^2(1 - y_\lambda) = 1/4(x_\lambda - y_\lambda)$, hence:

$$(1 - \lambda)(y_\lambda - x_\lambda) \geq 4\lambda(1 - y_\lambda). \tag{12}$$

2. The game at (a, a') :

In the same spirit, in the game at (a, a') , Player 2 has the option not to move, so we have:

$$x_\lambda \leq (1 - \lambda) \max_{\alpha \in I} (\alpha y_\lambda + (1 - \alpha - \alpha^2)x_\lambda + \alpha^2 z_\lambda).$$

Hence,

$$\lambda x_\lambda \leq (1 - \lambda) \max_{\alpha \in I} (\alpha(y_\lambda - x_\lambda) + \alpha^2(z_\lambda - x_\lambda)). \tag{13}$$

Also, by definition 2.12, x_λ is the value of the game (possibly played with mixed strategies), where Player 1 chooses α in I , Player 2 chooses β in J and the payoff to Player 1 is: $(1 - \lambda)g_\lambda(\alpha, \beta)$, where $g_\lambda(\alpha, \beta) =$

$$x_\lambda((1-\alpha-\alpha^2)(1-\beta-\beta^2)+\alpha\beta)+y_\lambda(\beta(1-\alpha-\alpha^2)+\alpha(1-\beta-\beta^2))+\beta^2(1-\alpha^2)1+\alpha^2(1-\beta^2)z_\lambda.$$

We want to prove that in this game, it is a dominant strategy for Player 2 not to move, that is to choose $\beta = 0$. We need to show that for all α and β , $g_\lambda(\alpha, 0) \leq g_\lambda(\alpha, \beta)$. As a function of β , $g_\lambda(\alpha, \beta)$ can be written as a constant plus:

$$\beta(1 - 2\alpha - \alpha^2)(y_\lambda - x_\lambda) + \beta^2(-(1 - \alpha - \alpha^2)x_\lambda - \alpha y_\lambda + 1 - \alpha^2 - \alpha^2 z_\lambda).$$

So we want to show that for all α in I , β in J :

$$(1 - 2\alpha - \alpha^2 - \alpha\beta)(y_\lambda - x_\lambda) + \beta((1 - \alpha^2)(1 - x_\lambda) - \alpha^2 z_\lambda) \geq 0$$

Since the expression is decreasing in α , it is enough to prove it with $\alpha = 1/4$:

$$(7 - 4\beta)(y_\lambda - x_\lambda) + \beta(15(1 - x_\lambda) - z_\lambda) \geq 0.$$

This is true for $\beta = 0$, and will be true for all β in J if and only if it is true for $\beta = 1/4$, so we are left with proving:

$$24(y_\lambda - x_\lambda) + 15(1 - x_\lambda) - z_\lambda \geq 0. \tag{14}$$

Consider $\lambda \leq 1/32$, and recall that $z_\lambda \leq 16\lambda$. If $x_\lambda \leq 1/2$, then clearly (14) holds. Assume on the contrary that $x_\lambda \geq 1/2$, then $z_\lambda \leq x_\lambda$, and (13) gives: $\lambda x_\lambda \leq (1 - \lambda)\frac{1}{4}(y_\lambda - x_\lambda) + 0$, so $(y_\lambda - x_\lambda) \geq 2\lambda$ and (14) holds as well.

We have shown that in the λ -discounted game at (a, a') with $\lambda \leq 1/32$, Player 2 has a pure dominant strategy which is $\beta = 0$. Considering a pure best reply of Player 1 against this strategy implies that the game at (a, a') has a value in pure strategies satisfying $x_\lambda = (1 - \lambda) \max_{\alpha \in I} (\alpha y_\lambda + (1 - \alpha - \alpha^2)x_\lambda + \alpha^2 z_\lambda)$, i.e. equation (6) is proved.

3. Small discount factors: For the sake of contradiction, assume that $z_{\lambda_n} \geq x_{\lambda_n}$ for a vanishing sequence λ_n of discount factors. Then by equation (6), we have for each n :

$$\lambda_n x_{\lambda_n} = (1 - \lambda_n) \left(\frac{1}{4}(y_{\lambda_n} - x_{\lambda_n}) + \frac{1}{16}(z_{\lambda_n} - x_{\lambda_n}) \right).$$

Since z_{λ_n} converges to 0, so does x_{λ_n} and y_{λ_n} , and moreover $\frac{y_{\lambda_n} - x_{\lambda_n}}{\lambda_n}$ converges to 0. This is in contradiction with equation (12). We have shown (5).

4. The game at (b, a') again: We proceed as for the game at (a, a') and will show that in the game at (b, a') , it is a dominant strategy for Player 1 not to move. By definition, y_λ is the value of the game where Player 1 chooses α in I , Player 2 chooses β in J and the payoff is $\lambda + (1 - \lambda)h_\lambda(\alpha, \beta)$, with $h_\lambda(\alpha, \beta) =$

$$y_\lambda((1-\alpha-\alpha^2)(1-\beta-\beta^2)+\alpha\beta)+x_\lambda(\beta(1-\alpha-\alpha^2)+\alpha(1-\beta-\beta^2))+\beta^2(1-\alpha^2)1+\alpha^2(1-\beta^2)z_\lambda.$$

We want to show that $h_\lambda(0, \beta) \geq h_\lambda(\alpha, \beta)$ for all α and β . That is, for all α and β ,

$$(x_\lambda - y_\lambda)(1 - 2\beta - \alpha\beta - \beta^2) + \alpha((1 - \beta^2)(z_\lambda - y_\lambda) - \beta^2) \leq 0.$$

For λ small enough, we have $z_\lambda \leq x_\lambda \leq y_\lambda$, and the above property is satisfied. Hence in the game at (b, a') it is dominant for Player 1 to choose $\alpha = 0$. Consequently, Player 2 has a pure optimal strategy and we can write:

$$y_\lambda = \lambda + (1 - \lambda) \min_{\beta \in J} (\beta x_\lambda + (1 - \beta - \beta^2)y_\lambda + \beta^2),$$

proving equation (7). And the proof of proposition 7.4 is complete.

8 Appendix B: uniform analysis

To study the uniform value, we restrict the analysis to idempotent games. The extension to the general case is open.

8.1 Definition of idempotent gambling games

Definition 8.1. *The gambling game is idempotent if $\Gamma \circ \Gamma = \Gamma$ and $\Lambda \circ \Lambda = \Lambda$.*

In that case, clearly $\Gamma = \Gamma^\infty$ and $\Lambda = \Lambda^\infty$, and vice versa, if $\Gamma = \Gamma^\infty$ then clearly $\Gamma \circ \Gamma = \Gamma$. Any state that could be reached in several stages can be reached immediately in a single stage. This holds true for instance in splitting games. Notice also that for any Γ , the multifunction Γ^∞ is idempotent. If $\Gamma \circ \Gamma = \Gamma$, then $\Gamma(x) = \tilde{\Gamma}^n(\delta_x) = \Gamma^\infty(x)$ for all n and x so if the gambling game is idempotent the notions of weak and strong acyclicity coincide.

An immediate corollary of theorem 3.1 is the following.

Corollary 8.2. *Consider a standard idempotent gambling game where a player has an acyclic gambling house. Then $\{v_\lambda\}$ converges uniformly to the unique function v in $\mathcal{C}(X \times Y)$ which is excessive, depressive and satisfies:*

$$Q1 : \forall (x, y) \in X \times Y, \exists p \in \Gamma(x), v(x, y) = v(p, y) \leq u(p, y),$$

$$Q2 : \forall (x, y) \in X \times Y, \exists q \in \Lambda(y), v(x, y) = v(x, q) \geq u(x, q).$$

8.2 Definition of uniform value and optimal strategies

In repeated and stochastic games, a stronger notion of limit value is given by the uniform value. As usual, a strategy of Player 1, resp. Player 2, is a measurable rule giving at every stage t , as a function of past and current states, an element in $\Gamma(x_t)$,

resp. of $\Lambda(y_t)$, where x_t and y_t are the states of stage t . A pair (x_1, y_1) of initial states and a pair of strategies (σ, τ) naturally define a probability on the set of plays $(X \times Y)^\infty$ (with the product σ -algebra, X and Y being endowed with their Borel σ -algebra), which expectation is written $\mathbb{E}_{(x_1, y_1), \sigma, \tau}$.

Definition 8.3. $w \in \mathcal{B}(X \times Y)$ is the uniform value of the gambling game and both players have optimal uniform strategies if:

There exists a strategy σ of Player 1 that uniformly guarantees w : for any $\varepsilon > 0$, there is N such that for any $n \geq N$ and initial states (x_1, y_1) , for any strategy τ of Player 2, $\mathbb{E}_{(x_1, y_1), \sigma, \tau} \left(\frac{1}{n} \sum_{t=1}^n u(x_t, y_t) \right) \geq w(x_1, y_1) - \varepsilon$.

And similarly, there exists a strategy τ of Player 2 that uniformly guarantees w : for any $\varepsilon > 0$, there is N such that for any $n \geq N$ and initial states (x_1, y_1) , for any strategy σ of Player 1, $\mathbb{E}_{(x_1, y_1), \sigma, \tau} \left(\frac{1}{n} \sum_{t=1}^n u(x_t, y_t) \right) \leq w(x_1, y_1) + \varepsilon$.

It is known [Sorin, 2002] that the above conditions imply similar inequalities for discounted payoffs: for Player 1, the same strategy σ is such that for any $\varepsilon > 0$, there is $\lambda_0 > 0$ satisfying: for any $\lambda \leq \lambda_0$, (x_1, y_1) and τ ,

$$\mathbb{E}_{(x_1, y_1), \sigma, \tau} \left(\lambda \sum_{t=1}^{\infty} (1 - \lambda)^{t-1} u(x_t, y_t) \right) \geq w(x_1, y_1) - \varepsilon.$$

And if the uniform value exists, it has to be $v = \lim_{\lambda} v_{\lambda} = \lim_n v_n$.

8.3 Definition of adapted strategies

Our main theorem 3.1 suggests particularly interesting strategies for the players. Consider again conditions $P1$ and $P2$, and fix a pair of states (x, y) . If $u(x, y) \geq v(x, y)$, the running payoff of Player 1 is at least as good as the payoff he should expect in the long run, so we may consider that Player 1 is “quite happy” with the current situation and in order to satisfy $P1$ it is enough for him not to move, i.e. to choose $p = \delta_x$. If on the contrary $u(x, y) < v(x, y)$, Player 2 is happy with the current situation and can choose $q = \delta_y$ to satisfy $P2$, whereas Player 1 should do something, and a possibility is to move towards a p satisfying $P1$. This looks interesting for Player 1 because if Player 2 does not react, eventually the distribution on states will approach (p, y) and (in expectation) Player 1 will be happy again with the current situation since $u(p, y) \geq v(p, y)$.

Definition 8.4. Let w in $\mathcal{B}(X \times Y)$.

A strategy of Player 1 is adapted to w if whenever the current state is (x, y) , it plays $p \in \Gamma(x)$ such that $w(x, y) \leq w(p, y) \leq u(p, y)$.

A strategy of Player 2 is adapted to w if whenever the current state is (x, y) , it plays $q \in \Gamma(y)$ such that $w(x, y) \geq w(x, q) \geq u(x, q)$.

If w satisfies $Q1$, resp. $Q2$, Player 1, resp. Player 2, has a strategy adapted to w (using a measurable selection theorem [Aliprantis and Border, 2006]). If moreover w is excessive, we have $w(x, y) = w(p, y) \leq u(p, y)$. Mertens Zamir [Mertens and Zamir, 1971] in repeated games with incomplete information and Oliu-Barton [Oliu-Barton, 2018] in splitting games used similar strategies derived from the MZ -characterization instead of the Q -characterization.

8.4 Bounded variation and existence of uniform adapted strategy

In repeated games with incomplete information or in splitting games, an important property is that any martingale on a simplex has bounded variations. This suggests the following.

Definition 8.5. *A gambling house Γ has vanishing L_1 -variation if for every $\varepsilon > 0$, there is N such that for all $n \geq N$ and any sequence (p_t) s.t. $p_{t+1} \in \tilde{\Gamma}(p_t)$, one has $\frac{1}{n} \sum_{t=1}^n d_{KR}(p_{t+1}, p_t) \leq \varepsilon$.*

The proof of the next result is inspired by Oliu-Barton [Oliu-Barton, 2018] in the framework of splitting games. He shows that Player 1 (resp. Player 2) can uniformly guarantee any excessive-depressive function satisfying $MZ1$ (resp. $MZ2$). Our proof is much shorter because it uses the new Q -characterization.

Proposition 8.6. *In a standard gambling game where Γ has vanishing L_1 -variation, if w in $\mathcal{B}(X \times Y)$ is excessive-depressive and satisfies $Q1$, then a strategy of Player 1 adapted to w uniformly guarantees w .*

Proof: Fix $\varepsilon > 0$. Because Lipschitz functions are dense in the set of continuous functions, there exists $K > 0$ and a K -Lipschitz function u_ε that is uniformly ε -close to u . Consider w in $\mathcal{B}(X \times Y)$ be an excessive-depressive function satisfying $Q1$, let σ be a strategy of Player 1 adapted to w and let τ be any strategy of Player 2. We fix the initial states and write $\mathbb{E} = \mathbb{E}_{(x_1, y_1), \sigma, \tau}$.

Then, the average payoff of the n -stage game is:

$$\begin{aligned} \frac{1}{n} \mathbb{E} \left(\sum_{t=1}^n u(x_t, y_t) \right) &= \frac{1}{n} \mathbb{E} \left(\sum_{t=1}^n u(x_t, y_t) - u(x_{t+1}, y_t) \right) + \frac{1}{n} \mathbb{E} \left(\sum_{t=1}^n u(x_{t+1}, y_t) \right) \\ &\geq \frac{1}{n} \mathbb{E} \left(\sum_{t=1}^n u_\varepsilon(x_t, y_t) - u_\varepsilon(x_{t+1}, y_t) \right) - 2\varepsilon + \frac{1}{n} \mathbb{E} \left(\sum_{t=1}^n u(x_{t+1}, y_t) \right) \\ &\geq -\frac{K}{n} \left(\sum_{t=1}^n d_{KR}(p_{t+1}, p_t) \right) - 2\varepsilon + \frac{1}{n} \mathbb{E} \left(\sum_{t=1}^n u(x_{t+1}, y_t) \right) \end{aligned}$$

Because the gambling game is of vanishing variation, there is N such that when $n \geq N$ one has $\frac{K}{n} \sum_{t=1}^n d_{KR}(p_{t+1}, p_t) \leq \varepsilon$. Because of $Q1$, $\mathbb{E}(u(x_{t+1}, y_t)) \geq \mathbb{E}(w(x_{t+1}, y_t)) =$

$\mathbb{E}(w(x_t, y_t))$. Since w is depressive, $\mathbb{E}(w(x_t, y_t)) \geq \mathbb{E}(w(x_t, y_{t-1}))$, so that $\mathbb{E}(w(x_{t+1}, y_t)) \geq \mathbb{E}(w(x_t, y_{t-1}))$, and this property holds for every t . Consequently, $\mathbb{E}(u(x_{t+1}, y_t)) \geq \mathbb{E}(w(x_2, y_1)) = w(x_1, y_1)$. We obtain finally $\frac{1}{n}\mathbb{E}(\sum_{t=1}^n u(x_t, y_t)) \geq w(x_1, y_1) - 3\varepsilon$, ending the proof.

Corollary 8.7. *If Γ and Λ have vanishing L_1 -variations then there is at most one excessive-depressive function in $\mathcal{B}(X \times Y)$ satisfying Q1 and Q2.*

Proof: If such a function exists, it can be guaranteed by both players. So it must be the uniform value of the game, which is unique whenever it exists.

Combining proposition 3.3, proposition 8.6 and corollary 8.7, we obtain the existence of the uniform value in a class of gambling games:

Theorem 8.8. *In a standard and idempotent gambling game where Γ and Λ have vanishing L_1 -variation, the uniform value v exists, and strategies adapted to v are uniformly optimal. Moreover, v is the unique excessive-depressive function in $\mathcal{B}(X \times Y)$ satisfying Q1 and Q2.*

Proof: From proposition 3.3, any accumulation point of v_λ satisfies P1 and P2, i.e. Q1 and Q2, so is unique and is the uniform value, as shown above.

Observe that acyclicity is not assumed in theorem 8.8. But vanishing L_1 -variation is a form of acyclicity (it rules out for example non-constant periodic orbits). A more formal link between acyclicity and vanishing variation is given in the next section.

8.5 Bounded L_2 -variation and acyclicity

Definition 8.9. *A gambling house Γ is of bounded L_2 -variation if there is $C > 0$ such that for every sequences $\{p_t\}$ satisfying $p_{t+1} \in \tilde{\Gamma}(p_t)$, one has*

$$\sum_{t=1}^{\infty} d_{KL}(p_{t+1}, p_t)^2 \leq C < +\infty$$

For example splitting games have bounded L_2 -variation.

Proposition 8.10. *If Γ is idempotent, non-expansive, leavable and of bounded L_2 -variation, then it is strongly acyclic and has vanishing L_1 -variation.*

Proof: Bounded L_2 -variation implies weak acyclicity because the real valued function:

$$\varphi(x) = \sup_{\{(p_t) \text{ s.t. } p_0=x \text{ and } p_{t+1} \in \tilde{\Gamma}(p_t)\}} \sum_{t=1}^{\infty} d_{KL}(p_{t+1}, p_t)^2,$$

is strictly decreasing along non-constant orbits of $\tilde{\Gamma}$ (i.e. $\arg \max_{p \in \Gamma(x)} \varphi(p) = \delta_x$). But for idempotent Γ , weak acyclicity and acyclicity coincide because $\Gamma^\infty = \Gamma$. Continuity of φ is a consequence of non-expansivity of Γ and the fact the bound C is uniform over the sequences $\{p_t\}$ satisfying $p_{t+1} \in \tilde{\Gamma}(p_t)$. Finally, that bounded L_2 -variation implies vanishing L_1 -variation is a consequence of Cauchy-Schwartz inequality ($\frac{1}{n} \sum_{t=1}^n d_{KL}(p_{t+1}, p_t) \leq \frac{1}{\sqrt{n}} \sqrt{\sum_{t=1}^n d_{KL}(p_{t+1}, p_t)^2}$).