

1 **Title: Extremely fast amelioration of plasmid fitness costs by multiple functionally-diverse**
2 **pathways**

3 **Authors:** James P. J. Hall^{1,2,3*} ORCID 0000-0002-4896-4592, Rosanna C. T. Wright² ORCID 0000-
4 0002-8095-8256, David Guymer³, Ellie Harrison² ORCID 0000-0002-2050-4631, Michael A. Brockhurst²
5 ORCID 0000-0003-0362-820X

6 **Affiliations:**

7 ¹Institute of Integrative Biology, University of Liverpool, Crown Street, Liverpool, L69 7ZB

8 ²Department of Animal and Plant Sciences, University of Sheffield, Western Bank, Sheffield, S10 2TN

9 ³Department of Biology, University of York, Wentworth Way, York, YO10 5DD

10 *Correspondence to: j.p.j.hall@liverpool.ac.uk, Institute of Integrative Biology, University of Liverpool,
11 Crown Street, Liverpool, L69 7ZB, United Kingdom. Tel: +44 (0)151 79 57691

12 **Author contributions:** Conceptualization: JPJH, EH, MAB. Investigation: JPJH, RCTW, DG. Formal
13 analysis: JPJH, RCTW, DG. Writing – original draft: JPJH. Writing – review and editing: JPJH, EH,
14 MAB.

15 **Data accessibility statement:** data supporting the results are in Supplementary Tables S1-S9, at
16 doi:10.17638/datacat.liverpool.ac.uk/953, and on the Short Read Archive, accession PRJEB32206.

17 **Conflicts of interest statement:** we have no conflicts of interest to declare.

18 **Keywords:** plasmids, horizontal gene transfer, compensatory evolution, amelioration, experimental
19 evolution, *Pseudomonas fluorescens*

20 **Word count:** 146 (abstract), 3441 (main text excluding references). **References:** 40

21 **Figures:** 2, plus 4 supplementary figures, 1 supplementary movie, and 9 supplementary tables.

22 **Abstract**

23 The acquisition of plasmids is often accompanied by fitness costs such that compensatory evolution is
24 required to allow plasmid survival, but it is unclear whether compensatory evolution can be extensive or
25 rapid enough to maintain plasmids when they are very costly. The mercury-resistance plasmid pQBR55
26 drastically reduced the growth of its host, *Pseudomonas fluorescens* SBW25, immediately after
27 acquisition, causing a small colony phenotype. However, within 48 hours of growth on agar plates we
28 observed restoration of the ancestral large colony morphology, suggesting that compensatory mutations
29 had occurred. Relative fitness of these evolved strains, in lab media and in soil microcosms, varied
30 between replicates, indicating different mutational mechanisms. Using genome sequencing we identified
31 that restoration was associated with chromosomal mutations in either a hypothetical DNA-binding protein
32 PFLU4242, RNA polymerase, or the GacA/S two-component system. Targeted deletions in *PFLU4242*,
33 *gacA*, or *gacS* recapitulated the ameliorated phenotype upon plasmid acquisition, indicating three distinct
34 mutational pathways to compensation. Our data shows that plasmid compensatory evolution is fast
35 enough to allow survival of a plasmid despite it imposing very high fitness costs upon its host, and indeed
36 may regularly occur during the process of isolating and selecting individual plasmid-containing clones.

37 **Introduction**

38 Plasmids are important vehicles for horizontal gene transfer (HGT), allowing bacteria to rapidly adapt to
39 new environments by transferring niche-adaptive traits (1). Plasmid acquisition can, however, disrupt
40 normal cellular function (2). The consequent fitness costs can limit plasmid survival because plasmid-
41 bearers will be outcompeted by plasmid-free cells that do not suffer the cost (2). Beneficial genes carried
42 by the plasmid cannot ensure its long-term persistence, as these genes can recombine onto the
43 chromosome (3,4). An important mechanism allowing plasmid survival is compensatory evolution
44 whereby mutations in chromosomal and/or plasmid genes ameliorate fitness costs (e.g. (5–7)). Where
45 transmission rates are too low to maintain plasmids by infectious transfer, plasmid survival effectively
46 becomes a race between the rate of compensatory evolution and the rate at which plasmid-bearers are
47 outcompeted (8).

48 Plasmid compensation is often explored using experimental evolution. In general, these experiments
49 involve the introduction of an initially costly plasmid to a strain, prolonged culture of plasmid bearers
50 (often through dozens of serial transfers), followed by assays on evolved plasmid-bearing strains and
51 often re-sequencing to identify underlying mutations. These studies have highlighted targets of loss-of-
52 function compensatory mutation, varying with the bacterial-plasmid pairing. Chromosomal accessory
53 helicases have been implicated with *Pseudomonas aeruginosa* and the small plasmid pNUK73 (7), and
54 with *Pseudomonas* sp. H2 and plasmid RP4 (6). Chromosomal global regulators have been identified: the
55 *fur* gene in *Shewanella oneidensis* MR-1 with pBP136 (9), and the *gacA/gacS* genes in *P. fluorescens*
56 SBW25 with pQBR103 (5). Compensatory mutations can also occur on plasmids, targeting replication
57 genes (10) or conjugation machinery (11–13). However, for very costly plasmids compensatory evolution
58 may be insufficient or too slow to enable persistence. For example, the mercury resistance plasmid
59 pQBR103 was lost from all populations of *P. aeruginosa* PAO1 even under mercury selection (3), and
60 plasmid pMS0506 was either lost or suffered large deletions when grown in *Acinetobacter baumannii*
61 ATCC19606 under selective (kanamycin) conditions (10).

62 In the current work, we show that several different routes of compensatory mutation, emerging during the
63 process of transconjugant colony growth, can overcome the heavy costs imposed when *P. fluorescens*
64 SBW25 newly acquires the conjugative plasmid pQBR55. pQBR55 is a member of the pQBR plasmid
65 collection, a set of relatively large (>130 kb) conjugative mercury resistance elements exogenously
66 isolated in the 1990s from Wytham Farm, Oxford. The sequenced pQBR plasmids, pQBR55, pQBR57,
67 and pQBR103, fall into different ‘groups’ based on RFLP fingerprint (14) (all of the sequenced pQBR
68 plasmids fall outwith the Enterobacteriaceae incompatibility typing scheme (15)) but DNA sequencing
69 indicated that pQBR57 and pQBR103 are distantly related to one another, and both are distantly related to
70 a family of IncP-2-related *Pseudomonas* megaplasmids (16,17). Besides mercury resistance, the
71 sequenced pQBR plasmids have few identifiable accessory traits, and no antibiotic resistance genes. Our
72 results indicate that compensatory evolution can rapidly and effectively facilitate the maintenance of
73 newly acquired mobile genetic elements.

74 **Materials and Methods**

75 **Strains and standard culture**

76 *Pseudomonas fluorescens* SBW25 was tagged with the mini-Tn7 streptomycin resistance (Sm^R) or
77 gentamicin resistance (Gm^R) cassette (5). For the 20 replicates used in the evolution experiment a *lacZ*-
78 expressing strain (18) was used, generating *P. fluorescens* SBW25 Sm^R -*lacZ* to aid identification of small
79 colonies by addition of X-gal (50 μ g/ml) to solid media. Strains were cultured in King’s B media at 28°C,
80 with addition of 1.2% w/v agar where appropriate. The pQBR55 donor strain, *P. putida*
81 UWC1(pQBR55), was a gift from Andy Lilley (KCL) via Andrew Spiers (University of Abertay).

82 **Establishing experimental lines**

83 Each replicate was established from an independent recipient colony. Overnight cultures of *P. fluorescens*
84 SBW25 Sm^R -*lacZ* and *P. putida* UWC1(pQBR55) were mixed in equal ratios and diluted 1:100 into 6 ml
85 KB broth which was incubated for 24 h at 180 rpm. Samples were spread on selective plates (250 μ g/ml

86 streptomycin, 20 mM HgCl₂, with 50 µg/ml X-gal) and colonies allowed to grow for 48 hours before
87 photographing (Figure S1). For each replicate, a colony was randomly selected as the one closest to a line
88 drawn down the centre of the plate, and was re-streaked twice onto selective media. A random colony was
89 again photographed, resuspended in KB + 20% w/v glycerol, and maintained at -80°C. Colonies were
90 tested for pQBR55 carriage by PCR using 1x GoTaq Green (Promega, WI USA) and 0.4 µM each of
91 primers pQBR55_0051_R1, pQBR55_0050_F1, merA_F, merA_R and thermocycling for 95°C 5', 30 x
92 (95°C 30", 58°C 30", 72°C 1'), 72°C 1' and in all cases were found to carry the plasmid. pQBR55_0050–
93 pQBR55_0051 adjoin the experimentally determined origin of pQBR55 replication, and thus their
94 presence is consistent with an independently-replicating plasmid (19). Primer sequences are in Table S3.
95 Glycerol stocks were used to establish subsequent experiments. Images were analysed with Imagemagick
96 (ImageMagick Studio LLC) and ImageJ (Rasband, W.S., ImageJ, U. S. National Institutes of Health,
97 Bethesda, Maryland, USA, 1997-2018). pQBR55 was conjugated into knockout strains from
98 UWC1(pQBR55) as described above, except selective plates included 30 µg/ml gentamicin and 20 mM
99 HgCl₂.

100 **Assays on experimental lines**

101 Phenotypic assays were performed largely according to Cheng et al. 2013 (20) on inocula from overnight
102 cultures. Production of extracellular protease was tested on nutrient agar supplemented with 5% w/v
103 powdered milk. Spots of 1:100 dilutions of culture (2.5 µl) were added and zones of clearing assessed
104 after 24 h incubation at room temperature. Viscosin production was assessed by drop collapse assay. A
105 sample (2 µl) of culture supernatant was spotted onto parafilm. The 'beading up' of the droplet indicated
106 a lack of viscosin. Flagellum-mediated motility was assessed by spotting cultures onto 'swim plates'
107 consisting of 5 g NaCl and 10 g tryptone per litre supplemented with 0.3% w/v agar, and assessing the
108 distance travelled after 24 h. Pectinase activity was assessed on slices of potato, which had been peeled,
109 soaked in 10% v/v bleach solution, and washed twice in deionised water (dH₂O). Approximately 8 mm
110 sections were placed in a petri dish and a depression cut into the surface, into which 50 µl culture was

111 pipetted. Potato slices were scored for soft rot after 4 days. Assays were repeated at least two times for
112 each evolved strain. All strains were tested alongside a *P. fluorescens* SBW25 wild-type positive control
113 and $\Delta gacA$ and $\Delta gacS$ negative controls.

114 Growth curves were carried out by subculturing overnight cultures and growing to OD600 ~ 0.4. Samples
115 of each culture were diluted 1:100 in KB and 5 μ l dilution used to inoculate 150 μ l KB in a 96-well
116 microtitre plate. Cultures were grown at 28°C, 180 rpm shaking in a Tecan M200 plate reader with
117 measurements taken every 15 minutes for 48 hours. Maximum growth rate across 3 timepoints was
118 estimated using Magellan (Tecan, Switzerland). Replica plating samples of endpoint culture onto 100 mM
119 HgCl₂ and testing by PCR showed no evidence of plasmid loss during the course of the experiment.
120 Rifampicin resistance of sequenced clones was assessed in a similar manner, except samples were added
121 to varying concentrations of KB + rifampicin (twofold dilutions from 0.2 to 200 μ g/ml). The threshold for
122 growth was set at OD600 > 0.03 after 48 h.

123 Phenotypic assays and growth curves were conducted on all 20 lineages, but six experimental lineages
124 either had a small colony morphology and/or slow or inconsistent growth, suggesting that they had not
125 ameliorated pQBR55 carriage. Competitions were therefore conducted only on the 14 remaining lineages.
126 Competitions were performed as described previously (21). Briefly, overnight cultures were washed in
127 M9 buffer (48 mM Na₂HPO₄, 22 mM KH₂PO₄, 8.55 mM NaCl, 19 mM NH₄Cl, pH 7.4, (22)), mixed with
128 plasmid-free *P. fluorescens* SBW25 Gm^R in approximately equal numbers, and samples serially diluted
129 and spread on KB + X-gal (50 μ g/ml) to enumerate starting cfus. Mixtures were diluted 1:100 into 6 ml
130 fresh KB media in a 30 ml glass universal and grown for 48 hours, or added 1:100 v/w to potting soil
131 microcosms pre-wetted with 900 μ l dH₂O and grown for 96 hours. Potting soil microcosms designed to
132 represent a more natural growth substrate of *P. fluorescens* consisted of 10 g unseived twice-autoclaved
133 John Innes #2 (manufactured by J. Arthur Bower, supplied by Vertigrow Ltd., UK) with ~25% w/v water
134 content as described previously (4,21). At the endpoint, samples of culture or soil wash were serially
135 diluted and spread on KB + X-gal and endpoint cfus were calculated. Relative fitness was calculated as

136 the ratio of Malthusian parameters (24). Endpoint colonies were replicated onto 100 mM HgCl₂ which
137 showed maintenance of mercury resistance in pQBR55-starting clones in all cases, and PCR analysis of
138 colonies from each sample showed no evidence of plasmid loss.

139 Photographs of transconjugant colonies were analysed with ImageJ using the Watershed and
140 AnalyseParticles tools. Measurements ± 2 standard deviations from the mean colony area for each image
141 were discarded as doublets or other errors, and the mean of the remaining measurements ($n \geq 8$ for each
142 sample) was analysed. To estimate the number of generations occurring during colony growth, three 48 h
143 old colonies were measured using ImageJ before each was dispersed in KB broth, serially diluted, and
144 plated to calculate cfu/colony. Ancestral plasmid-free colonies were calculated to contain 4.9×10^8 cells
145 with a standard error of 1.3×10^8 ($2.5 \pm 0.5 \times 10^7$ cells/mm²), whereas small colonies contained $2.2 \pm 0.8 \times$
146 10^6 cells ($3.3 \pm 1.1 \times 10^6$ cells/mm²). We therefore estimate an average of $\log_2(2.2 \times 10^6) = 21$
147 generations per small colony and $\log_2(4.9 \times 10^8) = 29$ generations per large colony. Assuming that each
148 lineage underwent two cycles of small colony and one of large colony morphology, we calculate ~ 70
149 generations.

150 **Sequencing**

151 Lineages were picked for genome resequencing based on *gac* and fitness phenotypes. Resequencing was
152 performed using Illumina technology (MiSeq/HiSeq) by MicrobesNG (<https://microbesng.uk>) which is
153 supported by the Biotechnology and Biological Sciences Research Council (BBSRC; grant number
154 BB/L024209/1), and reads are available at the ENA Short Read Archive with project accession number
155 PRJEB32206. Targeted sequencing of *gacA/S* and *PFLU4242* was performed by PCR amplifying the loci
156 with Phusion HF polymerase (NEB, MA USA) using 1x High-Fidelity buffer, 0.2 mM dNTPs, 0.5 μ M
157 each primer (see Table S3), and 1 μ l of glycerol stock as a template, with the following program: 98°C
158 30", 30 x (98°C 20", 63°C 30", 72°C 30"), 72°C 5'. Samples were run on agarose gels to ensure a single
159 product and amplicons purified using the QIAGEN PCR purification kit before sending for Sanger

160 sequencing. In the case of lineage 04, it was not possible to generate amplicons with any of six different
161 combinations of *PFLU4242* primers and thus this lineage was considered a *PFLU4242* deletion.

162 **Analysis of short-read sequencing**

163 Reads were mapped onto the *P. fluorescens* SBW25 chromosome (EMBL accession AM181176) and
164 pQBR55 (LN713927) using bwa mem version 0.7.17 (25) and variants called using gatk HaplotypeCaller
165 version 4.0.11.0 (26). Additionally, analysis with breseq version 0.33.0 (27) was carried out using the
166 default parameters. For each approach, predicted variants were compared between sequences and with the
167 plasmid-free ancestor to exclude those common to all sequenced strains. Repetitive regions prone to
168 spurious calling (28) were likewise masked from the analysis. The remaining predicted mutations were
169 examined manually for depth of coverage and consistency between reads using IGV (29), using a
170 threshold of mapping quality ≥ 60 and depth ≥ 10 . *PFLU4242* from lineages 19 and 20 was additionally
171 Sanger sequenced to verify this approach and confirm mutation presence.

172 To analyse plasmid copy number, coverage for plasmid and chromosome over 1 kb windows was
173 calculated. Windows with coverage ± 2 SD from the mean, due largely to poor mapping quality in
174 repetitive regions, were removed, and the mean across remaining windows was calculated. The ratio of
175 plasmid/chromosomal coverage in sequenced lineages ranged from 2.86 to 3.90 with a mean of 3.4,
176 suggesting modal pQBR55 copy number for all lineages of 3/cell. For pQBR57 the ratio in the ancestral
177 strain was 1.18 and for pQBR103 it was 1.36, suggesting modal copy number for these plasmids is 1/cell.

178 **Allelic replacement to generate knockout strains**

179 Strains lacking *gacA* or *gacS* were previously described by Harrison et al. (2015)(5). The *PFLU4242*
180 knockout was generated in a similar manner using a two-step process with the suicide vector pUIC3. Both
181 PCR and whole genome resequencing indicated that the gene of interest had been knocked out. The
182 $\Delta gacS$ mutant and the $\Delta PFLU4242$ mutant had no evidence of second-site mutations, however $\Delta gacA$ had
183 a single A3084294>G transition resulting in a F155L substitution in the putative integral membrane

184 protein PFLU2795, which was confirmed by Sanger sequencing. The consistent phenotypes between the
185 two different *gac* mutants and the fact that the $\Delta gacA$ mutant recapitulates the phenotype of lineage 13
186 suggests that this second-site mutation does not have a significant impact on our findings.

187 **Statistical analysis**

188 Fitness data for plasmid-bearers was analysed using a linear model with lineage and media as fixed
189 effects. Replicates where fitness could not be estimated due to lack of plasmid-bearer growth were
190 excluded. We did not detect an effect of marker (plasmid-free fitness no different from 1, KB $t[3] =$
191 0.485 , $p = 0.66$, soil $t[3] = 0.815$, $p = 0.475$), so to test whether plasmid-bearing strains were statistically
192 indistinguishable from plasmid-free we ran one-sample t-tests with $\mu = 1$. Logistic regression was used to
193 test association between *gac* activity and fitness, and non-parametric (Kruskal-Wallis) analysis was used
194 to test for the effect of different knockouts on colony size, due to heteroscedasticity. Analysis was
195 performed in R (R Core Team, Vienna).

196 **Results and discussion**

197 Acquisition of the plasmid pQBR55 caused a small-colony phenotype in *P. fluorescens* SBW25
198 transconjugants, a phenomenon not observed following acquisition of unrelated sympatric plasmids
199 pQBR57 or pQBR103 (Figure 1A) (21). The small colony phenotype was, however, transient: a further
200 48 h growth in liquid media or on agar plates (approx. 21 generations, Figures 1A, S1; Movie S1) restored
201 the ancestral (large) colony morphology. Re-emergence of large colony morphologies was not due to
202 plasmid loss as colonies remained mercury resistant and pQBR55-positive by PCR testing for *oriV*. Small
203 colony phenotypes in *P. aeruginosa* and *Staphylococcus aureus* reflect adaptations enabling prolonged
204 infection of humans and animals (30,31). However, as the pQBR55 small colony phenotype was
205 associated with plasmid acquisition and was rapidly lost, we hypothesized that it represented a
206 maladaptive response consistent with exceptionally high plasmid cost, and that restoration of the large
207 colony morphology was due to extremely fast compensatory evolution.

208 Unlike previous studies (e.g. (5–7,32)) it was not possible to conduct assays on transconjugants that had
209 just received the plasmid, as putatively compensated large colony variants emerged during assay
210 preparation. We therefore compared transconjugants that had restored the large morphology with one
211 another and with the ancestral plasmid-free strain. Twenty fresh independent *P. fluorescens*
212 SBW25(pQBR55) transconjugants were twice re-streaked on selective KB agar (~70 generations growth
213 in total). All remained pQBR55-positive, and 14/20 evolved a clear compensated (large colony)
214 morphology (Figures 1B, S1, S3). These 14 lineages varied in their fitness relative to the plasmid-free
215 ancestor when measured in nutrient-rich KB broth and in soil microcosms designed to approximate the
216 natural substrate of *P. fluorescens* (21) (Figures 1B, S2, Table S4). While most showed amelioration such
217 that fitness was similar to plasmid-free, five lineages still displayed low relative fitness, particularly in
218 soil (linear model, effect of lineage:media $F(10,74) = 11.94$, $p < 0.001$), where in some cases we failed to
219 detect competitive growth altogether. The fitness differences between evolved lineages suggested
220 occurrence of different compensatory mechanisms, as found previously (7,9), which could enhance
221 plasmid survival by increasing the supply of compensatory mutations.

222 To identify targets of compensatory mutation we chose 6 representative lineages (Figure 1B) and
223 performed whole genome resequencing. A single non-synonymous chromosomal mutation was detected
224 in each clone (Table S1), suggesting that the rapid emergence of compensation was not associated with
225 plasmid-induced hypermutation. No pQBR55 mutations were detected in any lineage. Lineages with the
226 least amelioration, 03 and 16, had mutations affecting RNA polymerase: lineage 03 had a 7 bp deletion in
227 the P26 partial terminator upstream of *rpoB* (*PFLU5534*) while lineage 16 had acquired a missense
228 mutation Pro584Ser in *rpoC* (*PFLU5533*). Lineages with intermediate amelioration, 13 and 18, had
229 mutations in *gacS* (*PFLU3777*) and *gacA* (*PFLU2189*), respectively (lineage 13 had an additional
230 synonymous C>T transition in *PFLU3410*). Lineage 19, which showed the most complete amelioration,
231 and 20, had mutations in *PFLU4242*, which encodes a hypothetical 527 amino acid protein with two
232 domains of unknown function (DUF262, DUF2081). The remaining evolved lineages phenotypically

233 resembled lineage 19 and targeted sequencing revealed each also had a mutation in *PFLU4242* (Figure
234 1B, Table S6). To test the ameliorative roles of *gacA/S* and *PFLU4242* mutations we generated pQBR55
235 transconjugants in *P. fluorescens* SBW25 $\Delta gacA$, $\Delta gacS$, and $\Delta PFLU4242$. Unlike wild-type, deletion-
236 mutant strains immediately and exclusively produced large-sized transconjugant colonies with fitness
237 similar to the evolved lineages (Figure 2, Figure S4, Table S7), suggesting that disrupting any of these
238 genes rendered *P. fluorescens* ‘pre-ameliorated’ for pQBR55 acquisition, and implicating these genes in
239 the high fitness costs of pQBR55.

240 GacA/GacS signaling positively regulates a suite of stationary phase phenotypes including flagella-
241 mediated motility and secreted products, suggesting that mutations affecting these genes have major
242 pleiotropic effects (20,33). Correspondingly, only strains with *gacA* or *gacS* mutations (13 and 18) had
243 lost the ability to produce extracellular protease, pectinase, surfactant (viscosin) and to swim (Figure 1B),
244 potentially restricting amelioration via this pathway in complex habitats like soil (34). Indeed, lineages
245 with the highest level of fitness in soil tended to retain GacA/GacS-regulated phenotypes (logistic
246 regression, effect of fitness in soil on protease result LRT $\text{Chisq}(1) = 15.8$, $p = 6.9e-5$). Mutation of
247 *gacA/gacS* was previously implicated in ameliorating the unrelated plasmid pQBR103, through
248 widespread effects on transcription (5). In *P. aeruginosa*, GacA/S controls small colony variants
249 generation through the activity of RsmA and the second messenger cyclic-di-GMP (31). Interestingly,
250 pQBR55 carries an RsmA homologue (PQBR55_0192), which might directly affect GacA/S or cyclic-di-
251 GMP signalling to cause small colonies in *P. fluorescens* (35).

252 Mutations to RNA polymerase are a common response to cellular stress, e.g. in rifampicin resistance (36)
253 and elevated temperature (37), and have previously been implicated in plasmid adaptation (6). Of the two
254 mutations detected here, one (RpoC Pro584Ser) probably affects specificity, whereas the other (deletion
255 of an attenuator upstream of *rpoB*) likely increases expression (36). These mutations may be adaptations
256 to modulate expression of specific disruptive genes, or to generally compensate for the transcriptional
257 burden of plasmid gene expression (38), and seem functionally distinct from those conferring rifampicin

258 resistance as we did not detect an increased rifampicin minimum inhibitory concentration (MIC) for
259 lineages 03 and 16 (Table S2).

260 The most complete amelioration occurred through *PFLU4242* mutation (Figure 1, Figure S4). *PFLU4242*
261 was likely acquired by *P. fluorescens* SBW25 via recent HGT as related strains *P. protegens* Pf-5 and *P.*
262 *fluorescens* Pf0-1 do not encode homologues whereas similar genes are found in distantly related genera
263 e.g. *Burkholderia* (WP_059533732.1, 80.5% amino acid identity), *Fischerella* (WP_016860471.1,
264 60.4%), and *Streptomyces* (WP_086730045.1, 50.8%). The GC-content of *PFLU4242* was 43.5%,
265 compared with 60.5% across the whole genome, also consistent with recent horizontal acquisition,
266 however we could not identify any neighbouring genes associated with mobility (e.g. transposases or
267 integrases), and the genes neighbouring *PFLU4242* homologues in these other species are not
268 homologous to those in *P. fluorescens* SBW25. The function of *PFLU4242* remains mysterious, but its
269 principal domain, DUF262, is a member of the ParB superfamily that includes nucleases as well as
270 partitioning systems (39). We did not detect any gross phenotypic effects of *PFLU4242* knockout on
271 colony morphology or growth in the absence of pQBR55. Further work is required to elucidate the
272 mechanistic basis of *PFLU4242* function, but our finding is consistent with a growing body of evidence
273 implicating horizontally-transferred DNA-binding proteins as key determinants of plasmid cost (6,7).

274 Re-examination of genome-resequencing data from previous experiments with *P. fluorescens* SBW25
275 implicates *PFLU4242* disruption in ameliorating unrelated plasmids pQBR103 (5) and pQBR57 (40),
276 suggesting a general mechanism behind plasmid costs in this host. Though *P. fluorescens* SBW25
277 isolated from the same site as the pQBR plasmids, it was originally plasmid-free (21), indicating it may
278 be a naive host, potentially vulnerable to conflicts between resident and newly-acquired genes.

279 Our observed rates of amelioration, occurring during the process of isolating individual plasmid-
280 containing clones, exceed those of previous studies and suggest a prominent role for compensation in
281 bacterial evolution. Similar mutations emerging during the preparation of plasmid-containing strains may
282 also influence the interpretation of experimental assays, leading, for example, to an underestimation of

283 plasmid carriage costs. In natural communities, rapid compensatory mutations could overcome the
284 oftentimes heavy metabolic and regulatory costs of plasmid-mediated horizontal gene transfer, and, in
285 combination with stability functions that expand the window of opportunity for such mutations to arise,
286 are likely to be a major factor promoting plasmid carriage and thus the maintenance and spread of
287 adaptive traits.

288 **Acknowledgements**

289 We would like to thank Alex Dunlop whose undergraduate work at the University of York provided some
290 preliminary data for this project but whom we were unfortunately unable to contact during the preparation
291 of this manuscript.


292 **Competing interests**

293 The authors have no competing financial interests to declare.

294 **Funding information**

295 This work was supported by ERC Grant Agreement 311490- COEVOCON (to M.A.B.) and a NERC
296 Standard Grant NE/R008825/1 (to J.P.J.H., E.H., and M.A.B.).

297



298

299 **Figure 1.** pQBR55-bearers rapidly ameliorate plasmid cost. **(A)** Transconjugants of pQBR55, but not

300 sympatric mercury resistance plasmids, consistently produce a small colony phenotype which is rapidly

301 resolved on restreaking. Scale bar indicates 1 mm. Lineage 20 is shown, photographs of all 20 lineages

302 are in Figure S1. Plasmid size and approximate modal copy number (CN) are provided for reference. **(B)**

303 Evolved lineages. Top: fitness relative to plasmid-free in KB (black) or in soil (grey). Four independent

304 replicates are shown, with mean and standard error overlaid. Where fitness could not be estimated (no


305 growth) it was set to zero. Control refers to competition between plasmid-free strains to assess any fitness

306 effects of the antibiotic markers. Middle: Gac phenotype assay results. Bottom: evolved loci. Colour

307 indicates target of locus-level parallel mutation, and the mutation is overlaid. del = deletion; P26 refers to

308 a partial attenuator upstream of *rpoB* (36). Strains subject to whole genome resequencing are marked with

309 triangles, the remaining *PFLU4242* sequences were determined by Sanger sequencing of amplicons.



311

312 **Figure 2.** pQBR55 transconjugant strains lacking in *gacA*, *gacS*, or *PFLU4242* do not display a small
313 colony morphology. Transconjugants of pQBR55 and pQBR57 were photographed and mean colony size
314 after 48 hours growth was measured. Four independent replicates are shown with mean and standard error
315 overlaid. Scale bar indicates 5 mm. We detected a significant effect of recipient on pQBR55 colony size
316 (Kruskal-Wallis $p = 0.016$) driven by the wild-type recipient (planned contrast Wilcoxon Test $p = 0.001$),
317 but not on pQBR57 (K-W $p = 0.061$; Wilcoxon $p = 0.8615$).

318

319 **References**

- 320 1. Norman A, Hansen LH, Sørensen SJ. Conjugative plasmids: Vessels of the communal gene pool.
321 *Philos Trans R Soc Lond B Biol Sci.* 2009 Aug;364(1527):2275–89.
- 322 2. San Millan A, MacLean RC. Fitness costs of plasmids: A limit to plasmid transmission. *Microbiol*
323 *Spectr.* 2017 Sep;5(5).
- 324 3. Kottara A, Hall JPJ, Harrison E, Brockhurst MA. Variable plasmid fitness effects and mobile genetic
325 element dynamics across pseudomonas species. *FEMS Microbiol Ecol.* 2018 Jan;94(1).
- 326 4. Hall JPJ, Wood AJ, Harrison E, Brockhurst MA. Source-sink plasmid transfer dynamics maintain gene
327 mobility in soil bacterial communities. *Proc Natl Acad Sci U S A.* 2016 Jul;113(29):8260–5.
- 328 5. Harrison E, Guymer D, Spiers AJ, Paterson S, Brockhurst MA. Parallel compensatory evolution
329 stabilizes plasmids across the parasitism-mutualism continuum. *Curr Biol.* 2015
330 Aug;25(15):2034–9.
- 331 6. Loftie-Eaton W, Bashford K, Quinn H, Dong K, Millstein J, Hunter S, et al. Compensatory mutations
332 improve general permissiveness to antibiotic resistance plasmids. *Nat Ecol Evol.* 2017
333 Sep;1(9):1354–63.
- 334 7. San Millan A, Toll-Riera M, Qi Q, MacLean RC. Interactions between horizontally acquired genes
335 create a fitness cost in pseudomonas aeruginosa. *Nat Commun.* 2015 Apr;6:6845.
- 336 8. Harrison E, Dytham C, Hall JPJ, Guymer D, Spiers AJ, Paterson S, et al. Rapid compensatory
337 evolution promotes the survival of conjugative plasmids. *Mob Genet Elements.* 2016
338 May;6(3):e1179074.
- 339 9. Stalder T, Rogers LM, Renfrow C, Yano H, Smith Z, Top EM. Emerging patterns of plasmid-host
340 coevolution that stabilize antibiotic resistance. *Sci Rep.* 2017 Jul;7(1):4853.

- 341 10. Sota M, Yano H, Hughes JM, Daughdrill GW, Abdo Z, Forney LJ, et al. Shifts in the host range of a
342 promiscuous plasmid through parallel evolution of its replication initiation protein. *ISME J.* 2010
343 Dec;4(12):1568–80.
- 344 11. De Gelder L, Williams JJ, Ponciano JM, Sota M, Top EM. Adaptive plasmid evolution results in host-
345 range expansion of a broad-host-range plasmid. *Genetics.* 2008 Apr;178(4):2179–90.
- 346 12. Turner PE, Williams ESCP, Okeke C, Cooper VS, Duffy S, Wertz JE. Antibiotic resistance correlates
347 with transmission in plasmid evolution. *Evolution.* 2014 Dec;68(12):3368–80.
- 348 13. Porse A, Schønning K, Munck C, Sommer MOA. Survival and evolution of a large multidrug
349 resistance plasmid in new clinical bacterial hosts. *Mol Biol Evol.* 2016 Nov;33(11):2860–73.
- 350 14. Lilley AK, Bailey MJ. The acquisition of indigenous plasmids by a genetically marked pseudomonad
351 population colonizing the sugar beet phytosphere is related to local environmental conditions.
352 *Appl Environ Microbiol.* 1997 Apr;63(4):1577–83.
- 353 15. Carattoli A, Zankari E, García-Fernández A, Voldby Larsen M, Lund O, Villa L, et al. In silico
354 detection and typing of plasmids using PlasmidFinder and plasmid multilocus sequence typing.
355 *Antimicrob Agents Chemother.* 2014 Jul;58(7):3895–903.
- 356 16. Xiong J, Alexander DC, Ma JH, Déraspe M, Low DE, Jamieson FB, et al. Complete sequence of
357 pOZ176, a 500-kilobase IncP-2 plasmid encoding IMP-9-mediated carbapenem resistance, from
358 outbreak isolate *pseudomonas aeruginosa* 96. *Antimicrob Agents Chemother.* 2013
359 Aug;57(8):3775–82.
- 360 17. Botelho J, Grosso F, Quinteira S, Mabrouk A, Peixe L. The complete nucleotide sequence of an IncP-
361 2 megaplasmid unveils a mosaic architecture comprising a putative novel bla VIM-2-harboring
362 transposon in *pseudomonas aeruginosa*. *J Antimicrob Chemother.* 2017;72(8):2225–9.

- 363 18. Zhang X-X, Rainey PB. Construction and validation of a neutrally-marked strain of *Pseudomonas*
364 *fluorescens* SBW25. *J Microbiol Methods*. 2007 Oct;71(1):78–81.
- 365 19. Turner SL, Lilley AK, Bailey MJ. Two *dnaB* genes are associated with the origin of replication of
366 pQBR55, an exogenously isolated plasmid from the rhizosphere of sugar beet. *FEMS Microbiol*
367 *Ecol*. 2002 Nov;42(2):209–15.
- 368 20. Cheng X, Bruijn I de, Voort M van der, Loper JE, Raaijmakers JM. The *gac* regulon of *Pseudomonas*
369 *fluorescens* SBW25. *Environ Microbiol Rep*. 2013 Aug;5(4):608–19.
- 370 21. Hall JPJ, Harrison E, Lilley AK, Paterson S, Spiers AJ, Brockhurst MA. Environmentally co-
371 occurring mercury resistance plasmids are genetically and phenotypically diverse and confer
372 variable context-dependent fitness effects. *Environ Microbiol*. 2015 Dec;17(12):5008–22.
- 373 22. Sambrook J, Fritsch EF, Maniatis T. *Molecular cloning: A laboratory manual*. Cold Spring Harbor
374 Laboratory; 1989.
- 375 23. Gómez P, Buckling A. Bacteria-phage antagonistic coevolution in soil. *Science*. 2011
376 Apr;332(6025):106–9.
- 377 24. Lenski RE, Rose MR, Simpson SC, Tadler SC. Long-Term experimental evolution in *Escherichia coli*.
378 I. Adaptation and divergence during 2,000 generations. *Am Nat*. 1991 Dec;138(6):1315–41.
- 379 25. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.
380 *Bioinformatics*. 2009 Jul;25(14):1754–60.
- 381 26. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et al. The genome
382 analysis toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data.
383 *Genome Res*. 2010 Sep;20(9):1297–303.

- 384 27. Deatherage DE, Barrick JE. Identification of mutations in laboratory-evolved microbes from next-
385 generation sequencing data using breseq. *Methods Mol Biol.* 2014;1151:165–88.
- 386 28. Hall JPJ, Williams D, Paterson S, Harrison E, Brockhurst MA. Positive selection inhibits gene
387 mobilisation and transfer in soil bacterial communities. *Nat Ecol Evol.* 2017 Sep;1(9):1348–53.
- 388 29. Thorvaldsdóttir H, Robinson JT, Mesirov JP. Integrative genomics viewer (IGV): High-performance
389 genomics data visualization and exploration. *Brief Bioinform.* 2013 Mar;14(2):178–92.
- 390 30. Proctor RA, Eiff C von, Kahl BC, Becker K, McNamara P, Herrmann M, et al. Small colony variants:
391 A pathogenic form of bacteria that facilitates persistent and recurrent infections. *Nat Rev*
392 *Microbiol.* 2006 Apr;4(4):295–305.
- 393 31. Malone JG. Role of small colony variants in persistence of pseudomonas aeruginosa infections in
394 cystic fibrosis lungs. *Infect Drug Resist.* 2015 Jul;8:237–47.
- 395 32. Hughes JM, Lohman BK, Deckert GE, Nichols EP, Settles M, Abdo Z, et al. The role of clonal
396 interference in the evolutionary dynamics of plasmid-host adaptation. *MBio.* 2012
397 Jul;3(4):e00077–12.
- 398 33. Lapouge K, Schubert M, Allain FH-T, Haas D. Gac/Rsm signal transduction pathway of gamma-
399 proteobacteria: From RNA recognition to regulation of social behaviour. *Mol Microbiol.* 2008
400 Jan;67(2):241–53.
- 401 34. Jousset A, Scheu S, Bonkowski M. Secondary metabolite production facilitates establishment of
402 rhizobacteria by reducing both protozoan predation and the competitive effects of indigenous
403 bacteria. *Funct Ecol.* 2008 Aug;22(4):714–9.

- 404 35. Madsen JS, Hylling O, Jacquiod S, Pécastaings S, Hansen LH, Riber L, et al. An intriguing
405 relationship between the cyclic diguanylate signaling system and horizontal gene transfer. *ISME*
406 *J.* 2018 Jun;
- 407 36. Qi Q, Preston GM, MacLean RC. Linking system-wide impacts of RNA polymerase mutations to the
408 fitness cost of rifampin resistance in *pseudomonas aeruginosa*. *MBio.* 2014 Dec;5(6):e01562.
- 409 37. Rodríguez-Verdugo A, Tenailon O, Gaut BS. First-Step mutations during adaptation restore the
410 expression of hundreds of genes. *Mol Biol Evol.* 2016 Jan;33(1):25–39.
- 411 38. Park C, Zhang J. High expression hampers horizontal gene transfer. *Genome Biol Evol.* 2012
412 Mar;4(4):523–32.
- 413 39. Machnicka MA, Kaminska KH, Dunin-Horkawicz S, Bujnicki JM. Phylogenomics and sequence-
414 structure-function relationships in the GmrSD family of type IV restriction enzymes. *BMC*
415 *Bioinformatics.* 2015 Oct;16:336.
- 416 40. Hall JPJ, Harrison E, Brockhurst MA. Competitive species interactions constrain abiotic adaptation in
417 a bacterial soil community. *Evol Lett.* 2018 Dec;2(6):580–9.
- 418

419 **Supplementary information for ‘Extremely fast amelioration of plasmid fitness costs by multiple**
420 **functionally-diverse pathways’**

421 **Authors:** James P. J. Hall^{1,2,3*} ORCID 0000-0002-4896-4592, Rosanna C. T. Wright² ORCID 0000-
422 0002-8095-8256, David Guymer³, Ellie Harrison² ORCID 0000-0002-2050-4631, Michael A. Brockhurst²
423 ORCID 0000-0003-0362-820X

424 **Affiliations:**


425 ¹Institute of Integrative Biology, University of Liverpool, Crown Street, Liverpool, L69 7ZB

426 ²Department of Animal and Plant Sciences, University of Sheffield, Western Bank, Sheffield, S10 2TN

427 ³Department of Biology, University of York, Wentworth Way, York, YO10 5DD

428 *Correspondence to: j.p.j.hall@liverpool.ac.uk, Institute of Integrative Biology, University of Liverpool,
429 Crown Street, Liverpool, L69 7ZB, United Kingdom. Tel: +44 (0)151 79 57691

430 **Data accessibility statement:** .csv tables of data supporting the results are in Supplementary Tables S1-
431 S9, at [doi:10.17638/datacat.liverpool.ac.uk/953](https://doi.org/10.17638/datacat.liverpool.ac.uk/953), and short read sequences are on the Short Read Archive,
432 accession PRJEB32206.




433

434 **Figure S1.** Colony morphologies of the 20 lineages. Where multiple colonies are visible, arrows mark the

435 one which was selected. Lineages marked with an asterisk were deemed to have recovered ancestral

436 phenotype based on growth curves and colony morphology.


437



438

439 **Figure S2.** Maximum growth rate of the 20 lineages. Three independent replicates were performed for
 440 each lineage. The ancestral plasmid-free clone for each lineage was also measured (n = 20, 'control').

441



442

443 **Figure S3.** The 20 lineages carry pQBR55. PCR for the *merA* resistance gene and the pQBR55 origin of


444 replication was performed on the stocked colony for each lineage. PCR using primers specific for *P.*

445 *fluorescens* SBW25 and *P. putida* KT2440 (from which the donor strain UWC1 was derived) were used

446 to ensure strains were transconjugants. 15a and 15b are additional colonies retrieved from lineage 15. M =

447 NEB 100 bp ladder.

448



449

450 **Figure S4.** Knockout transconjugants have relative fitness similar to evolved lines. Transconjugant
 451 colonies from Figure 2 were restreaked twice to isolate a single clone and competed against a plasmid-
 452 free competitor in KB. During this process, the wild-type small colony morphologies reverted to the large
 453 colony phenotype, and thus should be considered de novo mutants. We detected a significant effect of
 454 recipient (ANOVA $F_{3,12} = 9.6$, $p = 0.002$), driven largely by reduced amelioration by the $\Delta gacA$ mutant.

455

sample	chromosome	position	reference	alternative	depth	mapping quality	quality	name	locus tag	nucleotide mutation	amino acid mutation	GATK
ΔgacA	AM181176	3084294	A	G	21	60	786	PFLU_2795	PFLU_2795	c.463T>C	p.Phe155Leu	1:0,21:21:99:816,0
lineage 03	AM181176	6062127	ATCAGCCT	A	165	60	7242	rpoB-rpIL	PFLU_5534-PFLU_5535	n.6062128_6062134delTCAGCCT		1:0,163:163:99:7281,0
lineage 13	AM181176	4174597	T	G	94	60	3164	PFLU_3777	PFLU_3777	c.2078T>G	p.Val693Gly	1:0,94:94:99:3194,0
lineage 16	AM181176	6056066	G	A	204	60	7317	rpoC	PFLU_5533	c.1750C>T	p.Pro584Ser	1:0,204:204:99:7347,0
lineage 18	AM181176	2373295	G	A	72	60	2252	gacA	PFLU_2189	c.614G>A	p.Arg205His	1:0,72:72:99:2282,0
lineage 18	AM181176	3773830	C	T	73	60	2246	PFLU_3410	PFLU_3410	c.618G>A	p.Ser206Ser	1:0,73:73:99:2276,0
lineage 19	AM181176	4684561	T	C	63	60	2042	PFLU_4242	PFLU_4242	c.1369A>G	p.Asn457Asp	1:0,63:63:99:2072,0
lineage 20	AM181176	4685131	A	T	83	60	2820	PFLU_4242	PFLU_4242	c.799T>A	p.Leu267Met	1:0,83:83:99:2850,0

456

457 **Table S1.** Mutations detected in whole genome resequenced lineages. Results are shown from the
458 bwa/GATK pipeline, similar results were obtained using breseq.

459

lineage	replicate		
	A	B	C
13	12.5	12.5	12.5
16	6.25	6.25	12.5
18	12.5	12.5	12.5
19	12.5	12.5	12.5
20	12.5	12.5	12.5
03	12.5	12.5	12.5
plasmid-free	12.5	25	12.5

460

461 **Table S2.** Rifampicin minimum inhibitory concentrations (MIC) for sequenced evolved lineages. Values

462 are in $\mu\text{g/ml}$.

463

Primer name	Sequence	Purpose	
pQBR55_0051_R1	ATGGCTGCCCTAGACCTGG	Detecting pQBR55 origin of replication	
pQBR55_0050_F1	CAACGCCCGAACAAACGC		
merA_F	TGCAAGACACCCCTATTGGAC	Detecting merA presence	
merA_R	TTCGGCGACCAGCTTGATGAAC		
Δ PFLU4242_UF	TATAGGATCCACTAGTATGTTCCGTGGCTCCTC	Generating the PFLU4242 knockout	
Δ PFLU4242_UR	GAAATTATTTAACGACTCTTCCCTACTCGCATT		
Δ PFLU4242_DF	GGAAGAGTCGTTAAATAATTTTCGCTTTAATACCGTTAT		
Δ PFLU4242_DR	TATAGAATTCACTAGTGATTTTGGCCTTAAAGCAAAGC		
Δ PFLU4242_TESTF	GTCTCTACTACGATCAAGTATTGACAC		
Δ PFLU4242_TESTR	CTTGGGCCAAAAGCGGAC		
PFLU4242_FAMP	GAACTCGAGTACATTGGCGC	Amplifying and sequencing PFLU4242	
PFLU4242_RAMP	CGGGCCCTTTTACATAACGG		
PFLU4242_FINT	ACCTGGAGCGATGACTTTGA		
PFLU4242_RINT	TCCAGAAAAGCACGTACGC		
PFLU4242_intF_3	TTGCTGTGCGAAGGAATTTG		
PFLU4242_intR_5	TCCCAAACAAGCCACGC		
PFLU4242_FSEQ	AGGGAAGAGTCGTATGGAGT		
gacA_F	CAGCAAATAGAGCCGTCCGCCTC		Amplifying and sequencing gacA
gacA_R	CCAGCGCAGCTGTCTTGG		
gacA_400F	CGCCACGACCAAGTTGTTG		
gacA_600R	GGCTGGAAGACTTGATGGC		
gacS_F	CCAATCGCCGCCGGAC	Amplifying and sequencing gacS	
gacS_R	CAGGCCTGGCGGCAG		
gacS_600F	CTTGCTGCCGGTGTTTGG		
gacS_750R	GAGTTGATCGTACGGCTGATG		
gacS_1500F	AGGACGAGCAGGAAGACAG		
gacS_1650R	GATTTACCGCCCATCTGC		
gacS_2350F	TCGACTTGGTGTGATGGAC		
gacS_2550R	GGTCAGGTAATCGTCCATGC		
SBW25_F	ACTGCATTCAAACCTGACTGA		Distinguishing strains
SBW25_R	AATCACACCGTGGTAACCG		
KT2440_F	ATGGCAATGTCCGCAATCC		
KT2440_R	CGGAAGCCTCTGAACACG		

464

465 **Table S3.** Sequences of primers used in this study.

466

467 **Tables S4-S9 are provided as .csv files at [doi:10.17638/datacat.liverpool.ac.uk/953](https://doi.org/10.17638/datacat.liverpool.ac.uk/953)**

468 **Table S4.** Relative fitness plotted in Figure 1B.

469 **Table S5.** Gac assay results plotted in Figure 1B.

470 **Table S6.** Summary of genetics information plotted in Figure 1B.


471 **Table S7.** Colony size measurements plotted in Figure 2.

472 **Table S8.** Growth curve data plotted in Figure S2. Time is in seconds and rate is in Δ OD/s.

473 **Table S9.** Relative fitness of knockout strains plotted in Figure S4. Two replicate experiments were

474 performed, data from experiment 1 was plotted.

475



476

477 **Movie S1.** Breakout of compensated mutants from a small morphology colony. Small colonies restreaked
478 from lineage 15 were maintained at room temperature and photographed every 30 minutes. Multiple
479 independent mutants can be seen emerging from each colony. This movie is provided through the journal
480 website and at [doi:10.17638/datacat.liverpool.ac.uk/953](https://doi.org/10.17638/datacat.liverpool.ac.uk/953).

481

482