

ADIPOKINE EXPRESSION IN
MOUSE MODELS OF
ENDOTOXAEMIA AND SEPSIS

Thesis submitted in accordance
with the requirements of the
University of Liverpool for the
degree of MD

BY

Dr. Andrew Michael Rushton

December 2012

SUPERVISERS: Prof. Martin Leuwer
Prof. Paul Trayhurn

ACKNOWLEDGEMENTS

I would like to thank my wife Ruth for her patience and support during the work that went into this thesis.

I need to mention Mr. Robert Galvin and the Biomedical Services Unit at the University of Liverpool for their assistance in the setting up and conduct of the experimental models.

I would like to thank Prof. Martin Leuwer for his expert guidance in the production of this thesis.

TABLE OF CONTENTS

	PAGE
INTRODUCTION	1
RESEARCH QUESTION	4
MODELS OF SEPSIS	5
KEY FEATURES OF A MODEL:	5
MODELS USED:	6
AIMS:	6
LIMITATIONS OF MODELS	6
LEGAL REQUIREMENTS	8
1. PRELIMINARY CLP (SURGICAL SEPSIS) AND LPS (ENDOTOXAEMIA) WORK	8
METHOD	8
CLP SURGICAL PROTOCOL	9
ANAESTHESIA PROTOCOL	10
LPS PROTOCOL	11
SAMPLE COLLECTION	11
RESULTS	12
2. REFINED LPS MODEL (ENDOTOXAEMIA)	21
LPS METHOD	22
RNA EXTRACTION AND PCR	25
MEASUREMENT OF PROTEIN BY ELISA	27
DATA PRESENTATION AND STATS	27
RESULTS	28
Effect of LPS on adipokine expression 24 h after the induction of endotoxaemia	28

Effect of LPS on adipokine expression 4 h after the induction of endotoxaemia	31
White adipose tissue adipokine levels	35
Macrophage marker expression	37
HIF-1 α as an index of hypoxia	39
DISCUSSION	41
APPENDIX	48
REFERENCES	66

INTRODUCTION

Sepsis kills. With an estimated annual mortality of 30 -50 per 100,000 population, although survival has slightly increased in the past 10 - 15 years (1) (2). It affects 18 million people worldwide every year and ranks in the top 10 causes of death with annual treatment costs in the US alone of over \$16 billion (3).

SIRS	Two or more of: Temperature >38°C or <36°C Tachycardia >90 beats per minute Respiratory rate >20 breaths per minute or PaCO ₂ <4.3kPa White Blood cell count >12 x 10 ⁹ /L or <4 x 10 ⁹ /L or >10% immature (band) forms
Sepsis	SIRS due to infection
Severe Sepsis	Sepsis with evidence of organ hypoperfusion
Septic Shock	Severe sepsis with hypotension (systolic blood pressure < 90mmHg) despite adequate fluid resuscitation or the requirement for vasopressors / inotropes to maintain blood pressure

Table 1: Clinical definitions for the systemic inflammatory response syndrome (SIRS) and sepsis (4)

Sepsis, originally defined as putrefaction or decomposing of organic matter by micro-organisms (5), is the body's systemic response to infection. Consensus was reached with regards to clinically defining the condition in 1992 (table 1) (4) with the list of signs and symptoms increased in 2001 (6) and a simple haematological investigation has recently been shown to aid early diagnosis (7). However, these clinical criteria for the diagnosis of sepsis do not discriminate differences in underlying pathophysiology, thus hindering development and application of effective targeted therapies (8) (9) (10) (11).

The systemic inflammatory states of sepsis, septic shock and endotoxaemia affect most organs of the body including white adipose tissue with lipolytic activity and glucose metabolism particularly affected (12) (13). It is well established that glucose utilization is impaired and lipolysis altered in obesity, inflammation and sepsis (14) (15).

However, it has recently come to light that white adipose tissue (WAT), the body's main insulator and energy store, plays a mediatory role not confined to just glucose and fat metabolism and is a major endocrine secretory organ. WAT, by secreting protein mediators and signalling factors coined adipokines (16), contributes to inflammation related conditions in obesity such as the metabolic syndrome (17) (18).

Obesity is now a major health concern worldwide and has been described as an epidemic (19) (20). However, for obese ICU patients there appears to be a conundrum or even paradox with respect to outcomes. Obese patients were initially shown to have a poorer outcome from critical illness compared to non-obese patients (21-23) (24), but recent meta-analyses (25) (26) have suggested the reverse. But this is probably not the whole story, as a series of potential confounders may have affected these results (27) and indeed it may be the location of fat depots is a better predictor (28). Obese patients are still more difficult to manage due to difficulties and complications of ventilation and the high incidence of serious comorbidities such as ischaemic heart disease and diabetes.

As adipose tissue mass expands the concentration of a number of adipokines and proinflammatory markers expressed and secreted by adipocytes has been shown to rise. These include - cytokines (e.g. IL-6, IL-10), chemokines (e.g. IL-8, MCP-1, MIF) and acute-phase proteins (e.g. haptoglobin, serum amyloid A) as well as inflammation-

related adipokines such as leptin and nerve growth factor (NGF) (17) (18) (16) (29). These inflammatory responses may reflect hypoxia in the growing adipose tissue mass. Hypoxia is known to induce cytokine production in organs such as heart, kidneys and gut (30). Sepsis involves general microvascular dysfunction causing a decrease in oxygen delivery, producing tissue hypoxia and organ dysfunction (31). The recognition and appropriate response of cells to hypoxia is substantially mediated by the hypoxia inducible factor-1 (HIF-1) transcriptional system (32) (30, 33). Activation of HIF-1 during hypoxia, by stabilization of the HIF-1 α subunit, triggers expression of glycolytic enzymes and membrane glucose transporters, such as GLUT-1 (32). HIF-1 α was discovered more than 20 years ago when the expression of erythropoietin was being studied (34). It rapidly became clear that HIF-1 α is a key regulator of many other systems and it is now apparent that certain cytokines and growth factors can make use of this 'hypoxic signaling system' under normoxia (35). Some of these responses are amplified in the presence of hypoxia (36). There appears to be distinct signalling pathways, induced by hypoxia or Toll-like receptor 4 mediated, for HIF-1 α stabilisation leading to differential gene expression (37).

RESEARCH QUESTION

It was hypothesized that in endotoxaemia there is a major activation of the inflammatory response in adipose tissue, key inflammatory adipokines contributing to the metabolic dysregulation characteristic of endotoxaemia and sepsis. I used murine models of endotoxaemia and polymicrobial (surgical) sepsis to study the expression in both internal and subcutaneous adipose tissue of a series of inflammation-related adipokines. Evidence for the induction of adipose tissue hypoxia was sought through measurement of HIF-1 α .

MODELS OF SEPSIS

Sepsis is a complex disease, affecting multiple organ systems, and frequently requires intensive care. The average age of patients is 60 - 65 years and it is more common in men (38) with higher mortality with increasing age (3). Severe sepsis has long term (>6 months) effects on both mental and physical health (39).

As well as treating the underlying infection with appropriate antibiotics and surgery etc, the patient often needs haemodynamic and respiratory support. This includes the use of sedative, analgesic, inotropic, steroid and other medication. Frequent observation and continuous invasive monitoring of vital signs are used to guide treatment.

KEY FEATURES OF A MODEL:

Standardisation - in order to minimise individual differences and genetic polymorphisms responsible for the spectrum of responses seen in human sepsis (40) and ensure all subjects experience identical conditions. Hence, animals should be of the same strain, age, weight and experience identical husbandry conditions and the intervention should be of a standard dose for each arm of the study.

The procedures performed should be easily reproducible and hence reduce any operator variability. Effects of the model need to be clearly measurable which could be by direct observation, invasive / non-invasive monitoring and laboratory tests. Ultimately, the model needs to be valid - i.e. produce a definite septic response due to polymicrobial contamination or endotoxaemia etc. The actual model should also be as simple and cost effective as possible and use the least number of subjects to answer the question.

MODELS USED:

There are mainly two widely used models:

1. Injection, either intraperitoneal or intravenous, of bacteria or bacterial contents (e.g. LPS). These are simple methods that can be used to produce **endotoxaemia** or sepsis due to specific organisms (41).
2. The release of faeces into the peritoneal cavity by caecal ligation and puncture (CLP) or ascending colon stent peritonitis. These are the preferred methods to model the **polymicrobial** insult of **surgical / perioperative** sepsis (41) (42, 43).

AIMS:

To manage the complexity and demonstrate specific biological dysfunction in systems in order to gain a better understanding of the pathophysiology (Fig.1).

LIMITATIONS OF MODELS

Models by their nature are a controlled artificial environment with a set of specified procedures that in themselves can effect the outcome. Anaesthesia, which was a requirement of our licence, can effect onset and survival after LPS injection (44) and opiates, which were used in the CLP work, may alter the immune response (45). However, deep sedation incorporating opiates is used in the clinical management of patients with severe sepsis.




Fig 1: Aims of a model of sepsis

LEGAL REQUIREMENTS

IN ACCORDANCE WITH ANIMALS (SCIENTIFIC PROCEDURES) ACT 1986 THE SPECIFIC LICENCES UNDER WHICH WORK PERFORMED:

PROJECT LICENCE : PPL 40/2692

PERSONAL LICENCE : PIL 40/7646

In the first round of experiments both models have been implemented:

1. PRELIMINARY CLP (SURGICAL SEPSIS) AND LPS (ENDOTOXAEMIA) WORK

Caecal ligation and puncture produces a polymicrobial insult more akin to that seen in sepsis due to traumatic injury or surgery (42, 43).

METHOD

After obtaining a Home Office licence, we investigated thirty-two healthy, 8-10 week old, inbred, non-obese, male C57BL/6J mice (obtained from Charles River, UK) (body wt. 25-28 g). Eight animals served as controls; in eight animals, sepsis was induced by caecum ligation/puncture (CLP), in another eight animals by intra-peritoneal LPS injection (25 µg/g body wt) and eight animals were sham-operated. Twenty-four hours after induction of sepsis, the experimental animals were killed and RNA extracted from the epididymal fat pads (Fig.8). Real time PCR (see later experimental protocol) was used to quantify expression of IL-6, TNF α , IL-18, MCP-1, nerve growth factor (NGF), hypoxia-induced factor (HIF-1 α), adiponectin and leptin, in comparison to controls.

CLP SURGICAL PROTOCOL

The mouse was weighed and anaesthetised (see separate protocol).

The mouse was positioned on a cork board using loops of micropore tape (only need to tape lower limbs) and a heating mattress placed under the board. The abdomen was shaved (hovering the shaved fur) and cleaned with aqueous, not alcoholic, iodine solution.

A midline abdominal incision was made using small scissors and opened in 2 layers - being careful not to damage bowel.

The caecum was exposed - cotton buds are useful to aid retraction and move bowel (the abdomen of sham operated animals would be closed at this point as described below).

Once mobilised the caecum was tightly ligated with a silk tie at the base.

The caecum was punctured twice with a 21G needle, distal to the ties, and squeezed to expel some of its contents.

The abdomen was closed in 2 layers with 4/0 Dexon - peritoneum and musculature with a continuous suture and skin with interrupted sutures.

The mouse was given a subcutaneous injection of 1ml warmed 0.9% Saline.

Analgesia was provided, anaesthesia reversed, and post op care provided as described in the separate protocol.

ANAESTHESIA PROTOCOL

The mouse was placed in the anaesthetic chamber and general anaesthesia induced with 1-2% isoflurane (Abbott, uk) in nitrous oxide / oxygen mixture (N2O/O2/ISO).

Once anaesthetised the mouse was removed from chamber and injected with:

- Ketamine 75mg/kg - 0.2ml of 1:10 dilution
- Medetomidine 1mg/kg - 0.3ml of 1:10 dilution (reduce to 0.2ml if weighs <25g)

Supplemental oxygen was provided while the injectable anaesthetic took effect.

REVERSAL

The mouse was administered an Antisedan injection - 0.1ml sc of a 1:10 dilution.

POST-OP ANALGESIA

Analgesia was provided by an injection of Buprenorphine - 0.15ml sc of 1:20 dilution

POST-OP CARE

The mouse was given a fluid bolus of 1ml 0.9% Saline sc to the scruff and placed on bubble wrap mattress in an incubator with 2L /min O2. The incubator temperature was monitored - aiming to maintain ~30-32 C (heating mattress may need turning of and on) and the mouse turned frequently - ~ 30 minutely intervals. It was transferred to an individual cage when became mobile. Further fluid boluses 0.5ml 0.9% Saline sc were administered as required (if lethargic). The mice were monitored regularly by a trained anaesthetist as per license requirements.

LPS PROTOCOL

24Hr LPS

The mouse was weighed.

The mouse was placed in the anaesthetic chamber and general anaesthesia induced with 1-2% isoflurane (Abbott, uk) in nitrous oxide / oxygen mixture (N2O/O2/ISO).

Once deeply anaesthetised 600 µg of LPS (Escherichia coli O 111:B4, Sigma-Aldrich) was injected ip (0.3ml) ~ 25µg / g

To compensate for fluid losses 1ml of Saline was administered sc to the scruff.

Analgesia was provided by giving Buprenorphine injection sc 0.15ml of 1:20 dilution.

Post procedure the mice were housed in separate clean cages and maintained in the same temperature controlled (24-28°C) conditions with free access to standard laboratory chow and water.

Repeat fluid boluses were administered as indicated.

SAMPLE COLLECTION

24 hours post procedure all animals were killed by the schedule 1 method of rising concentration of carbon dioxide.



The fat deposits were removed and immediately frozen in liquid nitrogen and stored at -80°C until further analysis.

RESULTS

Twenty-four hours after induction of sepsis by LPS, real time PCR revealed substantial increases in the expression of IL-6 (500-fold), MCP-1 (55-fold), NGF (10-fold), TNF α (5-fold) and HIF-1 α (3-fold), while there were significant decreases in the expression of IL-18 (2.5-fold), adipsin (6-fold) and adiponectin (5-fold). The results in the CLP-group were qualitatively similar.



The work in these first set of experiments was presented as a poster (see appendix) at 2nd International Congress of the German Sepsis Society (GSS) in Weimar, Germany , September 7 - 10 , 2005.

RESULTS





*p<0.05 **p<0.01 ***p<0.001 ****p<0.0001

Fig. 2 Adipokine gene expression in sepsis and endotoxaemia. Twenty-four hours after induction of sepsis by CLP (blue bars) and endotoxaemia (red bars) or sham lapaparotomy (turquoise bars) surviving animals were killed by a rising concentration of CO₂ and the epididymal fat depots removed. The level of each adipokine mRNA was measured by real-time PCR and normalized to β -actin relative to the untreated control group (\cong 1). Real time PCR revealed significant decreases in the expression adipisin and adiponectin which were qualitatively similar in both the CLP and LPS groups. Results are given as mean values \pm SE for five to six mice in each group. * P <0.05, ** P <0.01, *** P <0.001 compared with controls.





*p<0.05 **p<0.01 ***p<0.001 ****p<0.0001

Fig. 3 T Adipokine gene expression in sepsis and endotoxaemia. Twenty-four hours after induction of sepsis by CLP (blue bars) and endotoxaemia (red bars) or sham lapaparotomy (turquoise bars) surviving animals were killed by a rising concentration of CO₂ and the epididymal fat depots removed. The level of each adipokine mRNA was measured by real-time PCR and normalized to β -actin relative to the untreated control group ($\equiv 1$). In the LPS group real time PCR revealed a substantial increase in the expression of MCP-1 (55-fold, while there was a significant decrease in the expression of IL-18 (2.5-fold). The results in the CLP-group were qualitatively similar. Results are given as mean values \pm SE for five to six mice in each group. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ compared with controls.



*p<0.05 **p<0.01 ***p<0.001 ****p<0.0001

Fig. 4 Adipokine gene expression in sepsis and endotoxaemia. Twenty-four hours after induction of sepsis by CLP (blue bars) and endotoxaemia (red bars) or sham lapaparotomy (turquoise bars) surviving animals were killed by a rising concentration of CO₂ and the epididymal fat depots removed. The level of each adipokine mRNA was measured by real-time PCR and normalized to β -actin relative to the untreated control group (\equiv 1). Real time PCR revealed substantial increases in the expression of IL-6 (500-fold), and HIF-1 α (3-fold) in the LPS group with qualitatively similar results in the CLP-group. Results are given as mean values \pm SE for five to six mice in each group. * P <0.05, ** P <0.01, *** P <0.001 compared with controls.



*p<0.05 **p<0.01 ***p<0.001 ****p<0.0001

Fig. 5 Adipokine gene expression in sepsis and endotoxaemia. Twenty-four hours after induction of sepsis by CLP (blue bars) and endotoxaemia (red bars) or sham lapaparotomy (turquoise bars) surviving animals were killed by a rising concentration of CO₂ and the epididymal fat depots removed. The level of each adipokine mRNA was measured by real-time PCR and normalized to β -actin relative to the untreated control group ($\equiv 1$). Real time PCR revealed substantial increases in the expression of NGF (10-fold) and TNF α (5-fold) in the LPS group with results in the CLP-group being qualitatively similar. Results are given as mean values \pm SE for five to six mice in each group. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ compared with controls.

2. REFINED LPS MODEL (ENDOTOXAEMIA)

Intraperitoneal injection of LPS was used which produces **endotoxaemia**. The efficacy had been proven from the preliminary work (see first set of experiments).

Further refinements were made to initial experimental model :

No opiates were used as they could effect inflammatory response (45).

The organs were harvested after cervical dislocation to minimise premortem hypoxia (compared to rising concentration of CO₂).

As no laparotomy was performed there was no potential influence from the surgical trauma.

Confounding factors were reduced by using a single sex (male to exclude the oestrous cycle), strain of mice, and a single operator performing all procedures. Baseline and control groups were included to assess if the stress of the new environment for the mice had an effect. A mixture of procedures performed each day - as mice for different days could have been from different batches.

More fat depots were collected and analysed for more markers than in the previous set of experiments.

These experiments have led to a publication (46) in Pflugers Archiv : European journal of physiology which itself has been regularly cited - now (December 2012) by over 20 articles (see appendix for list and copy of paper) .

LPS METHOD

Male 8 - 10 week old, inbred, C57BL/6J mice (obtained from Charles River, UK) were used in all experiments.

Mice were kept in the animal house for at least one week prior procedures, as per departmental protocol, to assure fitness.

The care of the mice and all experimental procedures were approved by the UK Home Office and were conducted in accordance with the appropriate Project License.

The mice were weighed prior to the procedure.

Anaesthesia was induced with 1-2% isoflurane (Abbott, uk) in nitrous oxide / oxygen mixture (N₂O/O₂) and maintained with the same agents.

Prior to injection the proposed site was cleaned with iodine solution.

LPS (Escherichia coli O 111:B4, Sigma-Aldrich) was injected intraperitoneally (ip) under general anaesthesia at a dose of 25 mg/kg.

Control animals were administered equivalent volumes of normal saline, ip.

All animals received 1 ml of normal saline subcutaneously (sc) to compensate for fluid losses.

Post procedure the mice were housed in separate clean cages and maintained in the same temperature controlled (24-28°C) conditions with free access to standard laboratory chow and water.

The animals were exposed to the same standard 12 hour day / night conditions.

The mice were monitored regularly by a trained anaesthetist as per license requirements.

All mice were killed at 4 or 24 hours after the injection of LPS by cervical dislocation.

The epididymal, subcutaneous and perirenal fat deposits were removed and immediately frozen in liquid nitrogen and stored at -80°C till further analysis.




Fig.6: Flow diagram of procedures for the LPS model

LPS = Lipopolysaccharide, ip = intraperitoneal, sc = subcutaneous, N/S = Normal (0.9%) saline

RNA EXTRACTION AND PCR

Total RNA was isolated from the adipose tissues using Trizol reagent (Invitrogen, UK).

Extracted RNA was quantified spectrophotometrically at 260 nm. Ratio of extinction at 260 and 280nm was between 1.7 and 1.9, and the quality was assessed by visualization of 18s and 28s ribosomal bands on an agarose gel.

1 µg of DNase I-treated RNA was reverse transcribed using a Reverse-iT™ 1st Strand Synthesis Kit (Abgene, UK) in the presence of anchored oligo dT in a total volume of 20 µl. on a PCR Express thermal Cycler (Hybaid, Ashford, UK). Controls without reverse transcription were performed to exclude the possibility of DNA contamination.

Real-time polymerase chain reaction (PCR) reactions were carried out in a final volume of 12.5 µl consisting of 12.5-50 ng of reverse transcribed cDNA mixed with optimal concentrations of primers and probe and qPCR™ Core kit (Eurogentec, UK) in 96-well plates on a Mx3005P detector (Stratagene, USA). The primer and probe sets were designed using Primer Express software (Applied Biosystems) or Beacon Designer (Biosoft, USA) and synthesized commercially (Eurogentec). The sequence of primers and probes is shown in Table 2. Typically, the amplification started with 10 min at 95°C and then 40 cycles of the following: 15 s at 95°C and 1 min at 60°C. β-actin was used as an endogenous reference. Relative quantitation values were expressed using the 2-ΔΔCt method (see User bulletin #2, ABI Prism 7700, Applied Biosystems) as fold changes in the target gene normalized to the reference gene and related to the expression of the controls. The PCR efficiency in all runs was close to 100%, and all samples were analyzed in at least duplicate.

Gene	Sequence 5'-3'
Adiponectin	Forward GGCTCTGTGCTCCTCCATCT Reverse AGAGTCGTTGACGTTATCTGCATAG Probe CCCATACACCTGGAGCCAGACTTGGT
Adipsin	Forward ACGACCTCATTCTTTTTAAGCTATCC Reverse CCGGGTCCACTTCTTTGTC Probe TCCCCACGTGAGACCCCTACCCTT
β-actin	Forward CAAGAAGGAAGGCTGGAAAAG Reverse ACGGCCAGGTCATCACTATTG Probe ACGAGCGGTTCCGATGCCCTG
F4/80	Forward AAGACTTGATACTCCAAAGTGAGC Reverse GAAGGAAGCATAACCAAGATCCC Probe CCCTGCACTGCTTGGCATTGCTGT
HIF-1α	Forward CAAGTCAGCAACGTGGAAGGT Reverse CTGAGGTTGGTACTGTTGGTATCA Probe TTCACTGCACGGGCCATATTCATGTC
IL-6	Forward ACAACCACGGCCTTCCCTACTT Reverse CACGATTCCCAGAGAACATGTG Probe TTCACAGAGGATACCACTCCCAACAGAACCT
IL-18	Forward GGCCTCTATTTGAAGATATGACTGATT Reverse CCTCTAGGCTGGCTATCTTTATACATACT Probe TGA CTGTAGAGATAATGCACCCCGGACC
Leptin	Forward CATCTGCTGGCCTTCTCCAA Reverse ATCCAGGCTCTCTGGCTTCTG Probe AGCTGCTCCCTGCCTCAGACCAGTG
MAC-1	Forward GAATGGATTGTGCTATTTGTTCCGG Reverse CGGAGCCATCAATCAAGAAGAC Probe TCCAACCTGCTGAGGCCGCC
MCP-1	Forward GGCTCAGCCAGATGCAGTTAA Reverse CCTACTCATTGGGATCATCTTGCT Probe CCCCACTCACCTGCTGCTACTCATTCA
MT2	Forward CTTCTTGTCGCTTACACCGTT Reverse AGCAGCACTGTTGCTCACTTC
NGF	Forward GCCAAGGACGCAGCTTTCTAT Rev AGTGATCAGAGTGTAGAACAACATGGA Probe CTGGCCGCAGTGAGGTGC
TNFα	Forward CCCAGACCCTCACACTCAGATC Reverse GCCACTCCAGCTGCTCCTC Probe TAGCCCACGTCGTAGCAAACCAACCAAG

Table 2. Primer and probe sequences used in real time PCR (amended) (46)

MEASUREMENT OF PROTEIN BY ELISA

The total amount of HIF-1 α in adipose tissue homogenates was measured by enzyme-linked immunosorbent assay (ELISA) (R & D Systems, UK) as previously described. The adiponectin, IL-6 and MCP-1 content of adipose tissue homogenates was also measured using commercial ELISAs, according to the manufacturer's protocol. For adiponectin (R & D Systems), the stated minimal detectable amount was 3 pg/ml, and the interassay coefficient of variation was 6%. For IL-6 (RayBiotech, USA), the minimal detectable quantity was 6 pg/ml and the interassay coefficient of variation <12%, while for MCP-1 (RayBio- tech), the values were <3 pg/ml and <12%, respectively.

DATA PRESENTATION AND STATS

Data are expressed as the mean \pm standard error of mean. The statistical significance of differences between treated and control groups was assessed with two-tailed Student's t test for independent samples. A *p* value <0.05 was considered to be statistically significant.

RESULTS

Effect of LPS on adipokine expression 24 h after the induction of endotoxaemia

In order to induce severe endotoxaemia, a high dose of lipopolysaccharide was administered; the survival rate at 4 h was 100%, while the 24 h survival rate was 61% [14/23]. In the first experiments, the mice were treated with LPS for 24 h and the epididymal, perirenal and subcutaneous adipose tissue depots examined. Tissues were only analysed for those animals that survived. LPS administration had no significant effect on leptin mRNA level in the epididymal WAT; nor was there any change in metallothein-2 mRNA (Fig.7). However, LPS induced a fivefold fall in adiponectin mRNA level, and there was a similar reduction in the level of both adipisin and IL-18 mRNA (Fig.7). In contrast to these genes, there were increases in mRNA level for IL-6, MCP-1, NGF and TNF α (Fig.7). The increase in NGF mRNA was small (twofold), but MCP-1 and TNF α mRNA levels increased by 24- and 18-fold, respectively, while IL-6 mRNA level increased as much as 250-fold.




Fig. 7 Adipokine gene expression in endotoxaemia in mice (46). Mice were injected with LPS to induce severe sepsis (*closed bars*) and adipose tissue depots removed 24 h later; control mice (*open bars*) received a saline injection. The level of each adipokine mRNA was measured in epididymal (*Epi*), perirenal (*Peri*) and subcutaneous (*Sub*) depots by real-time PCR and normalized to β -actin relative to the untreated control group ($\equiv 1$). *IL-6* Interleukin 6, *IL-18* interleukin 18, *MCP-1* monocyte chemoattractant protein 1, *MT2* metallothionein 2, *NGF* nerve growth factor, *TNF α* tumour necrosis factor α . Results are given as mean values \pm SE for five to six mice in each group. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ compared with controls.

Adiponectin gene expression in epididymal fat pads previously published in M.Phil thesis by Honggwang Bao (47)

Examination of the perirenal adipose tissue depot gave broadly similar results, with reductions in adipisin and adiponectin mRNA levels (Fig.7). However, the changes were less substantial than with the epididymal depot, the decreases in IL-18 and adiponectin mRNA levels being statistically insignificant. Leptin was again unchanged following treatment with LPS. TNF α mRNA level increased by eightfold, MCP-1 by 40-fold and the largest increase (120-fold) was with IL-6 (Fig.7).

The results obtained with the subcutaneous adipose tissue were less dramatic than with either the epididymal or perirenal depots. Adiponectin mRNA level fell fivefold, but there were no significant changes with leptin, adipisin, IL-18, NGF, TNF α and MT-2 (Fig.7). The lack of change in TNF α mRNA is noteworthy given the increases observed in the other two fat depots. MCP-1 and IL-6 mRNA levels were, however, elevated in subcutaneous fat in response to LPS, the increases being four- and 20-fold, respectively (Fig.7).

Effect of LPS on adipokine expression 4 h after the induction of endotoxaemia

To assess the speed of the response to LPS, studies were next undertaken at 4 h after injection. Epididymal, perirenal and subcutaneous adipose tissue depots were again examined. In the epididymal depot at 4 h, there were no significant changes in adipisin, adiponectin, IL-18 and MT-2 mRNA levels (Fig.8). There were, however, small, but statistically significant, increases in leptin (fourfold) and NGF (3.8-fold) mRNAs. Substantial increases were observed for TNF α (20-fold) and particularly MCP-1 (250-fold) and IL-6 (500-fold) mRNA. A broadly similar pattern was observed in the perirenal depot, although in this case, there was no change in leptin

mRNA level. The largest changes were again with MCP-1 and IL-6 mRNA, the levels of which were increased by 35- and 78-fold, respectively (Fig.8). TNF α level increased 23-fold, and a modest increase (fourfold) was observed with NGF.




Fig. 8 Acute effects of endotoxaemia on adipokine gene expression in mice (46). Mice were injected with LPS to induce severe sepsis (*closed bars*) and adipose tissue depots removed 4 h later; control mice (*open bars*) received a saline injection. The level of each adipokine mRNA was measured in epididymal (*Epi*), perirenal (*Peri*) and subcutaneous (*Sub*) depots by real-time PCR and normalized to β -actin relative to the untreated control group ($\equiv 1$). Results are given as mean values \pm SE for five to six mice in each group. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ compared with controls.

Adiponectin and TNF α gene expression in epididymal fat pads previously published in M.Phil thesis by Honggwang Bao (47)

In the subcutaneous adipose tissue, there were no changes in mRNA levels for IL-18, leptin and MT-2 (Fig.8). There was, however, an approximately 50% reduction in mRNA levels for adipsin and adiponectin. NGF and TNF α mRNA levels increased moderately, by four- and 17-fold, respectively. Very major changes were observed with MCP-1, the mRNA level of which increased 250-fold, and with IL-6 where the increase was >1,000-fold (Fig.8).

White adipose tissue adipokine levels

The adipokine protein content was measured by ELISA in the epididymal and subcutaneous fat depots for the two inflammatory adipokines which exhibited the largest changes in mRNA level. IL-6, which showed the greatest elevation in mRNA, was increased in both the epididymal and subcutaneous fat at 4 and 24 h after the injection of LPS (Fig.9). The increase in the epididymal depot was between five- and eightfold, while in the subcutaneous tissue the rise was between three- and eightfold.




Fig. 9 Adipokine levels in adipose tissue in endotoxaemia in mice (46). Mice were injected with LPS to induce severe sepsis (*closed bars*) and adipose tissue depots removed 4 or 24 h later; control mice (*open bars*) received a saline injection. Adipokine proteins were measured in the epididymal (*Epi*) and subcutaneous (*Sub*) depots by ELISA. Results are given as mean values \pm SE for three to six mice in each group. * $P < 0.05$, ** $P < 0.01$ compared with controls.

In contrast to IL-6, there was no significant increase in MCP-1 protein in either tissue at 24 h. However at 4 h, the amount of MCP-1 was approximately eightfold greater in the epididymal fat and twofold greater in the subcutaneous depot (Fig. 9). For adiponectin, only in the epididymal depot after 24 h of treatment with LPS was there a significant fall in the protein, reflecting the changes in the mRNA level. However, paradoxically the adiponectin level in the subcutaneous depot was actually increased, rather than decreased, at both 4 and 24 h in the LPS-treated mice.

Macrophage marker expression

To assess whether the increases in the expression of key inflammatory adipokine genes, particularly MCP-1, reflected an infiltration of macrophages into adipose tissue, the mRNA level of two macrophage markers was examined. The markers were F4/80 and MAC-1, and these were examined in 4 and 24 h LPS-treated mice and in all three WAT depots. In no case was there a statistically significant increase in F4/80 and MAC-1 mRNA level (Fig.10). Indeed, in one case, the perirenal tissue treated for 24 h, the level of both mRNAs was significantly decreased. F4/80 mRNA level was also decreased in the epididymal depot at 24 h.




Fig. 10 Expression of macrophage markers in adipose tissue in endotoxaemia in mice (46). Mice were injected with LPS to induce endotoxaemia (*closed bars*) and adipose tissue depots removed 4 or 24 h later; control mice (*open bars*) received a saline injection. The level of each macrophage marker mRNA was measured in epididymal (*Epi*), perirenal (*Peri*) and subcutaneous (*Sub*) depots by real-time PCR and normalized to β -actin relative to the untreated control group ($\equiv 1$). Results are given as mean values \pm SE for five to six mice in each group. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ compared with controls.

HIF-1 α as an index of hypoxia

In the final investigation, the effect of LPS treatment on the key molecular index of cellular oxygen deficiency, the hypoxia-sensitive transcription factor HIF-1 α , was examined. The expression and protein level of HIF-1 α was determined in both the epididymal and subcutaneous adipose tissue depots. At 4 h after the treatment with LPS, there was a three to fourfold increase in HIF-1 α mRNA in each depot, and importantly, this was accompanied by a two to threefold increase in HIF-1 α protein (Fig.11). HIF-1 α mRNA level was also twofold higher in both depots at 24 h, but at this later time point, there was no significant change in HIF-1 α protein (Fig.11).




Fig. 11 HIF-1 α gene expression and protein levels in adipose tissue in endotoxaemia in mice (46). Mice were injected with LPS to induce endotoxaemia (*closed bars*) and adipose tissue depots removed 4 or 24 h later; control mice (*open bars*) received a saline injection. HIF-1 α mRNA (**a**) and protein levels (**b**) were measured in the epididymal (*Epi*) and subcutaneous (*Sub*) depots by real-time PCR, the level being normalized to β -actin relative to the untreated control group ($\equiv 1$), and by ELISA, respectively. Results are given as mean values \pm SE for five to six mice in each group. * $P < 0.05$, ** $P < 0.01$ compared with controls.

DISCUSSION

This study demonstrates that there are major changes in the expression of key adipokine genes in each of the main white adipose tissue depots in mice following the induction of endotoxaemia. The expression of several major inflammatory-related adipokines was examined, and by far the largest response was observed with IL-6 and MCP-1. Adipose tissue produces large amounts of IL-6 and is considered to be a substantial contributor to circulating levels of this cytokine (48, 49). Plasma IL-6 levels are increased in endotoxaemia as well as sepsis, and it seems likely that adipose tissue contributes directly to this (50-52), particularly since in this study the amount of IL-6 in the fat depots was increased alongside the increase in mRNA level. Nevertheless, it should be noted that the increase in protein was much less than the changes in mRNA level, although this could reflect rapid release of IL-6 from the tissue once synthesised.

The chemokine MCP-1 shows increased expression in obesity, like IL-6, and is linked to the recruitment of macrophages within tissues, including adipose tissue. Indeed, in obesity there is a major infiltration of macrophages into adipose tissue (53) (14). Since the mRNA level of two macrophage markers—F4/80 and MAC-1—did not increase in any of the three adipose tissue depots examined, either at 4 or 24 h after the administration of LPS, it is unlikely that there was an infiltration of immune cells during the duration of endotoxaemia in the present work. The increased MCP-1 expression suggests that macrophage recruitment into adipose tissue may well occur long-term; however, in contrast to IL-6, no sustained elevation in MCP-1 protein levels was evident in adipose tissue homogenates following LPS

treatment.

In addition to macrophage infiltration, MCP-1 has been associated with the development of insulin resistance (54). This is also the case with IL-6 and TNF α (55) (56-58). Thus, the major stimulation of the expression of these three factors in response to LPS suggests that adipose tissue may contribute to insulin insensitivity and the dysregulation of glucose homeostasis in infection. TNF α is a pleiotrophic pro-inflammatory cytokine, and its actions in adipose tissue include the stimulation of lipolysis and of apoptosis (59) (60). It also has a major effect on the expression of other inflammation-related adipokines including the strong upregulation of IL-6 and MCP-1 production (61, 62).

In marked contrast, adiponectin expression was downregulated in all three adipose tissue depots 24 h after the administration of LPS. There was also a significant reduction in adiponectin mRNA in the subcutaneous depot at 4 h. Adiponectin, which is a key adipocyte-derived hormone, has multiple physiological effects, and these include anti-inflammatory and insulin sensitising actions (63-66). Thus, a fall in adiponectin expression may add to the upregulation of IL-6, MCP-1 and TNF α expression in leading to a major pro-inflammatory response and insulin resistance in endotoxaemia. However, although adiponectin protein level decreased in the epididymal depot at 24 h (the absence of a change at 4 h may reflect the time delay between alterations in gene transcription and subsequent protein synthesis), in the case of the subcutaneous depot, there was a paradoxical increase in the amount of the hormone. We have no satisfactory explanation for this anomaly, although it may indicate that adiponectin is less readily released from the subcutaneous adipose tissue in endotoxaemia.

Furthermore, low adiponectin levels observed in obesity (67) as a state of low-grade chronic inflammation are associated with increased insulin resistance (63, 66). Low adiponectin levels on admission to the intensive care unit have also been identified as an independent predictor of survival (68). Stress-induced hyperglycaemia frequently occurs in infection and is associated with an adverse outcome (69) (70). Elevated TNF α levels, increased MCP-1 production as well as low adiponectin may act in concert to induce and to maintain hyperglycaemia in critical illness. Although this study does not reveal the extent to which adipose tissue might contribute quantitatively to alterations in circulating levels of cytokines and chemokines in endotoxaemia, overall, the results suggest a role for the tissue in the changes in glucose metabolism associated with infection. It is emphasised that since in the 24-h studies there was a distinct mortality rate (39%) and that only tissues from surviving animals were investigated, there could be a link between the gene expression profile and the protective survival response.

Expression of the complement-related factor, adipsin, also fell during endotoxaemia. Leptin, on the other hand, showed no changes in gene expression. This is in contrast with previous studies in experimental animals and in humans where an upregulation has been described (50, 71) (72). However, the previously reported changes in leptin mRNA were very small, particularly in comparison with the dramatic changes in IL-6 and MCP-1 mRNA level.

Small increases in NGF mRNA level were observed in the epididymal and perirenal depots but not in the subcutaneous tissue. This target-derived neurotrophin is also involved in the inflammatory response, its expression and secretion from both murine and human adipocytes being stimulated by TNF α (73) (61). The low molecular weight

cysteine-rich, metal-binding protein metallothionein showed little response to LPS except in the perirenal depot where there was a small increase at 24 h. Similarly, there was little change in the expression of IL-18; this pro-inflammatory cytokine plays an important role in innate and acquired immunity, and its expression in adipocytes (74, 75) has been shown to be strongly upregulated by TNF α (75).

Certain depot-specific responses to LPS were evident in this study. Of the three adipose tissue sites examined, the largest response was observed in the epididymal depot in the case of both IL-6 and TNF α (at 24 h). Indeed, the subcutaneous depot showed no increase in TNF α mRNA level at 24 h. In this context, it has been shown that subcutaneous adipose tissue is affected to a lesser degree by sepsis-induced malperfusion than the splanchnic area and does not contribute to lactate production during critical illness (15).

In addition to depot differences, there is the important question of which cells within adipose tissue may be responsible for alterations in adipokine/cytokine production in endotoxaemia. White fat consists of two major components, mature adipocytes and the stromal-vascular fraction. The stromal-vascular fraction includes macrophages, fibroblasts, vascular endothelial cells and preadipocytes. Adipocytes are able to express each of the inflammation-related signals examined in the present study (see (16) (29)). However, while the key cytokines and chemokines are also expressed by macrophages, adipocytes alone are responsible for adiponectin and leptin production.

It could be hypothesized that during endotoxaemia, macrophages infiltrate adipose tissue, causing further inflammatory damage and substantially contributing to mediator production. This would imply

that immigrating immune cells are mainly responsible for the changes observed. However, as noted earlier, the mRNA level of two specific macrophage markers did not increase in WAT and even decreased in the case of the perirenal depot at 24 h. This suggests that mature adipocytes may be the key site of the response to LPS. Indeed, adipocytes express the Toll-like receptor 4 through which LPS exerts its inflammatory effect (76) (77). Irrespective of the cell type involved, adipose tissue can be regarded as a key inflammatory organ, which may contribute to the systemic response observed during systemic infection by generating and releasing inflammatory mediators into the circulation. However, the possibility that resident macrophages are activated and may contribute to adipokine/cytokine production in the various adipose tissue depots in endotoxaemia, without increased immune cell infiltration, cannot be excluded. It would be useful to investigate the adipose tissue using immunohistochemistry techniques that are able to preserve the architecture of the tissue to help answer the question.

It has recently been proposed that the inflammation that occurs in adipose tissue in obesity may, at least in part, be a response to relative hypoxia in groups of adipocytes distant from the vasculature (16). Evidence in support of this hypothesis is emerging (78) (79) (80), and hypoxia has been shown to directly increase the expression and release of inflammation-related adipokines such as leptin, IL-6, MIF and VEGF in murine and human adipocytes, while adiponectin with its anti-inflammatory action is decreased (81) (78) (82) (83) (80). We were therefore interested in whether hypoxia might underlie the changes in adipokine expression in endotoxaemia and hence fat depots were harvested after cervical dislocation to minimise premortem hypoxia (compared to rising concentration of CO₂). Increases in the mRNA encoding the hypoxia-sensitive transcription factor, HIF-1 α ,

were evident following administration of LPS, and increased HIF-1 α protein was found at 4 h. This suggests that at least in the early stages of endotoxaemia, hypoxia may occur in adipose tissue. The changes at the protein level were not sustained by 24 h, implying that any hypoxia may have been transitory. However, direct HIF-1 α gene activation by LPS, in a Toll-like receptor 4 manner, has been shown in macrophages (84) (85). Further work is therefore warranted on hypoxia in adipose tissue in severe infection and whether it underlies the changes in adipokine expression or if it is by direct stimulation by LPS / bacterial components. It may be possible to investigate this using gene arrays to look for specific gene expression profiles (37) at different time intervals.

A high dose of LPS was employed to induce endotoxaemia, and it may also reflect sepsis. However, the LPS challenge may not adequately reproduce the complexity of human sepsis. In particular, investigators with an interest in peri-operative sepsis have increasingly used caecal ligation and puncture (CLP) as the model of choice. In the series of experiments with CLP in the same strain of mice, it was possible to reproduce the changes in IL-6 mRNA levels found in the epididymal fat depot after LPS challenge. Here IL-6 mRNA levels were 87-fold higher than in control mice and sixfold higher than in sham-operated mice at 24 h. Apparently, CLP leads to qualitatively similar changes in IL-6 expression as LPS, suggesting that white adipose tissue is activated in septic shock as in endotoxaemia.

Clinical sepsis is commonly polymicrobial, but experimentation involving bacterial products, such as LPS, may serve to characterize response patterns directed against one particular organism (42). LPS was therefore used as the model of choice to examine the potential

role of WAT during severe infection. The dose employed, although high, is within the range used in previous studies and elicited a severe, acute, systemic inflammatory response in mice with all the symptoms of sepsis and septic shock (86). Importantly, the substantial increase in IL-6 and MCP-1 gene expression in the subcutaneous adipose tissue at 4 h indicates a systemic response in WAT rather than an effect local to the i.p. site of injection. The mortality rate of 39% in the 24-h series of experiments reflects acute life-threatening gram-negative infection of a severity similar to that occurring in human sepsis (3). Further investigations using various models of sepsis are warranted to specify the role of WAT in infection caused by different organisms.

In conclusion, the results demonstrate that an LPS challenge sufficient to mimic endotoxaemia leads to major changes in the expression of key adipokine genes in different adipose tissue depots of mice. Thus, white adipocytes (and preadipocytes) may well contribute directly to the most prominent features of infection – systemic inflammation and metabolic dysregulation. If this could be corroborated in studies involving human patients and their various fat depots, it would allow a new interpretation of the correlation between body mass index and sepsis mortality as well as the development of novel approaches to treating insulin resistance in severe infection. It is evident from our results that not all fat is equal and it would be valuable to investigate the differences between depots in the human population with samples obtained at laparotomy in patients with sepsis or chronic inflammatory states such as pancreatitis. It may also be possible to look for hypoxic responses in these tissues using gene arrays. Quantification of the size of various fat depots in septic ITU patients from their CT images and their possible relation to outcome would also be interesting.

APPENDIX

LPS MOUSE EXPERIMENTS

MOUSE NUMBER	WEIGHT	EXPT LENGTH	PROCEDURE	SURVIVAL	OTHER
1	24g	24	LPS	Y	
2	25.4g	24	LPS	Y	
3	25.5g	24	LPS	Y	
4	24.6g	24	SALINE	Y	
5	24.6g	24	SALINE	Y	
6	25g	24	SALINE	Y	
7	22.8g	24	CONTROL	Y	
8	24.5g	24	CONTROL	Y	
9	25.8g	24	CONTROL	Y	
10		24	BASLINE	Y	
11		24	BASLINE	Y	
12		24	BASLINE	Y	
13	27.5g	24	LPS	Y	
14	23g	24	LPS	Y	
15	24.5g	24	LPS	N	DIED 11HOURS AFTER THE INJECTION
16	26.5g	24	SALINE	Y	
17	24g	24	SALINE	Y	
18	23g	24	SALINE	Y	
19	22g	24	CONTROL	Y	
20	24g	24	CONTROL	Y	
21	25g	24	CONTROL	Y	
22	26g	24	BASLINE	Y	
23	26.4g	24	BASLINE	Y	
24	23.4g	24	BASLINE	Y	
25	26.6g	24	LPS	Y	
26	23.9g	24	LPS	N	DIED 21HOURS AFTER THE INJECTION
27	23.5g	24	LPS	Y	
28	25.4g	24	LPS	N	DIED 19 HOURS AFTER THE INJECTION
29	24.5g	24	SALINE	Y	
30	26g	24	SALINE	Y	
31	24.9g	24	CONTROL	Y	
32	24g	24	CONTROL	Y	
33		24	BASLINE	Y	
34		24	BASLINE	Y	
35	27g	4	LPS	Y	
36	22.4g	4	LPS	Y	
37	25.4g	4	SALINE	Y	
38	24.9g	4	SALINE	Y	
39	26.7g	4	CONTROL	Y	
40	24.5g	4	CONTROL	Y	
41	23.5	4	LPS	Y	
42	26.5	4	LPS	Y	
43	21.5	4	SALINE	Y	
44	24	4	SALINE	Y	
45	22.5	4	CONTROL	Y	
46	24.5	4	CONTROL	Y	

47	25.5	4	LPS	Y	4 Hours later weight 25.9
48	24	4	LPS	Y	4 Hours later weight 23.3
49	23	4	SALINE	Y	4 Hours later weight 22.9
50	26	4	SALINE	Y	4 Hours later weight 26.2
51	25	4	CONTROL	Y	4 Hours later weight 24.4
52	24.5	4	CONTROL	Y	4 Hours later weight 23.4
		Schedule 1			DREW BLOOD BY CARDIAC PUNCTURE UNDER GA
		Schedule 1			DREW BLOOD BY CARDIAC PUNCTURE UNDER GA
53	24.5	24	LPS	N	DIED 20 HOURS AFTER INJECTION
54	26.7	24	LPS	Y	
55	24.8	24	LPS	Y	
56	24.8	24	LPS	N	DIED 17.5 HOURS AFTER INJECTION
57	26.4	24	LPS	Y	
58	24.5	24	LPS	N	DIED 18 HOURS AFTER INJECTION
59	28.7	24	SALINE	Y	Large cystic tumour in abdomen - SAMPLES NOT TAKEN
60	25	24	SALINE	Y	
61	25.1	24	SALINE	Y	
62	26	24	SALINE	Y	
63	26.4	24	LPS	N	DIED 14 HOURS AFTER INJECTION
64	26.4	24	LPS	Y	
65	25.3	24	LPS	Y	
66	24.1	24	LPS	Y	
67	26.4	24	LPS	N	DIED 19.5 HOURS AFTER INJECTION
68	26.9	24	LPS	Y	
69	26.5	24	LPS	N	DIED 17 HOURS AFTER INJECTION
70	25	24	SALINE	Y	
71	26.5	24	SALINE	Y	
72	25.1	24	SALINE	Y	

European Journal of Anaesthesiology:
May 2005 - Volume 22 - Issue - p 177
Intensive Care Medicine

A-683

A possible role of white adipose tissue in sepsis

M. Leuwer, G. Marx, A. Rushton, A. Axon, G. Masterson, L. Hunter, P. Trayhurn
*Department of Anaesthesia, The University of Liverpool, Liverpool,
United Kingdom*

Background and Goal of Study: It has recently been established that white adipose tissue, besides its metabolic role, storing and releasing fatty acids, has a major role in secreting a variety of hormones and mediators which are referred to as adipokines.⁽¹⁾ In particular, it has been suggested that adipose tissue releases pro-inflammatory cytokines, e.g. IL-6 in response to hypoxia. Thus we felt tempted to investigate how adipokine expression relates to sepsis in an animal model.

Methods: After obtaining a Home Office licence under the Animal Rights Act, we investigated sixteen healthy, non-obese male mice (body weight 25–28 g). Eight animals served as controls, in another eight animals, sepsis was induced by intra-peritoneal LPS injection. Twenty-four hours after induction of sepsis, the experimental animals were killed and RNA extracted from epididymal fat was investigated by real time PCR, in order to quantify expression of IL-6, IL-18, nerve growth factor (NGF), hypoxia-induced factor (HIF), adiponectin and adiponectin in comparison to controls.

Results: Twenty-four hours after induction of sepsis, real time PCR revealed significant increases in the expression of IL-6 (500 fold), NGF (10 fold), TNF (5 fold) and HIF (3 fold), while there were significant decreases in the expression of IL-18 (0.4 fold), adiponectin (0.15 fold) and adiponectin (0.2 fold). **Conclusions:** These results show that white adipose tissue is an organ system which actually suffers from hypoxia during sepsis. It responds by a marked increase in the expression of pro-inflammatory cytokines. At the same time, the expression of anti-inflammatory cytokines as well as the expression of adiponectin is reduced. Thus, we propose the hypothesis that white adipose tissue may be an important contributor to the pathophysiology of sepsis.

Reference:

1 Trayhurn P and Wood IS. Adipokines: inflammation and the pleiotropic role of white adipose tissue. *British Journal of Nutrition* 2004; 92: 347–355.



School of Clinical
Science
University of
Liverpool, UK

ADIPOKINE EXPRESSION IN A MURINE MODEL OF SEPSIS

*Rushton A, Leuwer M, Axon A,
Marx G, Masterson G, Hunter L,
Trayhurn P.

INTRODUCTION

It has recently been established that white adipose tissue, besides its metabolic function in storing and releasing fatty acids, has a major role in secreting a variety of hormones and protein factors which are referred to as adipokines. (1) In particular, it has been suggested that adipose tissue releases pro-inflammatory cytokines, e.g. IL-6, in response to hypoxia. Thus we felt tempted to investigate whether adipokine expression relates to sepsis in an animal model.

METHODS

After obtaining a Home Office licence, we investigated thirty-two healthy, non-obese male mice (body wt. 25–28 g). Eight animals served as controls; in eight animals, sepsis was induced by intra-peritoneal LPS injection (25 µg/g body wt), in another eight animals by caecum ligation/puncture (CLP) and eight animals were sham-operated. Twenty-four hours after induction of sepsis, the experimental animals were killed and RNA extracted from the epididymal fat pads. Real time PCR was used to quantify expression of IL-6, TNF α , IL-18, MCP-1, nerve growth factor (NGF), hypoxia-induced factor (HIF-1 α), adipsin, adiponectin and leptin, in comparison to controls.

RESULTS


Twenty-four hours after induction of sepsis by LPS, real time PCR revealed substantial increases in the expression of IL-6 (500-fold), MCP-1 (55-fold), NGF (10-fold), TNF α (5-fold) and HIF-1 α (3-fold), while there were significant decreases in the expression of IL-18 (2.5-fold), adipsin (6-fold) and adiponectin (5-fold). The results in the CLP-group were qualitatively similar.

CONCLUSIONS

These results show that white adipose tissue is an organ system which responds to sepsis by a marked increase in the expression of pro-inflammatory cytokines. At the same time, the expression of anti-inflammatory cytokines as well as the expression of adipsin and adiponectin (anti-inflammatory) is reduced. Thus, we propose the hypothesis that white adipose tissue may be an important contributor to the pathophysiology of sepsis, and that this may relate, at least in part, to hypoxia within the tissue.

REFERENCES

(1) Trayhurn P and Wood IS. Adipokines: inflammation and the pleiotropic role of white adipose tissue. *British Journal of Nutrition* 2004; 92: 347-355



*p<0.05 **p<0.01 ***p<0.001 ****p<0.0001

Poster presentation at 2nd International Congress of the
German Sepsis Society (GSS) in Weimar, Germany ,
September 7 - 10 , 2005.

Endotoxaemia leads to major increases in inflammatory adipokine gene expression in white adipose tissue of mice

Martin Leuwer · Ingeborg Welters · Gernot Marx ·
Andrew Rushton · Hongguang Bao · Leif Hunter ·
Paul Trayhurn

Received: 6 June 2008 / Revised: 17 July 2008 / Accepted: 25 July 2008 / Published online: 2 August 2008
© Springer-Verlag 2008

Abstract The proposition that white adipose tissue is involved in the inflammatory response and metabolic dysregulation of endotoxaemia has been examined. Mice were injected with lipopolysaccharide (LPS; 25 mg/kg) and epididymal, perirenal and subcutaneous adipose tissue removed 4 or 24 h later. The expression of genes encoding key inflammation-related adipokines was measured by real-time polymerase chain reaction. At 24 h after the administration of LPS, there was no change in leptin mRNA level, and adiponectin mRNA fell. However, major increases in TNF α , MCP-1 (up to 40-fold) and IL-6 (up to 250-fold) mRNA levels were evident; a substantial elevation in these mRNAs occurred by 4 h, and adipose tissue IL-6 protein also increased (three- to eightfold). At 24 h, the responses in the subcutaneous depot were much lower than in epididymal and perirenal adipose tissue, but at 4 h, the subcutaneous tissue showed major increases in IL-6, MCP-1 and TNF α gene expression. In contrast to the inflammatory adipokines, the mRNA level of two macrophage markers, F4/80 and MAC-1, was unaltered in adipose

tissue during endotoxaemia. Expression of the hypoxia-sensitive transcription factor, HIF-1 α , gene was increased at both 4 and 24 h, and HIF-1 α protein was elevated at 4 h, suggesting that the tissue was hypoxic. It is concluded that white adipose tissue may play an important role in the production of inflammatory mediators in endotoxaemia.

Keywords Adipokines · Adipose tissue · Endotoxaemia · IL-6 · Inflammation · Sepsis

Introduction

Sepsis, septic shock and endotoxaemia are systemic inflammatory states that affect most organs of the body, including white adipose tissue, and glucose metabolism and lipolytic capacity are particularly affected [21, 43]. It has been demonstrated that obese patients have a worse outcome from critical illness than non-obese individuals [33, 55]. To date, this has mainly been regarded as a result of difficulties in and complications of ventilation as well as an increase in the incidence of comorbidities such as diabetes and cardiac disease. While it is well established that, in obesity [54], as well as during endotoxaemia and sepsis, lipolysis is altered and glucose utilisation is impaired [25], it has recently become clear that the potential mediatory role of white adipose tissue (WAT) is not confined to glucose and fat metabolism. WAT has become recognized as a major endocrine and secretory organ which contributes to inflammation-related diseases in obesity, such as the metabolic syndrome, by the secretion of protein mediators and signalling factors called adipokines [16, 34, 46].

In obesity, the plasma concentration of a number of adipokines and proinflammatory markers expressed and

M. Leuwer (✉) · I. Welters · A. Rushton · H. Bao
Critical Care Research Unit, School of Clinical Sciences,
University of Liverpool, University Clinical Departments,
Duncan Building,
Liverpool L69 3GA, UK
e-mail: mleuwer@liverpool.ac.uk

L. Hunter · P. Trayhurn
Obesity Biology Research Unit, School of Clinical Sciences,
University of Liverpool, University Clinical Departments,
Duncan Building,
Liverpool L69 3GA, UK

G. Marx
University Clinic for Anaesthesiology and Intensive Care,
University of Jena,
Jena, Germany

released by adipocytes has been shown to rise as adipose tissue mass expands. These include cytokines (e.g. IL-6, IL-10), chemokines (e.g. IL-8, MCP-1, MIP) and acute-phase proteins (e.g. haptoglobin, serum amyloid A) as well as inflammation-related adipokines such as leptin and nerve growth factor (NGF) [16, 34, 45, 46]. It has been suggested that these inflammatory changes may reflect hypoxia within the growing adipose tissue mass [46], and it is well recognised that in organs such as heart, kidneys and gut, hypoxia can induce the synthesis of cytokines and inflammatory mediators [40]. Sepsis is considered to be a state of generalised microvascular dysfunction with a consequent decrease in oxygen delivery, which is associated with tissue hypoxia and organ dysfunction [10]. The key intracellular pathway in hypoxia-induced metabolic changes involves activation of the transcription factor, HIF-1 [7, 39, 40]. Activation of HIF-1, through stabilisation of the HIF-1 α subunit, during hypoxia triggers expression of glycolytic enzymes and membrane glucose transporters, such as GLUT-1 [7].

Previous studies have indicated that inflammation induced by lipopolysaccharide (LPS) and other agents leads to an increase in leptin expression and circulating levels in experimental animals [11, 12, 37], and increased circulating levels of the hormone have been observed in inflammatory conditions in humans [1, 4, 6, 44]. Evidence has also been presented for increases in IL-6 expression in humans during experimentally induced inflammation [1, 19]. In the present study, we hypothesized that in endotoxaemia there is a major activation of the inflammatory response in adipose tissue, key inflammatory adipokines contributing to the metabolic dysregulation characteristic of endotoxaemia and sepsis. A murine model of endotoxaemia was used to study the expression in both internal and subcutaneous adipose tissue of a series of inflammation-related adipokines. Evidence for the induction of adipose tissue hypoxia was sought through measurement of HIF-1 α .

Materials and methods

Animals and tissue collection

All experiments were carried out on 8- to 10-week-old male C57BL/6J mice (Charles River, UK). The care of the mice and all experimental procedures were approved by the UK Home Office and were conducted in accordance with the appropriate Project License. LPS (*Escherichia coli* O 111: B4, Sigma-Aldrich) was injected intraperitoneally (ip) under general anaesthesia (2% isoflurane in N₂O/O₂) at a dose of 25 mg/kg. Control animals were administered equivalent volumes of normal saline, ip. All animals

received 1 ml of normal saline subcutaneously (sc) to compensate for fluid losses. Mice were housed in separate cages post-procedures and maintained in the same temperature-controlled conditions with free access to standard laboratory chow and water. All mice were killed at 4 or 24 h after injection of LPS by cervical dislocation. A 24-h time point is commonly used in studies on LPS-induced endotoxaemia [12], providing a sufficient period for a severe response to be established, while 4 h [11, 37] was used to investigate acute effects. The epididymal, subcutaneous and perirenal fat depots were removed and immediately frozen in liquid nitrogen.

Real-time polymerase chain reaction

Total RNA was extracted from adipose tissues with Trizol reagent (Invitrogen, UK), and 1 μ g of DNase I-treated RNA was reverse transcribed using a Reverse-iTTM 1ST Strand Synthesis Kit (Abgene, UK) in the presence of anchored oligo dT in a total volume of 20 μ l.

Real-time polymerase chain reaction (PCR) reactions were carried out in a final volume of 12.5 μ l consisting of 12.5–50 ng of reverse transcribed cDNA mixed with optimal concentrations of primers and probe and qPCRTM Core kit (Eurogentec, UK) in 96-well plates on a Mx3005P detector (Stratagene, USA). The primer and probe sets were designed using Primer Express software (Applied Biosystems) or Beacon Designer (Biosoft, USA) and synthesized commercially (Eurogentec). The sequence of primers and probes is shown in Table 1.

Typically, the amplification started with 10 min at 95°C and then 40 cycles of the following: 15 s at 95°C and 1 min at 60°C. β -actin was used as an endogenous reference. Relative quantitation values were expressed using the 2^{- $\Delta\Delta$ C_t} method (see User bulletin #2, ABI Prism 7700, Applied Biosystems) as fold changes in the target gene normalized to the reference gene and related to the expression of the controls. The PCR efficiency in all runs was close to 100%, and all samples were analyzed in at least duplicate.

Measurement of protein by enzyme-linked immunosorbent assay

The total amount of HIF-1 α in adipose tissue homogenates was measured by enzyme-linked immunosorbent assay (ELISA) (R & D Systems, UK) as previously described [51]. The adiponectin, IL-6 and MCP-1 content of adipose tissue homogenates was also measured using commercial ELISAs, according to the manufacturer's protocol. For adiponectin (R & D Systems), the stated minimal detectable amount was 3 pg/ml, and the interassay coefficient of variation was 6%. For IL-6 (RayBiotech, USA), the minimal

Table 1 Primer and probe sequences used in real time PCR

Gene	Sequence 5'-3'	
Adiponectin	Forward	ACGACCTCAATCTTTTAAAGCTATCC
	Reverse	TCCCCACGTGAGACCCCTACCCTT
	Probe	CCGGTTCACACTTCTTTGGTC
Adipsin	Forward	ACGACCTCAATCTTTTAAAGCTATCC
	Reverse	CCGGTTCACACTTCTTTGGTC
	Probe	TCCCCACGTGAGACCCCTACCCTT
β -actin	Forward	CAAGAAGGAAGGCTGGAAAAG
	Reverse	ACGCCAGGTCATCACTATTG
	Probe	ACGAGCGGTTCCGATGCCCTG
F4/80	Forward	AAGACTTGATACTCCAAAGTGAGC
	Reverse	GAAGGAAGCATAACCAAGATCCC
	Probe	CCCTGCACCTGTTGGCATTGCTGT
HIF-1 α	Forward	CAAGTCAGCAACGTGGAAAGGT
	Reverse	CTGAGGTTGGTACTGTTGGTATCA
	Probe	TTCACCTGCACGGCCATATCATGTC
IL-6	Forward	CATCTGCTGGCCCTTCTCCAA
	Reverse	CAGGCTCTCTGGCTTCTG
	Probe	AGCTGCTCCCTGCCTCAGACCAAGT
IL-18	Forward	GGCCTCTATTGGAAGATATGAC
	Reverse	CTCTAGGCTGGCTATCTTATA
	Probe	TGACTGTAGAGATAATGCACCCC
Leptin	Forward	CATCTGCTGGCCCTTCTCCAA
	Reverse	ATCCAGGCTCTCTGGCTTCTG
	Probe	AGCTGCTCCCTGCCTCAGACCAAGT
MAC-1	Forward	GAATGGATTGTGCTATTTGTTCCGG
	Reverse	CGGAGCCATCAATCAAGAAGAC
	Probe	TCCAACCTGCTGAGGCCGCC
MCP-1	Forward	GGCTCAGCCAGATGCAGTAA
	Reverse	CCTACTATTGGGATCATCTTGCT
	Probe	CCCCACTCACCTGCTGCTACTCA
MT2	Forward	CTTCTGTGCGCTTACACCGTT
	Reverse	AGCAGCACTGTTGCTCACTTC
	Probe	GCCAAAGGACGCACTTCTAT
NGF	Forward	AGTGATCAGAGGTGAGAACAACA
	Rev	TGGA
	Probe	CTGGCCGCACTGAGGTTGC
TNF α	Forward	CCCAGACCCTCACACTCAGATC
	Reverse	GCCACTCCAGTGCCTC
	Probe	TAGCCCACGTCGTAGCAACCAC
		CAAG

detectable quantity was 6 pg/ml and the interassay coefficient of variation <12%, while for MCP-1 (RayBiotech), the values were <3 pg/ml and <12%, respectively.

Statistical analysis

The statistical significance of differences between treated and control groups was assessed with Student's *t* test for independent samples. *P*<0.05 was considered significant.

Results

Effect of LPS on adipokine expression 24 h after the induction of endotoxaemia

In order to induce severe endotoxaemia, a high dose of lipopolysaccharide was administered; the survival rate at 4 h was 100%, while the 24 h survival rate was 61%. In the first experiments, the mice were treated with LPS for 24 h and the epididymal, perirenal and subcutaneous adipose tissue depots examined. Tissues were only analysed for those animals that survived. LPS administration had no significant effect on leptin mRNA level in the epididymal WAT; nor was there any change in metallothein-2 mRNA (Fig. 1). However, LPS induced a fivefold fall in adiponectin mRNA level, and there was a similar reduction in the level of both adipsin and IL-18 mRNA (Fig. 1). In contrast to these genes, there were increases in mRNA level for IL-6, MCP-1, NGF and TNF α (Fig. 1). The increase in NGF mRNA was small (two-fold), but MCP-1 and TNF α mRNA levels increased by 24- and 18-fold, respectively, while IL-6 mRNA level increased as much as 250-fold.

Examination of the perirenal adipose tissue depot gave broadly similar results, with reductions in adipsin and adiponectin mRNA levels (Fig. 1). However, the changes were less substantial than with the epididymal depot, the decreases in IL-18 and adiponectin mRNA levels being statistically insignificant. Leptin was again unchanged following treatment with LPS. TNF α mRNA level increased by eightfold, MCP-1 by 40-fold and the largest increase (120-fold) was with IL-6 (Fig. 1).

The results obtained with the subcutaneous adipose tissue were less dramatic than with either the epididymal or perirenal depots. Adiponectin mRNA level fell fivefold, but there were no significant changes with leptin, adipsin, IL-18, NGF, TNF α and MT-2 (Fig. 1). The lack of change in TNF α mRNA is noteworthy given the increases observed in the other two fat depots. MCP-1 and IL-6 mRNA levels were, however, elevated in subcutaneous fat in response to LPS, the increases being four- and 20-fold, respectively (Fig. 1).

Effect of LPS on adipokine expression 4 h after the induction of endotoxaemia

To assess the rapidity of the response to LPS, studies were next undertaken at 4 h after injection. Epididymal, perirenal and subcutaneous adipose tissue depots were again examined. In the epididymal depot at 4 h, there were no significant changes in adipsin, adiponectin, IL-18 and MT-2 mRNA levels (Fig. 2). There were, however, small, but statistically significant, increases in leptin (fourfold) and NGF (3.8-fold) mRNAs. Substantial increases were observed for TNF α (20-

Fig. 1 Adipokine gene expression in endotoxaemia in mice. Mice were injected with LPS to induce severe sepsis (closed bars) and adipose tissue depots removed 24 h later; control mice (open bars) received a saline injection. The level of each adipokine mRNA was measured in epididymal (*Epi*), perirenal (*Peri*) and subcutaneous (*Sub*) depots by real-time PCR and normalized to β -actin relative to the untreated control group ($\equiv 1$). *IL-6* Interleukin 6, *IL-18* interleukin 18, *MCP-1* monocyte chemoattractant protein 1, *MT2* metallothionein 2, *NGF* nerve growth factor, *TNF α* tumour necrosis factor α . Results are given as mean values \pm SE for five to six mice in each group. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ compared with controls





Fig. 2 Acute effects of endotoxaemia on adipokine gene expression in mice. Mice were injected with LPS to induce severe sepsis (closed bars) and adipose tissue depots removed 4 h later; control mice (open bars) received a saline injection. The level of each adipokine mRNA was measured in epididymal (Epi), perirenal (Peri) and subcutaneous (Sub) depots by real-time PCR and normalized to β -actin relative to the untreated control group (=1). Results are given as mean values \pm SE for five to six mice in each group. * P <0.05, ** P <0.01, *** P <0.001 compared with controls



fold) and particularly MCP-1 (250-fold) and IL-6 (500-fold) mRNA. A broadly similar pattern was observed in the perirenal depot, although in this case, there was no change in leptin mRNA level. The largest changes were again with MCP-1 and IL-6 mRNA, the levels of which were increased by 35- and 78-fold, respectively (Fig. 2). TNF α level increased 23-fold, and a modest increase (fourfold) was observed with NGF.


In the subcutaneous adipose tissue, there were no changes in mRNA levels for IL-18, leptin and MT-2 (Fig. 2). There was, however, an approximately 50% reduction in mRNA levels for adiponectin and adiponectin. NGF and TNF α mRNA levels increased moderately, by four- and 17-fold, respectively. Very major changes were observed with MCP-1, the mRNA level of which increased 250-fold, and with IL-6 where the increase was >1,000-fold (Fig. 2).

White adipose tissue adipokine levels

The adipokine protein content was measured by ELISA in the epididymal and subcutaneous fat depots for the two inflammatory adipokines which exhibited the largest changes in mRNA level. IL-6, which showed the greatest elevation in mRNA, was increased in both the epididymal and subcutaneous fat at 4 and 24 h after the injection of LPS (Fig. 3). The increase in the epididymal depot was between five- and eightfold, while in the subcutaneous tissue the rise was between three- and eightfold.

In contrast to IL-6, there was no significant increase in MCP-1 protein in either tissue at 24 h. However at 4 h, the amount of MCP-1 was approximately eightfold greater in the epididymal fat and twofold greater in the subcutaneous depot (Fig. 3). For adiponectin, only in the epididymal depot after 24 h of treatment with LPS was there a

Fig. 3 Adipokine levels in adipose tissue in endotoxaemia in mice. Mice were injected with LPS to induce severe sepsis (closed bars) and adipose tissue depots removed 4 or 24 h later; control mice (open bars) received a saline injection. Adipokine proteins were measured in the epididymal (Epi) and subcutaneous (Sub) depots by ELISA. Results are given as mean values \pm SE for three to six mice in each group. * P <0.05, ** P <0.01 compared with controls



significant fall in the protein, reflecting the changes in the mRNA level. However, paradoxically the adiponectin level in the subcutaneous depot was actually increased, rather than decreased, at both 4 and 24 h in the LPS-treated mice.

Macrophage marker expression

To assess whether the increases in the expression of key inflammatory adipokine genes, particularly MCP-1, reflected an infiltration of macrophages into adipose tissue, the mRNA level of two macrophage markers was examined. The markers were F4/80 and MAC-1, and these were examined in 4 and 24 h LPS-treated mice and in all three WAT depots. In no case was there a statistically significant increase in F4/80 and MAC-1 mRNA level (Fig. 4). Indeed, in one case, the perirenal tissue treated for 24 h, the level of both mRNAs was significantly decreased. F4/80 mRNA level was also decreased in the epididymal depot at 24 h.

HIF-1 α as an index of hypoxia

In the final investigation, the effect of LPS treatment on the key molecular index of cellular oxygen deficiency, the hypoxia-sensitive transcription factor HIF-1 α , was examined. The expression and protein level of HIF-1 α was determined in both the epididymal and subcutaneous adipose tissue depots. At 4 h after the treatment with LPS, there was a three- to fourfold increase in HIF-1 α mRNA in each depot, and importantly, this was accompanied by a

two- to threefold increase in HIF-1 α protein (Fig. 5). HIF-1 α mRNA level was also twofold higher in both depots at 24 h, but at this later time point, there was no significant change in HIF-1 α protein (Fig. 5).

Discussion

The present study demonstrates that there are major changes in the expression of key adipokine genes in each of the main white adipose tissue depots in mice following the induction of endotoxaemia. The expression of several major inflammation-related adipokines was examined, and by far the largest response was observed with IL-6 and MCP-1. Adipose tissue produces large amounts of IL-6 and is considered to be a substantial contributor to circulating levels of this cytokine [9, 26]. Plasma IL-6 levels are increased in endotoxaemia as well as sepsis, and it seems likely that adipose tissue contributes directly to this [1, 13, 19], particularly since in the present study the amount of IL-6 in the fat depots was increased alongside the increase in mRNA level. Nevertheless, it should be noted that the increase in protein was much less than the changes in mRNA level, although this could reflect rapid release of IL-6 from the tissue once synthesised.

The chemokine MCP-1 shows increased expression in obesity, like IL-6, and is linked to the recruitment of macrophages within tissues, including adipose tissue. Indeed, in obesity there is a major infiltration of macro-

Fig. 4 Expression of macrophage markers in adipose tissue in endotoxaemia in mice. Mice were injected with LPS to induce endotoxaemia (closed bars) and adipose tissue depots removed 4 or 24 h later; control mice (open bars) received a saline injection. The level of each macrophage marker mRNA was measured in epididymal (Epi), perirenal (Peri) and subcutaneous (Sub) depots by real-time PCR and normalized to β -actin relative to the untreated control group (=1). Results are given as mean values \pm SE for five to six mice in each group. * P <0.05, ** P <0.01, *** P <0.001 compared with controls





Fig. 5 HIF-1 α gene expression and protein levels in adipose tissue in endotoxaemia in mice. Mice were injected with LPS to induce endotoxaemia (*closed bars*) and adipose tissue depots removed 4 or 24 h later; control mice (*open bars*) received a saline injection. HIF-1 α mRNA (a) and protein levels (b) were measured in the epididymal (*Epi*) and subcutaneous (*Sub*) depots by real-time PCR, the level being normalized to β -actin relative to the untreated control group (=1), and by ELISA, respectively. Results are given as mean values \pm SE for five to six mice in each group. * P <0.05, ** P <0.01 compared with controls



phages into adipose tissue [52, 54]. Since the mRNA level of two macrophage markers—F4/80 and MAC-1—did not increase in any of the three adipose tissue depots examined, either at 4 or 24 h after the administration of LPS, it is unlikely that there was an infiltration of immune cells during the duration of endotoxaemia in the present work. The increased MCP-1 expression suggests that macrophage recruitment into adipose tissue may well occur long-term; however, in contrast to IL-6, no sustained elevation in MCP-1 protein levels was evident in adipose tissue homogenates following LPS treatment.

In addition to macrophage infiltration, MCP-1 has been associated with the development of insulin resistance [38]. This is also the case with IL-6 and TNF α [17, 20, 27, 35]. Thus, the major stimulation of the expression of these three factors in response to LPS suggests that adipose tissue may contribute to insulin insensitivity and the dysregulation of glucose homeostasis in infection. TNF α is a pleiotropic pro-inflammatory cytokine, and its actions in adipose tissue include the stimulation of lipolysis and of apoptosis [31, 36]. It also has a major effect on the expression of other inflammation-related adipokines including the strong upregulation of IL-6 and MCP-1 production [49, 50].

In marked contrast, adiponectin expression was downregulated in all three adipose tissue depots 24 h after the administration of LPS. There was also a significant reduction in adiponectin mRNA in the subcutaneous depot at 4 h. Adiponectin, which is a key adipocyte-derived hormone, has multiple physiological effects, and these include anti-inflammatory and insulin sensitising actions [5, 28, 29, 56]. Thus, a fall in adiponectin expression may

add to the upregulation of IL-6, MCP-1 and TNF α expression in leading to a major pro-inflammatory response and insulin resistance in endotoxaemia. However, although adiponectin protein level decreased in the epididymal depot at 24 h (the absence of a change at 4 h may reflect the time delay between alterations in gene transcription and subsequent protein synthesis), in the case of the subcutaneous depot, there was a paradoxical increase in the amount of the hormone. We have no satisfactory explanation for this anomaly, although it may indicate that adiponectin is less readily released from the subcutaneous adipose tissue in endotoxaemia.

Furthermore, low adiponectin levels observed in obesity [3] as a state of low-grade chronic inflammation are associated with increased insulin resistance [5, 56]. Stress-induced hyperglycaemia frequently occurs in infection and is associated with an adverse outcome [14, 48]. Elevated TNF α levels, increased MCP-1 production as well as low adiponectin may act in concert to induce and to maintain hyperglycaemia in critical illness. Although the present study does not reveal the extent to which adipose tissue might contribute quantitatively to alterations in circulating levels of cytokines and chemokines in endotoxaemia, overall, our results suggest a role for the tissue in the changes in glucose metabolism associated with infection. It is emphasised that since in the 24-h studies there was a distinct mortality rate and that only tissues from surviving animals were investigated, there could be a link between the gene expression profile and the protective survival response.

Expression of the complement-related factor, adipsin, also fell during endotoxaemia. Leptin, on the other hand,

showed no changes in gene expression. This is in contrast with previous studies in experimental animals and in humans where an upregulation has been described [1, 12, 37]. However, the previously reported changes in leptin mRNA were very small, particularly in comparison with the dramatic changes in IL-6 and MCP-1 mRNA level.

Small increases in NGF mRNA level were observed in the epididymal and perirenal depots but not in the subcutaneous tissue. This target-derived neurotrophin is also involved in the inflammatory response, its expression and secretion from both murine and human adipocytes being stimulated by TNF α [30, 49]. The low molecular weight cysteine-rich, metal-binding protein metallothionein showed little response to LPS except in the perirenal depot where there was a small increase at 24 h. Similarly, there was little change in the expression of IL-18; this pro-inflammatory cytokine plays an important role in innate and acquired immunity, and its expression in adipocytes [41, 53] has been shown to be strongly upregulated by TNF α [53].

Certain depot-specific responses to LPS were evident in the present study. Of the three adipose tissue sites examined, the largest response was observed in the epididymal depot in the case of both IL-6 and TNF α (at 24 h). Indeed, the subcutaneous depot showed no increase in TNF α mRNA level at 24 h. In this context, it has been shown that subcutaneous adipose tissue is affected to a lesser degree by sepsis-induced malperfusion than the splanchnic area and does not contribute to lactate production during critical illness [25].

In addition to depot differences, there is the important question of which cells within adipose tissue may be responsible for alterations in adipokine/cytokine production in endotoxaemia. White fat consists of two major components, mature adipocytes and the stromal-vascular fraction. The stromal-vascular fraction includes macrophages, fibroblasts, vascular endothelial cells and preadipocytes. Adipocytes are able to express each of the inflammation-related signals examined in the present study (see [46, 47]). However, while the key cytokines and chemokines are also expressed by macrophages, adipocytes alone are responsible for adiponectin and leptin production.

It could be hypothesized that during endotoxaemia, macrophages infiltrate adipose tissue, causing further inflammatory damage and substantially contributing to mediator production. This would imply that immigrating immune cells are mainly responsible for the changes observed. However, as noted above, the mRNA level of two specific macrophage markers did not increase in WAT and even decreased in the case of the perirenal depot at 24 h. This suggests that mature adipocytes may be the key site of the response to LPS. Indeed, adipocytes express the Toll-like receptor 4 through which LPS exerts its inflam-

matory effect [22, 42]. Irrespective of the cell type involved, adipose tissue can be regarded as a key inflammatory organ, which may contribute to the systemic response observed during systemic infection by generating and releasing inflammatory mediators into the circulation. However, the possibility that resident macrophages are activated and may contribute to adipokine/cytokine production in the various adipose tissue depots in endotoxaemia, without increased immune cell infiltration, cannot be excluded.

It has recently been proposed that the inflammation that occurs in adipose tissue in obesity may, at least in part, be a response to relative hypoxia in groups of adipocytes distant from the vasculature [46]. Evidence in support of this hypothesis is emerging [15, 32, 57], and hypoxia has been shown to directly increase the expression and release of inflammation-related adipokines such as leptin, IL-6, MIF and VEGF in murine and human adipocytes, while adiponectin with its anti-inflammatory action is decreased [8, 15, 23, 51, 57]. We were therefore interested in whether hypoxia might underlie the changes in adipokine expression in endotoxaemia. Increases in the mRNA encoding the hypoxia-sensitive transcription factor, HIF-1 α , were evident following administration of LPS, and increased HIF-1 α protein was found at 4 h. This suggests that at least in the early stages of endotoxaemia, hypoxia may occur in adipose tissue. The changes at the protein level were not sustained by 24 h, implying that any hypoxia may have been transitory. Further work is warranted on hypoxia in adipose tissue in severe infection and whether it underlies the changes in adipokine expression.

A high dose of LPS was employed to induce endotoxaemia, and it may also reflect sepsis. However, the LPS challenge may not adequately reproduce the complexity of human sepsis. In particular, investigators with an interest in peri-operative sepsis have increasingly used caecal ligation and puncture (CLP) as the model of choice. In a preliminary series of experiments with CLP in the same strain of mice, we were able to reproduce the changes in IL-6 mRNA levels found in the epididymal fat depot after LPS challenge. Here IL-6 mRNA levels were 87-fold higher than in control mice and sixfold higher than in sham-operated mice at 24 h (unpublished data). Apparently, CLP leads to qualitatively similar changes in IL-6 expression as LPS, suggesting that white adipose tissue is activated in septic shock as in endotoxaemia.

Clinical sepsis is commonly polymicrobial, but experimentation involving bacterial products, such as LPS, may serve to characterize response patterns directed against one particular organism [18]. We therefore used LPS as the model of choice to examine the potential role of WAT during severe infection. The dose employed, although high, is within the range used in previous studies and elicited a

severe, acute, systemic inflammatory response in mice with all the symptoms of sepsis and septic shock [24]. Importantly, the substantial increase in IL-6 and MCP-1 gene expression in the subcutaneous adipose tissue at 4 h indicates a systemic response in WAT rather than an effect local to the i.p. site of injection. The mortality rate of 39% in our 24-h series of experiments reflects acute life-threatening gram-negative infection of a severity similar to that occurring in human sepsis [2]. Further investigations using various models of sepsis are warranted to specify the role of WAT in infection caused by different organisms.

In conclusion, our results demonstrate that LPS challenge sufficient to mimic endotoxaemia leads to major changes in the expression of key adipokine genes in different adipose tissue depots of mice. Thus, white adipocytes (and preadipocytes) may well contribute directly to the most prominent features of infection—systemic inflammation and metabolic dysregulation. If this could be corroborated in studies involving human patients, it would allow a new interpretation of the positive correlation between body mass index and sepsis mortality as well as the development of novel approaches to treating insulin resistance in severe infection.

Acknowledgement We gratefully acknowledge the receipt of funding from the Biotechnology and Biological Sciences Research Council (UK) to PT. The authors declare that they have no conflicts of interest with respect to this study. PT is a member of COST BM0602.

References

- Anderson PD, Mehta NN, Wolfe ML, Hinkle CC, Pruscino L, Comiskey LL, Tabita-Martinez J, Sellers KF, Rickels MR, Ahima RS, Reilly MP (2007) Innate immunity modulates adipokines in humans. *J Clin Endocrinol Metab* 92:2272–2279
- Angus DC, Linde-Zwirble WT, Lidicker J, Clermont G, Carcillo J, Pinsky MR (2001) Epidemiology of severe sepsis in the United States: analysis of incidence, outcome, and associated costs of care. *Crit Care Med* 29:1303–1310
- Arita Y, Kihara S, Ouchi N, Takahashi M, Maeda K, Miyagawa J, Hotta K, Shimomura I, Nakamura T, Miyaoka K, Kuriyama H, Nishida M, Yamashita S, Okubo K, Matsubara K, Muraguchi M, Ohmoto Y, Funahashi T, Matsuzawa Y (1999) Paradoxical decrease of an adipose-specific protein, adiponectin, in obesity. *Biochem Biophys Res Commun* 257:79–83
- Barbier M, Cherbut C, Aube AC, Blottiere HM, Galmiche JP (1998) Elevated plasma leptin concentrations in early stages of experimental intestinal inflammation in rats. *Gut* 43:783–790
- Berg AH, Combs TP, Du X, Brownlee M, Scherer PE (2001) The adipocyte-secreted protein Acrp30 enhances hepatic insulin action. *Nat Med* 7:947–953
- Bornstein SR, Preas HL, Chrousos GP, Sufredini AF (1998) Circulating leptin levels during acute experimental endotoxemia and antiinflammatory therapy in humans. *J Infect Dis* 178:887–890
- Brahimi-Horn MC, Pouyssegur J (2007) Oxygen, a source of life and stress. *FEBS Lett* 581:3582–3591
- Chen B, Lam KSL, Wang Y, Wu D, Lam MC, Shen J, Wong L, Hoo RLC, Zhang J, Xu A (2006) Hypoxia dysregulates the production of adiponectin and plasminogen activator inhibitor-1 independent of reactive oxygen species in adipocytes. *Biochem Biophys Res Commun* 341:549–556
- Coppack SW (2001) Pro-inflammatory cytokines and adipose tissue. *Proc Nutr Soc* 60:349–356
- Ellis CG, Jagger J, Sharpe M (2005) The microcirculation as a functional system. *Crit Care* 9(Suppl 4):S3–S8
- Faggioni R, Fantuzzi G, Fuller J, Dinarello CA, Feingold KR, Grunfeld C (1998) IL-1 beta mediates leptin induction during inflammation. *Am J Physiol Reg Integr Comp Physiol* 43:R204–R208
- Grunfeld C, Zhao C, Fuller J, Pollock A, Moser A, Friedman J, Feingold KR (1996) Endotoxin and cytokines induce expression of leptin, the *ob* gene-product, in hamsters—a role for leptin in the anorexia of infection. *J Clin Invest* 97:2152–2157
- Haek CE, De Groot ER, Felt-Bersma RJ, Nuijens JH, Strack Van Schijndel RJ, Eerenberg-Belmer AJ, Thijs LG, Aarden LA (1989) Increased plasma levels of interleukin-6 in sepsis. *Blood* 74:1704–1710
- Heuer JG, Sharma GR, Zhang T, Ding C, Bailey DL, Stephens EJ, Holmes KC, Grubbs RL, Fynboe KA, Chen YF, Jakubowski JA (2006) Effects of hyperglycemia and insulin therapy on outcome in a hyperglycemic septic model of critical illness. *J Trauma* 60:865–872
- Hosogai N, Fukuhara A, Oshima K, Miyata Y, Tanaka S, Segawa K, Furukawa S, Tochino Y, Komuro R, Matsuda M, Shimomura I (2007) Adipose tissue hypoxia in obesity and its impact on adipocytokine dysregulation. *Diabetes* 56:901–911
- Hotamisligil GS (2006) Inflammation and metabolic disorders. *Nature* 444:860–867
- Hotamisligil GS, Shargill NS, Spiegelman BM (1993) Adipose expression of tumor necrosis factor- α —direct role in obesity-linked insulin resistance. *Science* 259:87–91
- Hubbard WJ, Choudhry M, Schwacha MG, Kerby JD, Rue LW 3rd, Bland KI, Chaudry IH (2005) Cecal ligation and puncture. *Shock* 24(Suppl 1):S2–S7
- Krogh-Madsen R, Plomgaard P, Akerström T, Møller K, Schmitz O, Pedersen BK (2008) Effect of short-term intralipid infusion on the immune response during low-dose endotoxemia in humans. *Am J Physiol Endocrinol Metab* 294:E371–E379
- Lagathu C, Bastard J-P, Auclair M, Manchi M, Capeau J, Caron M (2003) Chronic interleukin-6 (IL-6) treatment increased IL-6 secretion and induced insulin resistance in adipocyte: prevention by rosiglitazone. *Biochem Biophys Res Commun* 311:372–379
- Lauza-Jacoby S, Sedkova N, Phetteplace H, Perrotti D (1997) Sepsis-induced regulation of lipoprotein lipase expression in rat adipose tissue and soleus muscle. *J Lipid Res* 38:701–710
- Lin Y, Lee H, Berg AH, Lisanti MP, Shupiro L, Scherer PE (2000) The lipopolysaccharide-activated toll-like receptor (TLR)-4 induces synthesis of the closely related receptor TLR-2 in adipocytes. *J Biol Chem* 275:24255–24263
- Lolmède K, Durand de Saint Front V, Galitzky J, Lafontan M, Bouloumié A (2003) Effects of hypoxia on the expression of proangiogenic factors in differentiated 3T3-F442A adipocytes. *Int J Obesity* 27:1187–1195
- Munnel DN (2007) Advances in sepsis research derived from animal models. *Int J Med Microbiol* 297:393–400
- Martinez A, Chioloro R, Bollman M, Revelly JP, Berger M, Cayeux C, Tuppy L (2003) Assessment of adipose tissue metabolism by means of subcutaneous microdialysis in patients with sepsis or circulatory failure. *Clin Physiol Funct Imaging* 23:286–292
- Mohamed-Ali V, Goodrick S, Rawesh A, Katz DR, Miles JM, Yudkin JS, Klein S, Coppack SW (1997) Subcutaneous adipose tissue releases interleukin-6, but not tumor necrosis factor- α , in vivo. *J Clin Endocrinol Metab* 82:4196–4200

27. Moller DE (2000) Potential role of TNF- α in the pathogenesis of insulin resistance and type 2 diabetes. *Trends Endocrinol Metab* 11:212–217
28. Ouchi N, Kihara S, Arita Y, Maeda K, Kuriyama H, Okamoto Y, Hotta K, Nishida M, Takahashi M, Nakamura T, Yamashita S, Funahashi T, Matsuzawa Y (1999) Novel modulator for endothelial adhesion molecules—adipocyte-derived plasma protein adiponectin. *Circulation* 100:2473–2476
29. Ouchi N, Kihara S, Arita Y, Okamoto Y, Maeda K, Kuriyama H, Hotta K, Nishida M, Takahashi M, Muraguchi M, Ohmoto Y, Nakamura T, Yamashita S, Funahashi T, Matsuzawa Y (2000) Adiponectin, an adipocyte-derived plasma protein, inhibits endothelial NF- κ B signaling through a cAMP-dependent pathway. *Circulation* 102:1296–1301
30. Peeraully MR, Jenkins JR, Trayhurn P (2004) NGF gene expression and secretion in white adipose tissue: regulation in 3T3-L1 adipocytes by hormones and inflammatory cytokines. *Am J Physiol Endocrinol Metab* 287:E331–E339
31. Prins JB, Niesler CU, Winterford CM, Bright NA, Siddle K, Ormally S, Walker NJ, Cameron DP (1997) Tumor necrosis factor- α induces apoptosis of human adipose cells. *Diabetes* 46:1939–1944
32. Rausch ME, Weisberg SP, Vardhana P, Tortoriciello DV (2008) Obesity in C57BL/6J mice is characterized by adipose tissue hypoxia and cytotoxic T-cell infiltration. *Int J Obesity* 32:451–463
33. Ray DE, Matchett SC, Baker K, Wasser T, Young MJ (2005) The effect of body mass index on patient outcomes in a medical ICU. *Chest* 127:2125–2131
34. Rosen ED, Spiegelman BM (2006) Adipocytes as regulators of energy balance and glucose homeostasis. *Nature* 444:847–853
35. Rotter V, Nagaev I, Smith U (2003) Interleukin-6 (IL-6) induces insulin resistance in 3T3-L1 adipocytes and is, like IL-8 and tumor necrosis factor- α , overexpressed in human fat cells from insulin-resistant subjects. *J Biol Chem* 278:45777–45784
36. Ryden M, Arvidsson E, Blomqvist L, Perbeck L, Dicker A, Arner P (2004) Targets for TNF- α -induced lipolysis in human adipocytes. *Biochem Biophys Res Commun* 318:168–175
37. Sarraf P, Frederich RC, Turner EM, Ma G, Jaskowiak NT, Rivet DJ, Flier JS, Lowell BB, Flier DL, Alexander HR (1997) Multiple cytokines and acute inflammation raise mouse leptin levels: potential role in inflammatory anorexia. *J Exp Med* 185:171–175
38. Sartipy P, Loskutov DJ (2003) Monocyte chemoattractant protein 1 in obesity and insulin resistance. *Proc Natl Acad Sci USA* 100:7265–7270
39. Schumacker PT (2005) Hypoxia-inducible factor-1 (HIF-1). *Crit Care Med* 33:S423–S425
40. Semenza GL, Agani F, Feldser D, Iyer N, Kotch L, Laughner E, Yu A (2000) Hypoxia, HIF-1, and the pathophysiology of common human diseases. *Adv Exp Med Biol* 475:123–130
41. Skurk T, Kolb H, Muller-Scholze S, Rohrig K, Hauner H, Herder C (2005) The proatherogenic cytokine interleukin-18 is secreted by human adipocytes. *Eur J Endocrinol* 152:863–868
42. Song MJ, Kim KH, Yoon JM, Kim JB (2006) Activation of Toll-like receptor 4 is associated with insulin resistance in adipocytes. *Biochem Biophys Res Commun* 346:739–745
43. Sugawara K, Miyata G, Shintani R, Satomi S (2003) The lipolytic responsiveness to endotoxin in subcutaneous adipose tissue is greater than mesenteric adipose tissue. *Tohoku J Exp Med* 199:171–179
44. Torpy DJ, Bornstein SR, Chrousos GP (1998) Leptin and interleukin-6 in sepsis. *Horm Metab Res* 30:726–729
45. Trayhurn P (2005) Endocrine and signalling role of adipose tissue: new perspectives on fat. *Acta Physiol Scand* 184:285–293
46. Trayhurn P, Wood IS (2004) Adipokines: Inflammation and the pleiotropic role of white adipose tissue. *Br J Nutr* 92:347–355
47. Trayhurn P, Wood IS (2005) Signalling role of adipose tissue: adipokines and inflammation in obesity. *Biochem Soc Trans* 33:1078–1081
48. van den Berghe G, Wouters P, Weekers F, Verwaest C, Bruyninckx F, Schetz M, Vlasselaers D, Ferdinande P, Lauwers P, Bouillon R (2001) Intensive insulin therapy in the critically ill patients. *N Engl J Med* 345:1359–1367
49. Wang B, Jenkins JR, Trayhurn P (2005) Expression and secretion of inflammation-related adipokines by human adipocytes differentiated in culture: integrated response to TNF- α . *Am J Physiol Endocrinol Metab* 288:E731–E740
50. Wang B, Trayhurn P (2006) Acute and prolonged effects of TNF- α on the expression and secretion of inflammation-related adipokines by human adipocytes differentiated in culture. *Pflugers Archiv Eur J Physiol* 452:418–427
51. Wang B, Wood IS, Trayhurn P (2007) Dysregulation of the expression and secretion of inflammation-related adipokines by hypoxia in human adipocytes. *Pflugers Archiv Eur J Physiol* 455:479–492
52. Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL, Ferrante AW Jr (2003) Obesity is associated with macrophage accumulation in adipose tissue. *J Clin Invest* 112:1796–1808
53. Wood IS, Wang B, Jenkins JR, Trayhurn P (2005) The pro-inflammatory cytokine IL-18 is expressed in human adipose tissue and strongly upregulated by TNF α in human adipocytes. *Biochem Biophys Res Commun* 337:422–429
54. Xu H, Barnes GT, Yang Q, Tan G, Yang D, Chou CJ, Sole J, Nichols A, Ross JS, Tartaglia LA, Chen H (2003) Chronic inflammation in fat plays a crucial role in the development of obesity-related insulin resistance. *J Clin Invest* 112:1821–1830
55. Yaegashi M, Jean R, Zuriqat M, Noack S, Homel P (2005) Outcome of morbid obesity in the intensive care unit. *J Intensive Care Med* 20:147–154
56. Yamauchi T, Kamon J, Waki H, Terauchi Y, Kubota N, Hara K, Mori Y, Ide T, Murakami K, Tsuboyama-Kasaoka N, Ezaki O, Akanuma Y, Gavrilova O, Vinson C, Reitman ML, Kagechika H, Shudo K, Yoda M, Nakano Y, Tobe K, Nagai R, Kimura S, Tomita M, Froguel P, Kadowaki T (2001) The fat-derived hormone adiponectin reverses insulin resistance associated with both lipodystrophy and obesity. *Nat Med* 7:941–946
57. Ye J, Gao Z, Yin J, He Q (2007) Hypoxia is a potential risk factor for chronic inflammation and adiponectin reduction in adipose tissue of *ob/ob* and dietary obese mice. *Am J Physiol Endocrinol Metab* 293:E1118–E1128

Endotoxaemia leads to major increases in inflammatory adipokine gene expression in white adipose tissue of mice

Martin Leuwer & Ingeborg Welters & Gernot Marx & Andrew Rushton & Hongguang Bao & Leif Hunter & Paul Trayhurn

Above paper has been cited (as of December 2012) by :

Maury, E., and S. M. Brichard. "Adipokine dysregulation, adipose tissue inflammation and metabolic syndrome." *Molecular and cellular endocrinology* 314, no. 1 (2010): 1.

Redman, C. W. G., and I. L. Sargent. "Placental stress and pre-eclampsia: a revised view." *Placenta* 30 (2009): 38-42.

Bourlier, Virginie, and Anne Bouloumié. "Role of macrophage tissue infiltration in obesity and insulin resistance." *Diabetes & metabolism* 35.4 (2009): 251-260.

Balistreri, Carmela Rita, Calogero Caruso, and Giuseppina Candore. "The role of adipose tissue and adipokines in obesity-related inflammatory diseases." *Mediators of inflammation* 2010 (2010).

Lira, Fabio S., et al. "Endotoxin levels correlate positively with a sedentary lifestyle and negatively with highly trained subjects." *Lipids in health and disease* 9.1 (2010): 82.

Menzebach, Axel, et al. "A comprehensive study of survival, tissue damage, and neurological dysfunction in a murine model of cardiopulmonary resuscitation after potassium-induced cardiac arrest." *Shock* 33.2 (2010): 189.

Lolmède, Karine, et al. "Immune cells in adipose tissue: key players in metabolic disorders." *Diabetes & metabolism* 37.4 (2011): 283-290.

Langouche, Lies, et al. "Alterations in adipose tissue during critical illness: an adaptive and protective response?." *Critical Care* 14 (2010): 1-1.

Redman, C. W. G. "Preeclampsia: a multi-stress disorder." *La revue de médecine interne* 32.S1 (2011): 41-44.

Uji, Yoshitaka, et al. "Adiponectin deficiency promotes the production of inflammatory mediators while severely exacerbating hepatic injury in mice with polymicrobial sepsis." *Journal of Surgical Research* 161.2 (2010): 301-311.

V Stankov, Metodi, and Georg MN Behrens. "Contribution of inflammation to fat redistribution and metabolic disturbances in HIV-1 infected patients." *Current pharmaceutical design* 16.30 (2010): 3361-3371.

Robinson, Katherine, John Prins, and Bala Venkatesh. "Clinical review: Adiponectin biology and its role in inflammation and critical illness." *Critical Care* 15.2 (2011): 221.

TvariJonaviciute, Asta, et al. "Adiponectin and IGF-1 are negative acute phase proteins in a dog model of acute endotoxaemia." *Veterinary Immunology and Immunopathology* 140.1 (2011): 147-151.

Fain, John N., et al. "The inflammatory response seen when human omental adipose tissue explants are incubated in primary culture is not dependent upon albumin and is primarily in the nonfat cells." *Journal of Inflammation* 7.1 (2010): 4.

Asseldonk, Edwin JP, et al. "The Effect of the Interleukin-1 Cytokine Family Members IL-1F6 and IL-1F8 on Adipocyte Differentiation." *Obesity* 18.11 (2012): 2234-2236.

Thomas, Anthony P., et al. "Inflammatory Phenotyping Identifies CD11d as a Gene Markedly Induced in White Adipose Tissue in Obese Rodents and Women." *The Journal of nutrition* 141.6 (2011): 1172-1180.

Turner, J. J. O., et al. "Investigation of nuclear factor- κ B inhibitors and interleukin-10 as regulators of inflammatory signalling in human adipocytes." *Clinical & Experimental Immunology* 162.3 (2010): 487-493.

TvariJonaviciute, Asta, et al. "Effect of weight loss on inflammatory biomarkers in obese dogs." *The Veterinary Journal* (2012).

Hanses, Frank, et al. "Intracellular survival of *Staphylococcus aureus* in adipocyte-like differentiated 3T3-L1 cells is glucose dependent and alters cytokine, chemokine, and adipokine secretion." *Endocrinology* 152.11 (2011): 4148-4157.

Packer, M. J., et al. "Adipose tissue-derived adiponectin expression is significantly associated with increased post operative mortality in horses undergoing emergency abdominal surgery." *Equine Veterinary Journal* 43.s39 (2011): 26-33.

van Meurs, Matijs, et al. "Adiponectin diminishes organ-specific microvascular endothelial cell activation associated with sepsis." *Shock* 37.4 (2012): 392.

庞璨, and 贾伟平. "脂肪组织缺氧与肥胖相关性炎症的研究现状." *中华医学杂志* 88.36 (2009): 2588-2591.

Sugden, Mary C., and Mark J. Holness. "Autocrine Effects in White Adipose Tissue and Pancreatic Islets: Emergent Roles in the Regulation of Adipocyte and Pancreatic β -cell Function."

Pini, Maria, et al. "Obesity and IL-6 interact in modulating the response to endotoxemia in mice." *Cytokine* (2012).

Carmela Rita, Balistreri, Caruso Calogero, and Candore Giuseppina. "The Role of Adipose Tissue and Adipokines in Obesity-Related Inflammatory Diseases." *Mediators of Inflammation* 2010 (2010).

Ramsay, T. G., et al. "Peripheral tumor necrosis factor α regulation of adipose tissue metabolism and adipokine gene expression in neonatal pigs." *Veterinary Research Communications* (2012): 1-10.

詹莉莉, 杨志秋, and 傅正伟. "肥胖与慢性炎症的研究进展." *中国细胞生物学学报* 33.3 (2011): 297-305.

Henri, Marion. *Obésité et insulino-résistance: étude longitudinale avec un traceur du transport du glucose, le [125I]-6-déoxy-6-iodo-D-glucose*. Diss. Université de Grenoble, 2011.

REFERENCES

1. Harrison DA, Welch CA, Eddleston JM. The epidemiology of severe sepsis in England, Wales and Northern Ireland, 1996 to 2004: secondary analysis of a high quality clinical database, the ICNARC Case Mix Programme Database. Crit Care. 2006;10(2):R42.
2. Dombrovskiy VY, Martin AA, Sunderram J, Paz HL. Rapid increase in hospitalization and mortality rates for severe sepsis in the United States: a trend analysis from 1993 to 2003. Crit Care Med. 2007 May;35(5):1244-50.
3. Angus DC, Linde-Zwirble WT, Lidicker J, Clermont G, Carcillo J, Pinsky MR. Epidemiology of severe sepsis in the United States: analysis of incidence, outcome, and associated costs of care. Crit Care Med. 2001 Jul;29(7):1303-10.
4. Bone RC, Balk RA, Cerra FB, Dellinger RP, Fein AM, Knaus WA, et al. Definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. The ACCP/SCCM Consensus Conference Committee. American College of Chest Physicians/Society of Critical Care Medicine. Chest. 1992 Jun;101(6):1644-55.
5. Geroulanos S, Douka ET. Historical perspective of the word "sepsis". Intensive Care Med. 2006 Dec;32(12):2077.
6. Levy MM, Fink MP, Marshall JC, Abraham E, Angus D, Cook D, et al. 2001 SCCM/ESICM/ACCP/ATS/SIS International Sepsis Definitions Conference. Crit Care Med. 2003 Apr;31(4):1250-6.
7. Toh CH, Ticknor LO, Downey C, Giles AR, Paton RC, Wenstone R. Early identification of sepsis and mortality risks through simple, rapid clot-waveform analysis. Implications of lipoprotein-complexed C reactive protein formation. Intensive Care Med. 2003 Jan;29(1):55-61.
8. Marshall JC. Rethinking Sepsis: From Concepts to Syndromes to Diseases. Sepsis. 1999;3(1):5-10.

9. O'Brien JM, Jr., Ali NA, Aberegg SK, Abraham E. Sepsis. *Am J Med.* 2007 Dec;120(12):1012-22.
10. Chung TP, Laramie JM, Province M, Cobb JP. Functional genomics of critical illness and injury. *Crit Care Med.* 2002 Jan;30(1 Suppl):S51-7.
11. Tjardes T, Neugebauer E. Sepsis research in the next millennium: concentrate on the software rather than the hardware. *Shock.* 2002 Jan;17(1):1-8.
12. Lanza-Jacoby S, Sedkova N, Phetteplace H, Perrotti D. Sepsis-induced regulation of lipoprotein lipase expression in rat adipose tissue and soleus muscle. *J Lipid Res.* 1997 Apr;38(4):701-10.
13. Sugawara K, Miyata G, Shineha R, Satomi S. The lipolytic responsiveness to endotoxin in subcutaneous adipose tissue is greater than mesenteric adipose tissue. *Tohoku J Exp Med.* 2003 Mar;199(3):171-9.
14. Xu H, Barnes GT, Yang Q, Tan G, Yang D, Chou CJ, et al. Chronic inflammation in fat plays a crucial role in the development of obesity-related insulin resistance. *J Clin Invest.* 2003;112:1821-30.
15. Martinez A, Chioloro R, Bollman M, Revelly JP, Berger M, Cayeux C, et al. Assessment of adipose tissue metabolism by means of subcutaneous microdialysis in patients with sepsis or circulatory failure. *Clin Physiol Funct Imaging.* 2003 Sep;23(5):286-92.
16. Trayhurn P, Wood IS. Adipokines: Inflammation and the pleiotropic role of white adipose tissue. *Br J Nutr.* 2004;92:347-55.
17. Hotamisligil GS. Inflammation and metabolic disorders. *Nature.* 2006 Dec 14;444(7121):860-7.
18. Rosen ED, Spiegelman BM. Adipocytes as regulators of energy balance and glucose homeostasis. *Nature.* 2006;444:847-53.

19. Ogden CL, Carroll MD, Curtin LR, McDowell MA, Tabak CJ, Flegal KM. Prevalence of overweight and obesity in the United States, 1999-2004. *JAMA*. 2006 Apr 5;295(13):1549-55.
20. Kral JG, Kava RA, Catalano PM, Moore BJ. Severe Obesity: The Neglected Epidemic. *Obes Facts*. 2012 Apr 25;5(2):254-69.
21. Huttunen R, Syrjanen J. Obesity and the risk and outcome of infection. *Int J Obes (Lond)*. 2012 May 1.
22. Bercault N, Boulain T, Kuteifan K, Wolf M, Runge I, Fleury JC. Obesity-related excess mortality rate in an adult intensive care unit: A risk-adjusted matched cohort study. *Crit Care Med*. 2004 Apr;32(4):998-1003.
23. Ray DE, Matchett SC, Baker K, Wasser T, Young MJ. The effect of body mass index on patient outcomes in a medical ICU. *Chest*. 2005 Jun;127(6):2125-31.
24. Yaegashi M, Jean R, Zuriqat M, Noack S, Homel P. Outcome of morbid obesity in the intensive care unit. *J Intensive Care Med*. 2005 May-Jun;20(3):147-54.
25. Hogue CW, Jr., Stearns JD, Colantuoni E, Robinson KA, Stierer T, Mitter N, et al. The impact of obesity on outcomes after critical illness: a meta-analysis. *Intensive Care Med*. 2009 Jul;35(7):1152-70.
26. Oliveros H, Villamor E. Obesity and mortality in critically ill adults: a systematic review and meta-analysis. *Obesity (Silver Spring)*. 2008 Mar;16(3):515-21.
27. Kiraly L, Hurt RT, Van Way CW, 3rd. The outcomes of obese patients in critical care. *JPEN J Parenter Enteral Nutr*. 2011 Sep;35(5 Suppl):29S-35S.
28. Paolini JB, Mancini J, Genestal M, Gonzalez H, McKay RE, Samii K, et al. Predictive value of abdominal obesity vs. body mass index for determining risk of intensive care unit mortality. *Crit Care Med*. 2010 May;38(5):1308-14.

29. Trayhurn P, Wood IS. Signalling role of adipose tissue: adipokines and inflammation in obesity. *Biochem Soc Trans.* 2005;33:1078-81.
30. Semenza GL, Agani F, Feldser D, Iyer N, Kotch L, Laughner E, et al. Hypoxia, HIF-1, and the pathophysiology of common human diseases. *Adv Exp Med Biol.* 2000;475:123-30.
31. den Uil CA, Klijin E, Lagrand WK, Brugts JJ, Ince C, Spronk PE, et al. The microcirculation in health and critical disease. *Prog Cardiovasc Dis.* 2008 Sep-Oct;51(2):161-70.
32. Brahimi-Horn MC, Pouyssegur J. Oxygen, a source of life and stress. *FEBS Letters.* 2007;581:3582-91.
33. Kenneth NS, Rocha S. Regulation of gene expression by hypoxia. *Biochem J.* 2008 Aug 15;414(1):19-29.
34. Semenza GL, Wang GL. A nuclear factor induced by hypoxia via de novo protein synthesis binds to the human erythropoietin gene enhancer at a site required for transcriptional activation. *Mol Cell Biol.* 1992 Dec;12(12):5447-54.
35. Zhou J, Brune B. Cytokines and hormones in the regulation of hypoxia inducible factor-1alpha (HIF-1alpha). *Cardiovasc Hematol Agents Med Chem.* 2006 Jul;4(3):189-97.
36. Jantsch J, Chakravortty D, Turza N, Prechtel AT, Buchholz B, Gerlach RG, et al. Hypoxia and hypoxia-inducible factor-1 alpha modulate lipopolysaccharide-induced dendritic cell activation and function. *J Immunol.* 2008 Apr 1;180(7):4697-705.
37. Jantsch J, Wiese M, Schodel J, Castiglione K, Glasner J, Kolbe S, et al. Toll-like receptor activation and hypoxia use distinct signaling pathways to stabilize hypoxia-inducible factor 1alpha (HIF1A) and result in differential HIF1A-dependent gene expression. *J Leukoc Biol.* 2011 Sep;90(3):551-62.

38. Martin GS, Mannino DM, Eaton S, Moss M. The epidemiology of sepsis in the United States from 1979 through 2000. *N Engl J Med*. 2003 Apr 17;348(16):1546-54.
39. Hofhuis JG, Spronk PE, van Stel HF, Schrijvers AJ, Rommes JH, Bakker J. The impact of severe sepsis on health-related quality of life: a long-term follow-up study. *Anesth Analg*. 2008 Dec;107(6):1957-64.
40. Arcaroli J, Fessler MB, Abraham E. Genetic polymorphisms and sepsis. *Shock*. 2005 Oct;24(4):300-12.
41. Buras JA, Holzmann B, Sitkovsky M. Animal models of sepsis: setting the stage. *Nat Rev Drug Discov*. 2005 Oct;4(10):854-65.
42. Hubbard WJ, Choudhry M, Schwacha MG, Kerby JD, Rue LW, 3rd, Bland KI, et al. Cecal ligation and puncture. *Shock*. 2005 Dec;24 Suppl 1:52-7.
43. Wichterman KA, Baue AE, Chaudry IH. Sepsis and septic shock---a review of laboratory models and a proposal. *J Surg Res*. 1980 Aug;29(2):189-201.
44. Fuentes JM, Talamini MA, Fulton WB, Hanly EJ, Aurora AR, De Maio A. General anesthesia delays the inflammatory response and increases survival for mice with endotoxic shock. *Clin Vaccine Immunol*. 2006 Feb;13(2):281-8.
45. Carrigan KA, Saurer TB, Ijames SG, Lysle DT. Buprenorphine produces naltrexone reversible alterations of immune status. *Int Immunopharmacol*. 2004 Mar;4(3):419-28.
46. Leuwer M, Welters I, Marx G, Rushton A, Bao H, Hunter L, et al. Endotoxaemia leads to major increases in inflammatory adipokine gene expression in white adipose tissue of mice. *Pflugers Arch*. 2009 Feb;457(4):731-41.
47. Bao H. The expression of adiponectin in epididymal fat pads in a murine model of sepsis [M.Phil.]. Liverpool: Liverpool; 2006.

48. Coppack SW. Pro-inflammatory cytokines and adipose tissue. *Proc Nutr Soc.* 2001;60:349-56.
49. Mohamed-Ali V, Goodrick S, Rawesh A, Katz DR, Miles JM, Yudkin JS, et al. Subcutaneous adipose tissue releases interleukin-6, but not tumor necrosis factor-alpha, in vivo. *J Clin Endocrinol Metab.* 1997;82:4196-200.
50. Anderson PD, Mehta NN, Wolfe ML, Hinkle CC, Pruscino L, Comiskey LL, et al. Innate Immunity Modulates Adipokines in Humans. *J Clin Endocrinol Metab.* 2007;92:2272-9.
51. Hack CE, De Groot ER, Felt-Bersma RJ, Nuijens JH, Strack Van Schijndel RJ, Eerenberg-Belmer AJ, et al. Increased plasma levels of interleukin-6 in sepsis. *Blood.* 1989 Oct;74(5):1704-10.
52. Krogh-Madsen R, Plomgaard P, Akerstrom T, Moller K, Schmitz O, Pedersen BK. Effect of short-term intralipid infusion on the immune response during low-dose endotoxemia in humans. *Am J Physiol Endocrinol Metab.* 2008 Feb;294(2):E371-9.
53. Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL, Ferrante AW, Jr. Obesity is associated with macrophage accumulation in adipose tissue. *J Clin Invest.* 2003.;112(.):1796-808.
54. Sartipy P, Loskutoff DJ. Monocyte chemoattractant protein 1 in obesity and insulin resistance. *Proc Natl Acad Sci U S A.* 2003;100:7265-70.
55. Hotamisligil GS, Shargill NS, Spiegelman BM. Adipose expression of tumor necrosis factor-alpha - direct role in obesity-linked insulin resistance. *Science.* 1993;259:87-91.
56. Lagathu C, Bastard J-P, Auclair M, Maachi M, Capeau J, Caron M. Chronic interleukin-6 (IL-6) treatment increased IL-6 secretion and induced insulin resistance in adipocyte: prevention by rosiglitazone. *Biochem Biophys Res Commun.* 2003;311:372-9.

57. Moller DE. Potential role of TNF-alpha in the pathogenesis of insulin resistance and type 2 diabetes. *Trends Endocrinol Metab.* 2000;11:212-7.
58. Rotter V, Nagaev I, Smith U. Interleukin-6 (IL-6) induces insulin resistance in 3T3-L1 adipocytes and is, like IL-8 and tumor necrosis factor-alpha, overexpressed in human fat cells from insulin-resistant subjects. *J Biol Chem.* 2003;278:45777-84.
59. Prins JB, Niesler CU, Winterford CM, Bright NA, Siddle K, Orahilly S, et al. Tumor necrosis factor-alpha induces apoptosis of human adipose cells. *Diabetes.* 1997;46:1939-44.
60. Ryden M, Arvidsson E, Blomqvist L, Perbeck L, Dicker A, Arner P. Targets for TNF-[alpha]-induced lipolysis in human adipocytes. *Biochem Biophys Res Commun.* 2004;318:168-75.
61. Wang B, Jenkins JR, Trayhurn P. Expression and secretion of inflammation-related adipokines by human adipocytes differentiated in culture: Integrated response to TNF-{alpha}. *Am J Physiol Endocrinol Metab.* 2005;288:E731-E40.
62. Wang B, Trayhurn P. Acute and prolonged effects of TNF-alpha on the expression and secretion of inflammation-related adipokines by human adipocytes differentiated in culture. *Pflugers Archiv European Journal of Physiology.* 2006;452:418-27.
63. Berg AH, Combs TP, Du X, Brownlee M, Scherer PE. The adipocyte-secreted protein Acrp30 enhances hepatic insulin action. *Nat Med.* 2001;7:947-53.
64. Ouchi N, Kihara S, Arita Y, Maeda K, Kuriyama H, Okamoto Y, et al. Novel modulator for endothelial adhesion molecules - Adipocyte-derived plasma protein adiponectin. *Circulation.* 1999;100:2473-6.
65. Ouchi N, Kihara S, Arita Y, Okamoto Y, Maeda K, Kuriyama H, et al. Adiponectin, an adipocyte-derived plasma protein, inhibits endothelial NF-kappa B signaling

through a cAMP-dependent pathway. *Circulation*. 2000;102:1296-301.

66. Yamauchi T, Kamon J, Waki H, Terauchi Y, Kubota N, Hara K, et al. The fat-derived hormone adiponectin reverses insulin resistance associated with both lipotrophy and obesity. *Nat Med*. 2001;7:941-6.

67. Arita Y, Kihara S, Ouchi N, Takahashi M, Maeda K, Miyagawa J, et al. Paradoxical decrease of an adipose-specific protein, adiponectin, in obesity. *Biochem Biophys Res Commun*. 1999;257:79-83.

68. Koch A, Sanson E, Voigt S, Helm A, Trautwein C, Tacke F. Serum adiponectin upon admission to the intensive care unit may predict mortality in critically ill patients. *J Crit Care*. 2011 Apr;26(2):166-74.

69. Heuer JG, Sharma GR, Zhang T, Ding C, Bailey DL, Stephens EJ, et al. Effects of hyperglycemia and insulin therapy on outcome in a hyperglycemic septic model of critical illness. *J Trauma*. 2006 Apr;60(4):865-72.

70. van den Berghe G, Wouters P, Weekers F, Verwaest C, Bruyninckx F, Schetz M, et al. Intensive insulin therapy in the critically ill patients. *N Engl J Med*. 2001 Nov 8;345(19):1359-67.

71. Grunfeld C, Zhao C, Fuller J, Pollack A, Moser A, Friedman J, et al. Endotoxin and cytokines induce expression of leptin, the ob gene product, in hamsters. *J Clin Invest*. 1996 May 1;97(9):2152-7.

72. Sarraf P, Frederich RC, Turner EM, Ma G, Jaskowiak NT, Rivet DJ, et al. Multiple cytokines and acute inflammation raise mouse leptin levels: Potential role in inflammatory anorexia. *Journal of Experimental Medicine*. 1997;185:171-5.

73. Peeraully MR, Jenkins JR, Trayhurn P. NGF gene expression and secretion in white adipose tissue: regulation in 3T3-L1 adipocytes by hormones and inflammatory cytokines. *Am J Physiol Endocrinol Metab*. 2004;287:E331-E9.

74. Skurk T, Kolb H, Muller-Scholze S, Rohrig K, Hauner H, Herder C. The proatherogenic cytokine interleukin-18 is secreted by human adipocytes. *Eur J Endocrinol*. 2005 Jun;152(6):863-8.

75. Wood IS, Wang B, Jenkins JR, Trayhurn P. The pro-inflammatory cytokine IL-18 is expressed in human adipose tissue and strongly upregulated by TNF[alpha] in human adipocytes. *Biochem Biophys Res Commun*. 2005;337:422-9.

76. Lin Y, Lee H, Berg AH, Lisanti MP, Shapiro L, Scherer PE. The lipopolysaccharide-activated toll-like receptor (TLR)-4 induces synthesis of the closely related receptor TLR-2 in adipocytes. *Journal of Biological Chemistry*. 2000;275:24255-63.

77. Song MJ, Kim KH, Yoon JM, Kim JB. Activation of Toll-like receptor 4 is associated with insulin resistance in adipocytes. *Biochem Biophys Res Commun*. 2006;346:739-45.

78. Hosogai N, Fukuhara A, Oshima K, Miyata Y, Tanaka S, Segawa K, et al. Adipose Tissue Hypoxia in Obesity and Its Impact on Adipocytokine Dysregulation. *Diabetes*. 2007;56:901-11.

79. Rausch ME, Weisberg S, Vardhana P, Tortoriello DV. Obesity in C57BL/6J mice is characterized by adipose tissue hypoxia and cytotoxic T-cell infiltration. *Int J Obes (Lond)*. 2008 Mar;32(3):451-63.

80. Ye J, Gao Z, Yin J, He Q. Hypoxia is a potential risk factor for chronic inflammation and adiponectin reduction in adipose tissue of ob/ob and dietary obese mice. *Am J Physiol Endocrinol Metab*. 2007;293:E1118-28.

81. Chen B, Lam KSL, Wang Y, Wu D, Lam MC, Shen J, et al. Hypoxia dysregulates the production of adiponectin and plasminogen activator inhibitor-1 independent of reactive oxygen species in adipocytes. *Biochem Biophys Res Commun*. 2006;341:549-56.

82. Lolmede K, Durand de Saint Front V, Galitzky J, Lafontan M, Bouloumie A. Effects of hypoxia on the

expression of proangiogenic factors in differentiated 3T3-F442A adipocytes. *Int J Obesity*. 2003;27:1187-95.

83. Wang B, Wood IS, Trayhurn P. Dysregulation of the expression and secretion of inflammation-related adipokines by hypoxia in human adipocytes. *Pflugers Archiv Eur J Physiol*. 2007;455:479-92.

84. Blouin CC, Page EL, Soucy GM, Richard DE. Hypoxic gene activation by lipopolysaccharide in macrophages: implication of hypoxia-inducible factor 1alpha. *Blood*. 2004 Feb 1;103(3):1124-30.

85. Peyssonnaud C, Cejudo-Martin P, Doedens A, Zinkernagel AS, Johnson RS, Nizet V. Cutting edge: Essential role of hypoxia inducible factor-1alpha in development of lipopolysaccharide-induced sepsis. *J Immunol*. 2007 Jun 15;178(12):7516-9.

86. Mannel DN. Advances in sepsis research derived from animal models. *Int J Med Microbiol*. 2007 Sep;297(5):393-400.