

RUNNING TITLE PAGE

Early HbA1c: Predictors and trajectories

Corresponding author:

Dr Veena Mazarello Paes

Population, Policy and Practice Research and Teaching Department,

UCL Great Ormond Street Institute of Child Health,

1st floor, Wellcome Trust Building,

30 Guilford Street

London WC1N 1EH

UK

Email: veena.paes.14@ucl.ac.uk

TITLE PAGE

Factors predicting poor glycaemic control in the first two years of childhood onset type 1 diabetes in a cohort from East London, UK: Analyses using mixed effects fractional polynomial models

Veena Mazarello Paes^{1, 2}, Jessica K Barrett³, David B Dunger^{2, 4}, Evelien F Gevers^{5, 6}, David C Taylor-Robinson⁷, Russell M Viner^{1, 8}, Terence J Stephenson¹

¹Great Ormond Street Institute of Child Health, University College London, UK

²Department of Paediatrics, University of Cambridge, UK

³MRC Biostatistics Unit, University of Cambridge, UK

⁴Wellcome Trust - MRC Institute of Metabolic Sciences, University of Cambridge, UK

⁵Centre for Endocrinology, William Harvey Research Institute, Queen Mary University, UK

⁶Barts Health NHS Trust, Royal London Children's Hospital, UK

⁷University of Liverpool, UK

⁸The Royal College of Paediatrics and Child Health, London, UK

Authors details:

1. Veena Mazarello Paes MSc, MPH, MPhil, PhD

Population, Policy and Practice Research and Teaching Department,

UCL Great Ormond Street Institute of Child Health,

London WC1N 1EH, UK

Department of Paediatrics,

University of Cambridge,

Cambridge, UK

2. Jessica K Barrett MSc, PhD

MRC Biostatistics Unit,
University of Cambridge,
Cambridge, UK
Email: jkb23@medschl.cam.ac.uk

3. David B Dunger MD, FMedSci, FRCPCH
Department of Paediatrics,
University of Cambridge,
Cambridge, UK
Wellcome Trust - MRC Institute of Metabolic Sciences,
University of Cambridge,
Cambridge, UK
Email: dbd25@cam.ac.uk

4. Evelien F Gevers MD, PhD
Centre for Endocrinology, William Harvey Research Institute,
Queen Mary University,
London, UK
Department of Paediatric Endocrinology,
Barts Health NHS Trust,
Royal London Children's Hospital,
London, UK
Email: evelien.gevers@bartshealth.nhs.uk

5. David C Taylor-Robinson MB ChB, MPH, PhD, FFPH, MRCPCH
Department of Public Health and Policy
University of Liverpool,

Liverpool, UK

Email: david.taylor-robinson@liverpool.ac.uk

6. Russell M Viner MBBS, PhD, FRCPCH

Population, Policy and Practice Research and Teaching Department,

UCL Great Ormond Street Institute of Child Health,

London, UK

The Royal College of Paediatrics and Child Health

London, UK

Email: r.viner@ucl.ac.uk

7. Terence J Stephenson DM, FRCPCH

Population, Policy and Practice Research and Teaching Department,

UCL Great Ormond Street Institute of Child Health,

London, UK

Email: t.stephenson@ucl.ac.uk

Word count: abstract: 250 and main text: 4144

Number of tables: 2

Number of online supplemental tables: 6

Number of online supplemental figures: 3

Number of figures: 2

ABSTRACT & KEY WORD PAGE

Factors predicting poor glycaemic control in the first two years of childhood onset type 1 diabetes in a cohort from East London, UK: Analyses using mixed effects fractional polynomial models

Background/aims: Poor early glycaemic control in childhood onset type 1 diabetes (T1D) is associated with future risk of acute and chronic complications. Our aim was to identify the predictors of higher glycated hemoglobin (HbA1c) within 24 months of T1D diagnosis in children and adolescents.

Methods: Mixed effects models with fractional polynomials were used to analyse longitudinal data of patients <19 years of age, followed from T1D diagnosis for up to 2 years, at three diabetes clinics in East London, UK.

Results: A total of 2209 HbA1c observations were available for 356 patients (52.5% female; 64.4% non-white), followed from within 3 months of diagnosis during years 2005 to 2015, with a mean±SD of 6.2±2.5 HbA1c observations/participant. The mean age and HbA1c at diagnosis were 8.9±4.3 years and 10.7% ±4.3% (or expressed as mmol/mol HbA1c mean±SD 92.9±23.10 mmol/mol) respectively.

Over the 2 years following T1D diagnosis, HbA1c levels were mostly above the National Institute for Health, Care and Excellence (NICE), UK recommendations of 7.5% (<58mmol/mol). Significant ($p<0.05$) predictors of poorer glycaemic control were: Age at diagnosis (12-18 years), higher HbA1c at baseline (>9.5% i.e. >80mmol/mol), clinic site, non-white ethnicity and period (pre-year 2011) of diagnosis. Additionally in univariable analyses, frequency of clinic visits, HbA1c at diagnosis and type of insulin treatment regimen showed association with poor glycaemic control ($p<0.05$).

Conclusions: Major risk factors of poorer glycaemic control during 3-24 months following childhood onset T1D are: diagnosis prior to 2011, higher HbA1c levels at baseline, age at diagnosis, non-white ethnicity and clinic site.

Key words: Predictors, HbA1c, Type 1 diabetes mellitus, paediatric, longitudinal analyses

LIST OF ABBREVIATIONS

AIC: Akaike information criterion

BIC: Bayesian information criterion

BMI: Body mass index

DKA: Diabetic Ketoacidosis

HbA1c: Glycated haemoglobin A1c

IMD: Indices of multiple deprivations

NICE: National Institute for Health, Care and Excellence

T1D: Type 1 diabetes

-2LL: - 2 Log likelihood

INTRODUCTION

Evidence suggests that early glycaemic control in childhood onset Type 1 Diabetes (T1D) tracks into later life (1-4) and predicts risk of future vascular complications (5, 6). Studies in adults show a reduced risk of vascular complications and mortality in those who maintained lower HbA1c levels during early stages of the disease (7, 8). It is therefore important to identify if this is the case or whether other factors influence glycaemic trends from T1D diagnosis in children and adolescents (9, 10).

Poor glycaemic control is seen in children and adolescents soon after T1D diagnosis (11). But factors influencing these outcomes are poorly understood due to studies: with small sample size,

focused on older children and adolescents or investigated a limited number of demographic variables (12). This causes a difficulty in identifying and predicting a “risk signature” of poor control during the course of the disease in young T1D patients (13).

In order to investigate the predictors (socio-demographic, biological and clinical) affecting the glycaemic control within 24 months of T1D diagnosis in children and adolescents, we undertook a retrospective analysis of data collected prospectively from three diabetes clinics based in East London, UK.

METHODS

We investigated the predictors of glycaemic control (HbA1c levels) during 3 to 24 months from T1D diagnosis in young children and adolescents. Data included paediatric (<19 years of age at diagnosis) patients, recorded in the hospital database as newly diagnosed with T1D, in the period between 1 January 2005 and 31 December 2015 and receiving care from one of the three paediatric diabetes clinics of the Barts Health NHS Trust in East London, UK.

Due to increasing knowledge of more rare causes of diabetes, patients may have been misidentified during the study period as Type 1 Diabetes (default in the database), and may have had Type 2 Diabetes, Maturity Onset Diabetes or secondary diabetes. The records of three patients with very low HbA1c were looked at and these patients most likely did not have T1D and were excluded.

Each of the three clinics had a separate but unequal, multi-disciplinary diabetes specialist care team made up of clinicians, nurses, dieticians, psychologists and with access to interpreters. For example, a 24 hour paediatric Diabetes consultant on call service was available at only one of the clinics. The three clinics became part of a network in 2012 but care continued to be delivered by the respective care team at each clinic as per the updated care package recommended by the National Institute of Health and Care Excellence (NICE) (14). Type and methods for the electronic recording of anthropometric, socio-demographic and clinical data, collected at diagnosis and at each clinic visit

were not always uniform between the three clinics. For example, one of the three clinics recorded data of fewer patients in the first 3 months from T1D diagnosis (Supplementary figure panel 1a&b).

Variables

We used HbA1c, a measure of average blood glucose over the previous 8-12 weeks, as our marker of diabetes control and as the dependent variable in these analyses. Venous and finger HbA1c measured using BioRad Variant II Turbo (in laboratory) and Siemens/Bayer DCA 2000 (in clinic) respectively at diagnosis and each clinic visit. The HbA1c values recorded as percentages were converted to mmol/mol using the formula: $(\text{HbA1c value in \%} - 2.15) \times 10.929$. Baseline was defined as period within 3 months of T1D diagnosis and patients with baseline measurements were eligible. HbA1c levels at diagnosis (measurement within the first month of diagnosis) and at baseline (averaged HbA1c measurements within first 3 months of diagnosis) were assessed by tertiles or categorized into low (<7.5% or <58mmol/mol), moderate (7.5- 9.5% or 58-80 mmol/mol) and high (> 9.5% or >80mmol/mol) based on UK targets that were acceptable during the study period.

Based on our systematic review (12) and captured in the dataset, potential predictors of HbA1c which we used as independent variables in our analyses were sex, age at diagnosis, season/month/year/period of T1D diagnosis, ethnicity, index of multiple deprivation (IMD), clinic site, number of clinic visits, insulin regimen, body mass index (BMI), pH at diagnosis, HbA1c at diagnosis and baseline.

We tested the accuracy of the data in a number of patients and found that the insulin treatment may have been inaccurately documented in the database. Sometimes details of insulin regimen such as dosage, injection frequency and pump treatment were missing during the earlier years (prior to 2012) and before the individual clinics joined to form a network.

Age at diagnosis was treated as a categorical variable and stratified into the following groups: 0–5 (preschool), 6–11 (pre-pubertal), and 12-18 (pubertal/adolescent) years. Season of diagnosis was categorized into autumn (September to November), winter (December to February), spring (March to May) and summer (June to August). Period of diagnosis was categorized into pre-2011 and post-2011 to check for differences in outcomes between the two periods since the three diabetes clinics became part of the same network in 2012. Ethnicity was categorized into white and non-white for the main analysis. Socioeconomic status was measured using the UK IMD (2010), which gives a relative measure of neighborhood-level socioeconomic deprivation based on seven domains for small areas in England: income, employment, health and disability, education, skills and training, barriers to housing and services, living environment and crime. IMD scores were categorized into three groups with 3 being least deprived and 1 being most deprived (15). The number of clinic visits per patient was treated as a continuous variable.

Insulin treatment (between years 2005-2015) was categorized as continuous subcutaneous insulin infusion pump, 3 or more injections per day or 1 or 2 injections per day. BMI (obtained from the records of physical examination involving measurements of height and weight) measured as kg/m² at each clinic visit, was treated as a continuous variable and was standardized to BMI (WHO) z-scores. These were further categorized into normal weight, overweight/obese and thin (16).

Whole blood pH at diagnosis was treated as a continuous variable and in a sensitivity analysis was categorized into acidotic (pH<7.3), normal (pH 7.3 to 7.45) and alkalotic (pH>7.45).

Analyses

Baseline characteristics of the cohort were reported as means and standard deviations for continuous variables or counts and percentages for categorical variables. Baseline characteristics were compared across HbA1c categories at baseline using a global χ^2 test for categorical variables or a Wald test from linear regression for continuous variables.

Repeated measurements of the outcome HbA1c were modelled using mixed effects models. Fractional polynomials were used to account for nonlinear trajectories of HbA1c over time (17, 18). We estimated models for both 0 to 24 months (Model 1: with 253 T1D patients followed from diagnosis) and 3 to 24 months (Model 2: with 335 T1D patients followed from baseline), as data showed dramatic instability in HbA1c during the first three months after diagnosis, as expected. In this paper we report the results of the investigations of predictors of HbA1c levels from 3 to 24 months in the 335 patients who were followed from baseline. The trajectories and predictors of HbA1c during 0-3 months will be studied in depth and reported separately. The polynomial terms in the model were chosen by comparing model fit amongst all fractional polynomial models with one, two or three polynomial terms with any combination of the following polynomial powers of time: (-2 -1 -0.5 0 0.5 1 2 3), where 0 represents $\log(t)$ and a repeated power (p p) indicates the functions t^p and $\log(t)t^p$. The outcome model for 0 to 24 months (Model 1) was

$$HbA1c_i(t) = \beta_0 + \beta_1 \log(t) + \beta_2 t + \beta_3 t^2 + b_{0,i} + b_{1,i} \log(t) + b_{2,i} t + b_{3,i} t^2 + \varepsilon_i$$

and the outcome model for 3 to 24 months (Model 2) was

$$HbA1c_i(t) = \beta_0 + \beta_1 \sqrt{t} + \beta_2 \log(t) \sqrt{t} + b_{0,i} + b_{1,i} \sqrt{t} + b_{2,i} \log(t) \sqrt{t} + \varepsilon_i$$

where $HbA1c_i(t)$ is the outcome of interest for individual i . $\beta_0, \beta_1, \beta_2$ and β_3 are fixed effect coefficients describing the population average trajectory of the outcome and $b_{0,i}, b_{1,i}, b_{2,i}$ and $b_{3,i}$ are normally distributed random effects describing the deviation of individual i 's trajectory from the population average. The best model was selected within each class of one-term, two-term or three-term models. Fixed effects and random effects were included in the model selection procedure. However, the expectation of the random effects was assumed to be zero. Additionally, both the Akaike information criterion (AIC) and the Bayesian information criterion (BIC) were used to assess model fit (Supplementary tables S1 a & b give the AIC and BIC results from each class of models). Time-dependent predictors were not included in the model. However, mean of HbA1c and BMI measurements taken during the 0-3 month period from diagnosis were used as predictors during 3-24 month follow-up.

To study the predictors of glycaemic control between 3 and 24 months of diagnosis, predictor variables were included in the mixed effects model. The effect of the predictor variables represents a shift in the population average trajectory and can therefore be interpreted as a mean difference in HbA1c levels over time. Initially each predictor variable was separately added to the models 1 and 2, and significant predictor variables were then included in a multivariable model. Predictors showing significant interactions with sex, clinic site and age at diagnosis were also included in the multivariable model. All models include $\sqrt{\quad}$ and $\log^*\sqrt{\quad}$ functions of time as both fixed and random effects. Results for potential predictors were reported as mean differences and 95% confidence intervals (CI). Sensitivity analyses were conducted where appropriate.

Additionally, the above initial mixed model with each predictor variable were then fitted to include interactions between the predictor variable and the polynomial time functions, allowing the shape of the HbA1c trajectory to depend on the level of the predictor variable.

Analyses were performed using Stata 15, StataCorp. College Station, TX, 2015. The fractional polynomial models were fitted using the stata “fp” command.

RESULTS

A total of 356 T1D patients (52.5% female) with 2209 HbA1c observations (mean 6.2 ± 2.5 HbA1c observations/participant) and a follow-up of 2 years from T1D diagnosis at the 3 sites, during the study period (2005-2015) were eligible for inclusion in our study at baseline.

However, in the analyses of data covering the 3-24 months period from diagnosis, a total of 335 patients (53.4% female) with 1627 HbA1c observations (mean 7.4 ± 2.0 HbA1c observations/participant) were eligible. Of these, 253 (75.5%) patients were followed from and had HbA1c measurements within first month of T1D diagnosis (See Table 1 for baseline characteristics). Insulin regimen may have been inaccurately documented for some patients and data on insulin

dosage was missing, which restricted the analysis of this variable against our objectives. pH at diagnosis (mean \pm SD 7.27 \pm 0.16) was available for only 186/356 (52.25%) patients and 80 out of these 186 (43%) patients were acidotic. Mean BMI at diagnosis was 18.3 \pm 3.6 (BMI WHO z-score at diagnosis and at baseline was 0.46 \pm 1.2 and 0.53 \pm 1.3 respectively) and frequency of clinic visits from diagnosis ranged from 1 to 13. (See Supplementary Table S2 for baseline characteristics by HbA1c tertiles)

The fractional polynomial model 1 fitted to HbA1c measurements taken 0-24 months post T1D diagnosis revealed that the initial, high HbA1c levels of the cohort dropped rapidly after diagnosis due to initiation of insulin treatment. However, subsequently after around three months of diagnosis, the population mean HbA1c levels rose steadily above the then (2005-2015) NICE recommended targets and remained high throughout the entire duration of follow-up. The model 2 fitted to HbA1c measurements 3-24 months gave similar results for that period (Fig panel 1).

We then examined the association of various predictor variables with HbA1c trajectories from 3 to 24 months of T1D diagnosis. At the univariable level, pubertal age (12-18 years) at diagnosis, year of diagnosis pre 2011, non-white ethnicity, clinic site, frequency of clinic visits, insulin regimen type and higher HbA1c levels at diagnosis and during 0-3 months after diagnosis were significant predictors of higher mean HbA1c over the follow up period (Table 2).

After adjustment for predictors that were statistically significant in univariable analyses, the following predictors of glycaemic control remained statistically significant in a multivariable analysis (Table 2): age at diagnosis (12-18 years), earlier years of diagnosis (pre 2011), non-white ethnicity, HbA1c during the 1st three months after diagnosis) and clinic site.

The results were similar when the multivariable analyses were repeated in the sensitivity analyses with predictors: HbA1c tertiles at baseline (Supplementary table S3a) and HbA1c at diagnosis (supplementary table S3b).

Additional univariable and multivariable sensitivity analyses for predictor variables from 1-3 months after diagnosis consistently revealed the following as significant predictors of poor glycaemic control: year of diagnosis (pre-2011), clinic site, HbA1c and BMI (WHO z score) at 1-3 months after diagnosis.“

There was no effect modification when interaction with age at diagnosis, sex or clinic site were added to the mixed initial (with each predictor variable) and multivariable models. However, interaction with ethnicity by clinic site showed some significance in both univariable ($p=0.04$) and multivariable ($p=0.05$) analyses. The interaction year of diagnosis by gender seems to have an effect of borderline significance in multivariable analyses ($p=0.04$). Age at diagnosis by clinic interaction showed some significance ($p=0.05$) in the univariable analyses and $p=0.03$ in the sensitivity analyses when predictor HbA1c at diagnosis was added to the multivariable analyses (supplementary table S3b).

We then examined the following predictors by including interactions with polynomial terms in time since diagnosis in the initial mixed models with each predictor variable: sex, age at T1D diagnosis, year of diagnosis, ethnicity, clinic site, insulin regimen, HbA1c level at diagnosis and baseline (also tertiles at baseline) and BMI (WHO) z score at baseline (Supplementary table S4). The interaction with polynomial time terms allowed for the shapes of the HbA1c trajectories to differ between the compared groups. The following predictors had significant interactions with polynomial time terms: age at diagnosis, HbA1c at diagnosis (Figure panel 2) and HbA1c at baseline.

Interaction between age at diagnosis and polynomial time terms showed that pre-pubertal (6-11years) and pubertal children (12-18years) experienced a sharper increase in HbA1c after 3 months than preschool (0-5years) aged children, leading to higher levels throughout the study period ($p=0.001$) (Figure panel 2a).

There was a significant difference in HbA1c trajectories between those with low, moderate and high HbA1c levels at diagnosis (Figure panel 2b) and baseline ($p=0.001$ and $p=0.01$ respectively). All children had a similar starting point at 3 months but, the glycaemic control steadily deteriorated in those children with HbA1c levels above 7.5% (58 mmol/mol) at diagnosis or at baseline. But, there is some uncertainty in the HbA1c trajectory of those who had HbA1c levels $<7.5\%$ (<58 mmol/mol) at diagnosis or at baseline as there were only 13 (5.14%) and 38 (10.67%) patients in these groups at diagnosis and baseline respectively and the diagnosis details of these patients could not be verified. However, we found that the difference between HbA1c trajectories was of significance ($p=0.02$) when the analyses were repeated by HbA1c tertiles at baseline (Supplementary figure 2)

DISCUSSION

Our longitudinal, retrospective, cohort study of glycaemic control during 3-24 months after T1D diagnosis in 356 children and adolescents of multi-ethnic backgrounds, receiving care; between 1 January 2005 and 31 December 2015; at a network of three diabetes clinics located in East London, UK. We used fractional polynomial modelling to estimate the non-linear mean HbA1c trajectories to identify the predictors (socio-demographic, biological and clinical) of glycaemic control during the first 2 years of T1D diagnosis. We found that the partial remission period ends at around 3 months from diagnosis, and that independent risk factors of poorer glycaemic control during the 2005-2015 study period were higher levels of HbA1c at baseline, higher age at diagnosis (12-18 years), non-white ethnicity, year of diagnosis (pre- year 2011) and clinic factors.

It is important to ensure good glycaemic control from T1D onset, but lack of evidence on predictors of glycaemic control restricts prediction of future glycaemic trends and risk of complications in children and young people, thus hampering the proactive prognosis and management of glycaemic control in high risk patients. Furthermore, glycaemic control is an important outcome and an exposure and/or mediator of later-life outcomes. Previously, NICE recommended HbA1c targets below 7.5% (58mmol/mol) (19). However, since year 2015, NICE recommends achieving a target of HbA1c levels of 6.5% (48mmol/mol) or below including during the honeymoon phase (period shortly after T1D diagnosis) when the existing beta cells of the pancreas continue to produce some insulin to help control blood glucose. Our study found that it was challenging for the cohort to achieve recommended blood glucose and HbA1c levels after the honeymoon period and during the first 24 months of T1D diagnosis during the study period (2005-2015) as indicated by other studies (20, 21).

In this cohort of young children and adolescents with T1D, we found that higher HbA1c levels (>9.5% i.e. >80mmol/mol) at diagnosis/baseline were associated with higher HbA1c levels during the first 24 months of T1D diagnosis. Elevated HbA1c levels at diagnosis may be due to delayed access to medical treatment and/or delayed T1D diagnosis but evidence suggests that better glycaemic control can be achieved during this sensitive period through disease awareness and change of behaviors in diagnostic and care practice (22, 23).

Children diagnosed at an older age (12 to 18 year olds) had significantly poorer glycaemic control during the study period. Adolescence has proven to be yet another sensitive period which influences formation of enduring behaviours and diabetes self-management habits (24). This may relate to a complex interaction of factors such as increased insulin resistance, eating disorders, psychological disturbances and the psychosocial demands of adolescence leading to decreased adherence to management plans (25, 26).

Children diagnosed pre - 2011 had poorer glycaemic control which may be due to older diagnostic testing, treatment and recording strategies as reported by other studies (13, 27). Additionally, it may indicate that being part of a network improved the glycaemic outcomes of the patients registered at the three clinics, as the three clinics became part of a network only from 2012. However, this is a causal hypothesis which the currently reported data and analyses cannot address.

We also found that overweight or obese children had better glycaemic control, although this effect was statistically non-significant. This is likely to reflect the association of weight gain during T1D treatment with better adherence to insulin regimens and therefore BMI may represent an indirect marker of good glycaemic control. However, there is uncertainty in the HbA1c trajectory of underweight/thin children in our study as there were <10% children in this group. Evidence from other studies on the association of BMI with HbA1c is also inconclusive (28-30). Also, a small minority of patients may have been misclassified during the study period and may have had T2D, MODY or secondary diabetes.

Non-white ethnicity was associated with elevated HbA1c levels. This association is consistent with other studies and may be due to biological, cultural and lifestyle differences, sub-optimal treatment prescription or reluctance to try new therapy (31-33). Also, some ethnicities may have an inherent genetic predisposition to adverse health outcomes from T1D (34). The role of clinic in improving glycaemic control in children with T1D (particularly ethnic minorities) and whether more frequent clinic visits would result in better care and adherence to treatment among the T1D patients from ethnic minority remains unclear.

However, we did find that patients at one of the clinics had better glycaemic control than others and that more frequent clinic visits (univariable analyses) were associated with higher HbA1c levels. More frequent appointments for children with poor control may have been offered in some clinics. Other studies have indicated associations with glycaemic control and clinic related factors (2, 28,

35). Some studies have also highlighted the active role of diabetes teams in achieving glycaemic targets (36, 37). Frequency of clinic attendance may be high in those patients living closer to the diabetes clinics. Poor or non-attendance is associated with elevated diabetes complications risk due to non-adherence to treatment in the high risk groups (38).

Details of insulin regimen such as injection frequency, pump treatment, dosage and treatment changes were missing which meant that we were restricted in our investigations of this variable against our research objectives, although, pump regimen (univariable analyses) was associated with lower HbA1c levels. However, sensitivity analysis with and without adjusting for insulin regimen did not affect our conclusions. It would be useful to investigate the benefits of intensive insulin treatment earlier on in the course of the disease management and whether early HbA1c levels tracks in later years and reduces the risk of vascular complications.

In our study we found that sex showed no association (univariable analyses) with glycaemic control although interaction with year of diagnosis by sex (univariable and multivariable analyses) showed some association. This is contrary to the findings of some studies (5, 13, 30); whilst there are others who have reported no association (28, 35) between sex and HbA1c levels.

No association was seen between pH at diagnosis (proxy for DKA) and HbA1c levels during the 3-24 months from T1D diagnosis. Although this was also the conclusion of some studies (28, 30, 35), others have shown an association (39-41). It was not clear if the general practitioners (GP's) managed the T1D symptoms in any of these patients prior to referral to the diabetes clinics for specialist care. We did not have access to GP prescription data to confirm if any form of glycaemic control started prior to clinic diagnosis date although it would be extremely unusual in the UK for a GP to commence insulin treatment in a child with diabetes.

STRENGTHS AND LIMITATIONS

Historically, mixed effects modelling have been used to describe trajectories of repeated outcomes within and between clustered individuals. But since glycaemic trajectories are often non-monotonic and non-linear, they require more flexible models to accurately investigate the factors which influence the HbA1c levels or the outcomes it is associated with. To our knowledge this is the first longitudinal study to investigate predictors of HbA1c during the first 24 months of T1D duration using mixed effects multilevel models with fractional polynomial time terms which allowed flexibility and parsimony in the modelling of non-linear trajectories. The application of fractional polynomials requires the comparison of many models for model selection and represents a situation of multiple testing. The modest sample size was suitable for the two step approach using fractional polynomials in multilevel modelling. However, the sample size limitation precluded the enhancement of complexity of fitting models with all covariates. Assessment of predictors of HbA1c was performed in a two-step approach, with polynomial terms selected in the first step and predictor selection performed in the second step. The effect estimates of present models may be biased due to the two-step procedure, and no adjustment has been made for multiple testing. Data were abstracted from routinely collected electronic medical records, thus limiting selection and recall bias. It was not possible to ascertain that the primary outcome or exposure was not associated with loss to follow-up. The total sample size was modest but exceeded the minimum ($N > 100$) suggested for multilevel modelling and may have limited the power to detect weaker associations. The study cohort was rich in ethnic diversity and an unusually high proportion of socially deprived population, which enabled us to study these predictors in greater detail but was also a limitation since there was less contrast.

Data was limited to patients receiving care between 2005 and 2015 at three diabetes clinics which formed a network in 2012. So the methods of data collection and recording may have differed and changed over the study period and may not be generalizable to other populations. The data used were from a dataset maintained for ongoing clinical use across the 3 clinic sites, and, as with most clinical datasets, was not subject to the detailed data checking that would be expected in a research dataset. We verified data where possible in association with the clinic physicians. However it was not

practicable to examine and verify each individual clinical record. Additionally, it was not possible to investigate whether the disparity in the number of clinic visits and treatment across the study population was due to individual and/or clinic factors. This is a retrospective analysis of an observational study and residual confounding cannot be ruled out. Details of laboratory methods and instruments used across clinics between 2005 and 2015 could not be confirmed.

CONCLUSIONS

Our study identified higher HbA1c levels at baseline, non- white ethnicity, clinic site, BMI and year of diagnosis as major risk factors of poor glycaemic control during the first 24 months of children diagnosed with T1D. Since higher HbA1c at T1D diagnosis is associated with poorer subsequent HbA1c levels and is associated with increased risk of developing complications, these findings could help clinicians, policy makers and researchers better understand the characteristics of type 1 diabetes and encourage quick identification of high risk patients for consideration of appropriate individualized treatment strategy to meet the HbA1c targets during the 24 months following the diagnosis of T1D. These findings will also help explore the underlying explanations for poor glycaemic outcomes so further efforts are made to overcome these differences.

FUTURE RESEARCH

Further research into the reasons for differences in HbA1c outcomes is required. Also predictors of glycaemic control in the first 3 months of diagnosis and beyond 2 years of diagnosis need further investigation.

ACKNOWLEDGEMENTS

We thank the funders and colleagues from the University College London who have helped make this project a success, especially, GOS ICH R&D and UCL Ethics for approving the project, Dr Rakesh Amin, Dr Amal Khanolkar and Dr Tiara Paramita for initial discussions and help in the acquisition of Barts Health NHS Trust data. We also thank Dr Paramita Cifelli, Dr Abdul Moodambail,

Dr Jeremy Allgrove and Dr Anne Dawnay from Barts Health NHS Trust for providing us with the data - clarifying data collection methods and variable definitions. Our sincere thanks to Prof Tim Cole, Prof Mario Cortina Borja, Prof Allan Hackshaw, Dr Graham Wheeler and Dr Nicholas Counsell from UCL for initial discussions and ideas on the statistical analyses plans. We especially thank Prof Helen Roberts from UCL and Dr Justin Warner from Cardiff for commenting on the manuscript. We also thank Dr Loredana Marcovecchio and Dr Stephen Burgess from the University of Cambridge for sharing their experiences with similar data and discussions on analyses plans.

FUNDING

Reference: 109/0001: The Policy Research Unit in the Health of Children, Young People and Families is funded by the Department of Health and Social Care Policy Research Programme. This report is independent research commissioned and funded by the National Institute for Health Research Policy Research Programme. The views expressed in this publication are those of the authors and not necessarily those of the NHS, the National Institute for Health Research, the Department of Health and Social Care or its arm's length bodies, and other Government Departments. This research was supported by the National Institute for Health Research Biomedical Research Centre at Great Ormond Street Hospital for Children NHS Foundation Trust and University College London. Jessica Barrett is funded by the MRC Unit Programme (MC_UU_00002/5). David Taylor-Robinson is funded by the MRC on a Clinician Scientist Fellowship (MR/P008577/1)

CONFLICT OF INTEREST

No potential conflict of interest was reported by the authors.

CONTRIBUTION STATEMENT

VMP designed the study, analyzed the data and wrote the first draft of the manuscript. JB advised on the statistical plan and analyses. EG coordinated the acquisition of data and clarified data related queries. JB, DD, RV, EG, DTR and TS contributed to the study plan, analyses and critical revision

of the manuscript. All authors approved the final version. TS is the principal investigator on the funding from the National Institute for Health Research Policy Research Programme and lead for the Long Term Conditions theme of this work. TS and VMP are guarantors of this work, and as such had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analyses.

SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article.

A) Supplementary tables:

Supplementary Table S1a: Fractional polynomial three dimension model fit comparison algorithm for period 0-24 months post T1D diagnosis

Supplementary Table S1b: Fractional polynomial three-dimension model fit comparison algorithm for period 3-24 months post T1D diagnosis

Supplementary Table S2: Baseline characteristics of study population by HbA1c tertiles. Data are mean (\pm SD) or counts (%)

Supplementary Table S3a: Predictors of glycaemic control in children and adolescents from 3 to 24 months post-T1D diagnosis (MVA with predictor HbA1c tertiles at baseline)

Supplementary Table S3b: Predictors of glycaemic control in children and adolescents from 3 to 24 months post-T1D diagnosis (MVA with predictor HbA1c at diagnosis)

Supplementary Table S4: Analysis of interactions between the predictor variable and the polynomial time functions

B) Supplementary figures:

Supplementary figure panel 1a: Histograms showing patterns of visits to clinic during 0-3 months after T1D diagnosis

Supplementary figure panel 1b: Histograms showing patterns of visits to clinic after 3 months of T1D diagnosis

Supplementary figure 2: Predictor HbA1c tertiles at baseline showing significant differences between HbA1c trajectories over time

REFERENCES

1. Edge JA, James T, Shine B. Persistent individual tracking within overall improvement in HbA1c in a UK paediatric diabetes clinic over 15 years. *Diabetic medicine : a journal of the British Diabetic Association*. 2010;27(11):1284-8.
2. Lawes T, Franklin V, Farmer G. HbA1c tracking and bio-psychosocial determinants of glycaemic control in children and adolescents with type 1 diabetes: retrospective cohort study and multilevel analysis. *Pediatric diabetes*. 2014;15(5):372 - 83.
3. Nirantharakumar K, Mohammed N, Toulis KA, Thomas GN, Narendran P. Clinically meaningful and lasting HbA1c improvement rarely occurs after 5 years of type 1 diabetes: an argument for early, targeted and aggressive intervention following diagnosis. *Diabetologia*. 2018.
4. Hofer SE, Raile K, Frohlich-Reiterer E, Kapellen T, Dost A, Rosenbauer J, et al. Tracking of metabolic control from childhood to young adulthood in type 1 diabetes. *J Pediatr*. 2014;165(5):956-61.e1-2.
5. Samuelsson U, Steineck I, Gubbjornsdottir S. A high mean-HbA1c value 3-15 months after diagnosis of type 1 diabetes in childhood is related to metabolic control, macroalbuminuria, and retinopathy in early adulthood--a pilot study using two nation-wide population based quality registries. *Pediatric diabetes*. 2014;15(3):229-35.
6. Cutfield SW, Derraik JG, Reed PW, Hofman PL, Jefferies C, Cutfield WS. Early markers of glycaemic control in children with type 1 diabetes mellitus. *PloS one*. 2011;6(9):e25251.
7. Fullerton B, Jeitler K, Seitz M, Horvath K, Berghold A, Siebenhofer A. Intensive glucose control versus conventional glucose control for type 1 diabetes mellitus. *Cochrane Database Syst Rev*. 2014;2:Cd009122.
8. Kahler P, Grevstad B, Almdal T, Gluud C, Wetterslev J, Lund SS, et al. Targeting intensive versus conventional glycaemic control for type 1 diabetes mellitus: a systematic review with meta-analyses and trial sequential analyses of randomised clinical trials. *BMJ open*. 2014;4(8):e004806.
9. Rohan JM, Rausch JR, Pendley JS, Delamater AM, Dolan L, Reeves G, et al. Identification and prediction of group-based glycemic control trajectories during the transition to adolescence. *Health psychology : official journal of the Division of Health Psychology, American Psychological Association*. 2014;33(10):1143-52.
10. Hilliard ME, Guilfoyle SM, Dolan LM, Hood KK. Prediction of adolescents' glycemic control 1 year after diabetes-specific family conflict: the mediating role of blood glucose monitoring adherence. *Archives of pediatrics & adolescent medicine*. 2011;165(7):624-9.
11. Petitti DB, Klingensmith GJ, Bell RA, Andrews JS, Dabelea D, Imperatore G, et al. Glycemic control in youth with diabetes: the SEARCH for diabetes in Youth Study. *J Pediatr*. 2009;155(5):668-72.e1-3.
12. Mazarello Paes V, Charalampopoulos D, Edge J, Taylor-Robinson D, Stephenson T, Amin R. Predictors of glycemic control in the first year of diagnosis of childhood onset type 1 diabetes: A systematic review of quantitative evidence. *Pediatric diabetes*. 2017; doi:10.1111/peidi.12530.
13. Clements MA, Lind M, Raman S, Patton SR, Lipska KJ, Fridlington AG, et al. Age at diagnosis predicts deterioration in glycaemic control among children and adolescents with type 1 diabetes. *BMJ open diabetes research & care*. 2014;2(1):e000039.
14. Beckles ZL, Edge JA, Mugglestone MA, Murphy MS, Wales JK. Diagnosis and management of diabetes in children and young people: summary of updated NICE guidance. *Bmj*. 2016;352:i139.
15. Ramsay SE, Morris RW, Whincup PH, Subramanian SV, Papacosta AO, Lennon LT, et al. The influence of neighbourhood-level socioeconomic deprivation on cardiovascular disease mortality in older age: longitudinal multilevel analyses from a cohort of older British men. *Journal of epidemiology and community health*. 2015;69(12):1224-31.
16. Cole TJ, Lobstein T. Extended international (IOTF) body mass index cut-offs for thinness, overweight and obesity. *Pediatric obesity*. 2012;7(4):284-94.

17. Royston P, Altman DG. Regression Using Fractional Polynomials of Continuous Covariates: Parsimonious Parametric Modelling. *Journal of the Royal Statistical Society Series C (Applied Statistics)*. 1994;43(3):429-67.
18. Sauerbrei W, Meier-Hirmer C, Benner A, Royston P. Multivariable regression model building by using fractional polynomials: Description of SAS, STATA and R programs. *Computational Statistics & Data Analysis*. 2006;50(12):3464-85.
19. NICE Guidance CG15 Type 1 diabetes in children and young people: full guideline. National Institute for Health and Clinical Excellence. 2004, [<http://guidance.nice.org.uk/CG15/Guidance/pdf/English>].
20. RCPCH. National Paediatric Diabetes Audit Report 2015-16. Care processes and outcomes 2017. Available from: https://www.rcpch.ac.uk/system/files/protected/page/Complete%20NPDA%202015-16%20report%20FINAL_0.pdf.
21. Miller KM, Foster NC, Beck RW, Bergenstal RM, DuBose SN, DiMeglio LA, et al. Current state of type 1 diabetes treatment in the U.S.: updated data from the T1D Exchange clinic registry. *Diabetes care*. 2015;38(6):971-8.
22. Salardi S, Porta M, Maltoni G, Cerutti F, Rovere S, Iafusco D, et al. Ketoacidosis at diagnosis in childhood-onset diabetes and the risk of retinopathy 20years later. *Journal of diabetes and its complications*. 2016;30(1):55-60.
23. Diabetes UK: Do You Know The 4T's Of Type 1 Diabetes? 2016 <https://www.diabetes.org.uk/the4TS>.
24. King PS, Berg CA, Butner J, Drew LM, Foster C, Donaldson D, et al. Longitudinal trajectories of metabolic control across adolescence: associations with parental involvement, adolescents' psychosocial maturity, and health care utilization. *The Journal of adolescent health : official publication of the Society for Adolescent Medicine*. 2012;50(5):491-6.
25. Amiel SA, Sherwin RS, Simonson DC, Lauritano AA, Tamborlane WV. Impaired insulin action in puberty. A contributing factor to poor glycemic control in adolescents with diabetes. *The New England journal of medicine*. 1986;315(4):215-9.
26. Peveler RC, Bryden KS, Neil HA, Fairburn CG, Mayou RA, Dunger DB, et al. The relationship of disordered eating habits and attitudes to clinical outcomes in young adult females with type 1 diabetes. *Diabetes care*. 2005;28(1):84-8.
27. Hanberger L, Akesson K, Samuelsson U. Glycated haemoglobin variations in paediatric type 1 diabetes: the impact of season, gender and age. *Acta Paediatr*. 2014;103(4):398-403.
28. Viswanathan V, Sneeringer MR, Miller A, Eugster EA, DiMeglio LA. The utility of hemoglobin A1c at diagnosis for prediction of future glycemic control in children with type 1 diabetes. *Diabetes research and clinical practice*. 2011;92(1):65-8.
29. Redondo MJ, Rodriguez LM, Escalante M, O'Brian Smith E, Balasubramanyam A, Haymond MW. Beta cell function and BMI in ethnically diverse children with newly diagnosed autoimmune type 1 diabetes. *Pediatric diabetes*. 2012;13(7):564-71.
30. Akesson K, Hanberger L, Samuelsson U. The influence of age, gender, insulin dose, BMI, and blood pressure on metabolic control in young patients with type 1 diabetes. *Pediatric diabetes*. 2015;16(8):581-6.
31. Fredheim S, Delli A, Rida H, Drivvoll AK, Skriverhaug T, Bjarnason R, et al. Equal access to health care may diminish the differences in outcome between native and immigrant patients with type 1 diabetes. *Pediatric diabetes*. 2014;15(7):519-27.
32. Zuijdwijk CS, Cuerden M, Mahmud FH. Social determinants of health on glycemic control in pediatric type 1 diabetes. *J Pediatr*. 2013;162(4):730-5.
33. Carter PJ, Cutfield WS, Hofman PL, Gunn AJ, Wilson DA, Reed PW, et al. Ethnicity and social deprivation independently influence metabolic control in children with type 1 diabetes. *Diabetologia*. 2008;51(10):1835-42.
34. Bizzarri C, Paladini A, Benevento D, Fierabracci A, Cappa M. Family history and ethnicity influencing clinical presentation of type 1 diabetes in childhood. *Journal of endocrinological investigation*. 2015;38(10):1141-3.

35. Redondo MJ, Connor CG, Ruedy KJ, Beck RW, Kollman C, Wood JR, et al. Pediatric Diabetes Consortium Type 1 Diabetes New Onset (NeOn) Study: factors associated with HbA1c levels one year after diagnosis. *Pediatric diabetes*. 2014;15(4):294-302.
36. Swift PG, Skinner TC, de Beaufort CE, Cameron FJ, Aman J, Aanstoot HJ, et al. Target setting in intensive insulin management is associated with metabolic control: the Hvidoere childhood diabetes study group centre differences study 2005. *Pediatric diabetes*. 2010;11(4):271-8.
37. Raphael JL, Rueda A, Lion KC, Giordano TP. The role of lay health workers in pediatric chronic disease: a systematic review. *Academic pediatrics*. 2013;13(5):408-20.
38. Butalia S, Patel AB, Johnson JA, Ghali WA, Rabi DM. Association between diabetic ketoacidosis hospitalizations and driving distance to outpatient diabetes centres in adults with type 1 diabetes mellitus. *Canadian journal of diabetes*. 2014;38(6):451-5.
39. Fredheim S, Johannesen J, Johansen A, Lyngsoe L, Rida H, Andersen ML, et al. Diabetic ketoacidosis at the onset of type 1 diabetes is associated with future HbA1c levels. *Diabetologia*. 2013;56(5):995-1003.
40. Shalitin S, Fisher S, Yackbovitch-Gavan M, de Vries L, Lazar L, Lebenthal Y, et al. Ketoacidosis at onset of type 1 diabetes is a predictor of long-term glycemic control. *Pediatric diabetes*. 2018;19(2):320-8.
41. Duca LM, Reboussin BA, Pihoker C, Imperatore G, Saydah S, Mayer-Davis E, et al. Diabetic ketoacidosis at diagnosis of type 1 diabetes and glycemic control over time: The SEARCH for diabetes in youth study. *Pediatric diabetes*. 2019;20(2):172-9.

FIGURE PANEL 1: Estimated HbA1c trajectories based on fractional polynomials




Fig 1a: Estimated HbA1c trajectory of 253 patients followed from diagnosis




Fig 1b: Estimated HbA1c trajectory of 335 patients followed from baseline

FIGURE PANEL 2: Estimated HbA1c trajectories based on fractional polynomials by age at diagnosis or HbA1c at diagnosis




Fig 2a: Age at diagnosis




Fig 2b: HbA1c at diagnosis

Table 1: Baseline[#] characteristics of study population by HbA1c categories. Data are mean (±SD) or counts (%)

| Characteristic | Total | HbA1c levels at baseline (mmol/mol) | | | P value (for difference by HbA1c at baseline) |
|--|----------------------|-------------------------------------|-----------------------------|--------------------------------|--|
| | | <58mmol/mol | 58-80mmol/mol | >80mmol/mol | |
| Sex | N=356 | N=38 (10.67%) | N=117 (32.87%) | N=201 (56.46%) | 0.01* |
| Male | 169 (47.47%) | 25 (65.79%) | 63 (53.85%) | 81 (40.30%) | |
| Female | 187 (52.53%) | 13 (34.21%) | 54 (46.15%) | 120 (59.70%) | |
| Age at diagnosis (years) | N=356 8.9±4.3 | N=38 9.5±4.7 | N=117 8.07±4.5 | N=201 9.3±4.1 | 0.03* |
| Age at diagnosis | N=356 | N=38 | N=117 | N=201 | 0.14 [†] |
| 0-5yrs | 106 (29.78%) | 11 (28.95%) | 44 (37.61%) | 51 (25.37%) | |
| 6-11yrs | 143 (40.17%) | 13 (34.21%) | 40 (34.19%) | 90 (44.78%) | |
| 12-18yrs | 107 (30.06%) | 14 (36.84%) | 33 (28.21%) | 60 (29.85%) | |
| Period of T1D onset | N=356 | N=38 | N=117 | N=201 | 0.01* |
| Year 2005 to 2010 | 181 (50.84%) | 11 (28.95%) | 56 (47.86%) | 114 (56.72%) | |
| Year 2011 - 2015 | 175 (49.16%) | 27 (71.05%) | 61 (52.14%) | 87 (43.28%) | |
| Season of T1D onset | N=356 | N=38 | N=117 | N=201 | 0.02 [†] |
| Spring | 93 (26.12%) | 11 (28.95%) | 38 (32.48%) | 44 (21.89%) | |
| Summer | 75 (21.07%) | 7 (18.42%) | 13 (11.11%) | 55 (27.36%) | |
| Autumn | 82 (23.03%) | 7 (18.42%) | 32 (27.35%) | 43 (21.39%) | |
| Winter | 106 (29.78%) | 13 (34.21%) | 34 (29.06%) | 59 (29.35%) | |
| Ethnicity | N=351 | N=38 (10.82%) | N=116 (33.05%) | N=197 (56.13%) | 0.05 [†] |
| 1 White v/s | 125 (35.61%) | 14 (36.84%) | 51 (43.97%) | 60 (30.46%) | |
| 2 Non white | 226 (64.39%) | 24 (63.16%) | 65 (56.03%) | 137 (69.54%) | |
| Non-white sub groups | | | | | |
| a) Mixed | 35 (9.97%) | 2 (5.26%) | 13 (11.21%) | 20 (10.15%) | |
| b) Black | 79 (22.51%) | 9 (23.68%) | 27 (23.28%) | 43 (21.83%) | |
| c) Asian/other | 112 (31.91%) | 13 (34.21%) | 25 (21.55%) | 74 (37.56%) | |
| Indices of Multiple Deprivation | N=354 | N=38 (10.73%) | N=117 (33.05%) | N=199 (56.22%) | 0.39 [†] |
| Most deprived | 187 (52.82%) | 18 (47.37%) | 60 (51.28%) | 109 (54.77%) | |
| Moderate | 127 (35.88%) | 13 (34.21%) | 41 (35.04%) | 73 (36.68%) | |
| Least deprived | 40 (11.30%) | 7 (18.42%) | 16 (13.68%) | 17 (8.54%) | |
| No of patients at clinics | N=356 | N=38 | N=117 | N=201 | 0.01 [†] |
| 1 | 36 (10.11%) | 12 (31.58%) | 15 (12.82%) | 9 (4.48%) | |
| 2 | 112 (31.46%) | 9 (23.68%) | 36 (30.77%) | 67 (33.33%) | |
| 3 | 208 (58.43%) | 17 (44.74%) | 66 (56.41%) | 125 (62.19%) | |
| Insulin regimen | N=341 | N=37 (10.85%) | N=112 (32.84%) | N=192 (56.31%) | 0.60 [†] |
| 1-2 inj./day | 92 (26.98%) | 9 (24.32%) | 31 (27.68%) | 52 (27.08%) | |
| >=3 inj. per day | 233 (68.33%) | 25 (67.57%) | 74 (66.07%) | 134 (69.79%) | |
| Pump | 16 (4.69%) | 3 (8.11%) | 7 (6.25%) | 6 (3.13%) | |
| HbA1c at diagnosis (mmol/mol) | N=253 92.9 ±23.10 | N=13 (5.14%) 49.69±5.44 | N=62 (24.50%) 69.60±6.58 | N=178 (70.36%) 104.17±16.96 | 0.01* |
| HbA1c at baseline (mmol/mol) | N=356 | N=38 (10.67%) | N=117 (32.87%) | N=201 (56.46%) | 0.01* |

| | | | | | |
|---|-----------------------|-----------------------------|-----------------------------|-------------------------------|--------|
| | 85.79±23.75 | 49.89±6.03 | 68.98±6.29 | 102.36±16.89 | |
| HbA1c at 1 to 3 months of diagnosis (mmol/mol) | N=242 66.11 ±14.76 | N=29 (11.98%) 49.72±6.19 | N=91 (37.61%) 63.40±9.67 | N=122 (50.41%) 72.03±15.82 | 0.001* |
| HbA1c at 2 to 3 months of diagnosis (mmol/mol) | N=127 61.37 ±15.00 | N=19 (5.14%) 50.89±5.71 | N=44 (24.50%) 59.52±9.91 | N=64 (70.36%) 65.75±17.85 | 0.001* |
| pH at diagnosis | N=186 7.27±0.16 | N=9 (4.84%) 7.33±0.19 | N=40 (21.50%) 7.33±0.13 | N=137 (73.66%) 7.25±0.16 | 0.02* |
| pH at diagnosis | N=186 | N=9 | N=40 | N=137 | 0.01† |
| Normal 7.3-7.45 | 98 (52.69%) | 5 (55.56%) | 31 (77.50%) | 62 (45.26%) | |
| Acidotic <7.3 | 80 (43.01%) | 2 (22.22%) | 8 (20.00%) | 70 (51.09%) | |
| Alkalotic >7.45 | 8 (4.30%) | 2 (22.22%) | 1 (2.50%) | 5 (3.65%) | |
| BMI at diagnosis (kg/m²) | N=232 18.29±3.62 | N=11 (4.74%) 21.26±5.48 | N=56 (24.14%) 18.03±2.83 | N=165 (71.12%) 18.18±3.65 | 0.01* |
| BMI WHO z-score at baseline | N=330 | N=33 (10.00%) | N=109 (33.03%) | N=188 (56.97%) | 0.02† |
| Thin | 70 (21.21%) | 5 (7.14%) | 19 (27.14%) | 46 (65.71%) | |
| Normal weight | 103 (31.21%) | 6 (5.83%) | 30 (29.13%) | 67 (63.05%) | |
| Obese/overweight | 157 (47.58%) | 22 (14.01%) | 60 (38.22%) | 75 (47.77%) | |

#Baseline: Period within 3 months of T1D diagnosis; N: Number of patients; †: Global Wald test (χ^2); *: Wald test (χ^2); HbA1c: Glycated hemoglobin; pH: measure of acidity; BMI: Body mass index; WHO: World health organization


Table 2: Predictors of glycaemic control in children and adolescents from 3 to 24 months post-T1D diagnosis (MVA with predictor HbA1c at baseline)

| Predictor | Univariable analysis | | | Multivariable analysis (MVA) without interaction terms | | | MVA (with predictors showing significant interaction with sex, age at diagnosis or clinic) | | |
|---|----------------------|-------------------------|---------|--|------------------------|---------|--|-------------------------|---------|
| | N | Estimate (95%CI) | p-value | N | Estimate (95%CI) | p-value | N | Estimate (95%CI) | p-value |
| Sex (ref female) | 335 | 0.46 (-2.79 to 3.72) | 0.78* | | | | | | |
| Age at diagnosis (Ref 0-5yrs) | 335 | | 0.01† | 330 | | 0.02† | 330 | | 0.04† |
| 6-11yrs | | 0.96 (-2.92 to 4.84) | | | 0.59 (-3.11 to 4.28) | | | 0.44 (-3.23 to 4.11) | |
| 12-18yrs | | 5.84 (1.65 to 10.02) | | | 5.05 (1.07 to 9.03) | | | 4.49 (0.53 to 8.45) | |
| Year of Diagnosis (Ref pre 2011) | 335 | | 0.001* | 330 | | 0.001* | 330 | | 0.001* |
| 2011 – 2015 | | -7.47 (-10.63 to -4.31) | | | -6.09 (-9.17 to -3.01) | | | -9.43 (-13.63 to -5.23) | |
| Season of diagnosis (Ref Spring) | 335 | | 0.45† | | | | | | |
| Summer | | 1.71 (-3.08 to 6.51) | | | | | | | |
| Autumn | | 1.82 (-2.82 to 6.47) | | | | | | | |
| Winter | | 3.58 (-0.73 to 7.89) | | | | | | | |
| Ethnicity (Ref White) | 331 | | 0.002* | 330 | | 0.02* | 330 | | 0.01* |
| Non white | | 5.46 (2.07 to 8.85) | | | 3.89 (0.66 to 7.13) | | | 5.37 (1.28 to 9.45) | |
| IMD (Ref 1 most deprived) | 333 | | 0.59† | | | | | | |
| 2 (moderate) | | 0.89 (-2.65 to 4.42) | | | | | | | |
| 3 (least deprived) | | -2.06 (-7.47 to 3.34) | | | | | | | |
| Clinic (ref clinic 3) | 335 | | 0.001† | 330 | | 0.02† | 330 | | 0.01† |
| 1 | | -0.53 (-6.05 to 4.99) | | | 2.43 (-3.12 to 7.97) | | | 11.48 (2.42 to 20.55) | |


| | | | | | | | | | |
|---|-----|-------------------------|--------|-----|----------------------|--------|-----|--------------------------|--------|
| 2 | | 6.55 (3.08 to 10.02) | | | 4.66 (1.34 to 7.98) | | | 6.09 (0.12 to 12.06) | |
| Number of clinic visits¹ | 335 | 0.22 (0.03 to 0.41) | 0.02* | | | | | | |
| Insulin regimen(Ref >=3 inj. per day) | 333 | | 0.02† | | | | | | |
| 1-2 inj./day | | 0.62 (-2.13 to 3.36) | | | | | | | |
| Pump | | -7.64 (-13.18 to -2.10) | | | | | | | |
| HbA1c at diagnosis | 238 | 0.15 (0.07 to 0.23) | 0.001* | | | | | | |
| HbA1c at diagnosis (ref <58 mmol/mol) | 238 | | 0.02† | | | | | | |
| 58 to 80 | | 5.74 (-3.60 to 15.09) | | | | | | | |
| >80 mmol/mol | | 10.36 (1.59 to 19.14) | | | | | | | |
| HbA1c at baseline | 334 | 0.25 (0.17 to 0.33) | 0.001* | | | | | | |
| HbA1c at baseline (ref <58 mmol/mol) | 334 | | 0.001† | 330 | | 0.003† | 330 | | 0.001† |
| 58 to 80 | | 2.47 (-2.57 to 7.51) | | | 3.02 (-1.88 to 7.93) | | | 3.07 (-1.75 to 7.90) | |
| >80 mmol/mol | | 8.80 (3.84 to 13.76) | | | 7.55 (2.61 to 12.48) | | | 8.21 (3.30 to 13.12) | |
| HbA1c during 1 to 3 months after diagnosis | 230 | 0.37 (0.25 to 0.49) | 0.001* | | | | | | |
| HbA1c during 1 to 3 months after diagnosis (ref <58 mmol/mol) | 230 | | 0.001† | | | | | | |
| 58 to 80 | | 7.19 (3.01 to 11.36) | | | | | | | |
| >80 mmol/mol | | 14.26 (8.88 to 19.63) | | | | | | | |
| HbA1c at 2 to 3 months after diagnosis | 124 | 0.56 (0.40 to 0.71) | 0.001* | | | | | | |
| HbA1c at 2 to 3 months after diagnosis (ref <58 mmol/mol) | 124 | | 0.001† | | | | | | |
| 58 to 80 | | 12.87 (7.84 to 17.91) | | | | | | | |
| >80 mmol/mol | | 21.15 (13.05 to 29.26) | | | | | | | |
| pH at diagnosis (Ref normal pH 7.3-7.45) | 175 | | 0.71† | | | | | | |
| Acidotic <7.3 | | 1.80 (-2.84 to 6.44) | | | | | | | |
| Alkalotic >7.45 | | 2.65 (-8.43 to 13.74) | | | | | | | |
| BMI (WHO) z score at baseline (ref normal weight) | 309 | | 0.22† | | | | | | |
| Overweight/obese | | -2.27 (-6.26 to 1.73) | | | | | | | |
| Thin | | 1.41 (-3.38 to 6.20) | | | | | | | |
| Year of diagnosis # sex (Ref Females) | | | | | | | 330 | | 0.04† |
| Pre 2011 # Males | | | | | | | | -1.88 (-6.07 to 2.31) | |
| 2011 – 2015 # Males | | | | | | | | 5.38 (0.91 to 9.84) | |
| Ethnicity # clinic (Ref white # clinic 3) | | | | | | | 330 | | 0.05† |
| Non-white # Clinic 1 | | | | | | | | -13.84 (-24.94 to -2.75) | |
| Non-white # Clinic 2 | | | | | | | | -2.49 (-9.63 to 4.66) | |

†: Global Wald test (χ^2); *: Wald test (χ^2); Ref: reference group; IMD: Indices of multiple deprivation; ¹: clinic visits (range 2-13); HbA1c: Glycated hemoglobin; pH: measure of acidity; BMI: Body mass index; WHO: World health organization. All models include v and \log^*v functions of time as both fixed and random effects.

Supplementary figure panel 1a: Histograms showing patterns of visits to clinic during 0-3 months after T1D diagnosis



Supplementary figure panel 1b: Histograms showing patterns of visits to clinic after 3 months of T1D diagnosis



Supplementary figure 2: Predictor HbA1c tertiles at baseline showing significant differences between HbA1c trajectories over time




Fig 2: HbA1c tertiles within 3 months of diagnosis

Supplementary Table S1a: Fractional Polynomial three-dimension model fit comparison algorithm for period 0-24 months post T1D diagnosis

| Fractional Polynomial model | Index for comparison | | |
|-----------------------------------|----------------------|----------|----------|
| | -2LL | AIC | BIC |
| Linear model | 14398.42 | 14408.42 | 14435.43 |
| Model with one polynomial term | 13887.14 | 13897.14 | 13924.15 |
| Model with two polynomial terms | 13571.56 | 13585.56 | 13623.37 |
| Model with three polynomial terms | 13507.02 | 13525.02 | 13573.62 |

Supplementary Table S1b: Fractional Polynomial three-dimension model fit comparison algorithm for period 3-24 months post T1D diagnosis

| Fractional Polynomial model | Index for comparison | | |
|--------------------------------|----------------------|----------|----------|
| | -2LL | AIC | BIC |
| Linear model | 13221.92 | 13231.92 | 13258.89 |
| Model with one polynomial term | 13125.53 | 13135.53 | 13162.50 |

| | | | |
|-----------------------------------|----------|----------|----------|
| Model with two polynomial terms | 13106.62 | 13120.62 | 13158.38 |
| Model with three polynomial terms | 13100.65 | 13118.65 | 13167.20 |

Supplementary Table S2: Baseline[#] characteristics of study population by HbA1c tertiles. Data are mean (\pm SD) or counts (%)

| Characteristic | Total | HbA1c tertiles at baseline (mmol/mol) | | | P value (for difference by HbA1c tertiles at baseline) |
|--|---------------------------|---------------------------------------|-----------------------------------|--------------------------------------|--|
| | | Tertile 1 (<74mmol/mol) | Tertile 2 (74-95mmol/mol) | Tertile 3 (>95mmol/mol) | |
| Sex | N=356 | N=120 (33.71%) | N=120 (33.71%) | N=116 (32.58%) | 0.01* |
| Male | 169 (47.47%) | 66 (65.00%) | 61 (50.83%) | 42 (36.21%) | |
| Female | 187 (52.53%) | 54 (45.00%) | 59 (49.17%) | 74 (63.79%) | |
| Age at diagnosis (years) | N=356 8.9 \pm 4.3 | N=120 8.7 \pm 4.6 | N=120 8.2 \pm 4.5 | N=116 9.9 \pm 3.8 | 0.01* |
| Age at diagnosis | N=356 | N=120 | N=120 | N=116 | 0.003 [†] |
| 0-5yrs | 106 (29.78%) | 39 (32.50%) | 47 (39.17%) | 20 (17.24%) | |
| 6-11yrs | 143 (40.17%) | 42 (35.00%) | 42 (35.00%) | 59 (50.86%) | |
| 12-18yrs | 107 (30.06%) | 39 (32.50%) | 31 (25.83%) | 37 (31.90%) | |
| Period of T1D onset | N=356 | N=120 | N=120 | N=116 | 0.03* |
| Year 2005 to 2010 | 181 (50.84%) | 50 (41.67%) | 63 (52.50%) | 68 (58.62%) | |
| Year 2011 - 2015 | 175 (49.16%) | 70 (58.33%) | 57 (47.50%) | 48 (41.38%) | |
| Season of T1D onset | N=356 | N=120 | N=120 | N=116 | 0.03 [†] |
| Spring | 93 (26.12%) | 42 (35.00%) | 28 (23.33%) | 23 (19.83%) | |
| Summer | 75 (21.07%) | 14 (11.67%) | 30 (25.00%) | 31 (26.72%) | |
| Autumn | 82 (23.03%) | 26 (21.67%) | 29 (24.17%) | 27 (23.28%) | |
| Winter | 106 (29.78%) | 38 (31.67%) | 33 (27.50%) | 35 (30.17%) | |
| Ethnicity | N=351 | N=119 (33.90%) | N=119 (33.90%) | N=113 (32.19%) | 0.01 [†] |
| 1 White v/s | 125 (35.61%) | 53 (42.40%) | 43 (34.40%) | 29 (23.20%) | |
| 2 Non white | 226 (64.39%) | 66 (29.20%) | 76 (33.63%) | 84 (37.17%) | |
| Non-white sub groups | | | | | |
| a) Mixed | 35 (9.97%) | 10 (28.57%) | 17 (48.57%) | 8 (22.86%) | |
| b) Black | 79 (22.51%) | 29 (36.71%) | 21 (26.58%) | 29 (36.71%) | |
| c) Asian/other | 112 (31.91%) | 27 (24.11%) | 38 (33.93%) | 47 (41.96%) | |
| Indices of Multiple Deprivation | N=354 | N=120 (33.90%) | N=118 (33.33%) | N=116 (32.77%) | 0.01 [†] |
| Most deprived | 187 (52.82%) | 63 (52.50%) | 50 (42.37%) | 74 (63.79%) | |
| Moderate | 127 (35.88%) | 39 (32.50%) | 55 (46.61%) | 33 (28.45%) | |
| Least deprived | 40 (11.30%) | 18 (15.00%) | 13 (11.02%) | 9 (7.76%) | |
| No of patients at clinics | N=356 | N=120 | N=120 | N=116 | 0.01 [†] |
| 1 | 36 (10.11%) | 24 (20.00%) | 8 (6.67%) | 4 (3.45%) | |
| 2 | 112 (31.46%) | 38 (31.67%) | 36 (30.00%) | 38 (32.76%) | |
| 3 | 208 (58.43%) | 58 (48.33%) | 76 (63.33%) | 74 (63.79%) | |
| Insulin regimen | N=341 | N=116 (34.02%) | N=114 (33.43%) | N=111 (32.55%) | 0.07 [†] |
| 1-2 inj./day | 92 (26.98%) | 28 (24.14%) | 32 (28.07%) | 32 (28.83%) | |
| >=3 inj. per day | 233 (68.33%) | 79 (68.10%) | 75 (65.79%) | 79 (71.17%) | |
| Pump | 16 (4.69%) | 9 (7.76%) | 7 (6.14%) | 0 (0.00%) | |
| HbA1c at diagnosis (mmol/mol) | N=253 92.9 \pm 23.10 | N=53 (20.95%) 61.66 \pm 8.25 | N=89 (35.18%) 85.12 \pm 6.58 | N=111 (43.87%) 114.05 \pm 13.75 | 0.01* |
| HbA1c at baseline (mmol/mol) | N=356 | N=120 | N=120 | N=116 | 0.01* |

| | | | | | |
|--|----------------------|-----------------------------|------------------------------|------------------------------|--------|
| | 78.44±19.24 | 59.04±8.50 | 78.74±7.75 | 98.21±14.79 | |
| HbA1c at 2 to 3 months after diagnosis (mmol/mol) | N=127 61.37±15.00 | N=51 (40.16%) 55.08±8.36 | N=39 (30.71%) 65.05±14.57 | N=37 (29.13%) 66.16±19.29 | 0.001* |
| pH at diagnosis | N=186 7.27±0.16 | N=35 (18.82%) 7.32±0.14 | N=66 (35.48%) 7.27±0.16 | N=85 (45.70%) 7.25±0.17 | 0.07* |
| pH at diagnosis | N=186 | N=35 | N=66 | N=85 | 0.05† |
| Normal 7.3-7.45 | 98 (52.69%) | 26 (74.29%) | 31 (46.97%) | 41 (48.24%) | |
| Acidotic <7.3 | 80 (43.01%) | 7 (20.00%) | 32 (48.48%) | 41 (48.24%) | |
| Alkalotic >7.45 | 8 (4.30%) | 2 (5.71%) | 3 (4.55%) | 3 (3.53%) | |
| BMI at diagnosis (kg/m2) | N=232 18.29±3.62 | N=47 (20.26%) 18.80±3.78 | N=85 (36.64%) 17.74±3.14 | N=100 (43.10%) 18.52±3.89 | 0.18* |
| BMI WHO z-score at baseline | N=330 | N=109 (33.03%) | N=116 (35.15%) | N=105 (31.82%) | 0.10† |
| Thin | 70 (21.21%) | 18 (16.51%) | 29 (25.00%) | 23 (21.21%) | |
| Normal weight | 103 (31.21%) | 29 (26.61%) | 34 (29.31%) | 40 (38.10%) | |
| Obese/overweight | 157 (47.58%) | 62 (56.88%) | 53 (45.69%) | 42 (40.00%) | |

#Baseline: Period within 3 months of T1D diagnosis; N: Number of patients; †: Global Wald test (χ^2); *: Wald test (χ^2); HbA1c: Glycated hemoglobin; pH: measure of acidity; BMI: Body mass index; WHO: World health organization

Supplementary Table S3a: Predictors of glycaemic control in children and adolescents from 3 to 24 months post-T1D diagnosis (MVA with predictor HbA1c tertiles at baseline)

| Predictor | Univariable analysis | | | Multivariable analysis (MVA) without interaction terms | | | MVA (with predictors showing significant interaction with sex, age at diagnosis or clinic) | | |
|---|----------------------|-------------------------|---------|--|------------------------|---------|--|--------------------------|---------|
| | N | Estimate (95%CI) | p-value | N | Estimate (95%CI) | p-value | N | Estimate (95%CI) | p-value |
| Sex (Ref female) | 335 | 0.46 (-2.79 to 3.72) | 0.78* | | | | | | |
| Age at diagnosis (Ref 0-5yrs) | 335 | | 0.01† | 331 | | 0.01† | 331 | | 0.03† |
| 6-11yrs | | 0.96 (-2.92 to 4.84) | | | 1.15 (-2.60 to 4.90) | | | 1.05 (-2.68 to 4.78) | |
| 12-18yrs | | 5.84 (1.65 to 10.02) | | | 5.49 (1.50 to 9.49) | | | 5.10 (1.12 to 9.08) | |
| Year of Diagnosis (Ref pre 2011) | 335 | | 0.001* | 331 | | 0.001* | 331 | | 0.001* |
| 2011 – 2015 | | -7.47 (-10.63 to -4.31) | | | -6.36 (-9.47 to -3.26) | | | -9.16 (-13.40 to -4.92) | |
| Season of diagnosis (Ref Spring) | 335 | | 0.45† | | | | | | |
| Summer | | 1.71 (-3.08 to 6.51) | | | | | | | |
| Autumn | | 1.82 (-2.82 to 6.47) | | | | | | | |
| Winter | | 3.58 (-0.73 to 7.89) | | | | | | | |
| Ethnicity (Ref White) | 331 | | 0.001* | 331 | | 0.01* | 331 | | 0.004* |
| Non white | | 5.46 (2.07 to 8.85) | | | 4.12 (0.87 to 7.38) | | | 6.03 (1.90 to 10.15) | |
| IMD (Ref 1 most deprived) | 333 | | 0.59† | | | | | | |
| 2 (moderate) | | 0.89 (-2.65 to 4.42) | | | | | | | |
| 3 (least deprived) | | -2.06 (-7.47 to 3.34) | | | | | | | |
| Clinic (Ref clinic 3) | 335 | | 0.001† | 331 | | 0.01† | 331 | | 0.01† |
| 1 | | -0.53 (-6.05 to 4.99) | | | 1.96 (-3.59 to 7.51) | | | 11.08 (2.11 to 20.04) | |
| 2 | | 6.55 (3.08 to 10.02) | | | 5.14 (1.77 to 8.50) | | | 7.35 (1.24 to 13.45) | |
| Number of clinic visits ¹ | 335 | 0.22 (0.03 to 0.41) | 0.02* | | | | | | |
| Insulin regimen(Ref >=3 inj. per day) | 333 | | 0.02† | | | | | | |
| 1-2 inj./day | | 0.62 (-2.13 to 3.36) | | | | | | | |
| Pump | | -7.64 (-13.18 to -2.10) | | | | | | | |
| HbA1c tertiles at baseline (Tertile 1 mmol/mol) | 335 | | 0.01† | 331 | | 0.07† | 331 | | 0.03† |
| Tertile 2 | | 3.80 (-0.16 to 7.75) | | | 3.55 (-0.27 to 7.37) | | | 3.96 (0.17 to 7.75) | |
| Tertile 3 | | 5.79 (1.88 to 9.71) | | | 4.32 (0.38 to 8.26) | | | 5.02 (1.07 to 8.96) | |
| pH at diagnosis (Ref normal pH 7.3-7.45) | 175 | | 0.71† | | | | | | |
| Acidotic <7.3 | | 1.80 (-2.84 to 6.44) | | | | | | | |
| Alkalotic >7.45 | | 2.65 (-8.43 to 13.74) | | | | | | | |
| BMI (WHO) z score at baseline (Ref normal weight) | 309 | | 0.22† | | | | | | |
| Overweight/obese | | -2.27 (-6.26 to 1.73) | | | | | | | |
| Thin | | 1.41 (-3.38 to 6.20) | | | | | | | |
| Year of diagnosis # sex (Ref Females) | | | | | | | 331 | | 0.11† |
| Pre 2011 # Males | | | | | | | | -1.68 (-5.90 to 2.55) | |
| 2011 – 2015 # Males | | | | | | | | 4.45 (-0.04 to 8.93) | |
| Ethnicity # clinic (Ref white # clinic 3) | | | | | | | 331 | | 0.04† |
| Non-white # Clinic 1 | | | | | | | | -14.09 (-25.10 to -3.09) | |
| Non-white # Clinic 2 | | | | | | | | -3.49 (-10.75 to 3.78) | |

†: Global Wald test (χ^2); *: Wald test (χ^2); Ref: reference group; IMD: Indices of multiple deprivation; ¹: clinic visits (range 2-13); HbA1c: Glycated hemoglobin; pH: measure of acidity; BMI: Body mass index; WHO: World health organization. All models include v and log*v functions of time as both fixed and random effects

Supplementary Table S3b: Predictors of glycaemic control in children and adolescents from 3 to 24 months post-T1D diagnosis (MVA with predictor HbA1c at diagnosis)

| Predictor | Univariable analysis | | | Multivariable analysis (MVA) without interaction terms | | | MVA (with predictors showing significant interaction with sex, age at diagnosis or clinic) | | |
|--|----------------------|-------------------------|---------|--|-------------------------|---------|--|-------------------------|---------|
| | N | Estimate (95%CI) | p-value | N | Estimate (95%CI) | p-value | N | Estimate (95%CI) | p-value |
| Sex (ref female) | 238 | 0.82 (-3.09 to 4.73) | 0.68* | | | | | | |
| Age at diagnosis (Ref 0-5yrs) | 238 | | 0.05† | 234 | | 0.04† | 234 | | 0.003† |
| 6-11yrs | | 1.99 (-2.64 to 6.64) | | | 1.72 (-2.75 to 6.19) | | | 2.38 (-2.81 to 7.56) | |
| 12-18yrs | | 6.30 (1.16 to 11.45) | | | 6.05 (1.11 to 10.99) | | | 9.65 (3.84 to 15.47) | |
| Year of Diagnosis (Ref pre 2011) | 238 | | 0.001* | 234 | | 0.001* | 234 | | 0.001* |
| 2011 – 2015 | | -8.05 (-11.82 to -4.28) | | | -6.84 (-10.58 to -3.09) | | | -7.25 (-10.95 to -3.55) | |
| Season of diagnosis (Ref Spring) | 238 | | 0.42† | | | | | | |
| Summer | | 2.41 (-3.19 to 8.01) | | | | | | | |
| Autumn | | 4.37 (-1.17 to 9.91) | | | | | | | |
| Winter | | 3.51 (-1.69 to 8.70) | | | | | | | |
| Ethnicity (Ref White) | 234 | | 0.005* | 234 | | 0.01* | 234 | | 0.01* |
| Non white | | 5.96 (1.85 to 10.07) | | | 5.24 (1.34 to 9.14) | | | 5.46 (1.63 to 9.28) | |
| IMD (Ref 1 most deprived) | 236 | | 0.75† | | | | | | |
| 2 (moderate) | | -1.13 (-5.35 to 3.09) | | | | | | | |
| 3 (least deprived) | | -2.29 (-9.24 to 4.65) | | | | | | | |
| Clinic (ref clinic 3) | 238 | | 0.008† | 234 | | 0.17† | 234 | | 0.03† |
| 1 | | 1.83 (-7.64 to 11.30) | | | 2.70 (-7.01 to 12.40) | | | -9.11 (-30.94 to 12.72) | |
| 2 | | 6.75 (2.52 to 10.97) | | | 3.93 (-0.19 to 8.05) | | | 9.96 (2.05 to 17.87) | |
| Insulin regimen(Ref >=3 inj. per day) | 236 | | 0.08† | | | | | | |
| 1-2 inj./day | | -0.14 (-3.33 to 3.04) | | | | | | | |
| Pump | | -7.20 (-13.53 to -0.86) | | | | | | | |
| HbA1c at diagnosis | 238 | 0.15 (0.07 to 0.23) | 0.001* | | | | | | |
| HbA1c at diagnosis (ref <58 mmol/mol) | 238 | | 0.02† | 234 | | 0.05† | 234 | | 0.15† |
| 58 to 80 | | 5.74 (-3.60 to 15.09) | | | 5.69 (-3.35 to 14.74) | | | 4.13 (-4.81 to 13.06) | |
| >80 mmol/mol | | 10.36 (1.59 to 19.14) | | | 9.05 (0.50 to 17.61) | | | 7.06 (-1.44 to 15.56) | |
| pH at diagnosis (Ref normal pH 7.3-7.45) | 175 | | 0.71† | | | | | | |
| Acidotic <7.3 | | 1.80 (-2.84 to 6.44) | | | | | | | |
| Alkalotic >7.45 | | 2.65 (-8.43 to 13.74) | | | | | | | |
| BMI (WHO) z score at diagnosis (Ref normal weight) | 217 | | 0.68† | | | | | | |
| Overweight/obese | | -1.59 (-6.38 to 3.19) | | | | | | | |
| Thin | | 0.54 (-5.29 to 6.37) | | | | | | | |
| Age at diagnosis # clinic (Ref 0-5yrs # clinic 3) | | | | | | | 234 | | 0.03† |
| 6-11yrs # Clinic 1 | | | | | | | | 29.87 (0.71 to 59.03) | |
| 6-11yrs # Clinic 2 | | | | | | | | -5.33 (-15.04 to 4.38) | |
| 12-18yrs # Clinic 1 | | | | | | | | -6.01 (-18.78 to 30.80) | |

| | | | | | |
|------|--|-------------------------|------------------------|-------------------------|--|
| Thin | | -2.26 (-46.01 to 41.48) | 52.17 (17.51 to 86.83) | -18.29 (-36.92 to 0.35) | |
|------|--|-------------------------|------------------------|-------------------------|--|

t=T1D duration; *Global Wald test (χ^2); ref: reference group; WHO: World Health Organization. All models include V and log*V functions of time as both fixed and random effects