

## Manuscript Details

<b>Manuscript number</b>	JAD_2019_2434_R1
<b>Title</b>	Effectiveness of the Thinking Healthy Programme for perinatal depression delivered through peers: pooled analysis of two randomized controlled trials in India and Pakistan
<b>Article type</b>	Research Paper

### Abstract

**Background:** The Thinking Healthy Programme (THP) is recommended to treat perinatal depression in resource-limited settings, but scale-up is hampered by a paucity of community health workers. THP was adapted for peer-delivery (THPP) and evaluated in two randomized controlled trials in India and Pakistan. Our aim was to estimate the effectiveness of THPP on maternal outcomes across these two settings, and evaluate effect-modification by country and other pre-defined covariates. **Methods:** Participants were pregnant women aged  $\geq 18$  years with depression (Patient Health Questionnaire (PHQ-9) score  $\geq 10$ ), randomized to THPP plus enhanced usual care (EUC) or EUC-only. Primary outcomes were symptom severity and remission (PHQ-9 score  $< 5$ ) 6 months post-childbirth. Secondary outcomes included further measures of depression, disability and social support at 3 and 6 months post-childbirth. **Results:** Among 850 women (280 India; 570 Pakistan), 704 (83%) attended 6-month follow-up. Participants in the intervention arm had lower symptom severity (PHQ-9 score adjusted mean difference  $-0.78$  (95% confidence interval  $-1.47, -0.09$ )) and higher odds of remission (adjusted odds ratio  $1.35$  (1.02, 1.78)) versus EUC-only. There was a greater intervention effect on remission among women with short chronicity of depression, and those primiparous. There were beneficial intervention effects across multiple secondary outcomes. **Limitations:** The trials were not powered to assess effect-modifications. 10-20% of participants were missing outcome data. **Conclusions:** This pooled analysis demonstrates the effectiveness, acceptability and feasibility of THPP, which can be scaled-up within a stepped-care approach by engaging with the existing health care systems and the communities to address the treatment gap for perinatal depression in resource-limited settings.

<b>Keywords</b>	Pregnancy; community health workers; depression; patient health questionnaire; Pakistan; India
<b>Corresponding Author</b>	Fiona Vanobberghen
<b>Order of Authors</b>	Fiona Vanobberghen, helen weiss, Daniela Fuhr, Siham Sikander, Ejma Afonso, Ikhlaq Malik, Najia Atif, Amina Bibi, Tayyaba Bibi, Samina Bilal, Aveena De Sa, Ethel D'souza, Akankasha Joshi, Priya Korgaonkar, Revathi Krishna, Anisha Lazarus, Rakshanda Liaqat, Maria Sharif, Benedict Weobong, Ahmed Zaidi, Shaffaq Zuliqar, Vikram Patel, Atif Rahman
<b>Suggested reviewers</b>	Alan Stein, Jane Fisher, Judy Bass, Prabha Chandra

## Submission Files Included in this PDF

### File Name [File Type]

- SHARE pooled cover letter JAD revision FINAL.pdf [Cover Letter]
- SHARE pooled paper JAD responses FINAL.docx [Response to Reviewers]
- SHARE pooled paper JAD highlights FINAL.docx [Highlights]
- SHARE pooled paper JAD abstract FINAL.docx [Abstract]
- SHARE pooled paper JAD title page.docx [Title Page (with Author Details)]
- SHARE pooled paper JAD revision FINAL.docx [Manuscript File]
- Fig1A.tif [Figure]
- Fig1B.tif [Figure]
- SHARE pooled paper JAD conflict of interest.docx [Conflict of Interest]
- SHARE pooled paper JAD author statement.docx [Author Statement]
- SHARE pooled supplement JAD FINAL.docx [Supplementary Material]

To view all the submission files, including those not included in the PDF, click on the manuscript title on your EVISE Homepage, then click 'Download zip file'.

14 November 2019

Dear Dr Brambilla

We thank you and the reviewer for the consideration of our manuscript entitled “**Effectiveness of the Thinking Healthy Programme for perinatal depression delivered through peers: pooled analysis of two randomized controlled trials in India and Pakistan**”, and for your helpful feedback.

We are pleased to submit our revised manuscript with tracked changes, along with a point-by-point response to the feedback.

We thank you for your continued consideration of our manuscript and we look forward to hearing from you.

Yours sincerely



Fiona Vanobberghen



**Effectiveness of the Thinking Healthy Programme for perinatal depression delivered through peers:  
pooled analysis of two randomized controlled trials in India and Pakistan**

**POINT BY POINT RESPONSES TO REVIEWERS**

**Comments from the editors and reviewers:**

**-Reviewer 1**

- Since THP/ THPP is the main intervention in this study, the authors should describe more (in details) about this type of intervention in the Introduction or summarize a protocol in a Table.

**Response: We have added further details of THPP to the methods (pages 2-3 of the revised manuscript) including citing the complete description in the published trial protocol paper and main trial papers (line 53) and in the intervention development paper (line 63). Manuals are available on request, as indicated (lines 63-64). We took this opportunity to restructure the first part of the methods section to improve readability, separating out the design/setting, interventions and outcomes.**

- In the Discussion, the authors should compare THPP to other standard psychological intervention in perinatal depression (e.g. CBT, IPT, etc) and may propose the mechanism by which THPP can help in decreasing chronicity and increasing remission of depression.

**Response: We have added discussion of other psychological interventions to the discussion (page 11, lines 274-279). The mechanism of THPP was assessed through a pooled mediation analysis of the two trials, which found that the key mechanism was through increase in behavioral activation and social support (Singla et al., 2019). We have added this reference to the manuscript (line 276) where we mention behavioural activation.**

***Singla, D.R., MacKinnon, D.P., Fuhr, D.C., Sikander, S., Rahman, A., Patel, V., 2019. Multiple mediation analysis of the peer-delivered Thinking Healthy Programme for perinatal depression: findings from two parallel, randomised controlled trials. Br J Psychiatry 1-8. <https://doi.org/10.1192/bjp.2019.184>***

**Effectiveness of the Thinking Healthy Programme for perinatal depression delivered through peers:  
pooled analysis of two randomized controlled trials in India and Pakistan**

**HIGHLIGHTS**

- In this pooled analysis of two randomized controlled trials, we found consistent benefits of the Thinking Healthy Programme adapted for delivery by Peers (THPP) on maternal outcomes
- THPP resulted in 35% higher odds of remission at 6 months post-childbirth compared to enhanced usual care only
- There was evidence of beneficial intervention effects across a number of secondary outcomes, and in repeated measures analyses, demonstrating feasibility and acceptability of this peer-delivered intervention, and external validity of our results

## **Effectiveness of the Thinking Healthy Programme for perinatal depression delivered through peers: pooled analysis of two randomized controlled trials in India and Pakistan**

### **ABSTRACT**

**Background:** The Thinking Healthy Programme (THP) is recommended to treat perinatal depression in resource-limited settings, but scale-up is hampered by a paucity of community health workers. THP was adapted for peer-delivery (THPP) and evaluated in two randomized controlled trials in India and Pakistan. Our aim was to estimate the effectiveness of THPP on maternal outcomes across these two settings, and evaluate effect-modification by country and other pre-defined covariates.

**Methods:** Participants were pregnant women aged  $\geq 18$  years with depression (Patient Health Questionnaire (PHQ-9) score  $\geq 10$ ), randomized to THPP plus enhanced usual care (EUC) or EUC-only. Primary outcomes were symptom severity and remission (PHQ-9 score  $< 5$ ) 6 months post-childbirth. Secondary outcomes included further measures of depression, disability and social support at 3 and 6 months post-childbirth.

**Results:** Among 850 women (280 India; 570 Pakistan), 704 (83%) attended 6-month follow-up. Participants in the intervention arm had lower symptom severity (PHQ-9 score adjusted mean difference  $-0.78$  (95% confidence interval  $-1.47, -0.09$ )) and higher odds of remission (adjusted odds ratio  $1.35$  (1.02, 1.78)) versus EUC-only. There was a greater intervention effect on remission among women with short chronicity of depression, and those primiparous. There were beneficial intervention effects across multiple secondary outcomes.

**Limitations:** The trials were not powered to assess effect-modifications. 10-20% of participants were missing outcome data.

**Conclusions:** This pooled analysis demonstrates the effectiveness, acceptability and feasibility of THPP, which can be scaled-up within a stepped-care approach by engaging with the existing health care systems and the communities to address the treatment gap for perinatal depression in resource-limited settings.

**Key words:** Pregnancy; community health workers; depression; patient health questionnaire;

Pakistan; India

**Trial registrations:** ClinicalTrials.gov: NCT02104232, NCT02111915

**Effectiveness of the Thinking Healthy Programme for perinatal depression delivered through peers:  
pooled analysis of two randomized controlled trials in India and Pakistan**

Fiona Vanobberghen, PhD‡ [1,2,3], Helen A Weiss, PhD [1], Daniela C Fuhr, DrPH [4], Siham Sikander, PhD [5,6], Ejma Afonso, MA [7], Ikhlaq Ahmad, MPhil [5], Najia Atif, PhD [5], Amina Bibi, MPhil [5], Tayyaba Bibi, M.Com [5], Samina Bilal, MSc [5], Aveena De Sa, MA, MSc [7], Ethel D'Souza, M.Com [7], Akankasha Joshi, MA [7], Priya Korgaonkar, MA [7], Revathi Krishna, MSc [7], Anisha Lazarus, MPhil [7], Rakshanda Liaqat, MSc [5], Maria Sharif, MPhil [5], Benedict Weobong, PhD [7,8], Ahmed Zaidi, MPhil [5], Shaffaq Zuliqar, MPhil [5], Vikram Patel, PhD\* [7,9], Atif Rahman, PhD\* [10]

\*Contributed equally as joint mentor authors.

[1] MRC Tropical Epidemiology Group, Faculty of Epidemiology and Population Health, Department of Infectious Disease Epidemiology, London School of Hygiene and Tropical Medicine, London, UK

[2] Swiss Tropical and Public Health Institute, Basel, Switzerland

[3] University of Basel, Basel, Switzerland

[4] Faculty of Public Health and Policy, Department of Health Services Research and Policy, London School of Hygiene and Tropical Medicine, UK

[5] Human Development Research Foundation, Islamabad, Pakistan

[6] Health Services Academy, Islamabad, Pakistan

[7] Sangath Centre, Socorro Village, Bardez-Goa, Goa, India

[8] Faculty of Epidemiology and Population Health, Department of Population Health, London School of Hygiene and Tropical Medicine, UK

[9] Department of Global Health and Social Medicine, Harvard Medical School, Boston, USA

[10] Institute of Psychology, Health and Society, University of Liverpool, Liverpool, UK

**Corresponding author:** Fiona Vanobberghen, Swiss Tropical and Public Health Institute, Socinstrasse 57, Basel, Switzerland; +41 61 284 87 16; [fiona.vanobberghen@swisstph.ch](mailto:fiona.vanobberghen@swisstph.ch)

1 **Effectiveness of the Thinking Healthy Programme for perinatal depression delivered through peers:**  
2 **pooled analysis of two randomized controlled trials in India and Pakistan**

3

4 **INTRODUCTION**

5 Depression is the main cause of disability globally, and its prevalence is increasing (Friedrich, 2017).  
6 More than one in ten women experience a major depressive episode in the vulnerable perinatal  
7 period (pregnancy or the year following delivery) in high income countries (Fisher et al., 2012;  
8 Hendrick et al., 1998), with prevalences in low- and middle-income countries (LMIC) of up to 20%  
9 (Fisher et al., 2012; Woody et al., 2017). Perinatal depression is associated with adverse maternal  
10 outcomes including suicidal ideation (Gelaye et al., 2016), pregnancy and birth complications, and  
11 poorer mental, motor and emotional development of the infant (Field et al., 2006).

12

13 Psychological therapies like cognitive behavior therapy or behavioral activation are recommended  
14 for the treatment of mild and moderate perinatal depression (Guille et al., 2013). Due to the paucity  
15 of mental health professionals in LMIC (Kakuma et al., 2011), some of these interventions are  
16 modified, and provided as low-intensity psychological interventions delivered by trained para-  
17 professionals to depressed mothers in need (Rahman et al., 2013). One such low-intensity  
18 psychological intervention is the Thinking Healthy Programme (THP), developed by Rahman and  
19 colleagues (Rahman, 2007) and recommended by the World Health Organization (WHO) for the  
20 treatment of perinatal depression in LMIC (World Health Organization, 2015). THP is delivered by  
21 community healthcare workers (CHW) and has shown to be highly effective, halving the risk of  
22 perinatal depression and significantly improving infant health outcomes in a study in Pakistan  
23 (Rahman et al., 2008). However, the public health impact of THP has been hampered by the high  
24 occupational load of CHWs (Haq et al., 2008), therefore impairing the scale up of this programme  
25 (Jaskiewicz and Tulenko, 2012). To address this challenge, the National Institute of Mental Health  
26 (NIMH), through the Collaborative Hubs for International Research on Mental Health (CHIRMH)  
27 initiative, commissioned two studies in India and Pakistan to adapt the Thinking Healthy Programme

28 for delivery by trained peers, namely local women with no prior experience of healthcare delivery  
29 (Atif et al., 2017; D. Singla et al., 2014). Randomized controlled trials were conducted in two distinct  
30 settings, in urban/peri-urban Goa, India and rural Rawalpindi, Pakistan (Sikander et al., 2015), to  
31 evaluate the effectiveness and cost-effectiveness of the Thinking Healthy Programme Peer-delivered  
32 (THPP) compared to enhanced usual care (EUC). There were some differences in the trial designs  
33 (outlined below), but both trials found moderate effects of THPP on depression symptom severity  
34 and remission at 3 months post-childbirth (Fuhr et al., 2019; Sikander et al., 2019).

35

36 Our aim was to estimate the overall effectiveness of THPP on maternal outcomes, and evaluate  
37 effect modifications by country and other pre-defined covariates. The increased power from the  
38 pooled analysis enables us to look for consistency of results across the two settings, to assess the  
39 external validity of the intervention effect, and hence the ability to generalize findings regarding  
40 acceptability and feasibility of this peer-delivered intervention to other settings.

41

## 42 **METHODS**

### 43 **Trial settings, designs, and participants**

44 Data were pooled from the Goa, India trial (hereafter, THPP-India) and the Rawalpindi, Pakistan trial  
45 (hereafter, THPP-Pakistan). The trial settings are diverse, with higher rates of poverty (60% versus  
46 4%), lower proportions of women educated (65% versus 84%), higher fertility rates (3.8 versus 1.8)  
47 and larger households (6.2 versus 4.2 persons per household) in rural Rawalpindi compared to urban  
48 Goa (D. Singla et al., 2014). The trial designs were similar (Table 1): both trials enrolled pregnant  
49 women aged  $\geq 18$  years with moderate to severe depression defined by scoring  $\geq 10$  on the nine-item  
50 Patient Health Questionnaire (PHQ-9). THPP-India was run through healthcare facilities with women  
51 individually-randomized, while THPP-Pakistan was conducted in a community setting with woman  
52 randomized in village clusters to avoid contamination. The trials have been described in full  
53 elsewhere (Fuhr et al., 2019; Sikander et al., 2019, 2015).

54

55 **Trial interventions**

56 ~~Participants were randomized to, and compared~~ THPP plus EUC ~~to or~~ EUC-only. ~~The~~  
57 ~~interventions~~ THPP consisted of 6-14 sessions over the prenatal period to six months post-childbirth,  
58 covering behavioral activation, active listening, collaboration with the family, guided discovery and  
59 homework ~~(Atif et al., 2017)~~. In India, THPP was delivered as 6-14 individual sessions, and in Pakistan,  
60 THPP was delivered as ten individual and four group sessions. Peers received classroom and field  
61 training, and received regular supervisions by the trainers. Emphasis was on behavior activation  
62 strategies to enable the delivery by peers with no previous experience of delivering healthcare.  
63 Further details of THPP have been published previously (Atif et al., 2017) and manuals are available  
64 from the authors on request. ~~THPP India was run through healthcare facilities with women~~  
65 ~~individually randomized, while THPP Pakistan was conducted in a community setting with woman~~  
66 ~~randomized in village clusters to avoid contamination.~~

67  
68 **Trial outcomes**

69 In both trials, the primary outcomes were symptom severity (PHQ-9 score) and remission (PHQ-9  
70 score <5) at 6 months post-childbirth. ~~The trials have been described in full elsewhere (Fuhr et al.,~~  
71 ~~2019; Sikander et al., 2019, 2015)~~. In this analysis, we included the primary and secondary maternal  
72 outcomes.

73  
74 **Statistical methods**

75 Baseline characteristics of the two trial populations were described, and a multivariable logistic  
76 regression model was used to assess factors independently associated with country. Process  
77 indicators were summarized by country, including peer characteristics and attendance to therapy  
78 sessions among women in the intervention arm.

79  
80 Outcome data were analyzed using logistic and linear regression models to estimate odds ratios and  
81 mean differences, respectively (with 95% confidence intervals). Generalized estimating equations

82 (GEEs) with an exchangeable correlation structure were used to account for the village-level  
83 clustering in THPP-Pakistan (Sikander et al., 2019), with individuals in THPP-India acting as their own  
84 clusters. Models were adjusted for all variables included in the main trials' analyses, because they  
85 were: pre-specified in the analysis plan, stratification factors in the randomization, unbalanced  
86 between groups at baseline, or associated with missing outcome data at 6 months. We also included  
87 country. The following baseline variables were adjusted for as fixed effects: country, recruitment site,  
88 residence (rural/urban), union council, symptom severity, treatment expectations, education,  
89 chronicity of depression, and time between screening and birth. Participants with missing values  
90 were omitted from the models.

91

92 For the primary outcomes, the following variables were assessed as a priori effect modifiers, by  
93 fitting interactions between the intervention and the covariate: country, age, chronicity of  
94 depression, baseline symptom severity, treatment expectations, and parity. We also assessed effect  
95 modification by country for the secondary outcomes. Repeated measures analyses were performed,  
96 combining the 3 and 6 months results, with assessment of group by time interactions. All models  
97 were adjusted for the covariates listed above.

98

99 Sensitivity analyses were performed for the primary outcomes, using GEEs with individuals grouped  
100 as one cluster, and mixed effects models. No adjustments were made for multiple testing; results  
101 were interpreted based on the strength of evidence of effect size and consistency of results across  
102 outcomes. Analyses were conducted in Stata (StataCorp, 2015), following intention-to-treat  
103 principles. Data are available on request (Fiona Vanobberghen et al., 2018; F Vanobberghen et al.,  
104 2018).

105

## 106 **Ethical considerations**

107 Ethical approval for the THPP-India trial was obtained from the Institutional Review Boards (IRBs) at  
108 the London School of Hygiene and Tropical Medicine (LSHTM), Sangath (the trial-implementing

109 institution in India), and the Indian Council of Medical Research. Ethical approval for the THPP-  
110 Pakistan trial was obtained from the IRBs at the University of Liverpool, LSHTM, and the Human  
111 Development Research Foundation (the trial-implementing institution in Pakistan). Participants in  
112 both trials provided written informed consent (or witnessed informed consent/audio-recordings for  
113 illiterate participants).

114

## 115 **RESULTS**

116 Overall, 280 women were enrolled in THPP-India and 570 in THPP-Pakistan (Table 2). After adjusting  
117 for confounders (Appendix Table A1), THPP-Pakistan participants were less likely than those in THPP-  
118 India to work (6% versus 15%, with these low rates reflecting the national populations where the  
119 majority of women are housewives) (WHO Country Office Pakistan, 2013; World Bank, 2013), had  
120 received more education, had higher expectations of the usefulness of counseling, had higher  
121 baseline symptom severity, had lower social support scores, were more likely to have had a previous  
122 miscarriage or stillbirth, and were less likely to report domestic violence. Data on chronicity of  
123 depression at baseline were missing for 30% of THPP-Pakistan and no THPP-India participants. There  
124 were some differences between the women who did and did not have chronicity data in THPP-  
125 Pakistan: those with missing data were somewhat less likely to work, had higher treatment  
126 expectations, had lower symptom severity, were more likely to be multiparous, were more likely to  
127 have had a previous non-live birth, and were less likely to report domestic violence. Data were  
128 missing on the time between screening and birth of the child for 15% of THPP-Pakistan and 4% of  
129 THPP-India participants, due to women being lost to follow-up after screening. Depression was more  
130 chronic among THPP-Pakistan than THPP-India participants and time between screening and the  
131 birth of the child was shorter.

132

133 Overall, 26 and 66 peers were trained and delivered at least one session in THPP-India and THPP-  
134 Pakistan, respectively, with corresponding mean ages of 38 and 30 years and mean years of  
135 education completed 12 years in both trials (Table 3). No peers were lost during the THPP-India trial,

136 whereas 23 (35%) were lost during the THPP-Pakistan trial. Attendance to supervisions was 67% and  
137 88% in THPP-India and THPP-Pakistan, respectively. Quality was assessed differently in the two trials.  
138 In THPP-India, 18 items were assessed on the Therapy Quality scale of 0-2 (higher better) (D. R. Singla  
139 et al., 2014); the overall score across 72 sessions rated by independent raters was 1.49 (standard  
140 deviation 0.33). In THPP-Pakistan, assessments were done by independent raters for each peer in  
141 three sessions at each of three time-points using an 18-item competency checklist; the overall  
142 average was 84%. These results indicate average to good therapy quality.

143

144 The mean number of sessions attended by participants in the intervention arm was slightly higher in  
145 THPP-Pakistan compared to THPP-India (10.9 versus 9.8); this was driven by higher attendance in the  
146 postnatal period (Table 3). In THPP-India, treatment completion was defined as attending at least six  
147 sessions with at least one session in each of four phases (prenatal, and 1-2, 3-4 and 5-6 months  
148 postnatal); 99/138 (72%) women completed treatment. In THPP-Pakistan, treatment completion was  
149 defined as attending at least ten sessions; 201/258 (78%) women completed treatment.

150

151 Average symptom severity decreased considerably over time in both groups (Appendix Figure A1).  
152 There was evidence of a beneficial effect of the intervention on both primary outcomes at 6 months,  
153 with participants in the intervention group having lower symptom severity (adjusted mean difference  
154 -0.78, 95% confidence interval -1.47,-0.09) and higher odds of remission (adjusted odds ratio 1.35,  
155 95% confidence interval 1.02,1.78) compared with those in the control group (Table 4, Figures 1A  
156 and 1B). There was a trend towards stronger intervention effects in THPP-India compared to THPP-  
157 Pakistan for remission, but the difference was not statistically significant ( $p=0.18$ , with  $p=0.77$  for  
158 symptom severity). Results were robust to sensitivity analyses (Appendix Table A2).

159

160 After adjusting for confounders including country, there was evidence of a greater intervention effect  
161 on remission at 6 months among women with shorter chronicity of depression at baseline, and those  
162 primiparous ( $p=0.03$  for both interactions, Figure 1B and Appendix Table A3). Similar trends were

163 observed for symptom severity with chronicity and parity, although the p values were large (p=0.49  
164 and 0.30, respectively; Figure 1A and Appendix Table A4). For both outcomes, there were non-  
165 significant trends towards greater intervention effects among younger women (p=0.14 and 0.27 for  
166 remission and symptom severity, respectively). There was no consistent evidence of effect-  
167 modification by baseline symptom severity or treatment expectations.

168  
169 After adjusting for confounders including intervention group, the following factors were associated  
170 with higher symptom severity and lower odds of remission at six months: country, higher baseline  
171 symptom severity, and lower level of education (Appendix Table A5).

172  
173 There was consistently strong evidence of beneficial intervention effects across a number of  
174 secondary outcomes (Table 4). For symptom severity and WHO-DAS score, there were slightly  
175 stronger intervention effects at 3 months than at 6 months. For remission and MSPSS score, results  
176 were broadly similar at 3 and 6 months. For the composite outcomes of recovery and response at  
177 both 3 and 6 months, there was consistent and strong evidence of a benefit of the intervention.  
178 There was no evidence of an intervention effect on number of days unable to work in the last month.  
179 There was no evidence of effect modification by country for any of the secondary outcomes.

180  
181 There was no statistical evidence of group by time interactions, and assuming a constant intervention  
182 effect at 3 and 6 months (Appendix Table A6), there were strong evidence of intervention effects on  
183 symptom severity, remission, WHO-DAS score and MSPSS score (3 and 6 months combined), in line  
184 with the results of the individual trials (Fuhr et al., 2019; Sikander et al., 2019). There was no  
185 evidence of an intervention effect on number of days unable to work in the last month.

186

## 187 **DISCUSSION**

188 We present results of a pooled analysis of two of the largest trials evaluating a psychological  
189 intervention for perinatal depression in LMIC delivered by peers. With the increased power from this

190 pooled analysis, we found small (Rahman et al., 2013) but important benefits of THPP across a range  
191 of maternal depression, disability and social support outcomes, with 35% higher odds of remission at  
192 6 months post-childbirth compared to EUC-only. These findings open the possibilities of peer-  
193 delivered THPP to be implemented in other settings where there are limited numbers of mental  
194 health professionals to tackle the treatment gap for perinatal depression.

195  
196 Historically, intervention studies of depression in LMIC have generally had small sample sizes, with  
197 different recruitment mechanisms of study participants and using varying methodological designs  
198 and outcome measures (Chowdhary et al., 2014; Rahman et al., 2013). Our studies represent a new  
199 generation of complementary trials of psychological therapies nested in the community using  
200 innovative delivery mechanisms based on lay care providers and peer supervisors. THPP-India was  
201 conducted in a peri-urban setting in Goa while THPP-Pakistan was embedded in rural communities of  
202 Rawalpindi. These contextual and socio-economic differences were reflected in the trial populations  
203 and delivery agents (peers), with the THPP-Pakistan participants having more severe and more  
204 chronic depression at baseline compared to those in the THPP-India trial. THPP-Pakistan participants  
205 reported on average more years of education than those in THPP-India, contrary to expected (D.  
206 Singla et al., 2014) and compared to lower nationwide literacy rates of 45-50% (Population Census  
207 Organization, 1998; World Bank, 2013). There may have been over-reporting of education among the  
208 participants in Pakistan due to social desirability. Participants in THPP-Pakistan also had poorer  
209 depression outcomes at 6 months compared to those in THPP-India, after adjustment for  
210 confounders including baseline severity and chronicity. Slightly different models of incentivization  
211 were used across the two trials: peers in urban Goa received financial incentives since this was  
212 identified as an important motivator, while peers in rural Rawalpindi were volunteers, devoting their  
213 time to THPP for altruistic motives (Fuhr et al., 2019; Sikander et al., 2019; D. Singla et al., 2014). This  
214 may have been one reason for a greater turnover of peers during the trial in Pakistan (35% compared  
215 to none in India), though attendance to supervision was excellent (88%, compared to 67% in India).  
216 Furthermore, most peers in Pakistan who left were able to identify their own replacements,

217 indicating that there was a pool of women interested in this work. The health systems in which peers  
218 were trained to operate THPP were also distinct. Peers in Rawalpindi worked alongside government  
219 lady health workers delivering care in the community, while peers in Goa worked more  
220 independently and within a tiered public healthcare system (D. Singla et al., 2014). There were some  
221 differences in the design of the intervention between the two trials (Sikander et al., 2015), although  
222 the mean number of sessions attended by intervention recipients was similar. Despite these  
223 contextual differences, the impact of THPP by country was very similar, demonstrating the external  
224 validity of the intervention. In both countries, peers were trained to competently deliver THPP, with  
225 high proportions of women in the intervention arms completing their treatment. This indicates that  
226 delivery by peers was not seen as stigmatizing, and demonstrates a high degree of acceptability of a  
227 psychological therapy delivered by peers in community settings to mothers with depression in both  
228 contexts. Further, the intervention was shown to offer an appreciable improvement in health at low  
229 cost in THPP-Pakistan (incremental cost-effectiveness ratio US\$ 15.50 over the whole trial period)  
230 and even cost saving in THPP-India (-US\$ 93.53) (Fuhr et al., 2019; Sikander et al., 2019), despite  
231 monetary incentivisation for the peers contributing to 12% of the cost in the latter trial (D. Singla et  
232 al., 2014).

233

234 While the benefits of THPP were consistent for remission across three and six months post-childbirth,  
235 we observed stronger intervention effects at three compared to six months for symptom severity  
236 (PHQ-9 score) and disability (WHO-DAS). Similar effects were found in the individual trials (Fuhr et  
237 al., 2019; Sikander et al., 2019). This may be explained by the intervention design which puts  
238 emphasis on the first three months after child-birth by front-loading sessions antenatally and soon  
239 after childbirth, and/or by the natural remission rates of untreated depression over time (Austin et  
240 al., 2008; Posternak and Miller, 2001; Rojas et al., 2007; Tandon et al., 2018; Whiteford et al., 2013).  
241 The benefits at three months remain important to reduce the duration of the depression episode at  
242 such a critical time-point in childcare. Considering effect modification of the intervention, we  
243 observed greater benefits of THPP among women with shorter chronicity of depression at baseline

244 and those primiparous. THPP may be best adopted as a first-step psychological intervention to be  
245 used in a stepped care system for maternal depression in LMIC, most suitable for lower-risk groups.  
246 The need for a stepped collaborative care model is highlighted by the fact that nearly half of women  
247 in the intervention arm across both countries did not achieve remission by 6 months. Future research  
248 should explore if peers are able to sign-post higher risk women, and refer them to primary or  
249 specialist care where they may receive additional interventions. However, we did not observe a  
250 difference in the benefits of THPP by baseline severity, in contrast to previous studies of THP (Patel et  
251 al., 2017).

252  
253 There are some limitations of this study. The control group received enhanced usual care in a well-  
254 resourced setting, thus the intervention effects might be greater in a health system where no such  
255 care is provided (Fuhr et al., 2019). The individual trials were not powered for assessing effect  
256 modifications; while the pooling of the data from the two trials provides more power we cannot rule  
257 out having missed some effect modifications, including those by country. Some baseline covariates  
258 were missing, most notably chronicity of depression in 30% of THPP-Pakistan participants. Outcome  
259 data were missing for approximately 10% of THPP-India participants, and 20% of THPP-Pakistan  
260 participants, although we adjusted for factors associated with missingness which yielded similar  
261 results to multiple imputation analyses in the individuals trials (Fuhr et al., 2019; Sikander et al.,  
262 2019).

263  
264 THPP-India and THPP-Pakistan were pragmatic trials with high internal validity delivered in  
265 community settings to mothers recruited from routine health-care settings. By aligning THPP with  
266 routine clinical practice and working alongside CHWs and clinicians, we were able to evaluate real-  
267 life treatment effects in two diverse South Asian contexts. This pooled analysis demonstrates the  
268 effectiveness, acceptability and feasibility of the THP intervention delivered by peers. This is  
269 important because although the theoretical model was the same in the two settings, there were  
270 contextual differences in implementation. The fact that the intervention effects were similar in the

271 two trials demonstrates the external validity of the intervention, and hence potential for  
272 generalizability to other settings and populations.  
273  
274 THP uses cognitive behaviour therapy techniques and was designed to be delivered by lady health  
275 workers (Atif et al., 2017). THPP represents a further adaptation, with focus on behaviour activation  
276 strategies to enable successful delivery by peers (Atif et al., 2017; Singla et al., 2019). Other  
277 interventions for perinatal depression include educational sessions led by healthcare professionals,  
278 or interpersonal therapy with problem-solving methods through group activities (Rahman et al.,  
279 2013) but to our knowledge none have been adapted for delivery by non-healthcare professionals.  
280 Although the benefits of THPP were relatively small (Rahman et al., 2013), it has proven to be cost-  
281 effective (Fuhr et al., 2019; Sikander et al., 2019). These results therefore open avenues for  
282 collaborative care, where peers can be the “missing” community agent or case manager to increase  
283 awareness and provide first-line interventions, and work alongside specialists in primary, secondary  
284 and tertiary care. Further, the delivery through peers provides opportunities for women in the  
285 community to develop skills and improve their chances of employability. To make this happen, there  
286 is a need in global mental health to advocate for accreditation of evidence-based and cost-effective  
287 low-intensity psychological interventions like THPP. We conclude that THPP is suitable for  
288 implementation as part of a stepped-care approach by engaging with the existing health-care  
289 systems and the communities to treat perinatal depression in other contexts.

290

## 291 REFERENCES

- 292 Atif, N., Krishna, R.N., Sikander, S., Lazarus, A., Nisar, A., Ahmad, I., Raman, R., Fuhr, D.C., Patel, V.,  
293 Rahman, A., 2017. Mother-to-mother therapy in India and Pakistan: adaptation and  
294 feasibility evaluation of the peer-delivered Thinking Healthy Programme. BMC Psychiatry 17,  
295 79. <https://doi.org/10.1186/s12888-017-1244-z>  
296 Austin, M.-P., Frilingos, M., Lumley, J., Hadzi-Pavlovic, D., Roncolato, W., Acland, S., Saint, K., Segal,  
297 N., Parker, G., 2008. Brief antenatal cognitive behaviour therapy group intervention for the  
298 prevention of postnatal depression and anxiety: a randomised controlled trial. J. Affect.  
299 Disord. 105, 35-44. <https://doi.org/10.1016/j.jad.2007.04.001>  
300 Chowdhary, N., Sikander, S., Atif, N., Singh, N., Ahmad, I., Fuhr, D.C., Rahman, A., Patel, V., 2014. The  
301 content and delivery of psychological interventions for perinatal depression by non-specialist

302 [health workers in low and middle income countries: A systematic review. Best Pract. Res.](#)  
303 [Clin. Obstet. Gynaecol. 28, 113–133. https://doi.org/10.1016/j.bpobgyn.2013.08.013](#)  
304 [Field, T., Diego, M., Hernandez-Reif, M., 2006. Prenatal depression effects on the fetus and newborn:](#)  
305 [a review. Infant Behav. Dev. 29, 445–455. https://doi.org/10.1016/j.infbeh.2006.03.003](#)  
306 [Fisher, J., Cabral de Mello, M., Patel, V., Rahman, A., Tran, T., Holton, S., Holmes, W., 2012.](#)  
307 [Prevalence and determinants of common perinatal mental disorders in women in low- and](#)  
308 [lower-middle-income countries: a systematic review. Bull. World Health Organ. 90, 139G-](#)  
309 [149G. https://doi.org/10.2471/BLT.11.091850](#)  
310 [Friedrich, M.J., 2017. Depression Is the Leading Cause of Disability Around the World. JAMA 317,](#)  
311 [1517. https://doi.org/10.1001/jama.2017.3826](#)  
312 [Fuhr, D.C., Weobong, B., Lazarus, A., Vanobberghen, F., Weiss, H.A., Singla, D.R., Tabana, H., Afonso,](#)  
313 [E., De Sa, A., D'Souza, E., Joshi, A., Korgaonkar, P., Krishna, R., Price, L.N., Rahman, A., Patel,](#)  
314 [V., 2019. Delivering the Thinking Healthy Programme for perinatal depression through peers:](#)  
315 [an individually randomised controlled trial in India. Lancet Psychiatry 6, 115–127.](#)  
316 [https://doi.org/10.1016/S2215-0366\(18\)30466-8](#)  
317 [Gelaye, B., Kajeepeta, S., Williams, M.A., 2016. Suicidal ideation in pregnancy: an epidemiologic](#)  
318 [review. Arch. Womens Ment. Health 19, 741–751. https://doi.org/10.1007/s00737-016-](#)  
319 [0646-0](#)  
320 [Guille, C., Newman, R., Fryml, L.D., Lifton, C.K., Epperson, C.N., 2013. Management of Postpartum](#)  
321 [Depression. J. Midwifery Womens Health 58, 643–653. https://doi.org/10.1111/jmwh.12104](#)  
322 [Haq, Z., Iqbal, Z., Rahman, A., 2008. Job stress among community health workers: a multi-method](#)  
323 [study from Pakistan. Int. J. Ment. Health Syst. 2, 15. https://doi.org/10.1186/1752-4458-2-15](#)  
324 [Hendrick, V., Altshuler, L., Cohen, L., Stowe, Z., 1998. Evaluation of mental health and depression](#)  
325 [during pregnancy: position paper. Psychopharmacol. Bull. 34, 297–299.](#)  
326 [Jaskiewicz, W., Tulenko, K., 2012. Increasing community health worker productivity and](#)  
327 [effectiveness: a review of the influence of the work environment. Hum. Resour. Health 10,](#)  
328 [38. https://doi.org/10.1186/1478-4491-10-38](#)  
329 [Kakuma, R., Minas, H., van Ginneken, N., Dal Poz, M.R., Desiraju, K., Morris, J.E., Saxena, S., Scheffler,](#)  
330 [R.M., 2011. Human resources for mental health care: current situation and strategies for](#)  
331 [action. Lancet Lond. Engl. 378, 1654–1663. https://doi.org/10.1016/S0140-6736\(11\)61093-3](#)  
332 [Patel, V., Weobong, B., Weiss, H.A., Anand, A., Bhat, B., Katti, B., Dimidjian, S., Araya, R., Hollon, S.D.,](#)  
333 [King, M., Vijayakumar, L., Park, A.-L., McDaid, D., Wilson, T., Velleman, R., Kirkwood, B.R.,](#)  
334 [Fairburn, C.G., 2017. The Healthy Activity Program \(HAP\), a lay counsellor-delivered brief](#)  
335 [psychological treatment for severe depression, in primary care in India: a randomised](#)  
336 [controlled trial. The Lancet 389, 176–185. https://doi.org/10.1016/S0140-6736\(16\)31589-6](#)  
337 [Population Census Organization, 1998. District census report of Rawalpindi. Islamabad, Pakistan.](#)  
338 [Posternak, M.A., Miller, I., 2001. Untreated short-term course of major depression: a meta-analysis](#)  
339 [of outcomes from studies using wait-list control groups. J. Affect. Disord. 66, 139–146.](#)  
340 [Rahman, A., 2007. Challenges and opportunities in developing a psychological intervention for](#)  
341 [perinatal depression in rural Pakistan – a multi-method study. Arch. Womens Ment. Health](#)  
342 [10, 211–219. https://doi.org/10.1007/s00737-007-0193-9](#)  
343 [Rahman, A., Fisher, J., Bower, P., Luchters, S., Tran, T., Yasamy, M.T., Saxena, S., Waheed, W., 2013.](#)  
344 [Interventions for common perinatal mental disorders in women in low- and middle-income](#)  
345 [countries: a systematic review and meta-analysis. Bull. World Health Organ. 91, 593-601l.](#)  
346 [https://doi.org/10.2471/BLT.12.109819](#)  
347 [Rahman, A., Malik, A., Sikander, S., Roberts, C., Creed, F., 2008. Cognitive behaviour therapy-based](#)  
348 [intervention by community health workers for mothers with depression and their infants in](#)  
349 [rural Pakistan: a cluster-randomised controlled trial. Lancet 372, 902–909.](#)  
350 [https://doi.org/10.1016/S0140-6736\(08\)61400-2](#)  
351 [Rojas, G., Fritsch, R., Solis, J., Jadresic, E., Castillo, C., González, M., Guajardo, V., Lewis, G., Peters,](#)  
352 [T.J., Araya, R., 2007. Treatment of postnatal depression in low-income mothers in primary-](#)  
353 [care clinics in Santiago, Chile: a randomised controlled trial. Lancet Lond. Engl. 370, 1629–](#)  
354 [1637. https://doi.org/10.1016/S0140-6736\(07\)61685-7](#)

355 [Sikander, S., Ahmad, I., Atif, N., Zaidi, A., Vanobberghen, F., Weiss, H.A., Nisar, A., Tabana, H., Ain,](#)  
356 [Q.U., Bibi, A., Bilal, S., Bibi, T., Liaqat, R., Sharif, M., Zulfiqar, S., Fuhr, D.C., Price, L.N., Patel,](#)  
357 [V., Rahman, A., 2019. Delivering the Thinking Healthy Programme for perinatal depression](#)  
358 [through volunteer peers: a cluster randomised controlled trial in Pakistan. \*Lancet Psychiatry\*](#)  
359 [6, 128–139. \[https://doi.org/10.1016/S2215-0366\\(18\\)30467-X\]\(https://doi.org/10.1016/S2215-0366\(18\)30467-X\)](#)

360 [Sikander, S., Lazarus, A., Bangash, O., Fuhr, D.C., Weobong, B., Krishna, R.N., Ahmad, I., Weiss, H.A.,](#)  
361 [Price, L., Rahman, A., Patel, V., 2015. The effectiveness and cost-effectiveness of the peer-](#)  
362 [delivered Thinking Healthy Programme for perinatal depression in Pakistan and India: the](#)  
363 [SHARE study protocol for randomised controlled trials. \*Trials\* 16, 534.](#)  
364 [https://doi.org/10.1186/s13063-015-1063-9](#)

365 [Singla, D., Lazarus, A., Atif, N., Sikander, S., Bhatia, U., Ahmad, I., Nisar, A., Khan, S., Fuhr, D., Patel, V.,](#)  
366 [Rahman, A., 2014. “Someone like us”: delivering maternal mental health through peers in](#)  
367 [two South Asian contexts. \*J. Affect. Disord.\* 168, 452–458.](#)  
368 [https://doi.org/10.1016/j.jad.2014.07.017](#)

369 [Singla, D.R., MacKinnon, D.P., Fuhr, D.C., Sikander, S., Rahman, A., Patel, V., 2019. Multiple mediation](#)  
370 [analysis of the peer-delivered Thinking Healthy Programme for perinatal depression: findings](#)  
371 [from two parallel, randomised controlled trials. \*Br. J. Psychiatry J. Ment. Sci.\* 1–8.](#)  
372 [https://doi.org/10.1192/bjp.2019.184](#)

373 [Singla, D.R., Weobong, B., Nadkarni, A., Chowdhary, N., Shinde, S., Anand, A., Fairburn, C.G.,](#)  
374 [Dimijdan, S., Velleman, R., Weiss, H., Patel, V., 2014. Improving the scalability of](#)  
375 [psychological treatments in developing countries: an evaluation of peer-led therapy quality](#)  
376 [assessment in Goa, India. \*Behav. Res. Ther.\* 60, 53–59.](#)  
377 [https://doi.org/10.1016/j.brat.2014.06.006](#)

378 [StataCorp, 2015. Stata Statistical Software Release 14. StataCorp LP, College Station, TX.](#)

379 [Tandon, S.D., Ward, E.A., Hamil, J.L., Jimenez, C., Carter, M., 2018. Perinatal depression prevention](#)  
380 [through home visitation: a cluster randomized trial of mothers and babies 1-on-1. \*J. Behav.\*](#)  
381 [Med. <https://doi.org/10.1007/s10865-018-9934-7>](#)

382 [Vanobberghen, F., Weiss, H., D'Souza, E., Patel, V., Rahman, A., Fuhr, D., Weobong, B., Lazarus, A.,](#)  
383 [2018. Data for Thinking Healthy Programme India trial \(THPP-India\). \*Lond. Sch. Hyg. Trop.\*](#)  
384 [Med. Lond. U. K. <https://doi.org/10.17037/DATA.00000793>](#)

385 [Vanobberghen, Fiona, Weiss, H., Zaidi, A., Patel, V., Rahman, A., Sikander, S., Fuhr, D., Ahmad, I.,](#)  
386 [2018. Data for Thinking Healthy Programme Pakistan trial \(THPP-Pakistan\). \*Lond. Sch. Hyg.\*](#)  
387 [Trop. Med. Lond. U. K. <https://doi.org/10.17037/DATA.00000794>](#)

388 [Whiteford, H.A., Harris, M.G., McKeon, G., Baxter, A., Pennell, C., Barendregt, J.J., Wang, J., 2013.](#)  
389 [Estimating remission from untreated major depression: a systematic review and meta-](#)  
390 [analysis. \*Psychol. Med.\* 43, 1569–1585. <https://doi.org/10.1017/S0033291712001717>](#)

391 [WHO Country Office Pakistan, 2013. National integrated feedback report. WHO Country Office](#)  
392 [Pakistan.](#)

393 [Woody, C.A., Ferrari, A.J., Siskind, D.J., Whiteford, H.A., Harris, M.G., 2017. A systematic review and](#)  
394 [meta-regression of the prevalence and incidence of perinatal depression. \*J. Affect. Disord.\*](#)  
395 [219, 86–92. <https://doi.org/10.1016/j.jad.2017.05.003>](#)

396 [World Bank, 2013. World development indicators. World Bank, Washington, DC.](#)

397 [World Health Organization, 2015. Thinking Healthy. A manual for psychological management of](#)  
398 [perinatal depression.](#)

399 [Atif, N., Krishna, R.N., Sikander, S., Lazarus, A., Nisar, A., Ahmad, I., Raman, R., Fuhr, D.C., Patel, V.,](#)  
400 [Rahman, A., 2017. Mother-to-mother therapy in India and Pakistan: adaptation and](#)  
401 [feasibility evaluation of the peer-delivered Thinking Healthy Programme. \*BMC Psychiatry\* 17,](#)  
402 [79. <https://doi.org/10.1186/s12888-017-1244-z>](#)

403 [Austin, M.-P., Frilingos, M., Lumley, J., Hadzi-Pavlovic, D., Roncolato, W., Acland, S., Saint, K., Segal,](#)  
404 [N., Parker, G., 2008. Brief antenatal cognitive behaviour therapy group intervention for the](#)  
405 [prevention of postnatal depression and anxiety: a randomised controlled trial. \*J. Affect.\*](#)  
406 [Disord. 105, 35–44. <https://doi.org/10.1016/j.jad.2007.04.001>](#)

407 Chowdhary, N., Sikander, S., Atif, N., Singh, N., Ahmad, I., Fuhr, D.C., Rahman, A., Patel, V., 2014. The  
408 content and delivery of psychological interventions for perinatal depression by non-specialist  
409 health workers in low and middle income countries: A systematic review. *Best Pract. Res.  
410 Clin. Obstet. Gynaecol.* 28, 113–133. <https://doi.org/10.1016/j.bpobgyn.2013.08.013>

411 Field, T., Diego, M., Hernandez-Reif, M., 2006. Prenatal depression effects on the fetus and newborn:  
412 a review. *Infant Behav. Dev.* 29, 445–455. <https://doi.org/10.1016/j.infbeh.2006.03.003>

413 Fisher, J., Cabral de Mello, M., Patel, V., Rahman, A., Tran, T., Holton, S., Holmes, W., 2012.  
414 Prevalence and determinants of common perinatal mental disorders in women in low- and  
415 lower-middle-income countries: a systematic review. *Bull. World Health Organ.* 90, 139G-  
416 149G. <https://doi.org/10.2471/BLT.11.091850>

417 Friedrich, M.J., 2017. Depression Is the Leading Cause of Disability Around the World. *JAMA* 317,  
418 1517. <https://doi.org/10.1001/jama.2017.3826>

419 Fuhr, D.C., Weobong, B., Lazarus, A., Vanobberghen, F., Weiss, H.A., Singla, D.R., Tabana, H., Afonso,  
420 E., De Sa, A., D'Souza, E., Joshi, A., Korgaonkar, P., Krishna, R., Price, L.N., Rahman, A., Patel,  
421 V., 2019. Delivering the Thinking Healthy Programme for perinatal depression through peers:  
422 an individually randomised controlled trial in India. *Lancet Psychiatry* 6, 115–127.  
423 [https://doi.org/10.1016/S2215-0366\(18\)30466-8](https://doi.org/10.1016/S2215-0366(18)30466-8)

424 Gelaye, B., Kajeepeta, S., Williams, M.A., 2016. Suicidal ideation in pregnancy: an epidemiologic  
425 review. *Arch. Womens Ment. Health* 19, 741–751. <https://doi.org/10.1007/s00737-016-0646-0>

426

427 Guille, C., Newman, R., Fryml, L.D., Lifton, C.K., Epperson, C.N., 2013. Management of Postpartum  
428 Depression. *J. Midwifery Womens Health* 58, 643–653. <https://doi.org/10.1111/jmwh.12104>

429 Haq, Z., Iqbal, Z., Rahman, A., 2008. Job stress among community health workers: a multi-method  
430 study from Pakistan. *Int. J. Ment. Health Syst.* 2, 15. <https://doi.org/10.1186/1752-4458-2-15>

431 Hendrick, V., Altshuler, L., Cohen, L., Stowe, Z., 1998. Evaluation of mental health and depression  
432 during pregnancy: position paper. *Psychopharmacol. Bull.* 34, 297–299.

433 Jaskiewicz, W., Tulenko, K., 2012. Increasing community health worker productivity and  
434 effectiveness: a review of the influence of the work environment. *Hum. Resour. Health* 10,  
435 38. <https://doi.org/10.1186/1478-4491-10-38>

436 Kakuma, R., Minas, H., van Ginneken, N., Dal Poz, M.R., Desiraju, K., Morris, J.E., Saxena, S., Scheffler,  
437 R.M., 2011. Human resources for mental health care: current situation and strategies for  
438 action. *Lancet Lond. Engl.* 378, 1654–1663. [https://doi.org/10.1016/S0140-6736\(11\)61093-3](https://doi.org/10.1016/S0140-6736(11)61093-3)

439 Patel, V., Weobong, B., Weiss, H.A., Anand, A., Bhat, B., Katti, B., Dimidjian, S., Araya, R., Hollon, S.D.,  
440 King, M., Vijayakumar, L., Park, A.-L., McDaid, D., Wilson, T., Velleman, R., Kirkwood, B.R.,  
441 Fairburn, C.G., 2017. The Healthy Activity Program (HAP), a lay counsellor-delivered brief  
442 psychological treatment for severe depression, in primary care in India: a randomised  
443 controlled trial. *The Lancet* 389, 176–185. [https://doi.org/10.1016/S0140-6736\(16\)31589-6](https://doi.org/10.1016/S0140-6736(16)31589-6)

444 Population Census Organization, 1998. District census report of Rawalpindi. Islamabad, Pakistan.

445 Posternak, M.A., Miller, I., 2001. Untreated short-term course of major depression: a meta-analysis  
446 of outcomes from studies using wait-list control groups. *J. Affect. Disord.* 66, 139–146.

447 Rahman, A., 2007. Challenges and opportunities in developing a psychological intervention for  
448 perinatal depression in rural Pakistan – a multi-method study. *Arch. Womens Ment. Health*  
449 10, 211–219. <https://doi.org/10.1007/s00737-007-0193-9>

450 Rahman, A., Fisher, J., Bower, P., Luchters, S., Tran, T., Yasamy, M.T., Saxena, S., Waheed, W., 2013.  
451 Interventions for common perinatal mental disorders in women in low- and middle-income  
452 countries: a systematic review and meta-analysis. *Bull. World Health Organ.* 91, 593–601.  
453 <https://doi.org/10.2471/BLT.12.109819>

454 Rahman, A., Malik, A., Sikander, S., Roberts, C., Creed, F., 2008. Cognitive behaviour therapy-based  
455 intervention by community health workers for mothers with depression and their infants in  
456 rural Pakistan: a cluster-randomised controlled trial. *Lancet* 372, 902–909.  
457 [https://doi.org/10.1016/S0140-6736\(08\)61400-2](https://doi.org/10.1016/S0140-6736(08)61400-2)

458 Rojas, G., Fritsch, R., Solis, J., Jadresic, E., Castillo, C., González, M., Guajardo, V., Lewis, G., Peters,  
459 T.J., Araya, R., 2007. Treatment of postnatal depression in low-income mothers in primary-

460 care clinics in Santiago, Chile: a randomised controlled trial. *Lancet Lond. Engl.* 370, 1629–  
461 1637. [https://doi.org/10.1016/S0140-6736\(07\)61685-7](https://doi.org/10.1016/S0140-6736(07)61685-7)

462 Sikander, S., Ahmad, I., Atif, N., Zaidi, A., Vanobberghen, F., Weiss, H.A., Nisar, A., Tabana, H., Ain,  
463 Q.U., Bibi, A., Bilal, S., Bibi, T., Liaqat, R., Sharif, M., Zulfiqar, S., Fuhr, D.C., Price, L.N., Patel,  
464 V., Rahman, A., 2019. Delivering the Thinking Healthy Programme for perinatal depression  
465 through volunteer peers: a cluster randomised controlled trial in Pakistan. *Lancet Psychiatry*  
466 6, 128–139. [https://doi.org/10.1016/S2215-0366\(18\)30467-X](https://doi.org/10.1016/S2215-0366(18)30467-X)

467 Sikander, S., Lazarus, A., Bangash, O., Fuhr, D.C., Weobong, B., Krishna, R.N., Ahmad, I., Weiss, H.A.,  
468 Price, L., Rahman, A., Patel, V., 2015. The effectiveness and cost-effectiveness of the peer-  
469 delivered Thinking Healthy Programme for perinatal depression in Pakistan and India: the  
470 SHARE study protocol for randomised controlled trials. *Trials* 16, 534.  
471 <https://doi.org/10.1186/s13063-015-1063-9>

472 Singla, D., Lazarus, A., Atif, N., Sikander, S., Bhatia, U., Ahmad, I., Nisar, A., Khan, S., Fuhr, D., Patel, V.,  
473 Rahman, A., 2014. “Someone like us”: delivering maternal mental health through peers in  
474 two South Asian contexts. *J. Affect. Disord.* 168, 452–458.  
475 <https://doi.org/10.1016/j.jad.2014.07.017>

476 Singla, D.R., Weobong, B., Nadkarni, A., Chowdhary, N., Shinde, S., Anand, A., Fairburn, C.G.,  
477 Dimijdan, S., Velleman, R., Weiss, H., Patel, V., 2014. Improving the scalability of  
478 psychological treatments in developing countries: an evaluation of peer-led therapy quality  
479 assessment in Goa, India. *Behav. Res. Ther.* 60, 53–59.  
480 <https://doi.org/10.1016/j.brat.2014.06.006>

481 StataCorp, 2015. Stata Statistical Software Release 14. StataCorp LP, College Station, TX.

482 Tandon, S.D., Ward, E.A., Hamil, J.L., Jimenez, C., Carter, M., 2018. Perinatal depression prevention  
483 through home visitation: a cluster randomized trial of mothers and babies 1-on-1. *J. Behav.*  
484 *Med.* <https://doi.org/10.1007/s10865-018-9934-7>

485 Vanobberghen, F., Weiss, H., D’Souza, E., Patel, V., Rahman, A., Fuhr, D., Weobong, B., Lazarus, A.,  
486 2018. Data for Thinking Healthy Programme India trial (THPP-India). *Lond. Sch. Hyg. Trop.*  
487 *Med. Lond. U. K.* <https://doi.org/10.17037/DATA.00000793>

488 Vanobberghen, Fiona, Weiss, H., Zaidi, A., Patel, V., Rahman, A., Sikander, S., Fuhr, D., Ahmad, I.,  
489 2018. Data for Thinking Healthy Programme Pakistan trial (THPP-Pakistan). *Lond. Sch. Hyg.*  
490 *Trop. Med. Lond. U. K.* <https://doi.org/10.17037/DATA.00000794>

491 Whiteford, H.A., Harris, M.G., McKeon, G., Baxter, A., Pennell, C., Barendregt, J.J., Wang, J., 2013.  
492 Estimating remission from untreated major depression: a systematic review and meta-  
493 analysis. *Psychol. Med.* 43, 1569–1585. <https://doi.org/10.1017/S0033291712001717>

494 WHO Country Office Pakistan, 2013. National integrated feedback report. WHO Country Office  
495 Pakistan.

496 Woody, C.A., Ferrari, A.J., Siskind, D.J., Whiteford, H.A., Harris, M.G., 2017. A systematic review and  
497 meta-regression of the prevalence and incidence of perinatal depression. *J. Affect. Disord.*  
498 219, 86–92. <https://doi.org/10.1016/j.jad.2017.05.003>

499 World Bank, 2013. World development indicators. World Bank, Washington, DC.

500 World Health Organization, 2015. Thinking Healthy. A manual for psychological management of  
501 perinatal depression.

503

504 **FIGURE LEGENDS**

505

506 **Figure 1. A) Remission (PHQ-9 score <5) and B) symptom severity (PHQ-9 score) at 6 months by**  
507 **potential effect modifiers.**

508 PHQ=Patient Health Questionnaire. Results are from linear or logistic generalized estimating equation models, adjusted for  
509 country, recruitment site, residence, union council, baseline symptom severity, treatment expectations, education,  
510 chronicity of depression, and time between screening and birth. The sizes of the squares indicating the point estimates are  
511 proportional to the number of participants in each category. These results are described further in Table 4 and Appendix  
512 Tables A3 and A4.

513

514

515 TABLES

516

517 Table 1. Summary of the trial designs.

	THPP-India	THPP-Pakistan
<b>Location</b>	Goa	Rawalpindi
<b>Study setting</b>	Antenatal clinics in two hospitals and three primary health centers	Community setting in rural sub-district
<b>Participants</b>	Pregnant women aged $\geq 18$ years with moderate to severe depression as defined by PHQ-9 score $\geq 10$	
<b>Intervention</b>	THPP delivered as 6-14 individual sessions, each lasting 30-45 minutes, in four phases (prenatal, up to two months post childbirth, 3-4 months after childbirth, and 5-6 months after childbirth), plus enhanced usual care	THPP delivered as ten individual and four group sessions, each lasting 30-45 minutes, from the third trimester of pregnancy to six months after childbirth, plus enhanced usual care
<b>Control</b>	Enhanced usual care: Standard care from the gynecologist plus (1) patients and gynecologists were informed of participants screening results; (2) gynecologists were given the adapted mental health Gap Action Programme (mhGAP) treatment guidelines for perinatal depression; (3) participants were provided with an information sheet on health-care during pregnancy and beyond	Enhanced usual care: Standard care from LHWs plus (1) all participants and LHWs were informed of participants screening results; (2) doctors and midwives at the primary healthcare centers were given the adapted mental health Gap Action Programme (mhGAP) treatment guidelines for perinatal depression; (3) participants were provided with an information sheet on healthcare during pregnancy and beyond
<b>Peers</b>	Women with children, a similar socio-demographic background as participants, and good communication skills. Recruited from the local community through word-of-mouth, particularly through key informants in women's self-help groups and CHWs. Financial incentives were paid based on sessions delivered, up to a maximum of Rs. 5900 (approximately US\$80) per trial participant, plus a monthly honorarium of Rs. 100 (approximately US\$1) to cover phone call costs	Women with children, a similar socio-demographic background as participants, and good communication skills. Identified through LHWs and elders, with recruitment and placement through primary health-care centers. Peers received a travel allowance for supervision meetings and a phone card/top-up to be able to stay in touch with their respective supervisors in case of support needed. No remunerations were paid for delivering sessions
<b>Randomization</b>	Individually-randomized (1:1 allocation ratio), stratified by place of residence (rural/urban)	Cluster-randomized (1:1 allocation ratio; 40 village clusters), stratified by 11 union councils
<b>Blinding</b>	Outcome assessors were blinded to participants' treatment allocation	
<b>Primary outcomes</b>	Symptom severity (PHQ-9 score) and remission (PHQ-9 $< 5$ ) at 6 months post-childbirth	
<b>Secondary outcomes<sup>a</sup></b>	Symptom severity (PHQ-9 score) at 3 months post-childbirth	
	Remission (PHQ-9 score $< 5$ ) at 3 months post-childbirth	
	Recovery (PHQ-9 score $< 5$ at both 3 and 6 months post-childbirth)	
	Response I (PHQ-9 score $< 10$ at both 3 and 6 months post-childbirth)	
	Response II ( $\geq 50\%$ reduction from baseline PHQ-9 score at both 3 and 6 months post-childbirth)	
	WHO-DAS score, at 3 and 6 months post-childbirth	
	Number of days unable to work in the last month, at 3 and 6 months post-childbirth	
	MSPSS score, at 3 and 6 months post-childbirth	
<b>Recruitment period and follow up</b>	October 2014-June 2017	October 2014-March 2017
<b>Number of participants</b>	280	570

518 The trial designs have been described previously in detail (Sikander et al., 2015). CHW=community health workers. LHW=  
519 Lady Health Worker. THPP=Thinking Healthy Programme Peer-delivered. PHQ=Patient Health Questionnaire.  
520 MSPSS=Multidimensional Scale of Perceived Social Support. WHO-DAS=World Health Organization Disability Assessment  
521 Schedule. <sup>a</sup>Maternal secondary outcomes only.  
522

523 **Table 2. Baseline characteristics by trial.**

	THPP-India	THPP-Pakistan	Total
<b>Number enrolled</b>	280	570	850
<b>Age, years</b> (mean (SD; range))	25 (4.6; 18-41)	27 (4.8; 18-45)	26 (4.8; 18-45)
<b>Level of education</b> (n (%))			
No formal education	34 (12%)	107 (19%)	141 (17%)
Up to primary	120 (43%)	39 (7%)	159 (19%)
Up to secondary	90 (32%)	333 (58%)	423 (50%)
Beyond secondary	36 (13%)	91 (16%)	127 (15%)
<b>Occupation</b> (n (%))			
Does not work	237 (85%)	533 (94%)	770 (91%)
Works	43 (15%)	37 (6%)	80 (9%)
<b>Chronicity of depression, weeks</b> (n (%)) <sup>a</sup>			
<12	173 (62%)	73 (18%)	246 (36%)
≥12	107 (38%)	326 (82%)	433 (64%)
<b>Symptom severity (PHQ-9 score)</b> (median (IQR))	12 (11,15)	14 (12,17)	14 (11,17)
<b>Symptom severity category</b> (n (%))			
Moderate (PHQ-9 score 10-14)	197 (70%)	312 (55%)	509 (60%)
Moderately severe (15-19)	67 (24%)	187 (33%)	254 (30%)
Severe (20-27)	16 (6%)	71 (12%)	87 (10%)
<b>MSPSS score</b> (mean (SD))	5 (1.1)	4 (1.4)	4 (1.4)
<b>Participant's expectation of usefulness of counseling</b> (n (%)) <sup>b</sup>			
Not useful	1 (0%)	4 (1%)	5 (1%)
A little	55 (20%)	26 (5%)	81 (10%)
somewhat useful	54 (19%)	114 (20%)	168 (20%)
Moderately useful	58 (21%)	246 (43%)	304 (36%)
Very useful	112 (40%)	178 (31%)	290 (34%)
<b>Parity</b> (n (%))			
Primiparous	119 (43%)	102 (18%)	221 (26%)
Multiparous	161 (57%)	468 (82%)	629 (74%)
<b>Previous miscarriage or still birth</b> (n (%))			
None	261 (93%)	377 (66%)	638 (75%)
One/more	19 (7%)	193 (34%)	212 (25%)
<b>Any domestic violence in last three months</b> (n (%)) <sup>c</sup>			
No	243 (87%)	486 (87%)	729 (87%)
Yes	37 (13%)	71 (13%)	108 (13%)
<b>Time between screening and birth of child, months</b> (mean (SD)) <sup>d</sup>	4 (1.6)	3 (1.2)	3 (1.5)

524 Percentages are of non-missing values. PHQ=Patient Health Questionnaire. MSPSS=Multidimensional Scale of Perceived  
 525 Social Support. SD=standard deviation. <sup>a</sup>Missing for 171 THPP-Pakistan participants. <sup>b</sup>Missing for 2 THPP-Pakistan  
 526 participants. <sup>c</sup>Missing for 13 THPP-Pakistan participants. <sup>d</sup>Missing for 11 THPP-India and 85 THPP-Pakistan participants.

527  
528

529 **Table 3. Process indicators.**


	THPP-India	THPP-Pakistan
<b>Peers</b>		
Peers trained and delivered at least one session (number)	26	66
Age of peers, years (mean, SD)	38 (7.5)	30 (5.5)
Years of education completed by peers (mean, SD)	12 (2.7)	12 (2.5)
Peers lost during the trial (number, %)	0	23 (35%)
Attendance to supervisions (%)	67%	88%
<b>Participants in the intervention arm</b>		
Number <sup>a</sup>	138	258
Sessions attended (mean, SD) <sup>b</sup>	9.8 (4.3)	10.9 (3.9)
Prenatal sessions attended (mean, SD) <sup>c</sup>	4.5 (1.9)	3.7 (1.7)
Postnatal sessions attended (mean, SD) <sup>d</sup>	5.4 (2.9)	7.3 (2.7)
Attended at least 6 sessions	113 (82%)	230 (89%)
Attended at least 10 sessions	87 (63%)	201 (78%)
Completed treatment <sup>e</sup>	99 (72%)	201 (78%)


530 SD=standard deviation. <sup>a</sup>Excluding 2 women in THPP-India and 25 in THPP-Pakistan who were discontinued as per protocol  
531 before month 3 due to child death, still birth or abortion. <sup>b</sup>Of possible 14 sessions in each trial. <sup>c</sup>Of possible 6 sessions in  
532 THPP-India and 5 in THPP-Pakistan. <sup>d</sup>Of possible 8 sessions in THPP-India and 9 in THPP-Pakistan. <sup>e</sup>Defined in THPP-India as  
533 attended at least 6 sessions, with at least 1 session in each of four phases (prenatal, and 1-2, 3-4 and 5-6 months postnatal);  
534 defined in THPP-Pakistan as attended at least 10 sessions.  
535

536 **Table 4. Primary and secondary outcomes.**

	Number of participants <sup>a</sup>		Mean (SE) or number (%) <sup>a</sup>		Intervention effect (adjusted mean difference or odds ratio; 95% CI) <sup>b</sup>	P value
	Control	Intervention	Control	Intervention		
<b>Primary outcomes</b>						
Symptom severity (PHQ-9 score) at 6 months						
Overall	355	349	6.0 (0.3)	5.1 (0.3)	-0.78 (-1.47,-0.09)	0.03
India	129	122	4.5 (0.4)	3.5 (0.4)	-0.95 (-2.31,0.41)	
Pakistan	226	227	6.8 (0.4)	6.0 (0.4)	-0.72 (-1.52,0.08)	
Remission (PHQ-9 score <5) at 6 months						
Overall	355	349	178 (50%)	201 (58%)	1.35 (1.02,1.78)	0.04
India	129	122	77 (60%)	89 (73%)	1.86 (1.08,3.20)	
Pakistan	226	227	101 (45%)	112 (49%)	1.20 (0.87,1.66)	
<b>Secondary outcomes</b>						
<b>Continuous variables</b>						
Symptom severity (PHQ-9 score) at 3 months	333	346	7.1 (0.4)	5.5 (0.3)	-1.84 (-2.43,-1.25)	<0.001
WHO-DAS score						
3 months	332	346	17.0 (1.0)	14.2 (0.9)	-3.17 (-5.73,-0.61) <sup>c</sup>	0.02 <sup>c</sup>
6 months	354	349	16.0 (1.0)	13.6 (0.9)	-2.11 (-4.77,0.55)	0.12
Number of days unable to work in last month						
3 months	332	346	1.7 (0.3)	1.6 (0.3)	-0.24 (-0.86,0.39)	0.46
6 months	354	349	1.6 (0.2)	1.6 (0.3)	0.02 (-0.51,0.55)	0.95
MSPSS score						
3 months	333	345	4.7 (0.1)	4.9 (0.1)	0.17 (0.03,0.31)	0.01
6 months	354	349	4.7 (0.1)	5.0 (0.1)	0.27 (0.11,0.43)	0.001
<b>Categorical variables</b>						
Remission (PHQ-9<5) at 3 months	333	346	155 (47%)	186 (54%)	1.34 (1.13,1.60)	0.001
Recovery (PHQ-9<5 at both 3 and 6 months)	317	325	100 (32%)	135 (42%)	1.60 (1.21,2.10)	0.001
Response I (PHQ-9<10 at both 3 and 6 months)	317	325	188 (59%)	224 (69%)	1.57 (1.16,2.10)	0.003
Response II (≥50% reduction from baseline PHQ-9 at both 3 and 6 months)	317	325	144 (45%)	190 (58%)	1.69 (1.27,2.24)	<0.001

537 PHQ=Patient Health Questionnaire. MSPSS=Multidimensional Scale of Perceived Social Support. WHO-DAS=World Health  
538 Organization Disability Assessment Schedule. SE=standard error. CI=confidence interval. <sup>a</sup>India and Pakistan trials combined  
539 for the secondary outcomes. <sup>b</sup>Linear or logistic GEE models, adjusted for country, recruitment site, residence, union council,  
540 baseline symptom severity, treatment expectations, education, chronicity of depression, and time between screening and  
541 birth (see methods). Baseline treatment expectations missing for one woman in Pakistan control group, therefore models  
542 are based on one fewer woman than the numbers indicated. Country-level intervention effects are based on models with  
543 interaction between group and country. <sup>c</sup>Using independent correlation matrix due to convergence problems when using  
544 exchangeable correlation structure.  
545





**Effectiveness of the Thinking Healthy Programme for perinatal depression delivered through peers:  
pooled analysis of two randomized controlled trials in India and Pakistan**

**Declaration of interest**

Declarations of interest: none.

## **Effectiveness of the Thinking Healthy Programme for perinatal depression delivered through peers: pooled analysis of two randomized controlled trials in India and Pakistan**

### **Contributors**

DCF, BW, AL, HAW, AR and VP designed the THPP-India trial. EA, ADS, AJ, PK, and RK were responsible for intervention implementation and data gathering instruments in the THPP-India trial. BW, AL, and VP were responsible for THPP-India trial conduct. ED'S designed and managed the database for the THPP-India trial. SS, IA, DCF, HAW, VP and AR designed the THPP-Pakistan trial. IA, NA, AB, TB, SB, RL, MS and SZ were responsible for intervention implementation and data gathering instruments in the THPP-Pakistan trial. SS, AZ, and AR were responsible for THPP-Pakistan trial conduct. AZ designed and managed the database for the THPP-Pakistan trial. FV and HAW performed the statistical analyses. VP and AR provided oversight. FV and HAW drafted the manuscript. All authors read and approved the manuscript.

### **Funding and role of the funding source**

This work was supported by the National Institute of Mental Health, USA (grant number 1U19MH095687). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute of Mental Health, the National Institute of Health, or the US Department of Health and Human Services. The funder of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report, except for LeShawndra Price who was a Scientific Collaborator for the individual trials, under the cooperative agreement that funded the research. We are grateful for support from the Directorate of Health Services and Directorate of Women and Child Development, Government of Goa, and Health Services Academy, Ministry of National Health Services Regulations & Coordination, Pakistan. FV and HAW were supported by a grant from the Medical Research Council (MRC) and the Department for International Development (DFID UK) under the MRC/DFID Concordat (K012126/1).

## **Acknowledgements**

We would like to thank Shreya Nagvekar, Sharmin Colaco, Seema Kanolkar and Kishori Mandrekar for their assistance and invaluable contribution to the THPP-India trial.

**Effectiveness of the Thinking Healthy Programme for perinatal depression delivered through peers:  
pooled analysis of two randomized controlled trials in India and Pakistan**

**SUPPLEMENT**

**Table A1. Factors associated with country.**

<b>Variable</b>	<b>Adjusted odds ratio (95% CI) for Pakistan versus India<sup>a</sup></b>	<b>P value</b>
<b>Age, per 10 years</b>	0.98 (0.50,1.91)	0.95
<b>Level of education</b>		<0.001
None	1 (reference)	
Primary	0.09 (0.04,0.24)	
Secondary	1.60 (0.73,3.50)	
Higher secondary/above	2.46 (0.96,6.28)	
<b>Occupation</b>		0.003
Does not work	1 (reference)	
Works	0.28 (0.12,0.65)	
<b>Chronicity of depression, weeks</b>		<0.001
<12	0.14 (0.08,0.25)	
≥12	1 (reference)	
<b>Symptom severity (PHQ-9 score)</b>		0.003
10-14	1 (reference)	
15-19	2.41 (1.32,4.40)	
≥20	3.73 (1.29,10.81)	
<b>MSPSS score, per unit</b>	0.40 (0.31,0.51)	<0.001
<b>Participant's expectation of usefulness of counselling</b>		<0.001
Not/a little useful	1 (reference)	
Somewhat useful	3.30 (1.14,9.55)	
Moderately useful	11.17 (3.97,31.43)	
Very useful	5.07 (1.89,13.60)	
<b>Parity</b>		0.56
Primiparous	1 (reference)	
Multiparous	1.22 (0.63,2.36)	
<b>Previous miscarriage or still birth</b>		<0.001
None	1 (reference)	
One or more	9.43 (4.05,21.95)	
<b>Domestic violence</b>		0.01
No	1 (reference)	
Yes	0.34 (0.15,0.79)	
<b>Time between screening and birth of child, per month</b>	0.54 (0.43,0.66)	<0.001

PHQ=Patient Health Questionnaire. MSPSS=Multidimensional Scale of Perceived Social Support. CI=confidence interval.

<sup>a</sup>Logistic regression model. N=589 (women with any missing data were excluded). Broadly similar results were obtained from models excluding chronicity of depression and time between screening and birth (which had the highest missingness, particularly in Pakistan).

**Table A2. Sensitivity analyses for the primary outcomes.**

	Number of clusters	Intervention effect (adjusted mean difference or odds ratio; 95% CI) <sup>a</sup>	P value	P value for effect modification by country
<b>Symptom severity (PHQ-9 score) at 6 months</b>				
Primary analysis as reported in Table 4 (GEEs with individuals in India acting as their own clusters)	291	-0.78 (-1.47,-0.09)	0.03	0.77
GEEs with individuals in India grouped as one cluster	41	-0.82 (-1.64,-0.01) <sup>b</sup>	0.05 <sup>b</sup>	0.81 <sup>b</sup>
Random effects models with individuals in India acting as their own clusters	291	-0.82 (-1.64,-0.01)	0.05	0.81
Random effects models with individuals in India grouped as one cluster	41	-0.82 (-1.64,-0.01)	0.05	0.81
<b>Remission (PHQ-9 score &lt;5) at 6 months</b>				
Primary analysis as reported in Table 4 (GEEs with individuals in India acting as their own clusters)	291	1.35 (1.02,1.78)	0.04	0.18
GEEs with individuals in India grouped as one cluster	41	1.39 (1.02,1.91) <sup>b</sup>	0.04 <sup>b</sup>	0.20 <sup>b</sup>
Random effects models with individuals in India acting as their own clusters	291	1.39 (0.02,1.91)	0.04	0.20
Random effects models with individuals in India grouped as one cluster	41	1.39 (1.02,1.91)	0.04	0.20

<sup>a</sup>Linear or logistic GEE models as described in the methods section, adjusted for country, recruitment site, residence, union council, baseline symptom severity, treatment expectations, education, chronicity of depression, and time between screening and birth (see methods). <sup>b</sup>Using independent correlation matrix due to convergence problems when using exchangeable correlation structure.

**Table A3. Remission (PHQ-9 score <5) at 6 months by potential effect modifiers.**

	Number of participants		Number with remission (%)		Intervention effect (adjusted odds ratio; 95% CI) <sup>a</sup>	P value for effect modification
	Control	Intervention	Control	Intervention		
Age, years						0.18
18-24	135	132	91 (69%)	74 (55%)	1.86 (1.08,3.20)	
≥25	220	217	110 (51%)	104 (47%)	1.20 (0.87,1.66)	
Chronicity of depression, weeks						0.03
<12	105	103	73 (71%)	55 (52%)	2.54 (1.39,4.65)	
≥12	172	178	88 (49%)	79 (46%)	1.10 (0.73,1.65)	
Missing <sup>b</sup>	78	68	40 (59%)	44 (56%)	-	
Baseline symptom severity (PHQ-9 score)						0.49
10-14	223	209	135 (65%)	124 (56%)	1.48 (1.01,2.17)	
≥15	132	140	66 (47%)	54 (41%)	1.17 (0.73,1.90)	
Treatment expectations <sup>c</sup>						0.85
None/little/somewhat	114	96	56 (58%)	58 (51%)	1.29 (0.73,2.26)	
Moderate/very useful	241	252	144 (57%)	120 (50%)	1.37 (0.98,1.93)	
Parity						0.03
Primiparous	91	95	72 (76%)	52 (57%)	2.59 (1.35,4.93)	
Multiparous	264	254	129 (51%)	126 (48%)	1.10 (0.80,1.52)	

PHQ=Patient Health Questionnaire. CI=confidence interval. <sup>a</sup>Logistic GEE models, adjusted for country, recruitment site, residence, union council, baseline symptom severity, treatment expectations, education, chronicity of depression, and time between screening and birth (see methods). Baseline treatment expectations missing for one woman in Pakistan control group, therefore models are based on one fewer women than indicated by the numbers indicated. These results are illustrated in Figure 1B. <sup>b</sup>Model includes missing chronicity category, but interaction effects are presented only among those with non missing chronicity. <sup>c</sup>Missing for one woman in Pakistan control group, who is omitted.

**Table A4. Symptom severity (PHQ-9 score) at 6 months by potential effect modifiers.**

	Number of participants		Mean symptom severity (PHQ-9 score; SE)		Intervention effect (adjusted mean difference; 95% CI) <sup>a</sup>	P value for effect modification
	Control	Intervention	Control	Intervention		
Age, years						0.27
18-24	135	132	5.9 (0.5)	4.4 (0.5)	-1.38 (-2.66,-0.10)	
≥25	220	217	6.0 (0.4)	5.5 (0.4)	-0.44 (-1.35,0.47)	
Chronicity of depression, weeks						0.49
<12	105	103	5.1 (0.5)	3.7 (0.5)	-1.67 (-3.16,-0.18)	
≥12	172	178	6.9 (0.5)	5.8 (0.4)	-0.99 (-2.02,0.03)	
Missing <sup>b</sup>	78	68	4.9 (0.6)	5.6 (0.7)	-	
Baseline symptom severity (PHQ-9 score)						0.80
10-14	223	209	5.1 (0.4)	4.4 (0.4)	-0.69 (-1.65,0.27)	
≥15	132	140	7.4 (0.6)	6.2 (0.5)	-0.91 (-2.14,0.31)	
Treatment expectations <sup>c</sup>						0.35
None/little/somewhat	114	96	6.3 (0.6)	4.5 (0.5)	-1.37 (-2.80,0.05)	
Moderate/very useful	241	252	5.8 (0.4)	5.3 (0.4)	-0.54 (-1.39,0.30)	
Parity						0.29
Primiparous	91	95	4.8 (0.5)	3.3 (0.5)	-1.51 (-3.05,0.04)	
Multiparous	264	254	6.4 (0.4)	5.8 (0.4)	-0.51 (-1.32,0.29)	

PHQ=Patient Health Questionnaire. SE=standard error. CI=confidence interval. <sup>a</sup>Linear GEE models, adjusted for country, recruitment site, residence, union council, baseline symptom severity, treatment expectations, education, chronicity of depression, and time between screening and birth (see methods). Baseline treatment expectations missing for one woman in Pakistan control group, therefore models are based on one fewer women than indicated by the numbers indicated. These results are illustrated in Figure 1A. <sup>b</sup>Model includes missing chronicity category, but interaction effects are presented only among those with non missing chronicity. <sup>c</sup>Missing for one woman in Pakistan control group, who is omitted.

**Table A5. Factors associated with the primary outcomes.**

	Symptom severity (PHQ-9 score) at 6 months		Remission (PHQ-9 score <5) at 6 months	
	Adjusted mean difference (95% CI) <sup>a</sup>	P value	Adjusted odds ratio (95% CI) <sup>a</sup>	P value
<b>Randomised group</b>		0.03		0.04
Control	0 (reference)		1 (reference)	
Intervention	-0.78 (-1.47,-0.09)		1.35 (1.02,1.78)	
<b>Country</b>		<0.001		0.001
India	0 (reference)		1 (reference)	
Pakistan	2.81 (1.23,4.39)		0.34 (0.18,0.64)	
<b>Recruitment site</b>		0.84		0.81
N/A (Pakistan) or Goa Medical College Hospital	0 (reference)		1 (reference)	
Asilo	0.32 (-1.15,1.79)		1.10 (0.62,1.96)	
Primary health centres	-0.46 (-3.93,3.02)		1.62 (0.37,7.20)	
<b>Residence</b>		0.67		0.85
N/A (Pakistan) or rural	0 (reference)		1 (reference)	
Urban	-0.38 (-2.09,1.33)		1.07 (0.54,2.09)	
<b>Union council</b>		0.002		0.03
N/A (India) or Guf	0 (reference)		1 (reference)	
Kallar	-0.61 (-2.22,0.99)		1.44 (0.76,2.75)	
Kanoha	0.06 (-1.70,1.83)		1.47 (0.71,3.02)	
Sakot	-1.67 (-3.81,0.46)		1.55 (0.66,3.65)	
Choa	-1.33 (-3.23,0.58)		1.00 (1.00,1.00)	
Smoot	-2.60 (-4.66,-0.53)		1.87 (0.86,4.09)	
Bewek	1.68 (0.15,3.21)		3.14 (1.32,7.47)	
Darkali	-0.92 (-2.38,0.54)		0.57 (0.29,1.11)	
Bishandot	0.49 (-1.19,2.17)		1.24 (0.69,2.24)	
Sagri	-0.84 (-2.39,0.71)		1.31 (0.67,2.56)	
Nallah	-2.57 (-4.75,-0.39)		1.52 (0.81,2.86)	
<b>Symptom severity (PHQ-9 score)</b>		0.004		0.02
10-14	0 (reference)		1 (reference)	
15-19	1.56 (0.61,2.51)		0.59 (0.41,0.85)	
≥20	1.39 (-0.10,2.87)		0.71 (0.40,1.26)	
<b>Participant's expectation of usefulness of counselling</b>		0.78		0.99
Not/a little useful	0 (reference)		1 (reference)	
Somewhat useful	-0.22 (-1.81,1.38)		0.99 (0.53,1.84)	
Moderately useful	-0.19 (-1.70,1.33)		0.95 (0.52,1.72)	
Very useful	-0.62 (-2.12,0.88)		1.01 (0.56,1.82)	
<b>Education status</b>		0.009		0.12
No formal education	0 (reference)		1 (reference)	
Up to primary	-0.37 (-1.83,1.09)		1.12 (0.64,1.98)	
Up to secondary	-0.47 (-1.65,0.71)		1.43 (0.91,2.27)	
Beyond secondary	-2.38 (-3.90,-0.86)		1.95 (1.07,3.55)	
<b>Chronicity of depression, weeks</b>		0.13		0.07

<12	-0.45 (-1.52,0.63)		1.06 (0.70,1.62)	
≥12	0 (reference)		1 (reference)	
Missing	-1.23 (-2.44,-0.01)		1.74 (1.09,2.79)	
<b>Time between screening and birth of child, per month</b>	-0.16 (-0.46,0.14)	0.30	1.08 (0.96,1.22)	0.19


PHQ=Patient Health Questionnaire. MSPSS=Multidimensional Scale of Perceived Social Support. CI=confidence interval. N/A=not applicable. <sup>a</sup>Linear or logistic GEE models as described in the methods section, adjusted for all variables shown in the table.

**Table A6. Analysis of repeated measures for the primary and secondary outcomes.**

	<b>P value for group by time interaction</b>	<b>Overall adjusted mean difference or odds ratio (95% CI)</b>	<b>P value for overall intervention effect</b>
<b>Primary outcomes</b>			
Symptom severity (PHQ-9 score)	0.17	-1.30 (-1.86,-0.74)	<0.001
Remission (PHQ-9 score <5)	0.98	1.39 (1.13,1.72)	0.002
<b>Secondary outcomes</b>			
WHO DAS score	0.81	-2.56 (-4.31,-0.81)	0.004
Number of days unable to work in last month	0.82	-0.12 (-0.55,0.32)	0.59
MSPSS score	0.37	0.25 (0.11,0.38)	<0.001

PHQ=Patient Health Questionnaire. WHO-DAS=World Health Organization Disability Assessment Schedule. MSPSS=Multidimensional Scale of Perceived Social Support. CI=confidence interval. Linear or logistic GEE models, adjusted for visit month, country, recruitment site, residence, union council, baseline symptom severity, treatment expectations, education, chronicity of depression, and time between screening and birth (see methods).

Figure A1. Symptom severity (PHQ-9 score) over time.



The central line shows the median score, the length of the box is the interquartile range, the lines are the most extreme values within 1.5 times the interquartile range, and points outside this range are shown individually.