

Recessive Z-linked lethals and the retention of haplotype diversity in a captive butterfly population

Ilik J. Saccheri, Samuel Whiteford, Carl J. Yung, Arjen E. van't Hof

Institute of Integrative Biology, University of Liverpool, Liverpool, UK

Abstract

Sex chromosomes are predicted to harbour elevated levels of sexually antagonistic variation due to asymmetries in the heritability of recessive traits in the homogametic vs heterogametic sex. This evolutionary dynamic may manifest as high recessive load specifically affecting the homogametic sex, and the retention of haplotype diversity in small populations. We tested the hypothesis that the Z chromosome in the butterfly *Bicyclus anynana* carries a high inbred load for male fertility and viability. Homozygosity of Z chromosome blocks was produced by daughter-father backcrosses, and inferred from marker loci positioned via a linkage map. Male sterility was, in general, unrelated to homozygosity in any region of the Z, but there was an extreme deficit of homozygous males within a 2cM interval in all families. In contrast, no corresponding skew in Z genotype was detected in their (hemizygous) sisters. The same pattern was observed in historically inbred lines, indicating a high frequency of recessive lethals in the ancestral population. Allele frequency changes between 1993 and 2006 (70 generations at $N_e \sim 160$) show that, despite the loss of many haplotypes, diversity was retained significantly above the neutral expectation. Effective overdominance in the lethal region can account for this effect locally but not in other parts of the chromosome, that are also associated with persistent linkage disequilibrium. These unexpected patterns suggest the operation of other factors, such as epistatic selection, recombination suppression, assortative mating, and meiotic drive. Our results highlight the role of balancing selection in maintaining the inbred load and linked genetic diversity.

Introduction

Understanding the factors that maintain the alleles responsible for inbreeding depression is relevant to explaining fitness variation in natural populations (Charlesworth, 2015), to predicting the risk of extinction of small populations (Saccheri *et al*, 1998; Bijlsma *et al*, 2000), and to interpreting changes in genetic marker diversity with respect to effective population size and selection (Zhao and Charlesworth, 2016). Mutational input of unconditionally deleterious variants, kept at low equilibrium frequencies by purifying selection, is the primary underlying process. However, it is increasingly acknowledged that a significant component of the inbred load is due to alleles maintained at intermediate frequencies by various forms of balancing selection (Charlesworth and Charlesworth, 1999; Charlesworth, 2015). The idea that overdominant loci, even if relatively few in number, could make a disproportionate contribution to fitness variation, has been around for a long time (reviewed in Crow, 1987) but lacks strong empirical support (Charlesworth and Willis, 2009; Liberatore *et al*, 2013). An alternative and perhaps more widespread balancing mechanism is antagonistic pleiotropy, where a single allele is beneficial through one trait but deleterious through another trait (Rose, 1982; Charlesworth and Hughes, 2000).

In species with separate sexes, sexually antagonistic pleiotropy (intralocus sexual conflict; Bonduriansky and Chenoweth, 2009) is predicted to affect many components of fitness (Connallon and Clark, 2014; Zajitschek and Connallon, 2018). This was borne out by the first genome-wide study of sexually antagonistic polymorphisms, which found many deeply conserved clusters of sexually antagonistic SNPs throughout the genome of *Drosophila melanogaster*, reflecting structural constraints to resolving functional conflicts (Ruzicka *et al*, 2019). Surprisingly, given some theoretical predictions (Rice, 1984), and earlier studies in *D. melanogaster* (Innocenti and Morrow, 2010) and humans (Lucotte *et al*, 2016), this study did not find a significant enrichment of these genes on the X chromosome, compared to autosomes. The major contribution from autosomes was attributed, in part, to sex-specific dominance effects, where the same allele confers a (partially) dominant benefit to one sex but a (partially) recessive cost to the opposite sex, which broaden the conditions (sexually antagonistic cost/benefit ratio) for the maintenance of autosomal polymorphisms (Fry, 2010).

Notwithstanding the likely importance of dominance modifiers for autosomal polymorphism (Spencer and Priest, 2016), Rice's (1984) single-locus model predicts that, in species with highly diverged heteromorphic sex chromosomes, intralocus sexual conflict at sex-linked loci may lead to asymmetries in the sex-specific inbred load. This is because a sexually antagonistic allele conferring a recessive cost to the homogametic sex (XX females or ZZ males) can be maintained through its beneficial effect to the heterogametic sex (XY males or ZW females), under the assumption that in the hemizygous state alleles are always expressed (i.e., dominance restrictions do not apply). In this model (Rice, 1984; Gibson *et al*, 2002), a relatively small pleiotropic benefit to the heterogametic sex can sustain low equilibrium frequencies of strongly deleterious recessives, whose homozygous cost is rarely exposed in the homogametic sex. Conversely, the conditions for maintaining alleles that are deleterious to the heterogametic sex are much more restrictive and rely on large dominant benefits to the homogametic sex, coupled with relatively mild costs to the heterogametic sex. Thus, we might expect a concentration of larger effect deleterious recessives affecting the homogametic sex on sex chromosomes. Sex chromosomes are also expected to be enriched for genes whose expression is biased towards the homogametic sex (Ellegren, 2011; Huylmans *et al*, 2017).

No clear pattern emerges as to the sex or gamety that is more sensitive to inbreeding (reviewed in Ebel and Phillips, 2016). For example, in *D. melanogaster*, inbreeding depression for fitness is greater for (XY) males than for (XX) females, due to male-male competition (Enders and Nunney, 2010; Mallet and Chippindale, 2011), whereas in *Caenorhabditis remanei*, (XX) females are much more sensitive to inbreeding than are (XO) males (Ebel and Phillips, 2016). Lack of a consistent sex chromosome effect is also provided by the pedigreed population of *Melospiza melodia*, where inbreeding depression for lifetime reproductive success was substantially greater for (ZW) females than (ZZ) males, predominantly through a reduction in the proportion of eggs hatching (Keller, 1998). Other factors, such as the genetic architecture of sex-specific traits, the contribution of these traits to total fitness, the strength of sexual selection and inbreeding imposed by life-history and mating system, and the mechanism of dosage compensation, will also influence sexual asymmetries in the inbred load (Janicke *et al*, 2013; Mank *et al*, 2014).

The physical distribution of the genetic load (*sensu lato*) within chromosomes, and the distribution of dominance effects, has major consequences for the pattern of linked molecular genetic diversity. In large populations, background selection against deleterious recessive alleles is expected to accelerate the rate of loss of linked neutral genetic variation, compared to the purely neutral case (Charlesworth *et al*, 1993). However, in small populations, linkage disequilibrium between two selected loci, each segregating deleterious recessives, and a linked (neutral) marker locus leads to apparent heterozygote advantage at the marker locus (associative overdominance; Ohta, 1971), even though the selected loci do not show true overdominance. Under certain conditions (low coefficient of dominance and weak selection) this type of association can retard the loss of genetic diversity due to genetic drift (Wang *et al*, 1999; Zhao and Charlesworth, 2016). This effect may account for some cases, typically in laboratory populations, where lower than expected declines in heterozygosity have been observed (e.g., Schou *et al*, 2017).

A laboratory population of the Afro-tropical butterfly *Bicyclus anynana* (Satyridae) has been maintained in captivity since 1988 and offers the opportunity to relate temporal changes in molecular genetic diversity to genetic (recessive) load in a male-homogametic (ZZ) species. Previous studies have revealed that *B. anynana* carries a very high inbred load (> 7 lethal equivalents per zygote), predominantly affecting male fertility and embryo viability (Saccheri *et al*, 1996; Saccheri *et al*, 2005). For comparison, the egg-adult inbred load in *Drosophila* spp. is of the order of 1-2 lethal equivalents (Dobzhansky *et al*, 1963; García *et al*, 1994). In *B. anynana*, sterility is strongly affected by the inbreeding status of males (inbreeding coefficient $F \sim 0.25$), whereas the degree of inbreeding of females appears to have essentially no effect on this trait (Saccheri *et al*, 2005). The qualitatively male-biased sterility effect and generally high inbred load, coupled with theoretical predictions of intralocus sexual conflict models (Rice, 1984; Gibson *et al*, 2002), suggested the presence of one or more Z-linked loci segregating recessive alleles with large deleterious effects. We therefore set out to test the null hypothesis that homozygosity (in diploid males) or haplotype identity (in hemizygous females) of Z chromosome segments (denoted by z) has no effect on fertility or viability. The W chromosome in this species is unusually small and degenerated (van't Hof *et al*, 2008), implying few if any functional regions that are homologous with the Z. (For practical reasons at the time of the experiments, the study was restricted to the Z chromosome, effectively ignoring any influence from the 27 autosomes.) In addition to seeking genotype-phenotype associations in segregating sibships, we used archival DNA from the ancestral base population to link changes in Z-haplotype diversity over 70 generations in captivity to selection and drift. A linkage map of

the Z chromosome was constructed and used to identify marker polymorphisms, which were then applied to segregating sibships and population samples.

Materials and methods

Overview of samples

This study combined analysis of experimental crosses (linkage mapping family and backcrosses) and bottlenecked lines, with population samples taken at two time points (1993 and 2006), all derived from the same initial laboratory population (Figure 1a).

Linkage map and genetic markers

Our objective was to obtain 8-10 evenly spaced markers across the Z chromosome; however, inconsistent gene order between *Bombyx mori* and *B. anynana* made it necessary to map a larger number of loci (34) to achieve this. The majority were initially targeted using *B. mori* Z chromosome scaffold orthologs (Wang *et al*, 2005). Ten Z-linked genes were already mapped in *B. anynana* (van't Hof *et al*, 2008; Beldade *et al*, 2009). Prior to the release of the *B. anynana* genome assembly (Nowell *et al*, 2017), gene sequences were obtained using degenerate primers based on conserved regions (Table S1), and expressed sequence tags (Beldade *et al*, 2006). Segregating polymorphisms were most commonly found by screening the father of the mapping family with intron-spanning PCR followed by Sanger sequencing. For some markers, exonic polymorphisms were used, and apterous A was mapped using a BAC sequence polymorphism outside the gene (genbank AC239117). Genotyping methods consisted of PCR-RFLP, indels producing distinct gel bands, and Sanger sequencing (PCR primers in Table S1). DNA was extracted from the head and thorax of frozen adults, following a standard phenol-chloroform protocol (Saccheri and Bruford, 1993). 1/100 dilution of this DNA extract was used as PCR template. PCR reactions were performed in 12 μ l volumes, containing 1X LongAmp buffer, 1.2 units of LongAmp polymerase (NEB), 0.4 μ M of each primer and 0.5 μ l of diluted DNA extract. A standard 3-step 40 cycle amplification was used with an annealing temperature (T_a) of 57°C for species-specific primers and T_a 50°C for degenerate primers. Most marker alleles consisted of multiple linked single nucleotide polymorphisms (SNPs) defining a 'microhaplotype'.

The mapping family is the same used for a previously published AFLP-based linkage map (van't Hof *et al*, 2008) with an additional 92 sibs included giving a total of 184 offspring. The large number of offspring and markers required a cost-effective mapping approach, allowing detailed mapping around recombination breakpoints combined with crude-mapping of non-recombining regions. A preliminary linkage map was constructed using indels or PCR-RFLP with distinct allelic bands on agarose gels ('cheap markers'). This revealed all recombination events in the mapping family. The next step was to roughly position the loci genotyped by Sanger sequencing ('expensive markers') relative to the preliminary map based on a subset of offspring. Subsequently, these loci were fine-tuned to their exact position by only sequencing individuals with recombination breakpoints near the 'expensive marker'. Missing genotypes inside non-recombinant chromosome sections (undisrupted inherited paternal haplotype sections) can then be reliably inferred using their surrounding 'cheap markers'. Errors caused by undetected double recombinants on the preliminary map are unexpected due to the restrictions of interference, i.e. the absence of double recombinations within small intervals (Muller, 1916). Markers near chromosome ends were sequenced in all offspring because their genotypes cannot be inferred reliably. The map was produced with Joinmap 4.0 (Kyazma) using the Haldane mapping method. The procedures to account for absence of female

recombination in Lepidoptera only apply to autosomes and were therefore not implemented for the Z chromosome mapping.

Backcross and egg hatching

Female Lepidoptera are heterogametic (ZW), receiving the Z chromosome from their father only, whereas homogametic (ZZ) males inherit one Z copy from each parent. Furthermore, there is no crossing-over in females. A daughter-father backcross (father Z_1Z_2 ; daughter either Z_1W or Z_2W) is therefore expected to produce a 1:1 ratio of Z_1Z_1 (homozygous) : Z_1Z_2 (heterozygous) sons (for the Z_1W daughter case) and a 1:1 ratio of Z_1W : Z_2W daughters. In reality, recombination breaks down the original Z_1 and Z_2 haplotypes, such that the above expectation holds, not for the whole chromosome, but for large chromosome segments, denoted by z (Figure 1b). For autosomal genes, the expected probability of identity by descent is 0.25 for both sexes. Fourteen daughter-father backcrosses were established from a laboratory stock population originally derived from Nkhata Bay, Malawi (our lab population had been in captivity for ~90 generations at the time of the experiments). Husbandry was standard for this species: larvae were reared on potted maize plants at 27°C; adults were fed on mashed banana. Potential backcross fathers were held at 20°C to retard senescence while their daughters developed into adults. Eggs were collected from mated daughters individually on cuttings of their wild host-plant, *Oplismenus compositus*. Backcross larvae were reared to adults in sleeve enclosures and 11-45 sons per backcross mated to (relatively) unrelated outbred females (F_1 hybrids of inter-line crosses). Egg batches from these females were used to estimate the frequency of sterile sons produced by each backcross (proportion of eggs that did not hatch or show visible signs of being fertilized, such as a head capsule). To keep sequencing costs down, only five of the backcross families were selected for genotyping, on the basis of having intermediate levels of sterility (maximising statistical power), with the exception of one family that was 100% sterile, and the presence of paternal polymorphisms at marker loci. In each of these, a sample of sons (mean $n = 26$) and daughters ($n = 16$, in one case 14) was genotyped. Different combinations of five or six Z-linked loci were used for different families because some fathers lacked polymorphisms in some regions.

Population samples

Archival DNA representing a random sample of the ancestral (base) population in 1993 ($n = 112$ males, 112 females), ~20 generations into captivity, was used to estimate the base population frequencies of Z microhaplotypes at eight marker loci (C2745, Kettin, Tpi, Catalase, C5197, Masc, Shaker, Ldh). The 1993 male sample was used to estimate the observed frequencies of homozygotes at these same loci. As a temporal comparison, a sample of 48 females from the parental (backcross) generation, separated in time from the archival sample by ~70 generations in captivity (assuming an average generation time of 10 weeks), was also genotyped for the full set of eight loci. Archival DNA was additionally available for six 'bottleneck' lines, each established from a single pair of randomly mated individuals from the same ancestral base population in 1993 (Saccheri *et al*, 1996; Saccheri *et al*, 1999), consisting of the founders and a small sample of the descendant F_3 . This material was used to assess genotype frequency distortion due to inbreeding of the outbred base population after relatively few generations in captivity.

Analysis of homozygosity, haplotype diversity, and linkage disequilibrium in the ancestral and descendant populations

The complexity and diversity of the marker sequences in the ancestral or base (Leiden-1993) population made it impractical to fully deconvolute heterozygous chromatographs into the constituent microhaplotypes; therefore, microhaplotype frequencies were estimated from the

female sample only. It was, however, possible to identify microhaplotype homozygotes in the male sample (those producing clean chromatographs). For each marker locus, the observed number of homozygotes was compared to its Hardy-Weinberg expectation, $n \sum q_i^2$, where q_i is the frequency of each allele (microhaplotype) and n is the size of the male sample. Chromosomal haplotypes spanning the eight marker loci were determined for each female sample (1993 and 2006). Microhaplotype identification numbers were assigned in order of their frequency in the 1993 sample.

Haplotype diversity (H_d), equivalent to expected heterozygosity, at each marker locus and time point (1993 and 2006) was estimated from the female allele (microhaplotype) frequency spectra as $(1 - \sum q_i^2) \frac{n}{n-1}$, where n is the size of the female sample (Nei, 1987). Standard error of H_d was calculated as the square root of the variance, as defined by Nei (1987) equation 8.12, replacing $2n$ by n for the haploid female sample. Allelic richness (the number of microhaplotypes, H_n) for each locus, as implemented in R package hierfstat (version 0.04; Goudet and Jombart, 2015), was calculated to account for differing n in 1993 and 2006. A point estimate of the expected H_d for each locus in 2006 due only to genetic drift was calculated as haplotype diversity in 1993 multiplied by $(1 - (1/1.5N_e))^t$, where the effective population size (N_e) was assumed to be 250 for (t) 40 generations (in Leiden), followed by 100 for 30 generations (in Liverpool). These effective population sizes are approximate as population sizes were not accurately recorded throughout the period but census numbers of adults (N) were generally kept at 400-500 in Leiden and 150-300 in Liverpool. We applied a $N_e:N$ ratio of 0.6, previously estimated for equivalent caged populations of *B. anynana* (Brakefield *et al*, 2001).

Simulation, implemented in simuPOP (Peng and Kimmel, 2005), was used to generate distributions of diversity and LD statistics under neutrality and in the presence of effective overdominance (high frequency and diversity of recessive lethals) in the C5197-Masc region. Population size was set to 450 and 200 for generations 1-40 and 41-70, respectively. Poisson variance in family size of 5, broadly consistent with empirical estimates (Brakefield *et al*, 2001), produces neutral behaviour expected for N_e of 250 for 40 generations, followed by N_e of 100 for 30 generations. The initial population was created from the observed 1993 haplotype frequencies. Chromosomal crossovers occurred in males only, at recombination rates between neighbouring pairs of loci as specified in the linkage map (Figure 1a). Females were hemizygous and non-recombining. For simplicity, a high frequency of recessive lethals in the C5197-Masc region was specified by assigning a fitness of zero to all C5197 homozygotes. The simulation was run 1000 times.

The R package boot (version 1.3) was used to generate distributions of the observed H_d in 1993 and 2006, with 1000 replicates for each sample (Davison and Hinkley, 1997; Canty and Ripley, 2019). Distributions of the change in heterozygosity at each locus were calculated from pairs of values from the two (1993 and 2006) heterozygosity distributions. All calculations were performed in R (version 3.3.3; R Core Team, 2017). The simulations were also used to estimate the joint probability of obtaining the observed, or greater, combination of H_n and H_d in the 2006 (Liverpool) sample.

Entropy-based linkage disequilibrium (eLD), an index of linkage disequilibrium for multiallelic loci (Okada, 2018), was estimated among all pairs of loci from the ancestral and descendant female samples. To examine eLD in the simulated populations, we generated genotype files for 100 replicates of the forward simulation and calculated the mean and standard deviation of eLD across these populations. To visualise allelic associations among marker loci, alluvial

diagrams displaying the structure and distribution of whole chromosome haplotypes in the 1993 and 2006 populations were generated with Disentangler (Kumasaka *et al*, 2011). Missing data was coded as null haplotypes indicated by the label "00".

Results

Linkage map

All 34 markers mapped together, with strong log of odd values (LOD > 10), to a single linkage group of 65 cM (Figure 2a). The segregation pattern in the mapping family was consistent with female hemizyosity: daughters inherited either of the paternal alleles but not the maternal allele, whereas sons inherited the (single) maternal allele plus one of the paternal alleles. All of the mapped orthologs of *B. mori* Z-genes are also on the *B. anynana* Z chromosome. However, the evolutionary conservation of synteny does not apply to the ordering of the genes, which is very different in the two species. The order in *B. anynana* also differs from that in *Biston betularia* (van't Hof *et al*, 2013). The gene Z-MOF (BaC1211), previously flagged as inconsistent between *B. mori* and *B. anynana* (Beldade *et al*, 2009), was confirmed to be Z-linked in *B. mori*, which has an autosomal paralog and a Z-ortholog (not included in an earlier gene annotation).

Genotype-phenotype associations in backcrosses

(i) Viability

Five or six marker loci were scored in each of the five backcross families (2BF₁, 8BF₁, 9BF₁, 11BF₁, 13BF₁) (Table S3). There was a clear and significant deficit of homozygous adult males (sons) within and close to the interval containing marker loci C5197, Masc and 6PGD (Table 1). In three of the five backcross families (8BF₁, 9BF₁, 11BF₁) there were no homozygotes in the C5197 to 6PGD interval. In the other two families (2BF₁ and 13BF₁) there were only one or two homozygous sons at 6PGD and Masc, respectively. The homozygous fraction tends to increase with increasing distance from this region, though it rarely reaches the null expectation of equality with heterozygotes, likely due to only one generation of recombination (in fathers only) to decouple local heterozygosity from heterozygosity in the lethal region. In contrast, both paternal alleles occur at approximately their expected ratio of 1:1 in the daughters of all backcrosses, and there is no striking deviation within the C5197-6PDG interval. The one exception to this overall pattern is for Tpi in 13BF₁, where all sons and all daughters carry the same paternal allele, which being the same as the maternal allele results in no heterozygous sons. The identity of the lethal core haplotype in the C5197-6PGD interval is different in four out of five families (9BF₁ and 13BF₁ share the same lethal haplotype).

(ii) Male fertility

The egg batches produced by the F₁ hybrid females mated to unrelated backcross sons showed a wide range of sterility and egg hatching rates (Table 2), but no association with *z* homozygosity of fathers at any of the marker loci (Table S4). The average proportion of sterile matings in the 2006 backcross families (0.62, s.d. 0.29) is somewhat higher than it was in similarly inbred parents (inbreeding coefficient = 0.25), produced by full-sib matings from the same population in 1999 and 2002 (0.43, s.d. 0.21; see Table 2 in Saccheri *et al*, 2005).

Homozygous lethal region in the ancestral population

There was a significant deficit of homozygotes at locus C5197 in the ancestral (Leiden-1993) population, but not at the Masc intronic marker. This pattern contrasts with the close to expected number of homozygotes at Tpi, Catalase, Shaker and Ldh (Table 3). Surprisingly, there was a highly significant excess of homozygotes at C2745. This was due to higher than

expected homozygotes for most of the less common haplotypes, whereas homozygotes for the most common haplotype (H1) were close to expected (15 observed vs 17 expected). In the partially inbred bottleneck lines (F_3 derived from Leiden-1993), there was a similar pattern of homozygote deficit in the C5197-Masc-6PGD interval as observed in the backcross families, and close to expected representation of founder alleles in the female sample (Table 4). The allelic identities of the missing homozygotes are diverse. The apparent deficit of homozygotes at Tpi (line 1.1) is in the opposite direction to the observation in 13BF₁, and possibly related to long-range linkage disequilibrium with the lethal region, generated through an interaction between drift and selection.

Changes in haplotype diversity and linkage disequilibrium

Comparison of the allele (microhaplotype) frequency distributions in the base (1993) and descendant (2006) populations (Figure 2b) reveals that the number of haplotypes declined substantially at every marker locus (45% average loss), but that H_d declined much less than the expected 26% at all marker loci, with the exception of Tpi (Table 5, Figure 3a, Table S5). At C2745, Catalase and C5197, H_d in 2006 was marginally higher than in 1993; at Masc, Shaker and Ldh it was ~10% lower. The retention of H_d was achieved, despite the loss of many alleles, by balancing the frequencies of the remaining alleles. The big drop at Tpi results from the loss of six out of nine haplotypes and a disproportionate increase in H1 (from 29% to 71%), which is the same allele that was exclusively inherited in 13BF₁ (Table 1); the alternative paternal allele in this family was H7.

The pattern of linkage disequilibrium in the base population (Figure 3c, upper half matrix) reveals two blocks of loci with high eLD values: C2745-Tpi (18 cM) and C5197-Ldh (18.3 cM); the former block centred on C2745-Kettin, and the latter block made up of two smaller blocks (C5197-Masc, which are very close to each other, and Shaker-Ldh, which are 7.4 cM apart). Surprisingly, these LD blocks were essentially retained over 70 generations, with some modifications (Figure 3c, lower half matrix). The lower block expanded to include Catalase; the upper block weakened locally (C2745-Tpi) but strengthened globally (C2745-Ldh). Some of the LD patterns are visually apparent in the distribution of eight-locus haplotypes (Figure S2).

Comparison with the simulated changes in H_n and H_d (Figure S1) confirm that the observed diversity in Liverpool 2006 is significantly greater than predicted under neutrality at six of the eight loci (Table 5). Moreover, the simulation incorporating overdominance at C5197, can explain the excess diversity at C5197 and Masc, but not at C2745, Catalase, Shaker and Ldh, which remain significant outliers (Table 5, Figure 3a). Similarly, LD in the simulated populations with C5197 overdominance, recover a closely similar estimate to that observed between C5197 and Masc in 2006, but uniformly low LD everywhere else (Figure 3c, in parentheses).

Discussion

This experiment was originally motivated to test the prediction that the very high male-specific sterility load in *B. anynana* (Saccheri *et al*, 2005) is Z-linked, and maintained through some form of intralocus sexual conflict (Rice, 1984). Lack of any consistent association between homozygosity at a sample of loci spanning the entire chromosome and male sterility does not support this hypothesis. The implication, therefore, is that the male-specific sterility load must be due to autosomal loci. In contrast, the Z chromosome does harbour a large proportion of the male-specific viability load, in the form of recessive lethals, with implications for the retention of linked diversity.

The complete, or near complete, absence of adult males homozygous within the region of the Z chromosome spanning C5197-6PGD in the Liverpool stock (2006) backcrosses indicates a high frequency of recessive lethals in this laboratory population. As this population had been in captivity for some 90 generations this could be a consequence of genetic drift (the presence of at least four different lethal core haplotypes makes this unlikely). However, the deficit of homozygotes in the Leiden (1993) outbred base population, which will have been subject to minimal genetic drift since being established from the wild, and in the inbred (bottlenecked) lines derived from it, strongly suggests that the high frequency of recessive lethals in this region is a feature of the natural population. Our experiment only measured the adult stage, and therefore does not allow detection of when, during development, the homozygous lethal effect occurs. However, as larval and pupal mortality is very low under benign rearing conditions (Brakefield *et al*, 2001), sometime during embryo development prior to hatching is most likely.

The presence of a few apparent homozygotes within the core region, particularly in the 1993 stock, is likely to reflect the fact that intronic marker microhaplotypes are only loosely linked to extended haplotypes in population samples, so we predict that many of the homozygotes at C5197 and Masc are cryptic haplotype heterozygotes at the functionally lethal region. This interpretation is supported by the complete or near complete absence of C5197-6PGD homozygotes in the backcross and bottleneck material, where parental haplotypes could be traced. In addition to C5197, Masc and 6PGD, the lethal region includes Z-lethal(3)neo18, C5904, SREBP, and 1-Cys peroxiredoxin. The name Z-lethal(3)neo18 is derived from the *D. melanogaster* ortholog, of which *B. anynana* has an autosomal copy and a Z-linked copy. Despite its suggestive name it remains unclear whether this gene is involved in the observed lethal effect since its function in *D. melanogaster* is not documented. Based on the *B. anynana* reference genome (Nowell *et al*, 2017), several other genes exist in the lethal interval, but none are obvious candidates for lethality either.

As demonstrated by our simulation model, heterozygote advantage could account for the observed retention of linked diversity and LD within the lethal-containing region (C5197 and Masc) through *ca.* 70 generations of genetic drift, despite the complete loss of one-third and one-half of haplotypes, respectively. We cannot at this stage rule out associative overdominance, as opposed to single-locus overdominance (Charlesworth and Willis, 2009), which would require different but closely linked lethal-carrying loci in repulsion phase, rather than a single locus with many recessive lethal alleles. A model involving sexually antagonistic pleiotropy is less plausible, given that most, possibly all, alleles appear to be unconditionally deleterious to males, and we did not observe any obvious fitness benefits to females.

The retention of haplotype diversity in other regions of the Z chromosome (C2745, Shaker, Ldh, and to lesser extent Kettin and Catalase) is intriguing. There were no obvious deviations from the expected segregation patterns in the backcross families and bottleneck lines, and recombination distances to the male-lethal region are large. Moreover, the family used to create the linkage map, and the backcross families do not suggest unusual recombination or the presence of inversion polymorphisms. However, the very similar pattern in LD between the 1993 and 2006 populations across the lower third of the chromosome (C5197-Ldh) does imply the interaction of reduced recombination and/or epistatic selection (Phillips, 2008).

In the specific case of C2745, the excess haplotype diversity is coupled with a significant excess of homozygotes in the ancestral male sample. Assortative mating, with respect to this or nearby loci, could conceivably produce both of these patterns. The restrictive conditions required for stable polymorphism under a one trait, two-locus model (Hedrick, 2017), suggest

a more complex architecture, perhaps involving sexual antagonism (Arnqvist, 2011) and linkage between (male) attractiveness and (female) preference (Iyengar *et al*, 2002). In contrast to the C5197-Ldh region, LD in the upper third of the chromosome declined, albeit more slowly than expected. This may relate to the artificially high population density environment of the caged population, and the associated relaxation of the conditions promoting mate choice behaviour in nature.

One of the marker loci (Tpi) stands out as having lost more haplotype diversity than expected. The highly skewed pattern of inheritance, in both sexes, of the paternal alleles in one of the backcross families, together with the more than doubling in frequency of this specific allele between 1993 and 2006, is suggestive of meiotic drive (Lindholm *et al*, 2016). The loss of autosomal suppressor alleles, originally present in the ancestral population, due to genetic drift may explain its spread in captivity. This selective sweep is likely to have contributed to the 1993-2006 increase in LD between Tpi and the neighbouring Catalase, whose most common haplotype (H6) is fully associated with the putative driving haplotype (Tpi-H1; Figure S2). Interestingly, there is also a suggestion that this haplotype causes male sterility in the homozygous state (all the sons in family 13BF₁ were both homozygous for Tpi and sterile).

Our finding of significant retention of heterozygosity on the Z contrasts the near neutral behaviour over 20 generations of the X in *D. melanogaster* lines, attributed to low levels of X-linked recessive load (Schou *et al*, 2017). Greater efficacy of purifying selection is expected to apply to sex-linked genes whose expression is biased towards the heterogametic sex (XY males, ZW females). Conversely, recessive or partially recessive deleterious mutations at sex-linked genes whose expression is biased towards the homogametic sex should experience less efficient selection. The contrasting pattern in non-synonymous to synonymous nucleotide diversity (π_n/π_s) expected from this evolutionary dynamic has been confirmed between categories of sex-biased Z-linked genes in two butterfly species (Rousselle *et al*, 2016). In addition to gamety, differences in sex chromosome-linked load likely reflect the outcome of many other factors, including species demography, life history, phylogeny, and intra/inter-specific coevolution.

This exploratory study has raised more questions than it has answered. Future work should be directed at resolving the identity of the male lethals in the C5197-6PGD region, as well as the underlying causes of the joint retention of heterozygosity and LD in the lower third of the Z chromosome, and the homozygote excess at the top end, around C2745. Further experiments are required to confirm the behaviour and identity of the putative meiotic driver in the vicinity of Tpi. Finally, the search for the loci causing inbred male sterility needs to be extended to the autosomes. The overall impression is of a complex set of balancing selection mechanisms, whose net effect is the maintenance of an elevated genetic load resistant to purifying selection, making this species particularly susceptible to inbreeding depression.

Acknowledgements

This work was made possible by the enthusiasm of Paul Brakefield, establishing the original lab colony of *Bicyclus anynana* in 1988 and promoting its use as a model for evolutionary biology. Dave Pye assisted with DNA extractions and sequencing. Comments by three reviewers led to improvements of a previous version of the paper. This work was principally supported by NERC grants (NER/M/S/2001/00040 and NE/N015711/1) and Leverhulme Research Fellowship (RF-2016-654) to IJS.

References

- Arnqvist G (2011). Assortative mating by fitness and sexually antagonistic genetic variation. *Evolution* **65**: 2111-2116.
- Beldade P, Rudd S, Gruber JD, Long AD (2006). A wing expressed sequence tag resource for *Bicyclus anynana* butterflies, an evo-devo model. *BMC Genomics* **7**: 130.
- Beldade P, Saenko SV, Pul N, Long AD (2009). A gene-based linkage map for *Bicyclus anynana* butterflies allows for a comprehensive analysis of synteny with the lepidopteran reference genome. *PLoS Genet* **5**: e1000366.
- Bijlsma R, Bundgaard J, Boerema AC (2000). Does inbreeding affect the extinction risk of small populations?: predictions from *Drosophila*. *J Evol Biol* **13**: 502-514.
- Bonduriansky R, Chenoweth SF (2009). Intralocus sexual conflict. *Trends Ecol Evol* **24**: 280-288.
- Brakefield PM, El Filali E, Van der Laan R, Breuker CJ, Saccheri IJ, Zwaan B (2001). Effective population size, reproductive success and sperm precedence in the butterfly, *Bicyclus anynana*, in captivity. *J Evol Biol* **14**: 148-156.
- Canty A, Ripley B (2019). boot: bootstrap R (S-Plus) functions.
- Charlesworth B (2015). Causes of natural variation in fitness: Evidence from studies of *Drosophila* populations. *Proc Natl Acad Sci USA* **112**: 1662-1669.
- Charlesworth B, Charlesworth D (1999). The genetic basis of inbreeding depression. *Genet Res* **74**: 329-340.
- Charlesworth B, Hughes KA (2000). The maintenance of genetic variation in life-history traits. In: Singh RS and Krimbas CB (eds) *Evolutionary Genetics: from Molecules to Morphology*. Cambridge University Press. Vol. 1, pp 369-392.
- Charlesworth B, Morgan MT, Charlesworth D (1993). The effect of deleterious mutations on neutral molecular variation. *Genetics* **134**: 1289-1303.
- Charlesworth D, Willis JH (2009). The genetics of inbreeding depression. *Nat Rev Genet* **10**: 783.
- Connallon T, Clark AG (2014). Evolutionary inevitability of sexual antagonism. *Proc R Soc B* **281**: 20132123.
- Crow JF (1987). Muller, Dobzhansky, and overdominance. *J Hist Biol* **20**: 351-380.
- Davison AC, Hinkley DV (1997). *Bootstrap Methods and their Applications*. Cambridge University Press: Cambridge.
- Dobzhansky T, Spassky B, Tidwell T (1963). Genetics of natural populations. XXXII. Inbreeding and the mutational and balanced genetic loads in natural populations of *Drosophila pseudoobscura*. *Genetics* **48**: 361-373.
- Ebel ER, Phillips PC (2016). Intrinsic differences between males and females determine sex-specific consequences of inbreeding. *BMC Evol Biol* **16**: 36.
- Ellegren H (2011). Emergence of male-biased genes on the chicken Z-chromosome: Sex-chromosome contrasts between male and female heterogametic systems. *Genome Res* **21**: 2082-2086.
- Enders LS, Nunney L (2010). Sex-specific effects of inbreeding in wild-caught *Drosophila melanogaster* under benign and stressful conditions. *J Evol Biol* **23**: 2309-2323.
- Fry JD (2010). The genomic location of sexually antagonistic variation: some cautionary comments. *Evolution* **64**: 10.1111/j.1558-5646.2009.00898.x.
- García N, López-Fanjul C, García-Dorado A (1994). The genetics of viability in *Drosophila melanogaster*: effects of inbreeding and artificial selection. *Evolution* **48**: 1277-1285.
- Gibson JR, Chippindale AK, Rice WR (2002). The X chromosome is a hot spot for sexually antagonistic fitness variation. *Proc R Soc B* **269**: 499-505.
- Goudet J, Jombart T (2015). hierfstat: estimation and tests of hierarchical F-statistics.

- Hedrick PW (2017). Assortative mating and linkage disequilibrium. *G3* **7**: 55-62.
- Huylmans AK, Macon A, Vicoso B (2017). Global dosage compensation is ubiquitous in Lepidoptera, but counteracted by the masculinization of the Z chromosome. *Mol Biol Evol* **34**: 2637-2649.
- Innocenti P, Morrow EH (2010). The sexually antagonistic genes of *Drosophila melanogaster*. *PLoS Biol* **8**: e1000335.
- Iyengar VK, Reeve HK, Eisner T (2002). Paternal inheritance of a female moth's mating preference. *Nature* **419**: 830-832.
- Janicke T, Vellnow N, Sarda V, David P (2013). Sex-specific inbreeding depression depends on the strength of male–male competition. *Evolution* **67**: 2861-2875.
- Keller LF (1998). Inbreeding and its fitness effects in an insular population of song sparrows (*Melospiza melodia*). *Evolution* **52**: 240-250.
- Kumasaka N, Okada Y, Takahashi A, Kubo M, Nakamura Y, Kamatani N. (2011). *12th International Congress of Human Genetics*: Montreal, Canada.
- Liberatore KL, Jiang K, Zamir D, Lippman ZB (2013). Heterosis: the case for single-gene overdominance. In: Chen ZJ and Birchler JA (eds) *Polypliod and Hybrid Genomics*. John Wiley & Sons, Inc, pp 137-152.
- Lindholm AK, Dyer KA, Firman RC, Fishman L, Forstmeier W, Holman L *et al* (2016). The ecology and evolutionary dynamics of meiotic drive. *Trends Ecol Evol* **31**: 315-326.
- Lucotte EA, Laurent R, Heyer E, Ségurel L, Toupance B (2016). Detection of allelic frequency differences between the sexes in humans: a signature of sexually antagonistic selection. *Genome Biol Evol* **8**: 1489-1500.
- Mallet MA, Chippindale AK (2011). Inbreeding reveals stronger net selection on *Drosophila melanogaster* males: implications for mutation load and the fitness of sexual females. *Heredity* **106**: 994-1002.
- Mank JE, Hosken DJ, Wedell N (2014). Conflict on the sex chromosomes: cause, effect, and complexity. *Cold Spring Harbor Persp Biol* **6**: a017715.
- Muller HJ (1916). The mechanism of crossing-over. II. IV. The manner of occurrence of crossing-over. *Am Nat* **50**: 284-305.
- Nei M (1987). *Molecular Evolutionary Genetics*. Columbia University Press: New York.
- Nowell RW, Elsworth B, Oostra V, Zwaan BJ, Wheat CW, Saastamoinen M *et al* (2017). A high-coverage draft genome of the mycalesine butterfly *Bicyclus anynana*. *Gigascience* **6**: 1-7.
- Ohta T (1971). Associative overdominance caused by linked detrimental mutations. *Genet Res* **18**: 277-286.
- Okada Y (2018). eLD: entropy-based linkage disequilibrium index between multiallelic sites. *Human Genome Variation* **5**: 29.
- Peng B, Kimmel M (2005). simuPOP: a forward-time population genetics simulation environment. *Bioinformatics* **21**: 3686-3687.
- Phillips PC (2008). Epistasis--the essential role of gene interactions in the structure and evolution of genetic systems. *Nat Rev Genet* **9**: 855-867.
- R Core Team (2017). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Rice WR (1984). Sex chromosomes and the evolution of sexual dimorphism. *Evolution* **38**: 735-742.
- Rose MR (1982). Antagonistic pleiotropy, dominance, and genetic variation. *Heredity* **48**: 63-78.
- Rousselle M, Faivre N, Ballenghien M, Galtier N, Nabholz B (2016). Hemizygoty enhances purifying selection: lack of fast-Z evolution in two satyrine butterflies. *Genome Biol Evol* **8**: 3108-3119.

- Ruzicka F, Hill MS, Pennell TM, Flis I, Ingleby FC, Mott R *et al* (2019). Genome-wide sexually antagonistic variants reveal long-standing constraints on sexual dimorphism in fruit flies. *PLoS Biol* **17**: e3000244.
- Saccheri I, Kuussaari M, Kankare M, Vikman P, Fortelius W, Hanski I (1998). Inbreeding and extinction in a butterfly metapopulation. *Nature* **392**: 491-494.
- Saccheri IJ, Brakefield PM, Nichols RA (1996). Severe inbreeding depression and rapid fitness rebound in the butterfly *Bicyclus anynana* (Satyridae). *Evolution* **50**: 2000-2013.
- Saccheri IJ, Bruford MW (1993). DNA fingerprinting in a butterfly, *Bicyclus anynana* (Satyridae). *J Hered* **84**: 195-200.
- Saccheri IJ, Lloyd H, Helyar S, Brakefield PM (2005). Inbreeding uncovers fundamental differences in the genetic load affecting male and female fertility in a butterfly. *Proc R Soc B* **272**: 39-46.
- Saccheri IJ, Wilson IJ, Nichols RA, Bruford MW, Brakefield PM (1999). Inbreeding of bottlenecked butterfly populations: estimation using the likelihood of changes in marker allele frequencies. *Genetics* **151**: 1053-1063.
- Schou MF, Loeschcke V, Bechsgaard J, Schlötterer C, Kristensen TN (2017). Unexpected high genetic diversity in small populations suggests maintenance by associative overdominance. *Mol Ecol* **26**: 6510-6523.
- Spencer HG, Priest NK (2016). The evolution of sex-specific dominance in response to sexually antagonistic selection. *Am Nat* **187**: 658-666.
- van't Hof AE, Marec F, Saccheri IJ, Brakefield PM, Zwaan BJ (2008). Cytogenetic characterization and AFLP-based genetic linkage mapping for the butterfly *Bicyclus anynana*, covering all 28 karyotyped chromosomes. *PLoS ONE* **3**: e3882.
- van't Hof AE, Nguyen P, Dalíková M, Edmonds N, Marec F, Saccheri IJ (2013). Linkage map of the peppered moth, *Biston betularia* (Lepidoptera, Geometridae): a model of industrial melanism. *Heredity* **110**: 283-293.
- Wang J, Hill WG, Charlesworth D, Charlesworth B (1999). Dynamics of inbreeding depression due to deleterious mutations in small populations: mutation parameters and inbreeding rate. *Genet Res* **74**: 165-178.
- Wang J, Xia Q, He X, Dai M, Ruan J, Chen J *et al* (2005). SilkDB: a knowledgebase for silkworm biology and genomics. *Nucleic Acids Res* **33**: D399-D402.
- Zajitschek F, Connallon T (2018). Antagonistic pleiotropy in species with separate sexes, and the maintenance of genetic variation in life-history traits and fitness. *Evolution* **72**: 1306-1316.
- Zhao L, Charlesworth B (2016). Resolving the conflict between associative overdominance and background selection. *Genetics* **203**: 1315-1334.

Table 1. The ratio of homozygous to heterozygous sons, or of alternate paternal alleles in daughters, at a series of Z-linked loci (ordered according to their relative positions on the chromosome) in a sample of adult offspring produced by five daughter-father backcrosses. Grey shading highlights the region in which homozygous sons are almost entirely missing. The significance of deviations from the null expectation of 1:1 (homozygotes vs heterozygotes or allele 1 vs allele 2) is indicated by asterisks (Fisher's exact test).

loci	sons: homozygotes : heterozygotes					daughters: allele 1 : allele 2				
	2BF1 (n=24)	8BF1 (n=33)	9BF1 (n=29)	11BF1 (n=24)	13BF1 (n=20)	2BF1 (n=16)	8BF1 (n=16)	9BF1 (n=16)	11BF1 (n=16)	13BF1 (n=14)
C2745	9:15	17:16	11:18	6:18	10:10	3:13	8:8	8:8	8:8	8:6
Kettin	9:15	-	-	-	-	4:12	-	-	-	-
Tpi	-	13:20	6:23 *	-	20:0 ***	-	6:10	6:10	-	14:0 ***
Cyp303a1	3:21 *	-	-	1:23 ***	-	4:12	-	-	11:5	-
C5197	-	-	0:29 ***	0:24 ***	0:20 ***	-	-	5:11	9:7	4:10
Masc	0:24 ***	0:33 ***	0:29 ***	0:24 ***	2:18 *	6:10	7:9	5:11	10:6	3:11
6PGD	1:23 ***	0:33 ***	-	-	-	8:8	7:9	-	-	-
Ldh	4:20 *	2:31 ***	2:27 ***	18:6	3:17 *	7:9	6:10	6:10	5:11	4:10
Henna	-	2:31 ***	1:28 ***	-	-	-	6:10	5:11	-	-

Table 2. The proportion of completely unfertilized (sterile) egg batches, and the mean proportion of hatching eggs, produced by samples of outcrossed (F_1 hybrid) females mated to the sons of 14 daughter-father backcrosses (n_1 is the number of mated pairs from which the estimates were derived; n_2 is the mean number of eggs collected per pair).

Line	n_1	n_2	Sterility (se)	Hatching (se)
1BF1	29	48	0.79 (0.08)	0.12 (0.05)
2BF1	23	45	0.39 (0.10)	0.31 (0.07)
3BF1	45	65	0.22 (0.06)	0.51 (0.06)
4BF1	21	68	0.09 (0.06)	0.62 (0.06)
5BF1	28	47	0.21 (0.08)	0.53 (0.07)
6BF1	21	38	0.71 (0.10)	0.17 (0.06)
7BF1	21	48	1.00 (0.00)	0.00 (0.00)
8BF1	33	60	0.64 (0.08)	0.16 (0.05)
9BF1	29	48	0.69 (0.09)	0.13 (0.05)
10BF1	20	44	0.65 (0.11)	0.33 (0.07)
11BF1	24	38	0.54 (0.10)	0.21 (0.07)
12BF1	13	39	0.77 (0.12)	0.07 (0.04)
13BF1	24	39	1.00 (0.00)	0.00 (0.00)
14BF1	11	28	0.91 (0.09)	0.09 (0.05)

Table 3. Comparison of the observed number homozygous males in a sample (n) taken from the ancestral (Leiden-1993) population vs the number expected under HWE and using allele frequencies estimated from an equivalent female sample, at each of eight Z-linked loci.

loci	n	Obs	Exp	χ^2	p-value
C2745	112	42	23.13	19.39	<0.001
Kettin	109	45	36	3.35	ns
Tpi	112	15	18.36	0.73	ns
Catalase	108	39	40	0.01	ns
C5197	112	14	26.78	8.02	<0.005
Masc	111	17	10.99	3.64	ns
Shaker	108	20	17.64	0.37	ns
Ldh	111	13	14.45	0.16	ns

Table 4. The ratios of homozygous to heterozygous males, or of alternate alleles in females, at a series of Z-linked loci (ordered according to their relative positions on the chromosome) in a sample of F_3 adults descended from six laboratory populations (lines) each established from a single pair of founders collected from the outbred Leiden-1993 population ca. 20 generations into captivity. Depending on whether the founding pair shared an allele, loci in these lines may segregate for two or three alleles. Grey shading highlights the region in which homozygous males are almost entirely missing. The significance of deviations from the null expectation is indicated by asterisks (Fisher's exact test). (Line 1.3 Tpi $n=12$ for both sexes).

Line: loci	males: homozygotes : heterozygotes						females: allele 1 : allele 2 : allele 3					
	1.1 (n=9)	1.2 (n=9)	1.3 (n=16)	1.4 (n=9)	1.5 (n=9)	1.6 (n=9)	1.1 (n=9)	1.2 (n=9)	1.3 (n=16)	1.4 (n=9)	1.5 (n=9)	1.6 (n=9)
C2745	1:8	5:4	13:3	8:1	8:1	2:7	5:2:3	5:4	10:5:1	4:3:2	5:4	3:3:3
Tpi	0:9	-	2:10	3:6	1:8	4:5	6:2:1	-	6:3:3	6:3	5:4	6:3
C5197	0:9	1:8	0:16	0:9	0:9	3:6	5:4	6:3	8:5:3	6:3	5:4	5:4
Masc	0:9	0:9	0:16	0:9	0:9	0:9	5:4	5:4	8:5:3	6:3	5:4	5:4
Ldh	6:3	8:1	3:13	2:7	6:3	1:8	7:2	8:1	4:9:3	6:3:1	9:0	4:4:1

Table 5. Comparison of haplotype number (H_n) and diversity (H_d) at eight Z-linked loci estimated from female samples of size n taken from the base (Leiden) laboratory population of *Bicyclus anynana* in 1993 and a descendant (Liverpool) population in 2006. P is the probability of jointly obtaining these H_n and H_d in 2006 under a neutral scenario (fitness of C5197 homozygotes = 1), and the overdominant one (fitness of C5197 homozygotes = 0).

	Marker loci															
	C2745		Kettin		Tpi		Catalase		C5197		Masc		Shaker		Ldh	
n	1993	2006	1993	2006	1993	2006	1993	2006	1993	2006	1993	2006	1993	2006	1993	2006
H_n	111	48	112	46	112	48	112	47	112	48	109	45	112	46	111	48
H_d	0.80	0.82	0.68	0.59	0.84	0.43	0.64	0.65	0.77	0.83	0.91	0.82	0.84	0.77	0.87	0.78
H_d se	0.03	0.02	0.05	0.11	0.02	0.06	0.03	0.06	0.04	0.02	0.01	0.03	0.02	0.03	0.01	0.03
$P(W_{C5197}=1)$	0.001		0.332		0.809		0.013		<0.001		<0.001		0.008		0.013	
$P(W_{C5197}=0)$	0.001		0.351		0.778		0.019		0.168		0.067		0.007		0.011	

Figure 1. Population samples and backcross design. (a) Timeline of the *Bicyclus anynana* Leiden/Liverpool lab population, showing the samples genotyped in this study, their relationship to each other, and the order in which they were analysed (1 to 5). (b) Design of daughter-father backcross, highlighting the identify (by descent) of Z chromosome segments (denoted by z_i , where i refers to the identity of the paternal Z in generation P). The position of the recombination breakpoints, indicated by change in colour, is arbitrary, as is which of the daughters to backcross (recombination does not occur in female Lepidoptera). Z_F and Z_M refers to whether the Z copy came from the father (F) or mother (M). This design is expected to generate F_2 brothers that differ with respect to which portion of the Z is either homozygous (hom) or heterozygous (het). Autosomes (not shown) are expected to experience the same degree of inbreeding regardless of sex (probability of identity by descent ~ 0.25 , averaged across all offspring, assuming zero coancestry between P generation parents). To assess the fertility of F_2 males, they were mated to unrelated outbred females. The F_2 generation is referred to as BF_1 (backcross F_1).




Figure 2. Z-chromosome markers and haplotype frequency changes. (a) Linkage map of the *Bicyclus anynana* Z chromosome, highlighting in blue all the marker loci used in this study (recombination distances in centimorgans). A subset of these were used to genotype population samples. (b) The distribution of micro-haplotype frequencies at eight Z-linked loci in the base (Leiden) laboratory population of *Bicyclus anynana* in 1993 (black bars) and a descendant (Liverpool) population in 2006 (hatched bars). Haplotype numbers were assigned within each locus according to their frequency in the 1993 sample and do not indicate direct correspondence between loci. Sample sizes are given in Table 5.




Figure 3. Observed and expected patterns in haplotype diversity and linkage disequilibrium at eight Z-linked loci in *Bicyclus anynana*. (a) Distributions of the proportional change between 1993 and 2006 (~70 generations) in haplotype diversity (H_d) at corresponding loci shown on a simplified chromosome (b). Bootstrap distribution of the observed data (gold); simulated distribution with overdominance at C5197 (green); neutral expectation (dotted line). (c) Entropy-based linkage disequilibrium (eLD) among all pairs of loci. Values in the upper half-matrix are for the Leiden population in 1993, and in the lower half-matrix for the descendant Liverpool population in 2006. In parentheses, the mean and standard deviation of eLD in a sample of 100 simulations assuming overdominance at C5197. Colour gradient is proportional to the magnitude of eLD.

