

The role of climate-fuel feedbacks on Holocene biomass burning in upper-montane Carpathian forests

Vachel A. Carter^a, Přemysl Bobek^b, Alice Moravcová^a, Anna Šolcová^b, Richard C. Chiverrell^c, Jennifer L. Clear^{d,e}, Walter Finsinger^f, Angelica Feurdean^{g,h,i}, Ioan Tanțăuⁱ, Enikő Magyari^{j,k,l}, Thomas Brussel^m, Petr Kuneš^a

A Department of Botany, Faculty of Science, Charles University, Prague, Czech Republic

B Institute of Botany of the Czech Academy of Sciences, Průhonice, Czech Republic

C Department of Geography and Planning, University of Liverpool, Liverpool, United Kingdom

D Department of Geography and Environmental Science, Liverpool Hope University, Liverpool, United Kingdom

E Department of Forest Ecology, Czech University of Life Sciences, Prague, Czech Republic

F ISEM, University Montpellier, CNRS, EPHE, IRD, Montpellier, France

G Department of Physical Geography, Goethe University, Altenhöferallee 1, 60438 Frankfurt am Main, Germany

H Senckenberg Biodiversity and Climate Research Centre (BiK-F), Senckenberganlage, 25, 60325, Frankfurt am Main, Germany

I Department of Geology, Babeş-Bolyai University, 400084 Cluj-Napoca, Romania

J Department of Environmental and Landscape Geography, Eötvös Loránd University, Budapest, Hungary

K MTA-MTM-ELTE Research Group for Paleontology, Budapest, Hungary

L GINOP Sustainable Ecosystems Group, MTA Centre for Ecological Research, Tihany, Hungary

M Department of Geography, University of Utah, Salt Lake City, UT, USA

Citation: Vachel A. Carter, Přemysl Bobek, Alice Moravcová, Anna Šolcová, Richard C. Chiverrell, Jennifer L. Clear, Walter Finsinger, Angelica Feurdean, Ioan Tanțău, Enikő Magyari, Thomas Brussel, Petr Kuneš, The role of climate-fuel feedbacks on Holocene biomass burning in upper-montane Carpathian forests, *Global and Planetary Change*, Volume 193, 2020, <https://doi.org/10.1016/j.gloplacha.2020.103264>

Highlights

- First high-resolution Holocene fire record from the High Tatra Mountains.
- Forest composition and density influence burning in upper-montane forests.
- Burning primarily occurred with intermediate *Picea abies* cover.
- Future positive climate-fuel feedbacks may exist with increasing *Pinus* cover.

Abstract

Over the past few decades, mean summer temperatures within the Carpathian Mountains have increased by as much as 2 °C leading to a projected increased forest fire risk. Currently, there are no paleofire records from the Western Carpathians that provide the long-term range of natural variability to contextualise the response of upper-montane forests to future environmental change and disturbance regimes. We present the first high-resolution Holocene fire history record from the upper-montane ecotone from the High Tatra Mountains, Slovakia, as well as provide a regional synthesis of pan-Carpathian drivers of biomass burning in upper-montane forests. Our results illustrate that forest composition and density both greatly influence biomass burning, creating two different climate-fuel feedbacks. First, warmer conditions in the early Holocene, coupled with generally higher abundances of *Pinus* sp., either *P. cembra* and/or *P. mugo/sylvestris*, created a positive climate-fuel relationship that resulted in higher amounts of biomass burning. Second, cooler and wetter late Holocene conditions led to denser *Picea abies* upper-montane forests, creating a negative climate-fuel feedback that reduced biomass burning in upper-montane forests across the Carpathians. Given that warmer and drier conditions are expected across the entire Carpathian region in the future, our results illustrate how future climate change could potentially create a positive climate-fuel relationship within upper-montane forests dominated by *Picea abies* and *Pinus cembra* and/or *P. mugo/sylvestris*.

Keywords: Fire, Upper-montane, Carpathians, Holocene, Sedimentary charcoal, Pollen, Macrofossils, *Picea abies*, Forest composition, Forest density

1. Introduction

Mountain forests are important ecosystems providing an array of ecosystem goods and services, with roughly half of the global human population depending upon these services (Körner and Ohsawa, 2005). Mountain forests are also key hotspots for biodiversity, with higher levels of endemic richness as a result of high environmental heterogeneity and geographic isolation (Noroozi et al., 2018). In Europe, forests cover 41% of the total mountain area with the majority of forest cover occurring in the Alps, Carpathians and Pyrenees Mountains (Price et al., 2011). Unfortunately, mountain forests are among the most sensitive ecosystems to climate change, with high mountain environments experiencing more rapid changes in temperature than low-elevation environments (Pepin et al., 2015). Since 2001, all major European alpine systems have witnessed an increasing trend in warm-adapted species (Gottfried et al., 2012), illustrating their sensitivity to warmer temperatures. Given that climate influences significantly the structure, composition and function of high mountain ecosystems (Prentice et al., 1992), climate-induced changes are likely going to be more pronounced near the upper treeline (Harsch et al., 2009).

Norway spruce (*Picea abies*) is one of the most important economic trees in Europe, dominating the boreal forests in northern Europe and upper-montane/subalpine areas of the Alps and Carpathian Mountains (hereafter referred to as Carpathians) (Caudullo et al., 2016). Norway spruce is susceptible to both heat and

drought due to its shallow root systems and may be vulnerable to the increasing temperatures at high-elevations. In fact, *Picea abies* is currently listed as one of the most vulnerable tree species in the Carpathians due to increasing summer temperatures (Lindner et al., 2008), with both montane (1100–1500 m a.s.l.) and subalpine (>1500 m a.s.l.) *Picea abies* forests projected to decline in extent with climate change (Werners et al., 2014). Mean summer temperatures in the Carpathians have increased by as much as 2 °C over the past few decades (Alberton et al., 2017), twice the global mean (Auer et al., 2007), which raises the prospect of greater sensitivity to climate change than montane forests elsewhere in Europe (UNEP, 2007). This sensitivity to climate change will likely propagate through the 21st century, with mean summer temperatures projected to increase by a further 1–2 °C across the Carpathians by the end of the century (Anders et al., 2014) in conjunction with heat waves and droughts. Projected changes in precipitation across the Carpathians are more varied but suggest greater seasonality with generally drier summers (Anders et al., 2014). If conditions continue as projected, drier conditions could lead to either a collapse and/or dieback of *Picea abies*, compromising the provision of ecosystem services (Lévesque et al., 2013).

Many mountains forests including old growth *Picea abies* forests are declining and/or vanishing, especially within the Romanian arc of the Carpathians as a result of deforestation, poor management practices, and increased rates of forest disturbances linked to windthrow, bark beetle outbreak and fire events (Achard et al., 2009; Hlásny and Sitková, 2010; Feurdean et al., 2011; Knorn et al., 2012, Knorn et al., 2013; Temperli et al., 2013; Grindean et al., 2019). These factors illustrate the sensitivity of *Picea abies* forests to human intervention, disturbance regimes, and climate change. Given that there is a high likelihood of increased fire risk among central and southern European ecosystems by the end of the 21st century (Lung et al., 2013), and that air temperature is a key variable for establishment and growth of high mountain forests (Körner and Paulsen, 2004), the coupled threat of future climate change and increased likelihood of fires potentially threatens upper-montane *Picea abies* forests across the Carpathians. Future climate change and increased fire risk could affect their ability to provide ecosystem services (Albrich et al., 2018), including negative impacts on key transboundary biosphere reserves (UNESCO-WNBR, n.d). However, there is currently a limited understanding regarding the long-term importance of fire in high mountain forests, specifically upper-montane *Picea abies*-dominated forests. This information is critical for conservation and management as *Picea abies* is also highly susceptible to other forest disturbances such as windthrow and bark beetle outbreaks. What limited understanding we have about fire activity in upper-montane forests throughout the Carpathians is from research mostly from the Eastern and Southern Carpathians, which demonstrate the importance of long-term fire occurrence in these elevations (Geantă et al., 2014; Feurdean et al., 2012, Feurdean et al., 2017; Finsinger et al., 2018; Florescu et al., 2018). No long-term records of fire dynamics existed from the Western Carpathians until recent research, which is from the lower montane forest zone dominated by beech (*Fagus sylvatica*), silver fir (*Abies alba*), and an admixture of Norway spruce (Kołaczek et al., 2020).

Here, we present a new c.9500-year high-resolution paleoecological record from Popradské pleso located in the upper-montane forest zone in the High Tatra Mountains, Slovakia, Western Carpathians. We address the existing knowledge gap concerning the long-term history of fire in upper-montane forests in the Western Carpathians by comparing the new record with other high-resolution fire records from the upper-montane forest zone spanning the entire Carpathian region. Our objectives are to; 1) quantify the long-term variability in wildfire (biomass burning) regime and assess spatial differences across the wider Carpathian upper-montane forested ecosystem; 2) evaluate the respective roles of climate and humans as drivers of past biomass burning; 3) explore how forest composition and density modulates biomass burning in these upper-montane forest ecosystems; and 4) use the paleofire record to infer the potential impacts of future climate warming (1–2 °C). Our study provides critical data for understanding the role of fire

disturbance regimes in old growth spruce forests in the Carpathians, with utility for understanding the future dynamics of equivalent montane ecosystems elsewhere.

2. Study region

2.1. Popradské pleso in the Western Carpathians

Popradské pleso (pleso is Slovak for 'glacial lake') (49°09'13" N, 20°04'47" E, 1513 m a.s.l.) is located in the Mengusovská dolina valley, on the south slope of the central High Tatra Mountains (Fig. 1). The High Tatra Mountains are a west-to-east trending mountain range along the Slovakia-Poland border situated within the Western Carpathians. This mountain range was heavily glaciated during the Last Glacial Maximum with the Mengusovská dolina valley hosting the largest glacier on the south slope (Zasadni and Kłapyta, 2014). Popradské pleso has a large catchment to lake area ratio (67.5:1), and a maximum water depth of ~18 m with a surface area of ~6.3–6.9 ha (Schaffer and Stummer, 1933; Pacl, 1973). The lake catchment has a granite-granodiorite bedrock and is surrounded by the tallest mountain peaks in the entire High Tatra region including the tallest mountain peak, Rysy (2500 m a.s.l.). The lake has one inflowing stream, Ľadový potok, on the northeast side, and one outflowing stream, Krupá, on the south side.




Fig. 1. Map of Popradské pleso, High Tatra Mountains, Slovakia (this study), in relationship to other *Picea abies*-dominated sites discussed; Tăul Muced and Poiana Știol, Eastern Carpathians (Feurdean et al., 2017), and Lake Brazi, Southern Carpathians (Magyari et al., 2012).

Popradské pleso is located ~50 m downhill from the upper treeline in the upper-montane forest belt (Grodzińska et al., 2004), which is dominated by Norway spruce with an admixture of Swiss stone pine (*Pinus cembra*) and dwarf mountain pine (*Pinus mugo*). While beech (*Fagus sylvatica*) and Silver fir (*Abies alba*) are both common at these elevations in the Southern Carpathians, *F. sylvatica* is currently absent and

A. alba is very rare at these elevations in the High Tatras (Rybníčková and Rybníček, 2006). However, *F. sylvatica* is present at lower elevations near Štrbské Pleso (1358 m; see Kučera, 2012 and Flachbart, 2007).

Climate in the high Western Carpathian mountain zone is cold and moist, with the highest quantities of precipitation (2000 to 2400 mm yr⁻¹) in the Carpathians occurring in the High Tatra Mountains (UNEP, 2007). Mean annual temperature and precipitation from the closest meteorological station to Popradské pleso, Štrbské Pleso (Slovak Hydrometeorological Institute), are 3.7 °C and 1826 mm, respectively. The local west-to-east mountain orientation has resulted in a strong windward/leeward precipitation gradient with the north side receiving more precipitation than the south site (Konček, 1974). Mean annual temperatures range between 2 to -4 °C in the coniferous belt dominated by *Picea abies* forests (Hess, 1965, Hess, 1974). For the High Tatras, climate models suggest 2 °C warming alongside 5–8% decreases in summer precipitation, with ~10–15% increases in winter precipitation by the end of 21st century (Anders et al., 2014).

2.2. Eastern and Southern Carpathian study sites

Poiana Știol (47°35'14" N; 24°48'43" E, 1540 m a.s.l.; Feurdean et al., 2017) and Tăul Muced (47°34'26" N; 24°32'42" E, 1360 m a.s.l.; Feurdean et al., 2017) are both located in the Rodna National Park and Biosphere Reserve, Eastern Carpathians. Poiana Știol is a nutrient-poor fen located at the current treeline ecotone dominated by *Picea abies*, *Pinus mugo*, *P. cembra* and *Juniperus communis* ssp. *nana* (Tanțău et al., 2011; Feurdean et al., 2016). The surrounding forest is composed of ~80% *Picea abies* and ~20% *Abies alba* (Topographic Maps of Romania, 1986). Tăul Muced is an ombrotrophic bog within a closed conifer forest, with onsite woody vegetation consisting of *Picea abies* with *P. mugo*. The surrounding forest is comprised of ~40% *Picea abies* and *Abies alba* and ~10% *Fagus sylvatica* (Topographic Maps of Romania, 1986). The climate at Poiana Știol is slightly wetter than at Tăul Muced due to its higher altitude (1300 mm and 1200 mm, respectively). Mean annual temperature is 2–2.2 °C at both sites (Dragotă and Kucsisca, 2011). For the Southern Carpathians, climate models suggest 2 °C warming alongside 10–15% decreases in summer precipitation, with ~10–15% increases in winter precipitation by the end of 21st century (Anders et al., 2014).

Lake Brazi (45°23'47" N; 22°54'6" E, 1740 m a.s.l.; Magyari et al., 2012) is a small (~0.4 ha), shallow lake (maximum water depth 1.1 m), located in the Retezat National Park, Retezat Mountains, Southern Carpathians. The site is located about 110 m below the timberline in a mixed conifer forest dominated by *Picea abies* and *Pinus cembra*, with *P. mugo* being restricted to the floating *Sphagnum* bog found along the lakeshore. Annual precipitation ranges between 900 and 1800 mm, while the mean annual temperature is around 6 °C in the foothill zone and -2 °C at 2500 m a.s.l. in the Retezat Mountains (Bogdan, 2008). For the Southern Carpathians, climate models suggest 2 °C warming alongside a 10–15% decrease in summer precipitation and a ~2–5% decrease in winter precipitation by the end of 21st century (Anders et al., 2014).

3. Methods

3.1. Field work

In September 2015, a 28 cm (POP 15-GC1) and a 4 m-long core (POP 15-1, drives 1–4) were extruded using a floating platform from the centre of Popradské pleso (water depth, 11 m). The sediment water interface was collected using a short gravity corer (Boyle, 1995), while a Russian corer was used to collect the long-core sediments in 1.5 m overlapping sections. Sediments were stored under refrigeration until they were subsampled at high-resolution (contiguous 0.5 cm intervals) for dating and geochemical analyses.

3.2. Lead 210 and radiocarbon dating

A total of 12 Accelerator Mass Spectrometry (AMS) radiocarbon (^{14}C) ages and a radiometric ^{210}Pb series were obtained to construct an age-depth model for Popradské pleso (Table 1). Most ^{14}C ages were measured on terrestrial plant macrofossils, but in the section below 1371 cm below the water surface, ages were measured on bulk pollen concentrations because of the lack of macrofossils. Pollen extractions for bulk pollen concentrations were modified after Brown et al. (1989) using a dilute Schulze solution of 10% KClO_3 and 35% HNO_3 (Gray, 1965).

Depth (cm)	Core ID	^{14}C Age $\pm 1\sigma$	Assigned ^{210}Pb age (Year AD)	Assigned age (cal yr BP)	Material Dated	Lab ID Number
1100	POP 15GC1		2015 \pm 1	-65		Pb210_1
1101	POP 15GC1		2005 \pm 2	-55		Pb210_2
1102	POP 15GC1		1985 \pm 5	-35		Pb210_3
1103	POP 15GC1		1966 \pm 7	-16		Pb210_4
1104	POP 15GC1		1945 \pm 10	5		Pb210_5
1105	POP 15GC1		1924 \pm 13	26		Pb210_6
1106	POP 15GC1		1901 \pm 15	49		Pb210_7
1107	POP 15GC1		1879 \pm 18	71		Pb210_8
1159.5	POP 15-1-1	1665 \pm 35			<i>Picea abies</i> needle	Poz-81508
1200	POP 15-1-1	2470 \pm 50			<i>Pinus</i> sp. needle	Poz-100371
1229.5	POP 15-1-1	3240 \pm 30			<i>Picea abies</i> needles	Poz-80176
1264.5	POP 15-1-2	3900 \pm 50			<i>Picea abies</i> needle fragments, <i>Vaccinium</i> sp. seed	Poz-100372
1310	POP 15-1-2	5655 \pm 35			<i>Pinus</i> sp. needles	Poz-80177
1344.5	POP 15-1-2	7180 \pm 50			<i>Picea abies</i> twig	Poz-100373
1371	POP 15-1-2	8700 \pm 40			<i>Picea abies</i> twig	Poz-80178
1374.5	POP 15-1-3	12790 \pm 20			Pollen	UGAMS-28410
1432	POP 15-1-3	16270 \pm 45			Pollen	UGAMS-28996
1532	POP 15-1-4	18500 \pm 55			Pollen	UGAMS-28997
1634	POP 15-1-4	16740 \pm 20			Pollen	UGAMS-28411
1635	POP 15-1-4	8370 \pm 100			<i>Betula</i> sp. bud scales	Poz-81509

Table 1. Radiocarbon (^{14}C) and ^{210}Pb dates used to constrain the depth-age relation for the Popradské pleso sediment sequence. Depths are presented by depth in centimeters below the water surface.

Radiometric dating targeted the upper 8 cm of the gravity core (POP-15-GC1), analysing for ^{210}Pb , ^{226}Ra , ^{137}Cs and ^{241}Am by direct gamma assay in the Liverpool University Environmental Radioactivity Laboratory using Ortec HPGe GWL series well-type coaxial low background intrinsic germanium detectors (Appleby et al., 1986). ^{210}Pb was determined via its gamma emissions at 46.5 keV, and ^{226}Ra by the 295 keV and 352 keV-rays emitted by its daughter radionuclide ^{214}Pb following 3 weeks storage in sealed containers to allow radioactive equilibration. ^{137}Cs and ^{241}Am were measured by their emissions at 662 keV and 59.5 keV respectively. The absolute efficiencies of the detectors were determined using calibrated sources and sediment samples of known activity. Corrections were made for the effect of self-absorption of low energy-rays within the sample (Appleby et al., 1992).

Age-depth relationships were modeled in a Bayesian framework using 'BACON' (Blaauw and Christen, 2011), including both the ^{210}Pb and ^{14}C dates. All ^{14}C dates were calibrated using the IntCal13 dataset (Reimer et al., 2013).

3.3. Lithostratigraphy, magnetic susceptibility and geochemical analyses

Prior to subsampling, high resolution digital line-scan photographs of the Pop15–1 core were taken using the University of Liverpool Geotek Multi-sensor core logger (MSCL-XZ). Geochemical data were determined non-destructively on a wet sediment basis at 5 mm resolution using an Olympus Delta ED-XRF (4 W)

mounted to the Geotek. Cores were measured under Soil mode (40 kV, 40 kV (filtered) and 15 kV beam intensities) for heavier elements and MiningPlus mode (40 kV and 15 kV) for lighter elements. Each beam intensity ran for 20 s with total runs of 60 s (soil mode) and 40 s (mining plus). The Olympus XRF instrument is subject to daily consistency checks and regular testing using library of certified geochemical reference materials (Boyle et al., 2015). Element concentrations are expressed in ppm, but are subject to possible matrix effects induced by variable down-core water and organic content arising from measuring wet sediments. Magnetic susceptibility (XLF) was measured on a wet sediment basis at 5 mm intervals using a Bartington Instruments MS2E high resolution point sensor paired to the Bartington MS3 meter mounted on the Liverpool Geotek MSCL-XZ. The scan XLF data reveal changes between minerogenic (positive values) and organic or high-water content sediments (negative or diamagnetic values).

3.4. Plant macrofossils and pollen analysis

Pollen analysis was conducted contiguously at 1 cm resolution throughout the gravity core, and at 2.5 cm resolution throughout the long core from Popradské pleso. Pollen processing utilized standard acid-base rinse procedures (Faegri et al., 1989) using 0.5 cm³ of sediment per sample. A Lycopodium tablet was added to each sample prior-to pollen processing, and was used as an exotic tracer to calculate both pollen concentration (grains cm³) and pollen accumulation rates (pollen influx; grains cm⁻² yr⁻¹) (Stockmarr, 1972). A minimum of 500 terrestrial pollen grains were identified at a magnification of 400× with the aid of reference material and identification keys (Beug, 2004; Punt, 1976–1996). Pollen counts were converted into pollen percentages based on the abundance of each pollen type relative to the sum of all identified terrestrial pollen. The pollen profile was divided into pollen assemblage zones based on optimal splitting by sums-of-squares using the broken-stick model using the software, Psimpoll (Birks and Gordon, 1985; Bennett, 1996). Legacy pollen data from the wider Carpathian region were available from Poiana Ştiol, Tăul Muced and Lake Brazi (see Magyari et al., 2012; Tanţău et al., 2011; Grindean et al., 2019). To access changes in forest density, we calculated an index using the ratio of *Picea abies* pollen (a) to *Pinus* pollen (b) following Higuera et al. (2014).

Plant macrofossil analysis was conducted every 2.5 cm from Popradské pleso using standard procedures described by e.g. Birks (2007). Sediment volume was measured by water displacement in a measuring cylinder (1–5 ml), and then gently sieved through a 100 µm sieve. Plant macrofossils were analysed using a stereomicroscope at a magnification of 15–50×, and were identified with the aid of identification keys (Cappers et al., 2006; Bojňanský and Fargašová, 2007; Katz et al., 1977; Tomlinson, 1985) and the herbarium collection at Charles University.

3.5. Charcoal analysis, fire-event detection, and change point analysis

Macroscopic charcoal was analysed at 1 cm resolution in both the gravity core and long core from Popradské pleso. Sediment sub-samples of 1 ml were left for 12 h in 10 ml, 5% solution of potassium hydroxide (KOH), to disaggregate organic matter. In order to bleach non-charred material, 5 ml of sodium hypochlorite (10% NaOCl) was added to the samples. The required length of bleaching process was adjusted according to the results of test specimens taken from the same sediment core. Samples were screened using a 125 µm sieve and washed into petri dishes. The majority of the non-charred organic particles were manually removed using a stereomicroscope before capturing images of charcoal fragments using a custom-built scanning device (www.microspock.cz). Images were analysed using ImageJ to count and measure the area of charcoal particles (Schneider et al., 2012). Macroscopic charcoal counts were transformed to concentrations (particles/area cm⁻³) and accumulation rates (CHARC (the number of particles cm⁻² yr⁻¹)).

The site-specific fire history was determined using the CharAnalysis software (Higuera et al., 2009). First, the CHARC record was interpolated to a constant temporal resolution of 30 years, which is based on the median sediment accumulation rate. Then, the charcoal record was decomposed into a background component (CHARback), and a peak component (CHARpeak), which represents either a single fire event or several temporally clustered fire events within the charcoal-source area. CHARback was calculated using a moving median smoothed with a 400-year moving-window that ensured a sufficient signal-to-noise index (SNI >3) (Kelly et al., 2011). The residual series (CHARpeak) was obtained by subtracting CHARback from CHARC, as we assume additive processes are responsible for increased charcoal delivery to the sedimentary basin. Fire episodes were inferred from CHARpeak by fitting a Gaussian mixture model to each overlapping 400-year window while assuming two sub-populations: CHARnoise and fire. This enables modeling both the mean and variance of the CHARnoise distribution locally. The CHARpeak exceeding the 99th percentile of the modeled CHARnoise distribution were considered peak events. Each peak that exceeded the locally defined threshold was statistically screened using the minimum-count test ($p < .05$) (Higuera et al., 2010).

Change point analysis was used to determine significant changes in biomass burning (i.e. periods of increased/decreased fire activity) for all charcoal records based on variations in both the mean and variance of CHARC using a penalty (β) = $6 \log(n)$ (Killick and Eckley, 2014). We also tested the potential effect of sedimentation-rate change on change point detection following Finsinger et al. (2018) using the 'change point' v2.2.2 package (Killick and Eckley, 2014).

3.6. Regional biomass burning

To disentangle local and regional drivers of fire in upper-montane forests in the Carpathians, the use of multiple sites, elevations and quantitative inter-site comparisons are necessary (Gavin et al., 2006). To reconstruct regional variations in biomass burning over time, we compared the record from Popradské pleso to three previously analysed sedimentary charcoal profiles from Poiana Știol, Tăul Muced and Lake Brazi (Fig. 1). These three sites were chosen based on the following criteria; 1) all sites are currently located in *Picea abies*-dominated forests at/near treeline; 2) all records are temporally well-constrained with limited hiatuses (see Feurdean et al., 2015; Feurdean et al., 2016; Finsinger et al., 2014); and 3) pollen and high-resolution charcoal data are available from each site, and span the majority of the Holocene. All fire histories were determined using the CharAnalysis software (Higuera et al., 2009) and followed the same methodologies presented in section 3.5 to allow for regional comparisons of biomass burning histories.

3.7. Local climate model

To reconstruct the influence of broad-scale changes in climate on fire activity at Popradské pleso, we developed a locally-derived macrophysical climate model (MCM; Bryson, 2005) using meteorological data of the Slovak Hydrometeorological Institute from the closest weather station, Štrbské pleso. For more information about the MCM, please see Carter et al. (2018).

3.8. Generalized additive models

Generalized additive models (GAMs; Hastie and Tibshirani, 1986) use a link function to investigate the relationship between the mean of the response variable (i.e. dependent variable) and a smoothed function of the predictor variable (i.e. independent variable). In this study, we investigated the relationship between biomass burning (e.g. total CHARC from each of the four study sites) and *Picea abies*, *Pinus*, primary forest cover (i.e., the summed percentage values of *Picea abies* and *Pinus*), forest density (the ratio of *Picea*:*Pinus*), and human indicators. Given that many secondary human indicator taxa were naturally abundant in the early Holocene (Fyfe et al., 2015), we used a base period from 8000 cal yr BP to present in the model to remove any bias secondary indicator pollen taxa may have in the early Holocene before the

intensification of human activity (see SI Table 1 for complete list of indicator taxa). We used a quasi-Poisson distribution with a log link function using the 'mgcv' package (Wood, 2017) in R. The GAMs were fit using restricted maximum likelihood smoothness selection.

4. Results

4.1. Popradské pleso chronology

The gravity core (POP-15-GC1) shows a very low accumulation rate with $^{210}\text{Pb}/^{226}\text{Ra}$ equilibrium at a depth of around 7 cm (Fig. 2). The exponential decline in unsupported ^{210}Pb concentrations with depth suggests a relatively uniform sedimentation rate throughout the period of time spanned by the ^{210}Pb record. High ^{137}Cs concentrations in the uppermost 2 cm most probably record fallout from the 1986 Chernobyl accident. Although the ^{137}Cs record from nuclear weapons test fallout again appears to have been obscured by Chernobyl ^{137}Cs , the 1963 fallout maximum is recorded by the presence of significant concentrations of ^{241}Am between 2 and 4 cm. ^{210}Pb ages calculated using the CRS and CIC models (Appleby and Oldfield, 1978) both suggest a relatively uniform sedimentation rate since the middle of the 19th century with a mean value during that time of $0.0065 \pm 0.0008 \text{ g cm}^{-2} \text{ y}^{-1}$ (0.048 cm y^{-1}). These results place 1986 at a depth of between 1 and 2 cm and 1963 between 2 and 3 cm, in reasonable agreement with the depths suggested by the ^{137}Cs and ^{241}Am records. The base of the ^{210}Pb age-depth model is consistent with European increases in airfall stable Pb concentrations in the sediment, associated with industrialization from the 1850's onwards (Renberg et al., 2000).




Fig. 2. Radiometric chronology showing: a) concentrations of supported and unsupported ^{210}Pb , b) concentrations of ^{137}Cs and ^{241}Am ; c) the ^{210}Pb ages and the depths of the 1986 (Chernobyl) and post-1963 (weapons testing) ^{137}Cs and ^{241}Am markers. d) Airfall Pb pollutant concentrations (determined by XRF).

Previous research from Popradské pleso suggested the possibility of a hiatus at the transition between the Late Glacial and the Holocene (Rybníčková and Rybníček, 2006). We found an age difference of ca. 4000 ^{14}C years between depths 1374.5 cm and 1371 cm below the water surface (Table 1). As core lithology changes dramatically at depth 1371.5 cm, transitioning from inorganic glacial sediments into organic-rich

gyttja (Fig. 3), we confirm the sedimentation hiatus originally suggested by Rybničková and Rybniček (2006). Due to multiple age-reversals between the bottom of the core (depth 1635 cm) and depth 1373 cm, the age-depth model and the basal age for Popradské pleso are insufficiently constrained. Therefore, we will only present and discuss the section of the sedimentary profile reflecting continuous accumulation spanning the Holocene (Fig. 3).




Fig. 3. Age-depth model for Popradské pleso constructed using ^{210}Pb (light blue line near modern) and 7 ^{14}C radiocarbon dates (shown in red) using 'BACON' (Blaauw and Christen, 2011). The gray area represents the 95th confidence intervals, and the white line delineates the mean probability. High (μm) resolution digital line-scan image of the Pop15–1 core taken using the University of Liverpool Geotek Multi-sensor core logger (MSCL-XZ). Down core profiles for low frequency magnetic susceptibility (XLF), μXRF scan geochemical data for Zr (ppm), Rb (ppm) and Pb/Rb for the long core. Pb/Rb ratio data continues to the modern period using dried sample XRF measurements.

4.2. Reconstructed vegetation dynamics at Popradské pleso

Summarising the dominant canopy taxa and significant changes in understory at Popradské pleso, the pollen percentage profile was divided by six statistically significant pollen assemblage zones (SI Fig. 1). Zone 1 (age undetermined; depth 1380–1370 cm) probably captures the Younger Dryas/Holocene transition, but due to a hiatus we were unable to confirm the exact ages of this zone. Regardless, this zone is characterized by high percentages of *Pinus* (>50%) and non-arboreal pollen (NAP; >20%), and by low percentages of *Picea abies* (<10%) similar to that found by Rybničková and Rybniček (2006) (SI Fig. 2). Low percentages of low-elevation arboreal species such as *Corylus avellana* were also recorded in this zone (SI Fig. 1). No plant macrofossils were found in Zone 1. Zone 2 (9500–8500 cal yr BP; depth 1370–1355 cm) is characterized by an increase in *Picea abies* percentages (~15%), with relatively high values of *Picea abies* influx values (~1000 grains $\text{cm}^{-2} \text{yr}^{-1}$), the presence of *Picea abies* stomata, and high concentration of *Picea abies* macrofossils. *Pinus* percentages (~20%) and influx (>2000 grains $\text{cm}^{-2} \text{yr}^{-1}$) values both declined from the previous zone, however *Pinus* stomata and macrofossils, both *Pinus* sp. and *P. cembra* are present throughout this zone (Fig. 4). While total herbaceous percentages decreased to their lowest values, several herbaceous macrofossils were found in Zone 2, such as an *Alchemilla* sp. seed, a *Caltha palustris*/*Trollius altissimus* seed, and a *Poaceae* seed (SI Fig. 1). *Larix* pollen was not found in Zone 2, however, *Larix decidua* needles

were found. Zone 3 (8500–5600 cal yr BP; depth 1355–1293 cm) can be characterized by the local dominance of *Picea abies* and *Pinus* with pollen percentages averaging 15–20%, and pollen influxes averages 2823 grains cm⁻² yr⁻¹ and 3335 grains cm⁻² yr⁻¹. *P. cembra* needles decreased during this zone, while *Pinus*. sp. (most likely *P. mugo*) needles and bud scales increased. *Corylus avellana* percentages and influxes continued to decrease in this zone. High abundances of macrofossils were found at the onset of Zone 3; for example, seeds from *Rubus idaeus*, *Alchemilla* sp. and *Juncus* sp. and *Rumex acetosa* fruit. Zone 4 (5600–3900 cal yr BP; depth 1293–1248 cm) is notable in that both *Abies alba* and *Fagus sylvatica* pollen percentages and influxes slowly increase during this zone (SI Fig. 1). *Picea abies* pollen percentages and influxes also increase to their highest values (~37% and ~ 7400 grains cm⁻² yr⁻¹) of the entire Holocene record before beginning a long-term declining trend. Zone 5 (3900–800 cal yr BP; depth 1248–1133 cm) is characterized by a decreasing trend in both *Picea abies* pollen percentages and influxes, despite the high abundances of *P. abies* needles being recorded. *Pinus* pollen percentage values (~20%) were relatively stable, with high abundances of *P. cembra* needles being recorded in Zone 5. *Abies alba* and *Fagus sylvatica* reached their highest percentages (15% and 11%) and influx values (~1900 grains cm⁻² yr⁻¹ and 1160 grains cm⁻² yr⁻¹), and human impact indicator pollen percentages and influxes begin to increase during this zone. *Sphagnum* sp. capsules were also prominent in this zone. The final zone, Zone 6 (800 cal yr BP–present; depth 1133–1100 cm) is characterized by declining percentages of trees and shrubs, specifically *Picea abies*, *Fagus sylvatica*, and *Abies alba*. However, pollen influx for each of these species shows rapid variability instead of a declining trend. Pollen percentages and influx values of *Pinus* and of human indicators increase throughout Zone 6. Macrofossils were not analysed from Zone 6.




Fig. 4. Reconstructed fire, geochemical, vegetation, and climate at Popradské pleso. From top to bottom; Fire events are indicated with a red '+' symbol; less significant fire events are delineated with a gray circle; Significant peaks identified by CharAnalysis, but are not true fire events are delineated with a black circle; CHARC (particles per cm⁻² yr⁻¹) (gray curve), with BCHAR (red line); fire frequency (fires/1000 years)(filled colour curve); Magnetic susceptibility (XLF; gray filled curve); Zirconium (Zr; brown filled curve); Rubidium (Rb; reddish brown filled curve); Potassium (K; olive filled curve); Zr/Rb ratio (Zr = dark brown filled curve; Rb = reddish brown filled curve); Picea influx (Picea = dark green filled curve; Pinus = light green filled curve); Pinus influx (Pinus = light green filled curve; Picea = dark green filled curve); Human Indicators influx (Human Indicators = brown filled curve); Winter Mean Precipitation (mm) (solid blue line); Summer Mean Precipitation (mm) (dashed blue line); Winter Mean Temperature (°C) (solid blue line); Summer Mean Temperature (°C) (solid red line); Peak frequency fires/1000 years (filled colour curve); Total Pinus sp. & Pinus cembra Macrofossils (Total Pinus sp. = light green filled curve; Pinus cembra = dark green filled curve).

Rb = orange filled curve); *Picea abies* pollen percentages (filled dark green curve), *Picea abies* pollen influx (black line), and *Picea abies* macrofossil concentrations (black vertical bars); *Pinus* pollen percentages (filled light green curve), *Pinus* pollen influx (black line), and *Pinus* sp. (black vertical bars) and *Pinus cembra* (red vertical bars) macrofossil concentrations; Human indicator taxa (i.e. the sum of primary and secondary anthropogenic pollen taxa) pollen percentages (filled orange curve), and human impact indicator species pollen influx (black line); MCM reconstructed summer temperature (red line); MCM reconstructed winter temperature (blue line); MCM reconstructed summer precipitation (dark blue line), MCM reconstructed winter precipitation (light blue dotted line). Pink vertical bars reflect the zones of significant biomass burning as indicated by change point analysis.

4.3. Reconstructed fire and erosion history at Popradské pleso

Elevated CHARC values were recorded between 9000 and 7000 cal yr BP, with low values between 5500 and 1000 cal yr BP and then elevated values again in the last ~1000 years (Fig. 4). The change-point analysis shows three periods of high biomass burning; 9000–7000 cal yr BP, around 6800 cal yr BP, and the past 2000 years (Fig. 4). A total of 24 fire episodes were detected over the past 9500 years, with periods of increased fire frequency centred between 9000 and 8500 cal yr BP, 7000–6000 cal yr BP, and over the last 3000 years. During the past 200 years, one single event was identified in the CHARC record (see black circle, Fig. 4). However, based on documentary evidence, this is a false positive because no fire event has occurred at Popradské pleso over the last 200 years. The increase in CHARC is possibly due to the charcoal being produced from the Horský Hotel Popradské Pleso which was built on the shore of Popradské pleso in the mid-1800s. Fire frequency averages ~2.5 fires/1000 years (or a fire return interval of ~400 years) throughout the entire record.

XRF geochemical profiles for K, Ti, Rb and Zr (Boyle, 2001; Davies et al., 2015), alongside magnetic susceptibility (XLF) data (Dearing, 1992; Oldfield et al., 2003), show changes in the flux of lithogenic elements from the granitic bedrock catchment. The inorganic stratigraphy (Fig. 4) displays two types of behavior, 1) sharp and short-lived peaks and 2) changes in the background concentrations of these lithogenic markers. Ratios for the typically insoluble lithogenic elements, Zr and Rb, provide additional information on sediment grain size, with Zr typically coarse silt whereas Rb, Ti and K associate with finer fractions (Boyle, 2001; Davies et al., 2015). The peaks in lithogenic elements occur throughout the Holocene, but are larger in magnitude between 8500–7750, 6500–5500 and > 4500 cal yr BP. Coincidentally, these periods align with periods of increased biomass burning as detected by change point analysis. These peaks across all lithogenic elements reflect short-lived run-off extreme events (e.g. storms or rapid snowmelt), with coincident peaks in Zr/Rb reflecting the higher-energy riverine delivery of coarser sediments to the lake. For many layers, particularly >4500 cal. yr BP, the Zr/Rb peak often occurs early in the erosion event before finer grain size show declining energy in the event. The background quantities for these lithogenic markers and the associated general erosion regime also changes, with lower concentrations for the earlier Holocene, but increasing in both their concentration and grain size (Zr/Rb) after 4500 cal yr BP.

4.4. Modeling biomass burning-vegetation relationships across the Carpathians

At Lake Brazi, biomass burning significantly increases with increasing *Pinus* and *Picea abies* abundance ($p < .001$) and increasing primary forest cover ($p < .001$), and significantly decreases with forest density ($p < .001$) (Table 2). Human indicators have a negligible effect on biomass burning at Lake Brazi ($p = .682$) (Fig. 7). At Poiana Știol, biomass burning significantly increases with increasing forest density ($p < .001$) and primary forest cover ($p < .03$). Both *Pinus* ($p = .319$) and *Picea abies* ($p = .45$) abundance and human indicators ($p = .104$) all had insignificant effects on biomass burning. At Popradské pleso, *Pinus* and human

indicators were both found to positively ($p < .001$) influence biomass burning. Biomass burning decreased with increasing *Picea abies* ($p < .001$) and increasing forest density ($p < .001$) at Popradské pleso. Primary forest cover had a negligible effect on biomass burning at Popradské pleso ($p = .298$). Finally, at Tăul Muced, biomass burning significantly increases with increasing primary forest cover ($p < .002$) and *Picea abies* ($p < .003$) pollen. Forest density ($p = .369$), human indicators ($p = .389$), and *Pinus* pollen ($p = .624$) all had insignificant effects on biomass burning.

Site name	Independent variable	edf	ref.df	p-value	r2 adj	Deviance explained
Lake Brazi	<i>Pinus</i>	3.78	4.71	0.001	0.4	44%
Lake Brazi	Primary Cover	3.53	4.2	0.001	0.3	36%
Lake Brazi	Forest Density	2.95	3.71	0.001	0.24	34%
Lake Brazi	<i>Picea abies</i>	1	1	0.001	0.06	8%
Lake Brazi	Human Indicators	1.41	1.72	0.682	0	1%
Poiana Ştiol	Forest Density	7.63	8.43	0.001	0.52	46%
Poiana Ştiol	Human Indicators	3.01	3.81	0.104	0.13	24%
Poiana Ştiol	<i>Pinus</i>	4.25	5.28	0.426	0.06	23%
Poiana Ştiol	Primary Cover	1	1	0.032	0.004	11%
Poiana Ştiol	<i>Picea abies</i>	1	1	0.718	-0.01	0.3
Popradské pleso	<i>Pinus</i>	1.94	2.44	0.001	0.45	45%
Popradské pleso	Human Indicators	3.97	4.86	0.001	0.27	43%
Popradské pleso	Forest Density	2.3	2.9	0.001	0.4	42%
Popradské pleso	<i>Picea abies</i>	1	1	0.001	0.12	19%
Popradské pleso	Primary Cover	2.58	3.3	0.298	0.02	10%
Tăul Muced	Primary Cover	4.95	6.1	0.002	0.11	24%
Tăul Muced	<i>Picea abies</i>	4.85	6	0.003	0.11	23%
Tăul Muced	Forest Density	2.71	3.34	0.369	0.01	6%
Tăul Muced	Human Indicators	1.84	2.31	0.389	0	3%
Tăul Muced	<i>Pinus</i>	2.12	2.6	0.624	0	3%

Table 2. Correlation between the independent variables, represented by pollen percentage data (*Pinus*, *Picea abies*, primary cover (i.e., the summed percentage values of *Picea abies* and *Pinus*, forest density (the ratio of *Picea*:*Pinus*), and human indicator taxa) and the dependent variable (biomass burning; charcoal influx) at each of the four study sites: from highest to lowest elevation, Lake Brazi, Poiana Ştiol, Popradské pleso and Tăul Muced.

5. Discussion

The new data on biomass burning (CHARC) from the High Tatra Mountains follows similar burning patterns to those experienced at the other upper-montane Carpathian sites during the Holocene (Fig. 5). Both in the High Tatras as well as at other Carpathian sites, forests were dominated by *Picea abies*, a fire avoider species with minimal fire adaptations (Feurdean et al., 2017), and often associated with reduced fire activity (Ohlson et al., 2011). As a consequence, minimal biomass burning would be expected, yet biomass burning has been an agent of forest disturbance prevalent in the Carpathians through the Holocene.




Fig. 5. Pan-Carpathian biomass burning and vegetation change at each of the four study sites presented in this study; from top to bottom, Lake Brazi, elevation 1740 m a.s.l.; Poiana Ştiol, elevation 1540 m a.s.l.; Popradské pleso, elevation 1513 m a.s.l.; Tăul Muced, elevation 1360 m a.s.l. Biomass burning indicators are represented by both CHARC and BCHAR (gray filled curve and red line) and fire frequency (filled colour

curve). Vegetation change is represented by pollen percentage data. Pink vertical bars reflect the zones of significant biomass burning as indicated by change point analysis.

5.1. Holocene wildfire activity across the Carpathians

5.1.1. Early and mid-Holocene

In the upper-montane forest zone, higher rates of biomass burning (i.e. total CHARC) were recorded between 10,000–6000 cal yr BP at Popradské pleso, Lake Brazi and Poiana Știol, as indicated by both change point analysis and increased fire frequencies (Fig. 5). Higher CHARC is likely connected to broad-scale climate, with summer temperatures significantly warmer-than-present as a result of peak summer insolation (Berger and Loutre, 1991). Increased abundances of *Pinus cembra* and *P. mugo* macrofossils at Lake Brazi, (Finsinger et al., 2018), *P. cembra* and *Pinus* sp. (likely either *P. cembra* or *P. mugo*) macrofossils at Popradské pleso (Fig. 4), and *P. sylvestris*/*mugo* macrofossils at Poiana Știol (Feurdean et al., 2016) were found during the early and mid-Holocene suggesting a more open upper-montane landscape, which is supported by lower *Picea*:*Pinus* ratio values until ~8500 cal yr BP (Fig. 6). Warmer conditions in the early Holocene, coupled with generally higher abundances of *Pinus* sp. likely *P. mugo* or *P. cembra*, created a positive climate-fuel relationship that resulted in higher amounts of biomass burning. *P. sylvestris* may have potentially been present in the early Holocene, as its pollen is indistinguishable from *P. mugo*. Long-term paleoecological records indicate that the late-successional *P. cembra* is able to withstand moderately frequent surface fires, but not high anthropogenic fire activity (Ali et al., 2005; Genries et al., 2009; Blarquez and Carcaillet, 2010; Schwörer et al., 2015), while the woody shrub, *P. mugo*, is favored by fire disturbances under drier conditions (Stähli et al., 2006; Leys et al., 2014). *P. sylvestris* is an early-successional species, fire-resister species, with a moderate-severity fire regime (Agee, 1988; Feurdean et al., 2017). Modeled *Pinus* response curves demonstrate a significant relationship with biomass burning at Lake Brazi and Popradské pleso (Table 2). This suggests that forest composition with higher amounts of *P. mugo*/*sylvestris* and/or *P. cembra*, despite the prevalence of *Picea abies*, may have been especially important for influencing biomass burning, especially at Lake Brazi and Popradské pleso.




Fig. 6. Figure illustrating how biomass burning (CHARC) increases (decreases) at all sites when the forest cover is comprised with more Pinus (Picea); from top to bottom, Lake Brazi, elevation 1740 m a.s.l.; Poiana Știol, elevation 1540 m a.s.l.; Popradské pleso, elevation 1513 m a.s.l.; Tăul Muced, elevation 1360 m a.s.l. Pink vertical bars reflect the zones of significant biomass burning as indicated by change point analysis. Horizontal gray bars on the Picea:Pinus curve illustrate GAMS values at which biomass burning increases (see Fig. 7).

High biomass burning during the early and mid-Holocene is witnessed across Europe (Marlon et al., 2013; Feurdean et al., 2012, Feurdean et al., 2013, Feurdean et al., 2017; Clear et al., 2014; Vanni re et al., 2016; Dietze et al., 2018; Florescu et al., 2019), as well as throughout the Western Carpathian region. For example, in the Polish Western Carpathians, biomass burning was highest between 10,000–9500 cal yr BP and 7500–6000 cal yr BP at lower-montane forests (Ko aczek et al., 2020). At Belianske l ky, Slovakia's largest spring fen located near the foothills of the High Tatra Mountains approximately 30 km from Popradsk  pleso, high macrocharcoal abundances were recorded ~8500 cal yr BP as a result of drier conditions (H jkov  et al., 2012). These results from Belianske l ky generally agree with high CHARC values at Popradsk  pleso. However, at Popradsk  pleso, both CHARC and fire frequency decrease and remain low, while indicators for soil erosions (i.e. K, Ti, Rb and XLF) all increase between 8400 and 7000 cal yr BP (Fig. 3). Interestingly, this time period overlaps with a potential sediment hiatus in the Polish Carpathian Mountains as a result of wetter conditions which caused the loss of sediment deposits (Ko aczek et al., 2020). Increased erosion is also recorded in both the Beskid Makowski Mountains and Beskid Wyspowy Mountains during this same time period (Margielewski, 2006, Margielewski, 2018). The time period between 8400 and 7000 cal yr BP overlaps with a period of global rapid climate changes (Mayewski et al., 2004), that is well documented across the entire Carpathian region, either by an increase in moisture (H jkov  et al., 2016; Jamrichov  et al., 2018;  olcov  et al., 2018; Dabkowski et al., 2019), reoccurring fire events (P l et al., 2016, P l et al., 2018; Vincze et al., 2017; Florescu et al., 2019), or as a change in climatic conditions or erosion activity (Schnitchen et al., 2006; Feurdean et al., 2008; T th et al., 2015; Ga ka et al., 2020). Together, this data suggests climate as the ultimate driver of biomass burning in the Carpathians in the early and mid-Holocene.

However, archaeological evidence demonstrates human occupation and activity near the High Tatra Mountains and throughout the Carpathian Basin since at least the Mesolithic period ~10,000 cal yr BP (Valde-Nowak and Soj k, 2010; Kert sz, 2002 and references therein). Florescu et al. (2018) found an increase in biomass burning ~8500 cal yr BP in the Southern Carpathians which coincided with a slight increase in meadow and pasture pollen types, suggesting that Early-Middle Neolithic (~8000 cal yr BP) populations occupied montane forest-elevations in the Southern Carpathians. Both *Cerealia* and *Secale* were first identified at T ul Muced ~7300 cal yr BP (Feurdean et al., 2017) illustrating a long history of human occupation in the Southern Carpathians. Similarly, others have also documented, or hinted at the possible presence of humans at montane-elevations in the Southern Carpathians in the late Neolithic (F rca  et al., 2003, F rca  et al., 2013; Feurdean et al., 2016, Feurdean et al., 2017). However, because certain taxa typically classified as secondary human indicator species (e.g. *Rumex*, *Poaceae*, *Plantago*) are apophytes and were naturally and abundantly present during the early Holocene, these data were left out of the GAMS model prior to 8000 cal yr BP, and thus our results are only indicative of anthropogenic activity post 8000 cal yr BP. It may be that Mesolithic and Neolithic populations were foraging seasonally among the Carpathians both in the lowlands and at higher-elevations, however, palynological evidence does not illustrate large-scale changes in vegetation at any of the study sites (Fig. 5). While low density Mesolithic/Neolithic populations could have contributed to the overall rate of biomass burning in the early and mid-Holocene, we suggest climate was the ultimate driver of early and mid-Holocene burning in upper-montane forests.

5.1.2. Mid to late Holocene

Biomass burning was reduced in the regions' upper-montane forest zone from ~6000 to ~2000 cal yr BP (Fig. 5) as a result of cooler and wetter conditions. Cooler and wetter conditions led to denser *Picea abies* upper-montane forests (Fig. 6), which naturally modulated biomass burning by regulating understory microclimates (Chen et al., 1999; Davis et al., 2018). Thus, this coupled climate-fuel relationship led to an

overall reduction in biomass burning in upper-montane forests across the Carpathians (Feurdean et al., 2020). However, despite the dominance of *Picea abies*, fires were not excluded from this forest type, which is illustrated by relatively moderate fire frequencies (~2 fires/1000 years) at most sites (Fig. 5).

At Popradské pleso, a substantial change in the erosion regime occurred ~4000 cal. yr BP, with the gradual increase in the baseline concentrations for lithogenic markers (Zr, K, Ti, Rb and XLF) showing a persistent shift to greater erosion (Fig. 4). Increased frequency of coarser and mineral rich layers from ~4000 cal. yr BP suggest also an intensification of the run-off events (e.g. storms or rapid snowmelt) (Schillereff et al., 2014, Schillereff et al., 2019; Chiverrell et al., 2019). Together, the long-term baseline concentrations and events (i.e. peaks) shown in the erosion proxies both point to increases in moisture and mineral in-wash. Soil erosion processes are highly dependent upon the magnitude of rainfall events (Schumacher et al., 2016), which the MCM model demonstrates is highly variable over the last 4000 years at Popradské pleso (Fig. 4; summer precipitation curve). We therefore hypothesize that an increase in summer precipitation could have potentially caused the overall decrease in biomass burning in the Western Carpathians. Our interpretation of an increase in moisture across the region is supported by rising groundwater levels leading to forestless bogs between 4395 and 4224 and again at 3940–3050 cal yr BP in the Polish Carpathians (Krapiec et al., 2016), an increase in illuvial deposits in the Beskid Makowski and Beskid Wyspowy Mountains (Margielewski, 2006; Margielewski, 2018; Starkel et al., 2013), as well as a sediment hiatus in the Polish Western Carpathians between 4940 and 3550 cal yr BP (Kofaczek et al., 2020). The timing of this sediment hiatus in the Polish Western Carpathians is interesting in that it aligns perfectly with a significant reduction in fire frequency at Popradské pleso, thus supporting a climatically driven, regional-scale moisture event. However, this moisture event was not confined to the Western Carpathians. Increased fluvial activity and/or increased moisture was also recorded in the Dniester River valley, Eastern Carpathians (Kofaczek et al., 2018) and in the Rodna Mountains, Eastern Carpathians (Gałka et al., 2016, Gałka et al., 2020; Diaconu et al., 2017).

Despite cooler and wetter conditions documented in the late Holocene, the forest density index (i.e. the *Picea:Pinus* ratio) shows a transition from a dense *Picea*-dominated to a more open *Pinus*-abundant system beginning ~3500 cal yr BP at Popradské pleso. This transition is supported by higher abundances of *Pinus cembra* macrofossils ~4000 cal yr BP. A more open landscape may have contributed to the intensification of the run-off events already discussed. However, *P. cembra* and *P. mugo* both have extension root systems that help prevent soil erosion in alpine environments. Thus, the intensification of human activities in the High Tatra Mountain area may have significantly contributed to changes in the forest composition and in biomass burning. The first indication of intense human impact in the High Tatra Mountains is dated to ~4000 cal yr BP (Obidowicz, 1996), which aligns with the first recording of primary human indicator pollen types at Popradské pleso (SI Fig. 1). The modeled response curves for Popradské pleso demonstrate a significant positive relationship with biomass burning and human indicator species (Fig. 7). Together, this data suggests that intensified human activities at high elevation areas may have contributed to the change in forest density. However, while there is no large-scale change in CHARC at Popradské pleso, fire frequency increases at this time (Fig. 5), suggesting that humans may have been utilizing small-scale fires to alter upper-montane forest composition as early as ~3500 cal yr BP.




Fig. 7. Generalized Additive Models showing the relationship between biomass burning (y-axis; CHARC) and dominant bottom-up drivers (from top to bottom, Pinus, Picea, primary canopy cover (sum of Pinus and Picea), human indicator species, and forest density (ratio of Picea:Pinus) at each of the four study sites shown from highest in elevation (Lake Brazi, left) to lowest in elevation (Tăul Muced, right). Pointwise confidence intervals (95%) are indicated by the gray bands.

Regionally, *Cerealia* pollen is first recorded ~4400 cal yr BP in the Polish Western Carpathians (Czerwiński et al., 2019), while both *Cerealia* and *Secale* are present in subalpine sediments ~4200 cal yr BP in the Rodna Mountains, Eastern Carpathians (Geantă et al., 2014). Similarly, *Cerealia* pollen has been consistently present in the Beskid Wyspowy Mountains, Polish Western Mountains since 2880 cal yr BP along with an increase in fire activity illustrating the connection between land use and fire (Czerwiński et al., 2019). An increase in biomass burning during the Middle (~2600 cal yr BP) and Late Bronze Age (~2100 cal yr BP) may be connected with the utilization of high elevation areas by humans in the Carpathians (Schumacher et al., 2016; Vincze et al., 2017), similar to a well-documented process in the Alps where seasonal pastoralism has resulted in a lowered treeline (Dietre et al., 2016; Tinner, 2007). Feurdean et al. (2020) found that biomass burning increased dramatically in continental regions during the Late Bronze Age and Iron Age as a result of increased deforestation and the creation of arable land. Over the last 2000 years, biomass burning has increased considerably at both Popradské pleso and Tăul Muced. The high rates of biomass burning recorded in the High Tatra Mountains likely reflects regional burning at lower elevations as a result of human activities which is demonstrated by an increase in human indicator species over the last 2000 years (Fig. 5), and a concurrent increase in microcharcoal – an indicator of regional biomass burning (SI Fig. 1). These data agree with archaeological evidence showing humans first impacted the forests near Belianske lúky ~2000 cal yr BP (Jankovská, 1972). Western Carpathian forests were being affected by intense human

activities, specifically through forest clearance by early farmers who used fire as a landscape management tool (Kukulak, 2000; Jawor, 2016a, Jawor, 2016b). Summit grasslands in the nearby Hercynian mid-mountains of central Europe were also affected by human occupation and activity as early as the Iron Age ~ 1900 cal yr BP (Novák et al., 2010). Intense human activities and land-use change also affected local beetle communities since 1250 CE at Diera Hollow downslope from Popradské pleso (Schafstall et al., 2020). However, despite regional literature and pollen records discussed here supporting the notion that humans influenced landscapes regionally beginning ~4000 cal yr BP, the modeled response curves illustrate a weak connection between biomass burning and human indicator species at Lake Brazi, Poiana Știol and Tăul Muced (Fig. 7). This suggests that human activities (i.e. land use) were less extensive in the upper-montane forest zone in Southern and Eastern Carpathians, as previously suggested by Finsinger et al. (2018).

5.2. Forest composition and forest density influences biomass burning in Carpathian upper-montane forests

The interplay between forest composition and forest density significantly influenced the overall patterns in biomass burning found in upper-montane Carpathian forests. Specifically, there is a strong relationship between biomass burning and forest density (i.e., *Picea*:*Pinus* ratio); when forest density decreases, biomass burning increases (Fig. 6, Fig. 7). The forest density index illustrates that biomass burning is highest when landscapes are more open and abundant with *Pinus*, while biomass burning is greatly reduced when dense *Picea abies* forests form (Fig. 7). This relationship is also reflected in the modeled primary forest cover (i.e. the sum of *Pinus* and *Picea* pollen) response curves which show that burning increases at intermediate pollen values ranging between ~30–50%. Biomass burning decreases at most sites when primary forest cover is denser (>50% pollen; Fig. 7). This suggests that when forests reach a threshold of ~50% primary canopy pollen, forests may be too dense, with an understory either too damp to burn and/or comprising insufficient ground-level fuel to sustain fire spread. While the canopy threshold of ~50% is slightly lower than identified by Feurdean et al. (2020), who suggest biomass burning increases when total forest canopy pollen percentages reaches 60%, our results agree with their findings in that biomass burning is reduced when tree cover (deduced from pollen data) reaches certain threshold levels in these upper-montane forests.

Together, the modeled response curves illustrate the sensitivity of upper-montane forests to fire through alterations in both forest composition and density. While these results are important for forest management implications, additional research utilizing pollen productivity estimates (PPEs) would be invaluable for forest managers interested in mitigating against future climate change in these forests (see Trondman et al., 2015; Abraham et al., 2016). Although our study is unable to quantify canopy openness due to the lack of adequate sites in the Carpathians used to calculate PPEs, our study does illustrate the need for further fire-related research from the region.

5.3. Future climate change impact on upper-montane Carpathian forests

Future climate projections for the Carpathians forecast an additional 1–2 °C increase in summer mean temperatures by the end of the 21st century, leading to projected increases in heat waves, droughts and fire activity (Anders et al., 2014; Lung et al., 2013). Both summer and winter precipitation are projected to decrease in the Southern Carpathians, by as much as 10–15% in the summer and 2–5% in the winter (Anders et al., 2014). Whereas in the Western Carpathians, summer precipitation is projected to decrease ~10%, yet winter precipitation is projected to increase by 10% (Anders et al., 2014). These projections suggest that changes in vegetation composition and disturbance regimes may be more pronounced in the Southern Carpathians forests.

There is a high confidence that treeline will shift upward in response to warming temperatures (see Greenwood and Jump, 2014 and references therein; Field et al., 2014), which could add additional stress to upper-montane forests already experiencing temperature amplification (Pepin et al., 2015). Our results offer potential analogs with regards to ecosystem responses and the anticipated impact of future climate change and fire risk, hypothetically leading to two different climate-fuel scenarios for these upper-montane forests. First, an upward shift of treeline could result in an increase in *Picea abies*, thereby creating a negative feedback with dense *Picea abies* forests generally suppressing the overall risk of fire. This feedback is generally seen in northern European forests, where a decrease in biomass burning with *Picea abies* expansion has been documented (Ohlson et al., 2011). Second, as exemplified by the warmer and drier early to mid-Holocene, *Pinus cembra* and/or *P. mugo* (or potentially *P. sylvestris*) could become vulnerable to warming temperatures, thereby developing increased susceptibility to fire danger and constituting a positive climate-fuel feedback. Given that this second scenario resembles more-closely future climate projections for the region i.e. warmer and drier conditions, especially those projected for the Southern Carpathians, we suggest that upper-montane forests may be more vulnerable to fire risk than currently thought, especially upper-montane forests found in the Southern Carpathians.

The second scenario has been documented for the wider region. For example, in the Alps, *P. cembra* benefited from above modern mean July temperatures during the mid-Holocene (Colombaroli et al., 2010). Increased drought conditions and reduced precipitation in the mid-Holocene favored *Pinus sylvestris* over *Picea abies*, leading to an increase in fire frequency in the Ural Mountains, Russia (Barhoumi et al., 2019). In the Balkan Bulgarian Mountains, fires mostly occurred in *Pinus*-dominated systems over the last 600 years, however, the largest fire documented historically have recently been in *Picea abies*-dominated forests as a result of humans (Panayotov et al., 2017). Finally, simulations using LandClim from the Alps project an upward shift of vegetation will occur, with *Picea abies* being replaced by *Pinus* in response to higher drought incidence (Schumacher and Bugmann, 2006).

Given that our sites from the Carpathians all document a shift to more *Pinus* pollen relative to *Picea abies* at present (Fig. 6), and that *Picea abies* forests are projected to decline in extent with climate change (Werners et al., 2014), this second scenario is especially concerning. Our results highlight that biomass burning is a prevalent disturbance agent in upper-montane forests, but these systems may become more vulnerable to an increasing fire risk with climate change. However, other factors such as soil development (Henne et al., 2011) and aspect (Courtney-Mustaphi and Pisaric, 2013) may influence and/or inhibit changes in vegetation composition and disturbance events in these forests in the future. Additionally, intensified land-use practices coupled with increased fire frequency could also influence/inhibit vegetation composition, as witnessed from the Alps (Berthel et al., 2012). Additional research from the region, especially from the Ukrainian Carpathians, that utilize high-resolution, multi-proxy approaches are necessary to fully understand the mechanics and spatial distribution of how upper-montane forests may be impacted with future climate change.

6. Conclusion

Our study provides the first long-term fire record from the upper-montane forest zone from the High Tatra Mountains in the Western Carpathians. Our results demonstrate that fire has been a continuous disturbance agent at Popradské pleso throughout the entire Holocene period. Our results also demonstrate that biomass burning across other upper-montane Carpathian forests is influenced by forest composition and forest density, both of these are strongly-dependent on climate. In the early and mid-Holocene, a positive climate-fuel feedback developed that linked warmer conditions and abundant *Pinus* with higher biomass burning. When climate cooled and became wetter in the late Holocene, climate-fuel relationships

created a negative feedback with *Picea abies* modulating less fire activity at upper-montane elevations. However, despite the reduced risk of fire implicit in *Picea abies* forests, biomass burning continued in forests with an intermediate primary forest cover (i.e. 30–50%; sum of *Picea abies* and *Pinus* pollen). These findings suggest that more dense forests (>50% pollen) had microclimate conditions not favorable for fire activity.

As temperatures continue to increase across the Carpathian region, so will the rate at which natural disturbances (i.e. windthrow, fire, bark beetle outbreaks) occur, potentially threatening these upper-montane forest ecosystems (see e.g. Seidl et al., 2014). In addition, treeline is expected to migrate upslope in response to increasing temperatures (see Greenwood and Jump, 2014 and references therein), which could alter natural disturbance regimes. Understanding how upper-montane forests and natural disturbance regimes (i.e. biomass burning) may be impacted with future climate change is important as these forests provide many ecosystem services. We hypothesize two potential future scenarios based on our results. First, an upward migration of treeline could result in an increase in *Picea abies* thereby creating a negative feedback with dense *Picea abies* forests modulating any fire risk. Second, as seen in the early and mid-Holocene when climate was warmer and drier than present, patches of *Pinus cembra* and/or *P. mugo/sylvestris* may potentially be vulnerable to warming temperatures and more susceptible to fire. Our study highlights how forest composition and forest density may influence future fire risk.

Funding

This work was supported by the Czech Science Foundation [EUROPIA project no. 16-06915S], the German Research Foundation [grant no. FE-1096/6-1], the GINOP project [no. 2.3.2-15-2016-00019], the Czech Academy of Sciences long-term research development project [no. RVO 67985939], and the Ministry of Education-UEFISCDI, Romania (project PN-III-P4-ID-PCE-2016-0711).

Acknowledgement

We would like to thank the High Tatra National Park administration for allowing us to core the Popradské pleso, and Peter Fleischer for his support in the field. We also thank John Boyle, Ladislav Hamerlík, Fiona Russell, Daniel Schillereff, Jolana Tátosová, and Daniel Vondrák for their assistance during coring, and Peter Appleby and Gayane Piliposian of the Environmental Radioactivity Research Centre at the University of Liverpool who provided the Radiometric (²¹⁰Pb, ²²⁶Ra, ¹³⁷Cs and ²⁴¹Am) dating.

References

- Abraham, V., Kuneš, P., Petr, L., Svitavská-Svobodová, H., Kozáková, R., Jamrichová, E., Švarcová, M.G., Pokorný, P., 2016. A pollen-based quantitative reconstruction of the Holocene vegetation updates a perspective on the natural vegetation in the Czech Republic and Slovakia. *Preslia* 88, 409–434.
- Achard, F., Eva, H., Mollicone, D., Popatov, P., Stibig, H.-J., Turubanova, S. & Yaroshenko, A., 2009. Detecting intact forests from space: hot spots of loss, deforestation and the UNFCCC. In: *Old-Growth Forests*, ed. C. Wirth, G. Gleixner & M. Heimann, Berlin and Heidelberg, Germany: Springer. pp. 411–427.
- Alberton, M., Andresen, M., Citadino, F., Egerer, H., Fritsch, U., Götsch, H., Hoffmann, C., Klemm, J., Mitrofanenko, A., Musco, E., Noellenburg, N., Pettita, M., Renner, K., Zebisch, M., 2017. Outlook on climate change adaptation in the Carpathian Mountains. United Nations Environment Programme, GRID-Arendal and Eurac Research, Nairobi, Vienna, Arendal and Bolzano.

Albrich, K., Rammer, W., Thom, D., Seidl, R., 2018. Trade-offs between temporal stability and level of forest ecosystem services provisioning under climate change. *Ecological Applications* 28 (7), 1884–1896.

Ali, A.A., Carcaillet, C., Talon, B., Roiron, P. and Terral, J-F., 2005. *Pinus cembra* L. (arolla pine), a common tree in the inner French Alps since the early Holocene and above the present tree line: a synthesis based on charcoal data from soils and travertines. *Journal of Biogeography* 32, 1659–1669.

Anders, I., Stagl, J., Ingeborg, A., Pavlik, D., 2014. Climate Change in Central and Eastern Europe. In: Rannow S., Neubert M. (eds) *Managing Protected Areas in Central and Eastern Europe Under Climate Change*. *Advances in Global Change Research*, vol 58. Springer, Dordrecht.

Appleby, P.G., Oldfield, F., 1978. The calculation of lead-210 dates assuming a constant rate of supply of unsupported 210Pb to the sediment. *Catena* 5, 1–8.

Appleby, P.G., Nolan, P.J., Gifford, D.W., Godfrey, M.J., Oldfield, F., Anderson, N.J., Batterbee, R.W., 1986. 210Pb dating by low background gamma counting. *Hydrobiologia* 143, 21–27.

Appleby, P.G., Richardson, N., Nolan, P.J., 1992. Self-absorption corrections for well-type germanium detectors. *Nuclear Instruments and Method in Physics Research Section B* 71, 228–233.

Auer, I., Böhm, R., Jurkovic, A., Lipa, W., Orlik, A., Potzmann, R., Schöner, W., Ungersböck, M., Matulla, C., Briffa, K., Jones, P.D., Efthymiadis, D., Brunetti, M., Nanni, T., Maugeri, M., Mercalli, L., Mestre, O., Moisselin, J-M., Begert, M., Müller-Westermeier, G., Kveton, V., Bochnicek, O., Stastny, P., Lapin, M., Szalai, S., Szentimrey, T., Cegnar, T., Dolinar, M., Gajic-Capka, M., Zaninovic, K., Majstorovic, Z., Nieplova, E., 2007. HISTALP—Historical instrumental climatological surface time series of the greater Alpine region 1760–2003. *International Journal of Climatology* 27, 17–46.

Barhoumi, G., Peyron, O., Joannin, S., Subetto, D., Kryshen, A., Drobyshev, I., Girardin, M., Brossier, B., Paradis, L., Pastor, T., Alleaume, S., Ali, A.A., 2019. Gradually increasing forest fire activity during the Holocene in the northern Ural region (Komi Republic, Russia). *The Holocene* 29(12), 1906–1920.

Bennett, K.D., 1996. Determination of the number of zones in a biostratigraphical sequence. *New Phytologist* 132 (1), 155–170.

Berger, A., Loutre, M.F., 1991. Insolation value for the climate of the last 10 million years. *Quaternary Science Reviews* 10, 297–317.

Beug, H.-J., 2004. *Leitfaden der Pollenbestimmung für Mitteleuropa und angrenzende Gebiete*. Verlag Dr Friedrich Pfeil, Munich.

Birks, H.J.B., Gordon, A.D., 1985. *Numerical Methods in Quaternary Pollen Analysis*. Academic Press, London.

Birks, H.H., 2007. Plant macrofossil introduction. In: Elias, S.A. (ed.) *Encyclopedia of Quaternary Science*, Volume 3. Elsevier, Amsterdam, 2266–2288.

Blaauw, M., and Christen, J.A., 2011. Flexible paleoclimate age-depth models using an autoregressive gamma process. *Bayesian Analysis* 6, 457–474.

Blarquez, O., and Carcaillet, C., 2010. Fire, fuel composition and resilience threshold in subalpine ecosystem. *PLoS ONE* 5(8), e13480.

- Bogdan, O., 2008. Carpații Meridionali. Clima, hazardele meteo - Climatice și impactul lor asupra dezvoltării turismului. In: Academia Română., Institutul de Geografie. Univ., Lucian Blaga, Sibiu, Romania (In Romanian), pp. 1–314.
- Bojňanský, V., Fargašová, A., 2007. Atlas of Seeds and Fruits of Central and East-European Flora. The Carpathian Mountains Region. Springer, Dordrecht.
- Boyle, J.F., 1995. A simple closure mechanism for a compact, large-diameter, gravity corer. *Journal of Paleolimnology* 13 (1), 85–87.
- Boyle, J.F., 2001. Inorganic geochemical methods in palaeolimnology. In: Last WM, Smol JP (eds) Tracking environmental change using lake sediments. Volume 2: physical and geochemical methods. Kluwer Academic Publishers, Dordrecht, pp 83–141.
- Boyle, J.F., Chiverrell, R.C., Schillereff, D.N., 2015. Approaches to water content correction and calibration for μ XRF core scanning: comparing x-ray scatter with simple regression of elemental concentrations, in: Rothwell, R.G., Croudace, I.W. (Eds.), *Developments in Palaeoenvironmental Research: Micro-XRF Studies of Sediment Cores*. Springer, Dordrecht.
- Brown, T.A., Nelson, D.E., Mathewes, R.W., Vogel, J.S., Southon, J.R., 1989. Radiocarbon dating of pollen by accelerator mass spectrometry. *Quaternary Research* 32, 205–212.
- Bryson, R., 2005. Archeoclimatology. *Encyclopedia of World Climatology*. Springer, Netherlands, pp. 58–63.
- Cappers, R.T.J., Bekker, R.M., Jans, J.E.A., 2006. *Digitale Zadenatlas van Nederland*, Groningen Archaeological Studies, vol. 4. Barkhuis Publishing & Groningen University Library, Groningen.
- Caudullo, G., Tinner, W., de Rigo, D., 2016. *Picea abies* in Europe: distribution, habitat, usage and threats. In: San-Miguel-Ayanz, J., de Rigo, D., Caudullo, G., Houston Durrant, T., Mauri, A. (Eds.), *European Atlas of Forest Tree Species*. Publ. Off. EU, Luxembourg.
- Chen, J., Saunders, S.C., Crow, T.R., Naiman, R.J., Broszofski, K.D., Mroz, G.D., Brookshire, B.L., Franklin, J.F., 1999. Microclimate in forest ecosystem and landscape ecology: variations in local climate can be used to monitor and compare the effects of different management regimes. *BioScience* 49, 288–298.
- Chiverrell, R. C., Sear, D., Warburton, J., Macdonald, N., Schillereff, D.N, Dearing, J.A., Bradley, J., 2019. Using lake sediment archives to improve understanding of flood magnitude and frequency: Recent extreme flooding in northwest UK. *Earth Surface Processes and Landforms*, 44(12), 2366–2376.
- Clark, J.S., Royall, P.D., 1995. Particle size evidence for source areas of charcoal accumulation in late Holocene sediments of eastern North American lakes. *Quaternary Research* 43, 80–89.
- Clear, J.L., Molinari, C., Bradshaw, R.H.W., 2014. Holocene fire in Fennoscandia and Denmark. *International Journal of Wildland Fire* 23, 781–789.
- Colombaroli, D., Henne, P.D., Kaltenrieder, P., Gobet, E., Tinner, W., 2010. Species response to fire, climate and human impact at tree line in the Alps as evidenced by palaeo-environmental records and a dynamic simulation model. *Journal of Ecology* 98, 1346–1357.

Czerwiński, S., Margielewski, W., Gałka, M., Kołaczek, P., 2019. Late Holocene transformations of lower montane forest in the Beskid Wyspowy Mountains (Western Carpathians, Central Europe): a case study from Mount Mogielica. *Palynology* 44(2), 355–368.

Davies, S., Lamb, H., Roberts, S. J., 2015. *Micro-XRF Studies of Sediment Cores: Applications of a non-destructive tool for the environmental sciences*. Springer Nature, Vol. 17. p. 189-226 38 p. (Developments in Paleoenvironmental Research; vol. 17).

Davis, K.T., Dobrowski, S.Z., Holden, Z.A., Higuera, P.E., Abatzoglou, J.T., 2018. Microclimate buffering in forests of the future: the role of local water balance. *Ecography* 42, 1–11.

Dearing, J.A., 1992. Sediment yields and sources in a Welsh upland lake-catchment during the past 800 years. *Earth Surf Process Landf* 17:1–22

Diaconu, A.-C., Tóth, M., Lamentowicz, M., Heiri, O., Kuske, E., Tanțău, I., Panait, A.-M., Braun, M., Feurdean, A., 2017. How warm? How wet? Hydroclimate reconstruction of the past 7500 years in northern Carpathians, Romania. *Palaeogeography, Palaeoclimatology, Palaeoecology* 482, 1–12.

Dietre, B., Walser, C., Kofler, W., Kothieringer, K., Hajdas, I., Lambers, K., Reitmaier, T., Haas, J.N., 2016. Neolithic to Bronze Age (4850-3450 cal. BP) fire management of the Alpine Lower Engadine landscape (Switzerland) to establish pastures and cereal fields. *The Holocene* 27, 181–196.

Dietze, E., Theuerkauf, M., Bloom, K., Brauer, A., Dörfler, W., Feeser, I., Feurdean, A., Gedminienė, L., Giesecke, T., Jahns, S., Karpińska-Kołaczek, M., Kołaczek, P., Lamentowicz, M., Latałowa, M., Marcisz, K., Obremaska, M., Pędziszewska, A., Poska, A., Rehfeld, K., Stančikaitė, M., Stivrins, N., Święta-Musznicka, J., Szal, M., Vassiljev, J., Veski, S., Wacnik, A., Weisbrodt, D., Wiethold, J., Vannièrè, B., Słowiński, M., 2018. Holocene fire activity during low-natural flammability periods reveals scale-dependent cultural human-fire relationships in Europe. *Quaternary Science Reviews* 201, 44–56.

Dragotă, C.S., and Kucsisca, G., 2011. Global climate change-related particularities in the Rodnei Mountains National Park. *Carpathian Journal of Earth and Environmental Sciences* 6 (1), 43–50.

Faegri, K., Kaland, P.E., Kzywinski, K., 1989. *Textbook of Pollen Analysis*. Wiley, New York. pp 1–3237.

Fărcaș, S., Tanțău, I., Bodnariuc, A., 2003. The Holocene human presence in Romanian Carpathians, revealed by the palynological analysis. *Wurzbürger Geographische Manuskripte*, 63, 111–128.

Fărcaș, S., Tanțău, I., Mîndrescu, M., Hurdu, B., 2013. Holocene vegetation history in the Maramureș Mountains (Northern Romanian Carpathians). *Quaternary International* 293, 92–104.

Feurdean, A., Klotz, S., Mosbrugger, V., Wohlfarth, B., 2008. Pollen-based quantitative reconstructions of Holocene climate variability in NW Romania. *Palaeogeography, Palaeoclimatology, Palaeoecology* 260, 494–504.

Feurdean, A., Spessa, A., Magyari, E.K., Veres, D., Hickler, T., 2012. Trends in biomass burning in the Carpathian region over the last 15,000 years. *Quaternary Science Reviews* 45, 111–125.

Feurdean, A., Liakka, J., Vannièrè, B., Marinova, E., Hutchinson, S.M., Mosbrugger, V., Hickler, T., 2013. 12,000-Years of fire regime drivers in the lowlands of Transylvania (Central-Eastern Europe): a data-model approach. *Quaternary Science Reviews* 81, 48–61.

Feurdean, A., Galka, M., Kuske, E., Tanțău, I., Lamentowicz, M., Florescu, G., Liakka, J., Hutchinson, S.M., Mulch, A., Hickler, T., 2015. Last Millennium hydro-climate variability in Central-Eastern Europe (Northern Carpathians, Romania). *The Holocene* 25, 1179–1192.

Feurdean, A., Gałka, M., Tanțău, I., Geantă, A., Hutchinson, S.M., Hickler, T., 2016. Tree and timberline shifts in the northern Romanian Carpathians during the Holocene and the responses to environmental changes. *Quaternary Science Reviews* 134, 100–113.

Feurdean, A., Florescu, G., Vanni re, B., Tanțău, I., O'Hara, R.B., Pfeiffer, M., Hutchinson, S.M., Gałka, M., Moskal-del Hoyo, M., Hickler, T., 2017. Fire has been an important driver of forest dynamics in the Carpathian Mountains during the Holocene. *Forest Ecology and Management* 389, 15–26.

Feurdean, A., Vanni re, B., Finsinger, W., Liakka, J., Panait, A., Warren, D., Connor, S., Forrest, M., Werner, C., Andri , M., Bobek, P., Carter, V.A., Davis, B., Diaconu, A., Dietze, E., Feeser, I., Florescu, G., Gałka, M., Giesecke, T., Jahns, S., Jamrichov, E., Kajukało, K., Kaplan, J., Karpińska-Kołaczek, M., Kołaczek, P., Kuneš, P., Kupriyanov, D., Lamentowicz, M., Lemmen, C., Maryari, E., Marcisz, K., Marinova, E., R sch, M., Słowiński, M., Stančikaitė, M., Szal, M., Święta-Musznicka, J., Tanțău, I., Theuerkauf, M., Tonkov, S., Valkó, O., Vassiljev, J., Veski, S., Vincze, I., Wacnik, A., Wiethold, J., Hickler, T., Niamir, A., Novenko, E., Obremaska, M., Pędziszewska, A., Pfeiffer, M., Poska, A., 2020. Fire risk modulation by forest structure and land use in Eastern and Central Europe. *Biogeosciences*. 71(5), 1213–1230.

Field, C.B., Barros, V.R., Mach, K.J., Mastrandrea, M.D., van Aalst, M., Adger, W.N., Arent, D.J., Barnett, J., Betts, R., Bilir, T.E., Birkmann, J., Carmin, J., Chadee, D.D., Challinor, A.J., Chatterjee, M., Cramer, W., Davidson, D.J., Estrada, Y.O., Gattuso, J.P., Hijioka, Y., Hoegh-Guldberg, O., Huang, H.Q., Insarov, G.E., Jones, R.N., Kovats, R.S., Romero-Lankao, P., Larsen, J.N., Losada, I.J., Marengo, J.A., McLean, R.F., Mearns, L.O., Mechler, R., Morton, J.F., Niang, I., Oki, T., Olwoch, J.M., Opondo, M., Poloczanska, E.S., P rtner, H.O., Redsteer, M.H., Reisinger, A., Revi, A., Schmidt, D.N., Shaw, M.R., Solecki, W., Stone, D.A., Stone, J.M.R., Strzepek, R.M., Suarez, A.G., Tschakert, P., Valentini, R., Vicuņa, S., Villamizar, A., Vincent, K.E., Warren, R., White, L.L., Wilbanks, T.J., Wong, P.P., Yohe, G.W., 2014. Technical summary. In: *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Field, C.B., V.R. Barros, D.J. Dokken, K.J. Mach, M.D. Mastrandrea, T.E. Bilir, M. Chatterjee, K.L. Ebi, Y.O. Estrada, R.C. Genova, B. Girma,

E.S. Kissel, A.N. Levy, S. MacCracken, P.R. Mastrandrea, and L.L.White (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 35-94.

Finsinger, W., Kelly, R., Fevre, J., Magyari, E.K., 2014. A guide to screening charcoal peaks in macrocharcoal-area records for fire-episode reconstructions. *The Holocene* 24(8), 1002–1008.

Finsinger, W., Fevre, J., Orbn, I., Pl, I., Vincze, I., Hubay, K., Birks, H.H., Braun, M., Tth, M., Magyari, E.K., 2018. Holocene fire-regime changes near the treeline in the Retezat Mts. (Southern Carpathians, Romania). *Quaternary International* 477, 94–105.

Flachbart, V., 2007 Bezbukov oblasti na Slovensku – skutonosť alebo fikcia? – In: RIZMAN I. (ed.): *Lesncka typolgia a zisťovanie stavu lesa vo vzbe na trvalo udržiateľn obhospodarovanie lesov : Zbornk prspevkov a prezentci z odbornho seminra konanho 3. 12. 2007 na NLC vo Zvolene v elektronickej forme [CD-ROM]. NLC – stav lesnch zdrojov a informatiky, Zvolen.*

Florescu, G., Vanni re, B., Feurdean, A., 2018. Exploring the influence of local controls on fire activity using multiple charcoal records from northern Romanian Carpathians. *Quaternary International* 488, 41–57.

Florescu, G., Brown, K.J., Carter, V.A., Kuneř, P., Veski, S., Feurdean, A., 2019. Holocene rapid climate changes and ice-rafting debris events reflected in high-resolution European charcoal records. *Quaternary Science Reviews* 222, 105877.

Fyfe, R.M., Woodbridge, J., Roberts, N., 2015. From forest to farmland: pollen-inferred land cover change across Europe using the pseudobiomization approach. *Global Change Biology* 21(3), 1197–1212.

Ga ka, M., Tan au, I., Ersek, V., Feurdean, A., 2016. A 9000 year record of cyclic vegetation changes identified in a montane peatland deposit located in the Eastern Carpathians (Central-Eastern Europe): autogenic succession or regional climatic influences? *Palaeogeography, Palaeoclimatology, Palaeoecology* 449, 52–61.

Ga ka, M., Tan au, I., Carter, V.A., Feurdean, A., 2020. The Holocene dynamics of moss communities in subalpine wetland ecosystems in the Eastern Carpathian Mountains, Central Europe. *The Bryologist* 123(1), 84–97.

Gavin, D.G., Hu, F.S., Lertzman, K., Corbett, P., 2006. Weak climatic control of standscale fire history during the late Holocene. *Ecology* 87, 1722–1732.

Geant , A., Ga ka, M., Tan au, I., Hutchinson, S., M ndrescu, M., Feurdean, A., 2014. High mountain region of the Northern Romanian Carpathians responded sensitively to Holocene climate and land use change: A multi-proxy analysis. *The Holocene* 24(8), 944–956.

Genries, A., Mercier, L., Lavoie, M., Muller, S.D., Radakovitch, O., Carcaillet, O., 2009. The effect of fire frequency on local Cembra Pine populations. *Ecology* 90(2), 476–486.

Gottfried, M., Pauli, H., Futschik, A., Akhalkatsi, M., Baran ok, P., Alonso, J.L.B., Coldea, G., Dick, J., Erschbamer, B., Calzado, M.R.F., Kazakis, G., Kraj i, J., Larsson, P., Mallaun, M., Michelsen, O., Moiseev, D., Moiseev, P., Molau, U., Merzouki, A., Nagy, L., Nakhutsrishvili, G., Pedersen, B., Pelino, G., Puscas, M., Rossi, G., Stanisci, A., Theurillat, J-P., Tomaselli, M., Villar, L., Vittoz, P., Vogiatzakis, I., Grabherr, G., 2012. Continent-wide response of mountain vegetation to climate change. *Nature Climate Change* 2, 111–115.

Gray, J., 1965 Extraction techniques. In Kummel, B. and Raup, G., eds., *Handbook of Palaeontological Techniques*. San Francisco, W. H. Freeman & Co.: 530–587.

Greenwood, S., Jump, A.S., 2014. Consequences of treeline shifts for the diversity and function of high altitude ecosystems. *Arctic Antarctic and Alpine Research* 46, 829–840.

Grindean, R., Tan au, I., Feurdean, A., 2019. Linking vegetation dynamics and stability in old-growth forests of Central Eastern Europe: Implications for forest conservation and management. *Biological Conservation* 229, 160–169.

Grodzi ska, K., Godzik, B., Fr czek, W., Badea, O., Oszl nyi, J., Postelnicu, D., Shparyk, Y., 2004. Vegetation of the selected forest stands and land use in the Carpathian Mountains. *Environmental Pollution* 130, 17–32.

H jkov , P., Grootjans, A.B., Lamentowicz, M., Rybn ikov , E., Madaras, M., Opravilov , V., Michaelis, D., H jek, M., Joosten, H., Wo ejko, L., 2012. How a *Sphagnum fuscum*-dominated bog changed into a

calcareous fen: the unique Holocene history of a Slovak spring-fed mire. *Journal of Quaternary Science* 27(3), 233–243.

Hájková, P., Pařil, P., Petr, L., Chattová, B., Matys Grygar, T., Heiri, O., 2016. A first chironomid-based summer temperature reconstruction (13–5 ka BP) around 49°N in inland Europe compared with local lake development. *Quaternary Science Reviews* 141, 94–111.

Harsch, M.A., Hulme, P.E., McGlone, M.S., Duncan, R.P., 2009. Are treelines advancing? A global meta-analysis of treeline response to climate warming. *Ecology Letters* 12(10), 1040–1049.

Hastie, T.J., Tibshirani, R.J., 1986. Generalized additive models. *Statistical Science* 1(3), 297–318.

Hess, M., 1965. Piętra klimatyczne w polskich Karpatach Zachodnich (Vertical climatic zones in the Polish Western Carpathians). *Zesz Nauk UJ Pr Geograficzne* 11, 1–267 (in Polish).

Hess, M., 1974 Piętra klimatyczne w Tatrach (Vertical climatic zones in the Tatra Mountains). *Czasopismo Geograficzne* 45, 75–95 (in Polish, English summary).

Higuera, P.E., Brubaker, L.B., Anderson, P.M., Hu, F.S., Brown, T., 2009. Vegetation mediated the impacts of postglacial climate change on fire regimes in the south-central Brooks Range, Alaska. *Ecological Monographs* 79, 201–219.

Higuera, P.E., Gavin, D.G., Bartlein, P.J., Hallett, D.J., 2010. Peak detection in sediment–charcoal records: impacts of alternative data analysis methods on fire-history interpretations. *International Journal of Wildland Fire*. 19, 996–1114.

Higuera, P.E., Briles, C.E., Whitlock, C., 2014. Fire-regime complacency and sensitivity to centennial through millennial-scale climate change in Rocky Mountain subalpine forests, Colorado, USA. *Journal of Ecology* 102, 1429–1441.

Hlásny, T., and Sitková, Z., 2010. Spruce forests decline in the Beskids. National Forest Centre—Forest Research Institute Zvolen, Czech University of Life Sciences Prague, Forestry and Game Management Research Institute Jíloviště–Strnady, Zvolen, Slovakia. pp. 1–165.

Jamrichová, E., Gálová, A., Gařpar, A., Horsák, M., Frodlová, J., Hájek, M., Hajnalová, M., Hájková, P., 2018. Holocene development of two calcareous spring gens at the Carpathian-Pannonian interface controlled by climate and human impact. *Folia Geobotanica* 53, 243–263.

Jawor, G., 2016a. Northern extent of settlement on the Wallachian law in medieval Poland. *Res Historica* 41, 35–49.

Jawor, G., 2016b. Seasonal pastoral exploitation of forests in the area of Subcarpathia in the 15th and 16th century. *Balcanica Posnaniensia* 23, 175–185.

Katz, N.J., Katz, S.V., Skobeeva, E.I., 1977. Atlas of Plant Remains in Peat. Nedra, Moscow.

Kelly, R.F., Higuera, P.E., Barrett, C.M., Hu, F.S., 2011. A signal-to-noise index to quantify the potential for peak detection in sediment–charcoal records. *Quaternary Research* 75, 11–17.

Kertész, R., 2002. Mesolithic hunter-gatherers in the northwestern part of the great Hungarian Plain. *Praehistoria* 3, 281–304.

Killick, R., Eckley, I.A., 2014. Change point: an R Package for change point analysis. *Journal of Statistical Software* 58(3), 1–19.

Knorn, J., Kuemmerle, T., Szabó, A., Mîndrescu, M., Keeton, W.S., Radeloff, V.C., Abrudan, I., Griffiths, P., Gancz, V., Hostert, P., 2012. Forest restitution and the protected area effectiveness in post-socialist Romania. *Biological Conservation* 146 (1), 204–212.

Knorn, J., Kuemmerle, T., Radeloff, V.C., Keeton, W.S., Gancz, V., Biriş, I.-A., Svoboda, M., Griffiths, P., Hagatis, A., Hostert, P., 2013. Continued loss of temperate old-growth forests in the Romanian Carpathians despite an increasing protected area network. *Environmental Conservation* 40, 182–193.

Kołaczek, P., Gałka, M., Apolinarska, K., Płóciennik, M., Gąsiorowski, M., Brooks, S.J., Hutchinson, S.M., Karpińska-Kołaczek, M., 2018. A multi-proxy view of exceptionally early postglacial development of riparian woodlands with *Ulmus* in the Dniester River valley, western Ukraine. *Review of Palaeobotany and Palynology* 250, 27–43.

Kołaczek, P., Margielewski, W., Gałka, M., Karpińska-Kołaczek, M., Buczek, K., Lamentowicz, M., Borek, A., Zernitskaya, V., Marcisz, K., 2020. Towards the understanding the impact of fire on the lower montane forest in the Polish Western Carpathians during the Holocene. *Quaternary Science Reviews* 229, 106137

Koncek, M., 1974. *Klima Tatier*. Vyd. Slovenskoj Akad. Vied., Bratislava, pp. 856.

Körner, C., and Paulsen, J., 2004. A world-wide study of high altitude treeline temperatures. *Journal of Biogeography* 31, 713–732.

Körner, C., and Ohsawa, M., 2005. *Mountain Systems. Ecosystems and Human Well-Being. Current States and Trends* (ed Millennium Ecosystem Assessment), pp. 681–716. Island Press, Washington, DC, USA.

Krąpiec, M., Margielewski, M., Korzeń, K., Szychowska-Krąpiec, E., Nalepka, D., Łajczak, A., 2016. Late Holocene palaeoclimate variability: The significance of bog pine dendrochronology related to peat stratigraphy. The Puścizna Wielka raised bog case study (Orwa – Norway Targ Basin, Polish Inner Carpathians). *Quaternary Science Reviews* 148, 192–208.

Kučera, P., 2012. Remarks on the intramontane continentality of the Western Carpathians defined by the absence of *Fagus sylvatica*. *Thaiszia Journal of Botany* 22 (1), 65–82.

Kukulak, J., 2000. Sedimentary record of early wood burning in alluvium of mountain streams in the Bieszczady range, Polish Carpathians. *Palaeogeography, Palaeoclimatology, Palaeoecology* 164, 167–175.

Lévesque, M., Saurer, M., Siegwolf, R., Eilmann, B., Brang, P., Bugmann, H., Rigling, A., 2013. Drought response of five conifer species under contrasting water availability suggest high vulnerability of Norway spruce and European larch. *Global Change Biology* 19(10), 3184–3199.

Leys, B., Carcaillet, C., Blarquez, O., Lami, A., Musazzi, S., Trevisan, R., 2014. Resistance of mixed subalpine forest to fire frequency changes: the ecological function of dwarf pine (*Pinus mugo* ssp. *mugo*). *Quaternary Science Reviews* 90, 60–68.

Lindner, M., Garcia-Gonzalo, J., Kolström, M., Green, T., Reguera, R., Maroschek, M., Seidl, R., Lexer, M. J., Netherer, S., Schopf, A., Kremer, A., Delzon, S., Barbati, A., Marchetti, M., Corona, P., 2008. Impacts of

Climate Change on European Forests and Options for Adaptation. European Forest Institute, Joensuu, Finland.

Lung, T., Lavalle, C., Hiederer, R., Dosio, A., Bouwer, L.M., 2013. A multi-hazard regional level impact assessment for Europe combining indicators of climatic and non-climatic change. *Global and Environmental Change* 23, 522–536.

Mayewski, P.A., Rohling, E.E., Stager, C.J., Karlén, W., Maasch, K.A., Meeker, D.L., Meyerson, E.A., Gasse, F., van Kreveld, S., Holmgren, K., Lee-Thorp, J., Rosqvist, G., Rack, F., Staubwasser, M., Schneider, R.R., Steig, E.J., 2004. Holocene climate variability. *Quaternary Research* 62, 243-255.

Magyari, E.K., Jakab, G., Balint, M., Kern, Z., Buczko, K., Braun, M., 2012. Rapid vegetation response to lateglacial and early Holocene climatic fluctuation in the South Carpathian Mountains (Romania). *Quaternary Science Reviews* 35, 116–130.

Margielewski, W., 2006. Records of the late glacial-Holocene palaeoenvironmental changes in landslide forms and deposits of the Beskid Makowski and Beskid Wyspowy Mts. Area (polish outer Carpathians). *Folia Quaternary* 76, 1-149.

Margielewski, W., 2018. Landslide fens as a sensitive indicator of paleoenvironmental changes since the late glacial: a case study of the Polish Western Carpathians. *Radiocarbon* 60, 1199-1213.

Marlon, J.R., Bartlein, P.J., Danialu, A-L., Harrison, S.P., Maezumi, S.Y., Power, M.J., Tinner, W, Vannièrè, B., 2013. Global biomass burning: a synthesis and review of Holocene paleofire records and their controls. *Quaternary Science Reviews* 65, 5–25.

Meier, E.S., Lischke, H., Schmatz, D.R., Zimmermann, N.E., 2012. Climate, competition and connectivity affect future migration and ranges of European trees. *Global Ecology and Biogeography* 21, 164–178.

Miezi, P., Jakuš, R., Pennerstorfer, J., Havašová, M., Škvarenina, J., Ferenčík, J., Bičárová, S., Bilčík, D., Blaženec, M., Netherer, S., 2017. Storms, temperature maxima and the Eurasian spruce bark beetle *Ips typographus* - An inferno trio in Norway spruce forests of the Central European High Tatra Mountains. *Agricultural and Forest Meteorology* 242, 85–95.

Noroozi, J., Talebi, A., Doostmohammadi, M., Rumpf, S.B., Peter Linder, H., Schneeweiss, G.M., 2018. Hotspots within a global biodiversity hotspot – areas of endemism are associated with high mountain ranges. *Scientific Reports* 8, 10345

Novák, J., Petr, L., Treml, V., 2010. Late Holocene human-induced changes to the extent of alpine areas in the East Sudetes, Central Europe. *The Holocene* 20(6), 895–905.

Obidowicz, A., 1996. A late glacial-Holocene history of the formation of vegetation belts in the Tatra Mountains. *Acta Palaeobotanica* 36(2), 159–206.

Ohlson, M., Brown, K.J., Birks, H.J.B., Grytnes, J.-A., Hörnberg, G., Niklasson, M., Seppä, H., Bradshaw, R.H.W., 2011. Invasion of Norway spruce diversifies the fire regime in boreal European forests: spruce invasion alters the fire regime. *Journal of Ecology* 99, 395–403.

Oldfield, F., Wake, R., Boyle, J., Jones, R., Nolan, S., Gibbs, Z., Appleby, P., Fisher, E., Wolff, G., 2003. The late-Holocene history of Gormire Lake (NE England) and its catchment: a multiproxy reconstruction of past human impact. *The Holocene* 13, 677–690.

Pacl, J., 1973. *Hydrologia Tatranského Národného Parku [Hydrology of the Tatra National Park]*. Zborník Prác o TANAPe 15:181–212.

Pál, I., Buczkó, K., Braun, M., Vincze, I., Pálffy, J., Molnár, M., Finsinger, W., Magyari, E.K., 2016. Small-scale moisture availability increase during the 8.2 ka climatic event inferred from biotic proxy records in the South Carpathians (SE Romania). *The Holocene* 26, 1382–1396.

Pál, I., Buczkó, K., Vincze, I., Finsinger, W., Braun, M., Biró, T., Magyari, E.K., 2018. Terrestrial and aquatic ecosystem responses to early Holocene rapid climate change (RCC) events in the South Carpathian Mountains, Romania. *Quaternary International* 477, 79–93.

Panayotov, M., Gogushev, G., Tsavkov, E., Vasileva, P., Tsvetanov, N., Kulakowski, D., Bebi, P., 2017. Abiotic disturbances in Bulgarian mountain coniferous forests – An overview. *Forest Ecology and Management* 388, 13–28.

Pepin, N., Bradley, R., Diaz, H., Baraer, M., Caceres, E.B., Forsythe, N., Fowler, H., Greenwood, G., Hashmi, M.Z., Liu, X.D., Miller, J.R., Ning, L., Ohmura, A., Palazzi, E., Rangwala, I., Schöner, W., Severskiy, I., Shahgedanova, M., Wang, M.B., Williamson, S.N., Yang, D.Q., 2015. Elevation-dependent warming in mountain regions of the world. *Nature Climate Change* 5, 424–430.

Prentice, I.C., Cramer, W., Harrison, S.P., Leemans, R., Monserud, R.A., Solomon, A.M., 1992. A global biome model based on plant physiology and dominance, soil properties and climate. *Journal of Biogeography* 19, 117–134.

Price, M.F., Gratzner, G., Duguma, L.A., Kohler, T., Maselli, D., Romeo, R., 2011. *Mountain Forests in a Changing World – Realizing Values, Addressing Challenges*. FAO/MPS and SDC, Rome, Italy.

Punt, W., 1976–1996. *The Northwest European Pollen flora 1–7*. Elsevier, Amsterdam.

Reimer, P.J., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Ramsey, C.B., Buck, C.E., Cheng, H., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Hafliðason, H., Hajdas, I., Hatté, C., Heaton, T.J., Hoffmann, D.L., Hogg, A.G., Hughen, K.A., Kaiser, K.F., Kromer, B., Manning, S.W., Niu, M., Reimer, R.W., Richards, D.A., Scott, E.M., Southon, J.R., Staff, R.A., Turney, C.S.M., van der Plicht, J., 2013. IntCal13 and Marine13 radiocarbon age calibration curves 0–50,000 years cal BP. *Radiocarbon* 55(4), 1869–1887.

Renberg, I., Brännvall, M-L., Bindler, R., Emteryd, O., 2000. Atmospheric lead pollution history during four millennia (2000 BC to 2000 AD) in Sweden. *Ambio* 29(3), 150–156.

Rybníčková, E., Rybníček, K., 2006. Pollen and macroscopic analyses of sediments from two lakes in the High Tatra Mountains, Slovakia. *Vegetation History and Archaeobotany* 15, 345–356.

Schaffer, J., and Stummer, F., 1933. *Atlas der Seen der Hohen Tatra, III*. Geographisches Institut der Deutschen Universität Prag. Blätter Popper See und Tschirmer See.

Schillereff, D. N., Chiverrell, R. C., Macdonald, N., Hooke, J. M., 2014. Flood stratigraphies in lake sediments: A review. *Earth Science Reviews* 135, 17–37.

<https://doi.org/10.1016/j.gloplacha.2020.103264>

Schillereff, D. N., Chiverrell, R. C., Macdonald, N., Hooke, J., Welsh, K. E., Piliposian, G., Croudace, I. W., 2019. Convergent human and climate forcing of late-Holocene flooding in Northwest England. *Global and Planetary Change*, 182, 102998.

Schumacher, S., Bugmann, H., 2006. The relative importance of climatic effects, wildfires and management for future forest landscape dynamics in the Swiss Alps. *Global Change Biology* 12, 1435–1450.

Schumacher, M., Schier, W., Schütt, B., 2016. Mid-Holocene vegetation development and herding-related interferences in the Carpathian region. *Quaternary International* 415, 253–267.

Schneider, C.A., Rasband, W.S., Eliceiri, K.W., 2012. NIH Image to ImageJ: 25 years of image analysis. *Nature Methods* 9, 671–675.

Schnitchen, C., Charman, D.J., Magyari, E., Braun, M., Grigorszky, I., Tóthmérész, B., Molnár, M., Szántó, Z., 2006. Reconstructing hydrological variability from testate amoebae analysis in Carpathian peatlands. *Journal of Paleolimnology* 36, 1–17.

Seidl, R., Schelhaas, M.-J., Rammer, W., Verkerk, P.J., 2014. Increasing forest disturbances in Europe and their impact on carbon storage. *Nature Climate Change* 4, 806–810.

Šolcová, A., Hájková, P., Petřík, J., Tóth, P., Rohovec, J., Bátora, J., Horsák, M., 2018. Early and middle Holocene ecosystem changes at the Western Carpathian/Pannonian border driven by climate and Neolithic impact. *Boreas* 47(3), 897–909.

Stähli, M., Finsinger, W., Tinner, W., Allgöwer, B., 2006. Wildland fire history and fire ecology of the Swiss National Park (Central Alps): New evidence from charcoal, pollen and plant macrofossils. *The Holocene* 16, 805–817.

Starkel, L., Michczyńska, D., Krąpiec, M., Margielewski, W., Nalepka, D., Pazdur, A., 2013. Progress in the Holocene chrono- climatostratigraphy of Polish territory. *Geochronometria*. 40(1), 1–21.

Stockmarr, J., 1972. Tablets with spores used in absolute pollen analysis. *Pollen and Spores* 13, 614–621.

Tanțău, I., Feurdean, A., de Beaulieu, J.L., Reille, M., Fărcaș, S., 2011. Holocene vegetation history in the upper forest belt of the Eastern Romanian Carpathians. *Palaeogeography, Palaeoclimatology, Palaeoecology* 309, 281–290.

Temperli, C., Bugmann, H., Elkin, C., 2013. Cross-scale interactions among bark beetles, climate change, and wind disturbances: a landscape modeling approach. *Ecological Monographs* 83, 383–402.

Tinner, W., Conedera, M., Gobet, E., Hubschmid, P., Wehrli, M., Ammann, B., 2000. A palaeoecological attempt to classify fire sensitivity of trees in the southern Alps. *The Holocene* 10, 565–574.

Tinner, W., 2007. Treeline studies, in: Elias, S.A. (Ed.), *Encyclopedia of Quaternary Science*. Elsevier B.V., Amsterdam, The Netherlands, pp. 2374–2384.

Tomlinson, P., 1985. An aid to the identification of fossil buds, bud-scales and catkin-bracts of British trees and shrubs. *Circaea* 3, 45–130.

Topographic maps of Romania, 1986. 1:25000.

<https://doi.org/10.1016/j.gloplacha.2020.103264>

Tóth, M., Magyari, E.K., Buczkó, K., Braun, M., Panagiotopoulos, K., Heiri, O., 2015. Chironomid-inferred Holocene temperature changes in the southern Carpathians (Romania). *The Holocene* 25, 569–582.

Troels-Smith, J., 1955. Karakterisering af løse jordarter. Geological Survey of Denmark, Copenhagen. pp. 1–73.

Trondman, A.-K., Gaillard, M.-J., Mazier, F., Sugita, S., Fyfe, R., Nielsen, A.B., Twiddle, C., Barratt, P., Birks, H.J.B., Bjune, A.E., Björkman, L., Broström, A., Caseldine, C., David, R., Dodson, J., Dörfler, W., Fischer, E., van Geel, B., Giesecke, T., Hultberg, T., Kalnina, L., Kangur, M., van der Knaap, P., Koff, T., Kuneš, P., Lagerås, P., Latałowa, M., Lechterbeck, J., Leroyer, C., Leydet, M., Lindbladh, M., Marquer, L., Mitchell, F.J.G., Odgaard, B.V., Peglar, S.M., Persson, T., Poska, A., Rösch, M., Seppä, H., Veski, S., Wick, L., 2015. Pollen-based quantitative reconstructions of Holocene regional vegetation cover (plant-functional types and land-cover types) in Europe suitable for climate modelling. *Global Change Biology* 21, 676–697.

UNEP, 2007. Carpathians Environmental Outlook. United Nations Environment Programme (UNEP). Geneva, Switzerland, pp. 236.

UNESCO-WNBR. <http://www.unesco.org/new/en/natural-sciences/environment/ecological-sciences/biosphere-reserves/europe-north-america/polandslovakia/tatra/>. Accessed April 29, 2020.

Valde-Nowak, P., and Soják, M., 2010. Contribution to the Mesolithic in the Slovak Carpathians. *Slovenská Archeológia*, 58, 1–12.

Vannière, B., Blarquez, O., Rius, D., Doyen, E., Brücher, T., Colombaroli, D., Connor, S., Feurdean, A., Hickler, T., Kaltenrieder, C., Leys, B., Massa, C., Olofsson, J., 2016. 7000-year human legacy of elevation-dependent European fire regimes. *Quaternary Science Reviews* 132, 206–212.

Vincze, I., Orbán, I., Birks, H.H., Pál, I., Finsinger, W., Hubay, K., Marinova, E., Jakab, G., Braun, M., Biró, T., Tóth, M., Dănuș, C., Ferencz, I.V., Magyari, E.K., 2017. Holocene treeline and timberline changes in the South Carpathians (Romania): Climatic and anthropogenic drivers on the southern slopes of the Retezat Mountains. *The Holocene* 27(11), 1613–1630.

Walther GR., Beißner S., Pott R., 2005. Climate Change and High Mountain Vegetation Shifts. In: Broll G., Keplin B. (eds) *Mountain Ecosystems*. Springer, Berlin, Heidelberg.

Werners, S.E., Bos, E., Civic, K., Hlásny, T., Hulea, O., Jones-Walters, L., Kópataki, E., Kovbasko, A., Moors, E., Nieuwenhuis, E., van de Velde, I., Zingstra, H., Zsuffa, I., 2014. Climate change vulnerability and ecosystem-based adaptation measures in the Carpathian region: Final Report - Integrated assessment of vulnerability of environmental resources and ecosystem-based adaptation measures. Wageningen, Alterra Wageningen UR (University & Research centre), Alterra report 2572. 132 pp.

Wood, S.N., 2017. *Generalized Additive Models: An Introduction with R* (2nd edition). Chapman and Hall/CRC. pp 1–476.

Zasadni, J and Kłapyta, P., 2014. The Tatra Mountains during the Last Glacial Maximum. *Journal of Maps* 10 (3), 440–456.