

1 **Production of a functionally active recombinant SARS-CoV-2 (COVID-19) 3C-Like**
2 **protease and a soluble inactive 3C-like protease-RBD chimeric in a prokaryotic**
3 **expression system**

4 Carolina De Marco Verissimo^{1*}; Jesús López Corrales¹; Amber L. Dorey¹; Krystyna
5 Cwiklinski¹; Richard Lalor¹; Nichola Eliza Davies Calvani¹; Heather L. Jewhurst¹; Andrew
6 Flaus²; Sean Doyle³; John P. Dalton¹.

7
8 1- Molecular Parasitology Lab (MPL) - Centre for One Health and Ryan Institute, School of
9 Natural Science, National University of Ireland Galway, Galway, Republic of Ireland.

10 2- Centre for Chromosome Biology, School of Natural Science, National University of Ireland
11 Galway, Galway, Republic of Ireland

12 3- Department of Biology – Maynooth University, Maynooth, Republic of Ireland.

13
14 *Corresponding author: carolina.verissimo@nuigalway.ie

28 **SUMMARY**

29 During the SARS-CoV-2 intracellular life-cycle, two large polyproteins, pp1a and pp1ab, are
30 produced. Processing of these by viral cysteine proteases, the papain-like protease (PLpro) and
31 the chymotrypsin-like 3C-like protease (3CL-pro) release non-structural proteins necessary for
32 the establishment of the viral replication and transcription complex (RTC), crucial for viral
33 replication. Hence, these proteases are considered prime targets against which anti-COVID-19
34 drugs could be developed. Here, we describe the expression of a highly soluble and functionally
35 active recombinant 3CL-pro using *Escherichia coli* BL21 cells. We show that the enzyme
36 functions in a dimeric form and exhibits an unexpected inhibitory profile because its activity
37 is potently blocked by serine rather than cysteine protease inhibitors. In addition, we assessed
38 the ability of our 3CL-pro to function as a carrier for the Receptor Binding Domain (RBD) of
39 the Spike protein. The co-expressed chimeric protein, 3CLpro-RBD, did not exhibit 3CL-pro
40 activity, but its enhanced solubility made purification easier and improved RBD antigenicity
41 when tested against serum from vaccinated individuals in ELISAs. Chimeric proteins
42 containing the 3CL-pro could represent an innovative approach to developing new COVID-19
43 vaccines.

44

45

46

47

48

49

50

51

52

53

54

55

56

57 INTRODUCTION

58 The severe acute respiratory syndrome coronavirus (SARS-CoV-2) was first identified in
59 Wuhan, China, in December 2019 and subsequently reported throughout the world [1-3].
60 Person-to-person transmission of the virus resulted in rapid distribution of SARS-CoV-2,
61 leading to the unprecedented pandemic of coronavirus disease 2019 (COVID-19), which up to
62 now has claimed >6 million lives [4]. The impact of this pandemic on the global health and
63 economy prompted the rapid action on the development, testing and approval of prophylactic
64 COVID-19 vaccines, followed by mass immunization programs [5].

65 SARS-CoV-2 is an enveloped virus that contains a single-strand of positive-sense
66 RNA. Infection begins when the virus attaches to cells via the angiotensin converting enzyme
67 2 (ACE2) receptor, mediated by the receptor binding domain (RBD) of the major glycoprotein
68 expressed on the virus surface, the Spike protein [6, 7]. Fusion of the viral membrane with the
69 lumen of the endosomal membrane leads to endocytosis, facilitating infection via entry of the
70 viral RNA into the cytosol. Applying new approaches and technologies, a multitude of vaccines
71 have been developed, four of which were licensed by the regulatory agencies [4] and have been
72 administered across the world, representing a relief and a unique opportunity to prevent the
73 deaths of millions of people and control the pandemic. In general, each of these four vaccines
74 induce antibodies against the Spike protein and bind to the RBD to block its interaction with
75 ACE2 [8, 9].

76 During the intracellular viral life cycle, two large polyproteins, pp1a and pp1ab, are
77 translated. Sixteen non-structural proteins (nsp) are co-translationally and post-translationally
78 released from pp1a and pp1ab upon proteolytic activity of two virus cysteine proteases, the
79 papain-like protease (PLpro) and the chymotrypsin-like 3C-like protease (3CL-pro), also
80 known as the main protease. These proteases allow the establishment of the viral replication
81 and transcription complex (RTC), which is crucial for virus replication inside the cells [10].
82 The 3CL-pro plays a prominent role on viral gene expression and replication [11, 12].
83 Moreover, recent studies comparing the protease from other coronaviruses, SARS-CoV and
84 Middle East Respiratory Syndrome CoV (MERS-CoV), or from picornaviruses, show that they
85 are all highly conserved in terms of proteolytic activity and structure [13-15] and revealed
86 important immunomodulatory properties for this enzyme. Amongst other mechanisms, during
87 viral infection 3CL-pro contributes to the delay of host anti-viral innate immune response by
88 cleaving or inactivating key elements of the Retinoic acid-inducible gene I (RIG-I) like
89 receptors (RLRs)-mediated Type I interferon (INF-I) signalling pathway, which allows
90 effective viral infection and contribute for disease progression and severity [15-17]. Thus, the

91 3CL-pro could be considered an attractive target for the development of future anti-COVID-
92 19 treatments.

93 Here we describe the production of a recombinant 3CL-pro in a prokaryotic expression
94 system and its purification as a highly soluble and functionally active protease. We also
95 generated a 3C-like protease-RBD gene construct that enabled the production of a chimeric
96 protein, named 3CLpro-RBD. This strategy proved useful to enhance the solubility and
97 antigenicity of the RBD, albeit the recombinant chimeric protein did not exhibit proteolytic
98 activity, understandable since the functional 3CL-pro functions as a dimer.

99

100

101 **METHODS**

102 **Ethical statement**

103 Human experimental work was conducted according to Human Research Ethics Committees.
104 Sera samples from individuals double-vaccinated with Pfizer/BioNTech (BNT162b2) vaccine
105 were obtained from healthy volunteers following ethical approval by the National University
106 of Ireland Galway, Ireland, research ethics committee (R20.Jun.06). The samples were pooled
107 and immediately stored at -80°C. All participants provided written informed consent prior to
108 the study. Negative control samples obtained from the Irish Blood Transfusion Service. These
109 blood samples were previously characterized by De Marco Verissimo et al. [18].

110

111 **Recombinant protein production in *Escherichia coli* cells and purification**

112 Sequences encoding the 3CL-pro and RBD proteins were codon optimized for expression in
113 *Escherichia coli* and cloned into the pET-28a(+) vector (Genscript Biotech). The chimeric
114 protein 3CLpro-RBD was produced by generating a gene construct that linked the 3CL-pro and
115 RBD genes by a bridge sequence that encoded for glycine-proline triple repeat (GPGPGP) (see
116 Figure 1). The recombinantly produced proteins contain a thrombin cleavage site followed by
117 a C-terminal His-tag. The synthesized vectors were transformed into BL21 competent *E. coli*
118 cells (ThermoFisher Scientific) following the manufacturer's instructions and stored in Luria
119 Bertani (LB) broth (Sigma-Aldrich) supplemented with 25% glycerol at -80°C. LB broth
120 supplemented with 50 µg/mL kanamycin was inoculated from the glycerol stock and incubated
121 shaking (200 rpm) at 37°C overnight. The culture was then diluted in fresh LB broth
122 supplemented with kanamycin, incubated at 37°C to OD₆₀₀ 0.6 and protein expression induced

123 with 1 mM isopropyl- β -D-1-thiogalactopyranoside (IPTG; ThermoFisher Scientific) for 4 hr
124 at 30°C (3CL-pro and RBD); 18 h at 16°C (3CLpro-RBD chimera). Following centrifugation at
125 10,000 x g for 10 min at 4°C, the bacterial pellets were re-suspended in 10 mL ST buffer (10
126 mM Tris, 150 mM NaCl, pH 8.0).

127 The bacteria pellets were treated with lysozyme (10 μ g/mL), sonicated on ice (6 x 10
128 seconds, 40% amplitude) and centrifuged 15,000 x g at 4°C for 30 min. The soluble
129 recombinant protein within the supernatant was purified and dialysed using the Profinia
130 Affinity Chromatography Protein Purification System (Bio-Rad), with the mini profinity
131 IMAC and mini Bio-Gel P-6 desalting cartridges (Bio-Rad). The protein concentration and
132 purity were verified by Bradford Protein Assay (Bio-Rad) and by 4-20% SDS-PAGE gels (Bio-
133 Rad) stained with Biosafe Coomassie (Bio-Rad), respectively. The gels were visualised using
134 a G:BOX Chemi XRQ imager (Syngene).

135 As RBD protein was found within the inclusion bodies, processing of the pellets,
136 protein purification and dialysis were performed as described by Schlager et al. [19] and
137 employed by us previously to extract recombinant SARS-CoV-2 proteins previously [18].
138 Briefly, 1% (w/v) SDS buffer (8 mM Na₂HPO₄, 286 mM NaCl, 1.4 mM KH₂PO₄, 2.6 mM KCl,
139 1% (w/v) SDS, pH 7.4) containing 0.1 mM DTT was added to the cell pellet to solubilize the
140 inclusion bodies. After sonication, the samples were centrifuged 15,000 x g at 4°C for 30 min
141 and the resulting supernatant containing the target protein was filtered and purified using a pre-
142 equilibrated Ni-NTA beads column (Qiagen). The recombinant protein was eluted using 4 mL
143 of elution buffer (8 mM Na₂HPO₄, 286 mM NaCl, 1.4 mM KH₂PO₄, 2.6 mM KCl, 0.1%
144 Sarkosyl (w/v), 250 mM imidazole, pH 7.4) and buffer-exchanged into 1x PBS containing
145 0.05% sarkosyl, pH 7.4.

146

147 **Size exclusion chromatography**

148 The purified recombinant 3CL-pro purified was additionally subjected to size-exclusion (gel
149 filtration) chromatography to resolve its dimerization state. The purification was performed
150 using a high performance Superdex 75 10/300 GL (Tricorn) column, with a flow rate of 400
151 μ L/min and eluted into 1x PBS. Three known proteins of different molecular sizes were
152 resolved in the column as standards, namely conalbumin (75 kDa), carbonic anhydrase (29
153 kDa) and aprotinin (6.5 kDa) (Sup Figure S1). Once the retention parameters were determined,
154 the r3CL-pro, at 1 mg/mL in PBS, was added to the column for purification. Aliquots of 200

155 μL of the sample were collected and stored at 4°C for further analysis using an activity assay
156 (see below).

157

158 **Fluorogenic assay to assess the enzymatic activity of the recombinant 3CL-pro**

159 The enzymatic activity of the recombinant r3CL-pro purified by affinity chromatography and
160 of the different fractions produced by gel filtration were verified using a fluorogenic assay
161 using the substrate LGS AVLQ-rhodamine 110-dp (BostonBiochem). Unless highlighted, all
162 the screening assays were performed at 37°C , in a $100\ \mu\text{L}$ reaction volume Hepes buffer (20
163 mM Hepes, 2 mM EDTA, pH 7.4). Initially, the reaction buffer was mixed with either of the
164 recombinant proteins, r3CL-pro (500 nM), rRBD (500 nM) or r3CLpro-RBD chimera (500 nM),
165 and incubated for 5 min at room temperature. The fluorogenic substrate ($20\ \mu\text{M}$) was added to
166 the wells and the proteolytic activity was measured at 37°C , over 1 h, as relative fluorescent
167 units (RFU) in a PolarStar Omega Spectrophotometer (BMG LabTech). All assays were carried
168 out in triplicate. Commercial broad-spectrum protease inhibitors, namely serine protease
169 inhibitors AEBSF (5 mM; Sigma-Aldrich) and Futhan-175 (FUT-175, $200\ \mu\text{M}$; BD-
170 Pharmingen-Bioscience), and the cysteine protease inhibitor E-64 ($200\ \mu\text{M}$; Sigma-Aldrich),
171 were added to the reaction, individually, for further characterization of the proteolytic activity
172 of the recombinant r3CL-pro.

173

174 **Assessment of the immunogenicity of the r3CL-pro, the rRBD and the chimeric r3CLpro-** 175 **RBD**

176 Seven weeks-old male and female CD1 outbred mice were used to assess the immunogenicity
177 of the recombinantly-produced proteins according to the schedule shown in Figure 2. All
178 animal experimental procedures were carried out by Eurogentec, BE, as follows: Group 1,
179 adjuvant control group (Montanide ISA 206VG, Seppic) (n= 9); Group 2, r3CLpro-RBD
180 chimera ($15\ \mu\text{g}$) formulated in the Montanide adjuvant (1:1 v/v) (n=10); Group 3, r3CL-pro ($15\ \mu\text{g}$)
181 formulated in the Montanide adjuvant (1:1 v/v) (n=7); Group 4, rRBD ($15\ \mu\text{g}$) formulated
182 in the Montanide adjuvant (1:1 v/v) (n=7); Group 5, r3CL-pro + rRBD ($7.5\ \mu\text{g}$ of each)
183 formulated in the Montanide adjuvant (1:1 v/v) (n=7).

184

185 **ELISA to assess antibodies against the recombinant SARS-CoV-2 proteins in serum of** 186 **vaccinated humans**

187 Flat-bottom 96 well microtitre plates (Nunc MaxiSorp, Biolegend) were coated with r3CL-pro,
188 r3CLpro-RBD chimer, rRBD or cmRBD as described above. After incubation in blocking
189 buffer (2% BSA in PBS 0.05% Tween-20 (v/v), pH 7.4, PBST) and washing steps, pooled
190 serum samples from (a) 10 vaccinated individuals (collected at least 10 days after the second
191 dose Pfizer/BioNTech (BNT162b2) vaccine), or from 10 negative controls individuals
192 (samples from the Irish Blood Transfusion Service obtained before COVID-19 pandemic) were
193 diluted 1:100 in blocking buffer and added to the plate. After 1 h incubation at room
194 temperature (RT), and washing five times with PBST, the secondary antibody HRP anti-
195 Human IgG (Fc specific) (Sigma-Aldrich) was added (1:15,000), and the plates incubated for
196 1 h at RT. After washing five times, TMB (3,3',5,5'-Tetramethylbenzidine Liquid Substrate
197 Supersensitive, Sigma-Aldrich) substrate was added to each well. Following a three-minute
198 incubation the reaction was stopped with 2 N sulphuric acid and plates read at 450 nm in a
199 PolarStar Omega Spectrophotometer. All samples were analysed in triplicate.

200

201 **Analysis of the immune response of mice to the recombinant 3CL-pro and r3CLpro-RBD** 202 **chimer by ELISA.**

203 The antibody response of individual mouse serum at day 0 and day 35 (Figure 2) was assessed
204 by ELISA using r3CL-pro and r3CLpro-RBD chimer as antigens. Flat-bottom 96 well
205 microtitre plates (Nunc MaxiSorp, Biolegend) were coated overnight at 4°C with either r3CL-
206 pro (2 µg/mL) or r3CLpro-RBD chimer (2 µg/mL) diluted in carbonate buffer (pH 9.6). After
207 incubation in blocking buffer (2% BSA in PBS-0.05% Tween-20 (v/v), pH 7.4; PBST) and
208 washing steps, mice serum diluted 1:100 in blocking buffer was added to the antigen-coated
209 wells and incubated for 1 hr at RT. After washing five times with PBST, the secondary antibody
210 HRP goat to mouse-anti-IgG (ThermoFisher Scientific) was added (1:10,000), and the plates
211 incubated for 1 h at RT. After washing five times, TMB substrate was added to each well.
212 Following a three-minute incubation the reaction was stopped with 2 N sulphuric acid and
213 plates read at 450 nm in a PolarStar Omega Spectrophotometer. All samples were analysed in
214 triplicates.

215

216

217 **RESULTS**

218 **Production of SARS-CoV-2 recombinant proteins in *E. coli***

219 The 3C-like protease (r3CL-pro) was readily produced as a recombinant protein in *E. coli*;
220 analysis of bacterial lysate showed that it was a prominent protein that separated into the
221 soluble fraction making it easy to isolate by affinity chromatography. The purified protein
222 resolved at the expected molecular size of ~34 kDa, as a highly soluble protein, and our
223 purification yielded 5.3 mg enzyme per litre of bacterial culture (Figure 3A).

224 By marked contrast, we found that rRBD did not extract with the solubilisation buffers
225 used but remained in the insoluble pellet, presumably in inclusion bodies. Accordingly, we
226 employed an alternative means of solubilisation that included the chaotropic detergent sodium
227 dodecyl sulphate (SDS) in the buffer, which proved successful in extracting the protein from
228 the pellet [18, 19]. After this extraction procedure, the recombinant RBD could be isolated by
229 NTA-affinity chromatography (Figure 3B). The purified ~29 kDa protein remained soluble
230 after dialysis against PBS containing 0,05% sarkosyl to remove the SDS detergent. This yielded
231 ~1.5 mg of protein per litre of bacterial culture.

232 By expressing the 3CL-pro and RBD proteins as a chimera (Figure 1), 3CLpro-RBD
233 (~60 kDa), we found that the recombinant protein exhibits intermediate solubility to the protein
234 expressed alone and, therefore, we were able to purify the chimera using the same automated
235 protocol adopted with the r3CL-pro. It provided a yield of ~1.2 mg per litre of bacterial culture
236 (Figure 3B).

237 Further confirmation that we had purified the targeted proteins was obtained by western
238 blot analysis, where we probed the purified proteins with antibody to the His-tag present on all
239 three recombinant proteins (Figure 3C).

240

241 **Recombinant 3CL-pro of SARS-CoV-2 is functionally active in dimeric form, while the** 242 **chimeric protein does not exhibit activity**

243 Once we successfully produced a soluble recombinant 3CL-pro, we proceeded to check its
244 proteolytic activity using an appropriate substrate. Since the cleavage site of 3CL-pro is highly
245 unique the commercially available LGS AVLQ-rhodamine 110-dp substrate could be utilized
246 to specifically assay the activity of the recombinant enzyme. The assay revealed that our
247 recombinant enzyme is a functionally active protease at 37°C in neutral pH (Figure 4).
248 Unexpectedly, however, despite 3CL-pro being described as a cysteine protease, we found that
249 the enzyme was not inhibited by the cysteine proteases inhibitors such as E-64, but was
250 susceptible to two broad-spectrum serine protease inhibitors, AEBSF and Futhan-175; the

251 enzyme was completely inhibited by these latter compounds at concentrations of 5 mM and
252 200 μ M, respectively (Figure 4).

253 Using gel filtration to further purify the functionally active recombinant r3CL-pro we
254 were able to determine the presence of a mixture of dimers and oligomers within the product
255 purified by affinity chromatography (Peak 1 and 2, respectively; Figure 5). The importance of
256 the dimerization for proteolytic activity of the r3CL-pro was determined by assaying the
257 individual fractions within the two main protein peaks detected during purification. Our data
258 revealed that the r3CL-pro is only functionally active when in its dimeric form, which
259 represents the predominant peak in the chromatogram obtained (Figure 5).

260 In order to examine if the expression of the 3CL-pro in a chimeric format with the RBD,
261 3CLpro-RBD chimer, had an effect on its proteolytic activity, we also assayed the activity of
262 the recombinant chimeric protein and rRBD produced. Neither protein exhibited enzymatic
263 activity when assayed in the same conditions used with the r3CL-pro (Figure 4).

264

265 **Antibodies in serum of naturally infected and vaccinated individuals recognize r3CLpro-** 266 **RBD chimeric protein.**

267 In order to determine if our recombinant SARS-CoV-2 proteins had common epitopes with
268 those present in the virus or with the viral proteins expressed upon vaccination, we performed
269 ELISA tests with sera from vaccinated humans using our recombinant proteins r3CL-pro,
270 r3CLpro-RBD chimer and rRBD as target antigens. In parallel, we used the commercial RBD
271 recombinantly produced in mammalian cells (cmRBD), which is commonly used in
272 immunological and functional assays [20]. Our results show that, when compared to negative
273 control samples, sera from vaccinated individuals contain antibodies that recognise the
274 r3CLpro-RBD chimer. These individuals also recognized the rRBD, albeit with lower intensity
275 (Figure 6). Surprisingly, a discrete antibody response against the r3CL-pro was also observed
276 with the vaccinated group (Figure 6). In addition, comparing the antibody response of
277 individuals naturally infected using nucleocapsidic (Npro) and S2Frag proteins as target
278 antigens in parallel, we verified that 50% of the individuals analysed mounted a significant
279 antibody response to 3CL-pro (Supplementary Fig S2).

280

281 **r3CL-pro and r3CLpro-RBD chimer induce antibody response in vaccinated mice.**

282 In order to assess and compare the immunogenicity of the r3CL-pro and r3CLpro-RBD chimera,
283 we immunized outbred CD1 mice with each protein in a regime similar to that initially
284 recommended for the available COVID-19 vaccines (i.e., initially the recommendation for
285 Pfizer and Oxford vaccines was 2 doses administered 3 weeks apart, Figure 2) [8, 9]. Groups
286 of CD1 mice were immunized with each of the proteins (15 µg) or with adjuvant Montanide
287 ISA 206VG alone, as base-line controls, which was used to formulate the preparations with the
288 recombinant proteins. Pre-immune (Day 0) and immune sera (Day 35) of each animal was
289 assessed for antibodies using ELISA tests.

290 Our ELISA results show that those animals immunized with adjuvant alone did not react to any
291 recombinant protein (Figure 7). Mice immunized with r3CL-pro responded with high levels of
292 antibodies against the r3CL-pro but their antibodies did not bind well to the r3CLpro-RBD
293 chimeric (Figure 7A and B). Conversely, mice immunized with the r3CLpro-RBD chimeric
294 produced with antibodies that bound to the chimeric protein in ELISA, but elicited a low level
295 response to the r3CL-pro (Figure 7A).

296

297

298 **DISCUSSION**

299 The 3C-like protease is regarded as a prime target for therapeutic drug treatment of
300 COVID-19 due to its unique specificity for cleaving peptide bonds that are absent in human
301 proteins [14, 21, 22]. This protease plays a central role during viral replication, being
302 responsible for the cleavage of 11 sites within the polyprotein 1ab, ultimately releasing 13 non-
303 structural proteins involved in SARS-CoV-2 replication inside the host's cells [23, 24]. Such
304 activity is associated with the ability of this protease to recognize and cleave the unique peptide
305 sequence Leu/Phe/Met-Gln ↓ Gly/Ser/Ala (↓ denotes the cleavage site). We were able to
306 demonstrate that the r3CL-pro produced in the present study using bacterial expression systems
307 has the same requirement for proteolytic activity, including for a glutamine (Gln) at the P1
308 position, as the recombinant enzyme was able to cleave specifically the LGSAVLQ-Rh110
309 substrate. Since 3CL-pro is reported to be functionally active as a dimer [25, 26], we further
310 investigated its molecular state using size-exclusion chromatography and determined that the
311 purified product was dominated by r3CL-pro dimers. Together with the results of the activity
312 assays, our data indicate that the production of this enzyme in the *E. coli* system employed here
313 allowed for protein fold and dimerization similar to that of the native form. The unexpected
314 inhibition of this protease by broad-spectrum serine protease inhibitors, but not by cysteine

315 protease inhibitors, does not support the claim that this protease is a cysteine peptidase, but is
316 more in keeping with the reported chymotrypsin-like protease activity of 3CL-pro [10, 27].
317 Perhaps this observation could be exploited in anti-3CL-pro novel drug designing.

318 In our previous study aimed at improving COVID-19 diagnostics [18] by using the
319 Npro and S2Frag proteins as parallel target antigens on ELISA tests, we also evaluated the
320 antibody response of individuals naturally infected with SARS-CoV-2 against the r3CL-pro.
321 Our data showed that, although higher antibodies titres are generated against Npro and S2frag,
322 50% of the individuals analysed had significant levels of antibodies against 3CL-pro,
323 indicating that this protease is naturally immunogenic during COVID-19 infection
324 (Supplementary Fig S2). As expected, individuals vaccinated against COVID-19 with the
325 currently available vaccines do not raise a significant antibody response against 3CLpro
326 (Figure 6). Nonetheless, in the present study, we demonstrated that the recombinant version of
327 this protein is immunogenic and induces a strong and specific antibody response in mice.
328 Together, this data indicates the potential application of the 3CL-pro as a vaccine target against
329 COVID-19. Inducing antibodies to the 3CL-pro could provide broader antibody and cellular
330 immunity to individuals and thus induce a stronger protection against SARS-CoV-2 infection
331 [15, 16].

332 Given the difficulty in producing recombinant soluble SARS-CoV-2 proteins in
333 prokaryotic systems [18, 28, 29], we considered that the highly soluble 3CL-pro could serve as
334 a carrier protein to produce and deliver RBD. Thus, we designed a construct to generate a
335 recombinant chimeric protein, 3CLpro-RBD. The two proteins were linked using a GP-linker
336 that allows separation and flexibility between them, so that each molecule is stable and can
337 function separately. This approach improved the solubility and production of the RBD.
338 However, we observed that the functional activity of the r3CL-pro chimeric was lost, likely
339 due to incorrect protease folding or the inability of the 3CL-pro to form dimers when linked to
340 the RBD.

341 Surprisingly, our ELISA tests revealed that CD1 mice immunized with the chimeric
342 protein developed antibodies against the 3CL-pro portion of the antigen, but not to the RBD
343 part. Nonetheless, antibodies in serum from individuals fully-vaccinated against COVID-19
344 recognized the chimeric protein more efficiently than the rRBD alone, indicating that the
345 presentation of the RBD is improved when produced in the chimeric format. Furthermore, the
346 ability of vaccinated individuals to recognise the commercial RBD produced in mammalian
347 cells (cmRBD), but not our rRBD produced in *E. coli* cells, suggests that glycosylated epitopes

348 are important in antibody recognition. Based on current data, antibodies that bind to the RBD
349 can neutralize the SARS-CoV-2 virus by preventing it binding to ACE2 receptors and
350 consequent host cells invasion and infection [22, 30-32]. A chimeric protein carrying the RBD
351 certainly could help to circumvent the low cellular immunogenicity problem faced with
352 vaccines, and the combination of RBD with more immunogenic molecules has been
353 demonstrated to be a useful strategy to create alternative vaccine targets to fight COVID-19
354 [28, 29, 33]. This is the first time that another SARS-CoV-2 molecule, and specifically the
355 3CL-pro, has been considered as an option to fuse with the antigenic RBD, which could
356 represent a future strategy to develop COVID-19 vaccine targets, as the 3CL-pro could have a
357 superior ability to induce long-term neutralizing antibody responses, as well as potent cellular
358 immunity, and also overcome the problems with the SARS-CoV-2 variants that are mainly
359 associated high plasticity of the Spike protein [34-36]. The concept of chimerization of SARS-
360 CoV-2 molecules that we introduced in this study might open new avenues on the discovery of
361 drug and vaccine targets to fight COVID-19.

362

363

364 **CONCLUSIONS**

365 Here we present a straightforward, efficient and cheap method to express and purify a
366 highly soluble and functional 3C-like protease, which is regarded as a main drug target at which
367 to develop therapies against SARS-CoV-2 infection. The enzyme could be useful in the
368 development of high-throughput assays for the screening and isolation of new anti-COVID-19
369 compounds. In addition, given its solubility and potential for triggering a cellular immune
370 response to fight infection, we introduced the idea of using the 3CL-pro as a carrier to other
371 SARS-CoV-2 proteins, in this case RBD, to improve their expression and delivery as potential
372 vaccines. Co-expressing 3CL-pro and RBD in a chimeric format resulted in loss of the 3CL-
373 pro activity (likely due to the inability of the enzyme domain to dimerise), but enhanced the
374 solubility of the RBD expressed alone, and improved its antigenic properties. Chimeric proteins
375 containing the 3CL-pro could represent a new approach to engender next generation protein-
376 subunit COVID-19 vaccine candidates.

377

378

379 **FINANCIAL SUPPORT**

380 This work was supported by the Science Foundation Ireland (SFI) COVID-19 Rapid Response
381 Funding Call, proposal ID 20/COV/0023 and 20/COV/0048.

382

383 **CONFLICT OF INTEREST**

384 None.

385

386 **DATA AVAILABILITY AND SUPPLEMENTARY FILE**

387 The data that support the findings of this study are all included in the publication.
388 Supplementary file data archive available on the Cambridge University Press - Cambridge Core
389 website.

390

391 **REFERENCES**

- 392 1. Sun B, et al Kinetics of SARS-CoV-2 specific IgM and IgG responses in COVID-19 patients.
393 *Emerging Microbes & Infections*. 2020;9(1):940-48.
- 394 2. Meo SA, et al. Omicron SARS-CoV-2 new variant: global prevalence and biological and
395 clinical characteristics. *European Review for Medical and Pharmacological Sciences*.
396 2021;25(24):8012-18.
- 397 3. Aleem A, Akbar Samad AB, Slenker AK Emerging Variants of SARS-CoV-2 And Novel
398 Therapeutics Against Coronavirus (COVID-19). In: *StatPearls*. Treasure Island (FL):
399 StatPearls Publishing Copyright © 2022, StatPearls Publishing LLC., 2022.
- 400 4. World Health Organization. Coronavirus disease (COVID-19) pandemic
401 (<https://covid19.who.int/>). 2022. Accessed May 20th 2022.
- 402 5. Coronavirus (COVID-19) Vaccinations: Our World in Data, 2022.
403 (<https://ourworldindata.org/covid-vaccinations>). 2022. Accessed May 22nd 2022.
- 404 6. Letko M, Marzi A, Munster V Functional assessment of cell entry and receptor usage for
405 SARS-CoV-2 and other lineage B betacoronaviruses. *Nature Microbiology*. 2020;5(4):562-69.
- 406 7. Holmes KV SARS-associated coronavirus. *New England Journal of Medicine*.
407 2003;348(20):1948-51.
- 408 8. Polack FP, et al. Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *New*
409 *England Journal of Medicine*. 2020;383(27):2603-15.
- 410 9. Voysey M, et al Safety and efficacy of the ChAdOx1 nCoV-19 vaccine (AZD1222) against
411 SARS-CoV-2: an interim analysis of four randomised controlled trials in Brazil, South Africa,
412 and the UK. *Lancet*. 2021;397(10269):99-111.
- 413 10. V'Kovski P, et al Coronavirus biology and replication: implications for SARS-CoV-2.
414 *Nature Reviews Microbiology*. 2021;19(3):155-70.
- 415 11. Gordon CJ, et al The antiviral compound remdesivir potently inhibits RNA-dependent
416 RNA polymerase from Middle East respiratory syndrome coronavirus. *Journal of Biological*
417 *Chemistry*. 2020;295(15):4773-79.
- 418 12. Dai W, et al Structure-based design of antiviral drug candidates targeting the SARS-CoV-
419 2 main protease. *Science*. 2020;368(6497):1331-35.

- 420 13. Anand K, et al Coronavirus main proteinase (3CLpro) structure: basis for design of anti-
421 SARS drugs. *Science*. 2003;300(5626):1763-67.
- 422 14. Zhang CH, et al. Potent Noncovalent Inhibitors of the Main Protease of SARS-CoV-2 from
423 Molecular Sculpting of the Drug Perampanel Guided by Free Energy Perturbation
424 Calculations. *ACS Central Science*. 2021;7(3):467-75.
- 425 15. Ng CS, Stobart CC, Luo H. Innate immune evasion mediated by picornaviral 3C protease:
426 Possible lessons for coronaviral 3C-like protease? *Reviews in Medical Virology*. 2021;31(5):1-
427 22.
- 428 16. Blanco-Melo D, et al SARS-CoV-2 launches a unique transcriptional signature from in
429 vitro, ex vivo, and in vivo systems. *bioRxiv*. 2020:004655.
- 430 17. Lokugamage KG, et al Type I interferon susceptibility distinguishes SARS-CoV-2 from
431 SARS-CoV. *Journal of Virology*. 2020;94(23):e01410-20.
- 432 18. De Marco Verissimo C, et al. Improved diagnosis of SARS-CoV-2 by using nucleoprotein
433 and spike protein fragment 2 in quantitative dual ELISA tests. *Epidemiology and Infection*.
434 2021;149:e140.
- 435 19. Schlager B, Straessle A, Hafen E Use of anionic denaturing detergents to purify insoluble
436 proteins after overexpression. *BMC Biotechnology*. 2012;12:95.
- 437 20. Huang WC, et al. SARS-CoV-2 RBD Neutralizing Antibody Induction is Enhanced by
438 Particulate Vaccination. *Advanced Materials*. 2020;32(50):e2005637.
- 439 21. Jukič M, et al Prioritisation of Compounds for 3CL(pro) Inhibitor Development on SARS-
440 CoV-2 Variants. *Molecules (Basel, Switzerland)*. 2021;26(10):3003.
- 441 22. Chen YW, Yiu C-PB, Wong K-Y. Prediction of the SARS-CoV-2 (2019-nCoV) 3C-like
442 protease (3CL pro) structure: virtual screening reveals velpatasvir, ledipasvir, and other drug
443 repurposing candidates. *F1000Research*. 2020;9:129.
- 444 23. Froggatt HM, Heaton BE, Heaton NS Development of a Fluorescence-Based, High-
445 Throughput SARS-CoV-2 3CL(pro) Reporter Assay. *Journal of Virology*. 2020;94(22).
- 446 24. Jaskolski M, et al. Crystallographic models of SARS-CoV-2 3CL(pro): in-depth assessment
447 of structure quality and validation. *IUCrJ International Union of Crystallography*. 2021;8(Pt
448 2):238-56.
- 449 25. Chen S, et al Residues on the dimer interface of SARS coronavirus 3C-like protease: dimer
450 stability characterization and enzyme catalytic activity analysis. *Journal of Biochemistry*.
451 2008;143(4):525-36.
- 452 26. Goyal B, Goyal D Targeting the Dimerization of the Main Protease of Coronaviruses: A
453 Potential Broad-Spectrum Therapeutic Strategy. *ACS Combinatorial Science*. 2020;22(6):297-
454 305.
- 455 27. Molavi Z, et al Identification of FDA approved drugs against SARS-CoV-2 RNA
456 dependent RNA polymerase (RdRp) and 3-chymotrypsin-like protease (3CLpro), drug
457 repurposing approach. *Biomedicine and Pharmacotherapy*. 2021;138:111544.
- 458 28. Bellone ML, et al Production in *Escherichia coli* of recombinant COVID-19 spike protein
459 fragments fused to CRM197. *Biochemical and Biophysical Research Communications*.
460 2021;558:79-85.
- 461 29. Tan TK, et al A COVID-19 vaccine candidate using SpyCatcher multimerization of the
462 SARS-CoV-2 spike protein receptor-binding domain induces potent neutralising antibody
463 responses. *Nature Communications*. 2021;12(1):542-42.
- 464 30. Weinreich DM, et al REGN-COV2, a Neutralizing Antibody Cocktail, in Outpatients with
465 Covid-19. *New England Journal of Medicine*. 2020;384(3):238-51.
- 466 31. Hoffmann H-H, et al Functional interrogation of a SARS-CoV-2 host protein interactome
467 identifies unique and shared coronavirus host factors. *Cell Host & Microbe*. 2021;29(2):267-
468 80. e5.
- 469 32. Chen P, et al SARS-CoV-2 Neutralizing Antibody LY-CoV555 in Outpatients with Covid-
470 19. *New England Journal of Medicine*. 2020;384(3):229-37.


471 33. Hong SH, et al Immunization with RBD-P2 and N protects against SARS-CoV-2 in
 472 nonhuman primates. *Science Advances*. 2021;7(22):eabg7156.
 473 34. Gómez CE, Perdiguero B, Esteban M Emerging SARS-CoV-2 Variants and Impact in
 474 Global Vaccination Programs against SARS-CoV-2/COVID-19. *Vaccines (Basel)*. 2021;9(3).
 475 35. Plante JA, et al Spike mutation D614G alters SARS-CoV-2 fitness. *Nature*.
 476 2021;592(7852):116-21.
 477 36. Oude Munnink BB, et al The next phase of SARS-CoV-2 surveillance: real-time molecular
 478 epidemiology. *Nature Medicine*. 2021;27(9):1518-24.

479

480

481 **LEGENDS FOR FIGURES**


482 **Figure 1. Primary sequence of the SARS-CoV-2 proteins and schematic representation**
 483 **of the 3CLpro-RBD chimeric protein structure.** A: The amino acid sequence of the SARS-
 484 CoV-2 3C-like protease (3CL-pro) used for recombinant expression in *Escherichia coli*. B: The
 485 amino acid sequence of the receptor binding domain (RBD) (residues 319-542 of the full
 486 SARS-CoV-2 Spike protein). C: Schematic representation of the 3CLpro-RBD chimeric
 487 protein structure showing the unique GP linker. SARS-CoV-2 proteins, 3CL-pro and the RBD
 488 are linked by a GP triplet (Glycine, G, and Proline, P), allowing their expression as a stable
 489 chimeric protein. Thr: Thrombin cleavage site; 6His: Histidine tag added to the protein C-
 490 terminal.



491

492



493 **Figure 2. Graphical schematic showing the schedule for the immunization of CD1**
 494 **outbred mice using the recombinant SARS-CoV-2 proteins.** Red arrows indicate when the
 495 mice were immunized with adjuvant alone or with the recombinant 3CL-pro and 3CLpro-RBD
 496 chimer. Blue arrow indicates the end of the experiment and when the final bleeding was taken.



497

498


499 **Figure 3. Recombinant expression of the SARS-CoV-2 proteins, 3C-like protease,**
 500 **receptor binding domain (RBD), and 3CLpro-RBD chimer.** A: Purification of recombinant
 501 3C-like protease. The supernatant after bacterial pellet digestion (1); proteins that did not bind
 502 to the column in the run through (2); proteins in the wash (3); purified and dialysed recombinant
 503 protein (3CL-pro). B: The proteins were recombinantly expressed in the prokaryotic expression
 504 system, *E. coli*, purified and resolved in SDS-PAGE at the expected respective molecular size:
 505 RBD, ~29 kDa; 3CL-pro, ~34 kDa; 3Cpro-RBD chimer, ~60 kDa. C: Western blot of the
 506 recombinant proteins probed with the monoclonal anti-6Histidine tag antibody. M: Molecular
 507 weight in kilodaltons.



508

509


510 **Figure 4. Enzymatic activity of the SARS-CoV-2 recombinant proteins.** The enzymatic
 511 activity of the r3C-Like protease (r3CL-pro, 500 nM) was tested with or without various broad-
 512 spectrum protease inhibitor, namely the serine protease inhibitors AEBSF (5 mM) and Futhan-
 513 175 (FUT-175, 200 μ M), and the cysteine protease inhibitor E-64 (200 μ M). The activity of
 514 the r3CLpro-RBD chimer (500 nM) and of the receptor binding domain (rRBD; 500 μ M) was
 515 assessed in parallel using the same substrate, LGS AVLQ-Rh110 (20 μ M). Enzymatic activity
 516 presented as relative fluorescence units (RFU) at 485-535 nm. Error bars indicate standard
 517 deviation of three separate experiments.



518

519


520 **Figure 5. Gel filtration chromatography of the recombinant 3CL-pro.** The chromatogram
 521 of the purification of the r3CL-pro by gel filtration. Peak 1 (light blue), appeared at 11.918 mL
 522 was calculated to represent a protein of ~ 60.9 kDa, while the Peak 2 (light blue), at 12.74 mL,
 523 represents a protein of ~ 42 kDa, which indicates the presence of 3CL-pro as a dimer and a
 524 oligomer, respectively (for the protein standards data see Supplementary Figure S1). The
 525 enzymatic activity of each fraction within the peaks (n = 17, in red) was determined in relation
 526 to the activity detected with the r3CL-pro purified only by affinity chromatography. In yellow,
 527 the three fractions where enzymatic activity was detected. Black arrows indicate the retention
 528 (mL) for the standards conalbumin (11.37) and carbonic anhydrase (13.50). The complete
 529 chromatogram for the standards is presented in the Supplementary Fig S2.



530

531



532 **Figure 6. Immune recognition of the recombinant SARS-CoV-2 proteins by antibodies in**
533 **sera from COVID-19 fully-vaccinated individuals.** ELISA tests were performed to assess
534 the presence antibodies in serum of negative control individuals (NC) or COVID-19-vaccinated
535 individuals that bind r3CL-pro, r3CLpro-RBD chimer, rRBD or commercial RBD (cmRBD).
536 Results presented as the mean and standard deviation of OD 450 nm values of all the
537 individuals of the group.



538

539

540 **Figure 7. Evaluation of the antibody response induced by the recombinant 3CL-pro and**
541 **3CLpro-RBD chimer proteins in CD1 outbred mice.** Groups of CD1 outbred mice were
542 immunized with either r3CL-pro, r3CLpro-RBD chimer or Adjuvant only and evaluated for
543 specific antibodies using ELISA tests. A: ELISA using r3CL-pro as target antigen; B: ELISA
544 using r3CLpro-RBD chimer as target antigen to assess specificity immune response stimulated.
545 Results presented as the mean and standard deviation of OD 450 nm values of all the animals
546 of the group.



3C-Like protease
 3Cpro-RBD chimer
 Adjuvant

547


548

549

SUPPLEMENTARY MATERIAL


550

551



Peak Table - UV

| Peak | Retention ml | Area m ² mAU | Area % | Ext coeff. μ g ⁻¹ ml cm ⁻¹ | Fraction(s) | Volume ml | Conductivity mS/cm |
|--------|--------------|-------------------------|--------|--|---------------|-----------|--------------------|
| Peak A | 11.367 | 75.31 | | 22.38 | 5.B.9 - 5.C.5 | 1.801 | 14.81 |
| Peak B | 13.496 | 95.26 | | 28.3 | 5.C.7 - 5.D.5 | 2.200 | 14.80 |
| Peak C | 17.598 | 166.0 | | 49.32 | 5.D.9 - 5.F.4 | 4.001 | 14.79 |



552

553 **Supplementary Figure S1.** Protein standards optimization for size exclusion chromatography.

554

555

556

557

558

559


560

561

562

563

564



565

566

567 **Supplementary Figure S2. Antibody response of individuals naturally infected with**
 568 **SARS-CoV-2 to the 3C-like protease.** IgG antibodies against the r3C-like protease were
 569 detected in sera from 42 individuals confirmed positive for SARS-CoV-2 infection by RT-
 570 PCR. (A) Sera from the RT-PCR positive SARS-CoV-2 patients were tested for antibodies
 571 against 3C-like protease and compared to their antibodies levels against the Nucleoprotein
 572 (Npro). (A) Sera from the RT-PCR positive SARS-CoV-2 patients were tested for antibodies
 573 against 3C-like protease and compared to their antibodies levels against the Subunit 2 of the
 574 Spike protein (S2Frag). ELISA antibody tests developed in our previous study (De Marco
 575 Verissimo et al., 2021). ▲ sera were negative for antibodies against both Npro and S2frag by
 576 ELISA but positive for 3C-like protease; ● sera were negative for antibodies against any of the
 577 viral antigens tested; ● sera were positive for antibodies against Npro or S2frag by ELISA and
 578 for antibodies against 3C-like protease; ● sera were positive for antibodies against Npro or
 579 S2frag by ELISA but not for antibodies against 3C-like protease. Individual ELISA results are
 580 presented as Optical density (OD 450 nm) divided by the calculated cut-off (CO) to each
 581 ELISA test developed (considering the negative control group). The cut-off value for each
 582 antigen is indicated by the dotted line.

583

584

585