

**DEAE-Dextran enhances the lentiviral transduction of primary human mesenchymal stromal cells
from all major tissue sources without affecting their proliferation and phenotype**

Francesco Amadeo^{1,2}, Vivien Hanson¹, Patricia Murray², Arthur Taylor²

1. Cellular Therapies Laboratory, NHS Blood and Transplant, Liverpool, UK.

2. Department of Molecular Physiology and Cell Signalling, University of Liverpool, Liverpool, UK.

Corresponding author e-mail address: taylora@liverpool.ac.uk

Orcid IDs: Francesco Amadeo 0000-0002-3868-2348; Patricia Murray 0000-0003-1316-148X; Arthur Taylor 0000-0003-2028-6694

Abstract (100- 200 words)

Genetic engineering of mesenchymal stromal cells (MSCs) is a tool widely used to explore MSC properties in both *in vitro* and *in vivo* applications. Lentiviral infection with the use of polycations as an adjuvant is a method that is commonly used to generate stably transduced cells. However, it is known that some polycations can negatively affect primary MSCs and to date, no study has explored the effect of different polycations on the transduction efficiency and properties of all main types of MSCs, namely those derived from umbilical cord, bone marrow and adipose tissue. Here we explore a range of polycations, using transduction protocols with and without spinoculation, to produce stably transduced MSCs from these three tissue sources. We identified that an overnight incubation with diethylaminoethyl-dextran (DEAE-Dextran) is the protocol associated with the best transduction efficiency without compromising the viability of the cells, and which worked consistently with lentiviral particles encoding for different transgenes. Transduced and sorted MSC populations revealed no significant changes in proliferation, morphology and expression of MSC markers compared to naïve MSCs. Following this study, we conclude that DEAE-Dextran is a polycation that can be successfully used to enhance the transduction of MSCs from all major tissue sources.

Keywords: mesenchymal stromal cells, polycations, lentivirus, transduction, reporter genes.

Declarations:

Funding: This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 813839.

Conflicts of interest/Competing interests: The authors have no conflicts of interest to declare that are relevant to the content of this article.

Availability of data and material: All datasets from this study are publicly available on Zenodo, DOI: 10.5281/zenodo.6594236

Authors' contributions: V. H. provided the umbilical cord mesenchymal stromal cells. F.A. performed the *in vitro* experiments and the data analysis. F. A. produced the figures and the tables. F.A., A.T and P.M. wrote the paper. All authors contributed to the design of the experiments and to the interpretation of the results.

Acknowledgments: we thank Sandra Calcat-i-Cervera (National University of Ireland, Galway) for providing the bone marrow mesenchymal stromal cells; we thank Erika Erika and Eleonora Scaccia (University of Heidelberg) for providing the adipose derived mesenchymal stromal cells.

Introduction

Mesenchymal stromal cells (MSCs) are multipotent cells showing great promise in different pre-clinical studies and currently used in many clinical trials [1, 2]. Several properties and mechanisms of actions of these cells are still not completely understood and genetic engineering is a tool that can be used to gain a better understanding of their properties *in vitro* [3], help to investigate what happens to them when administered *in vivo* [4, 5], or even as a strategy to improve their therapeutic efficacy [6].

Lentiviral vectors (LV) are widely used as gene delivery vectors for various research applications because of their ability to infect both dividing and non-dividing cells. The interaction between the lentiviral particle and the cell of interest is receptor independent [7] and can be reduced or prevented by the presence of strong electrostatic repulsion between the negatively charged cell and the envelope of the virus [8, 9].

The addition of polycations, positively charged molecules, can reduce the charge of cell membranes, and increase the chance of an interaction between the lentivirus and the cell surface [9]. Polybrene (Pb) is the most commonly used polycation in lentiviral transduction and is associated with a very high transduction efficiency [10, 11]. Nevertheless, it has also been reported to have negative effects on some types of cell [12-14]. In particular, its use for the transduction of human endometrium-derived mesenchymal stromal cells has been described to negatively affect their proliferation, migration ability and differentiation potential [14]. Other polycations, such as diethylaminoethyl-dextran (DEAE-dextran) and protamine sulphate (Ps) have also been successfully used to transduce different cell lines [9, 15].

Another method reported to increase the binding of retroviruses to the surface of the cells is spinoculation [16-18]. Although various mechanisms have been proposed [16, 17], it is still unclear what process is responsible for this enhancement. During spinoculation, the cells are centrifuged at low speed (below 2000g) in the presence of the virus. Importantly, it has been reported that

spinoculation together with the optimal polycation can synergistically increase transduction efficiency [15].

A transduction method that is universally efficacious across all major types of MSC would facilitate research on their properties and mechanisms of action but to date, no studies have assessed how different protocols affect MSCs from different tissue sources and across different donors. Importantly, any protocol for the genetic modification of MSCs must not alter their properties, unless that is the specific intention such as in a gene knockdown study, as this would compromise the research findings. In this work, we optimise a transduction protocol with the goal of obtaining genetically engineered umbilical cord (UC), bone marrow (BM) and adipose-derived (A) MSCs that share the same properties as the unmodified (naïve) cells. We identify that a protocol based on the use of DEAE-dextran is capable of producing stable transduced mesenchymal stromal cell populations without affecting their properties.

Materials and Methods

Cell culture

Human MSCs from three different tissue sources were used. Umbilical cord (UC) MSCs were obtained from the NHS Blood and Transplant (NHSBT, Liverpool, UK) from three different healthy donors, licenced under the Human Tissue Act (HTA). The cells were isolated according to NHSBT good manufacturing practice (GMP) procedures: the tissue was halved horizontally, chopped into big pieces and cultured undisturbed for 7 days. The pieces were then removed, and the adherent cells were expanded for two passages under standard tissue culture conditions. Bone marrow (BM) MSCs were provided by the University of Galway (Ireland), after being purchased from Lonza (Basel, Switzerland). Adipose derived (A) MSCs were provided by the University of Heidelberg (Germany), after being isolated from lipoaspirates harvested after informed consent. Briefly, the tissue was digested with a solution of NB4 collagenase (Serva/Nordmark) under gentle agitation for 30 minutes. After a straining step to remove the undigested tissue, the cells were centrifuged, resuspended in complete medium

and expanded under standard tissue culture conditions. The Mannheim Ethics Commission II approved the study (vote 2011-215N-MA). Cells from three different donors were obtained for each tissue source, leading to a total of 9 MSC samples. The cells were cryopreserved and shipped to the Department of Molecular Physiology and Cell Signalling of the University of Liverpool, where all the cells were cultured following standard mammalian tissue culture protocols. The cells were grown in MEM- α containing GlutaMAX (Gibco) and supplemented with 10% foetal bovine serum (FBS; Gibco) and kept at 37 °C in a humidified incubator, with 5% CO₂.

Transduction

In the set of experiments aimed to explore different transduction protocols we infected the 3 types of MSCs with pHIV-Luc2-ZsGreen (Supplementary Figure 1a) with a multiplicity of infection (MOI) of 5. The pHIV-Luc2-ZsGreen vector was a gift from Bryan Welm (Addgene plasmid #39196). We seeded the cells at a density of 5×10^3 cells/cm² and transduced them 4 hours later in the presence of three different polycations: polybrene (Pb, 8 μ g/mL), protamine sulphate (Ps, 20 μ g/mL) or DEAE-dextran (6 μ g/mL). MSCs transduced without any adjuvant served as control of the basal transduction efficiency. To evaluate a possible effect of the polycations themselves on the cells, we also incubated the cells with either Pb, Ps or DEAE-dextran alone. Finally, we used MSCs alone (without any polycation or LV) as controls. We transduced the cells under two conditions: (a) overnight incubation at 37°C (“static” condition), replacing the medium containing the LV/adjuvant after 16h or (b) centrifugation of the cells for 1 hour at 750 g and room temperature, followed by incubation for 1 hour at 37°C, and then replacing the medium containing the LV/adjuvant (“centrifuged” condition). We assessed the transduction efficiency 7-10 days post transduction via flow cytometry using a FACScalibur (BD Biosciences) and used the CellTiter-Glo[®] Luminescent Cell Viability Assay (#G7571, Promega) to evaluate the effect of the transduction on the viability and proliferation of the cells.

Once we identified the optimal transduction protocol, we applied it for the transduction with 4 other different lentiviral particles. We seeded MSCs at a density of 5×10^3 cells/cm² and transduced them

with one of the following lentiviral particles (Supplementary Figure 1b): (i) pHIV-eGFP (a gift from Bryan Welm & Zena Werb; Addgene plasmid #21373), (ii) pHIV-dTomato (a gift from Bryan Welm; Addgene plasmid #21374), (iii) pCDH-EF1 α -Luc2-P2A-tdTomato (a gift from Kazuhiro Oka; Addgene plasmid #72486), (iv) pLV-mCherry (a gift from Pantelis Tsoulfas, Addgene plasmid #36084). All transductions were carried out using an MOI of 5 in the presence or absence of DEAE-dextran (6 μ g/mL) applying the static protocol. We assessed the transduction efficiency via flow cytometry after 7 days of culture.

In the set of experiments aimed to evaluate the effect of the transduction on the proliferation, expression of MSC markers and morphology of the cells, we transduced the MSCs with the pHIV-Luc2-ZsGreen construct (MOI of 5) in the presence of 6 μ g/mL of DEAE-dextran applying the static protocol. Following expansion, we sorted the cells based on the ZsGreen expression using a FACSaria II (BD Biosciences). Then we compared these cells to untransduced controls in terms of proliferation, morphology and expression of MSC markers. Table 1 summarises the passage of each donor sample at each study step.

Proliferation and doubling time

After sorting and expansion for one passage, Luc2-ZsGreen⁺ and naïve MSCs were seeded at their optimal seeding density (3x10² cells/cm² for adipose cells and 3x10³ cells/cm² for umbilical cord and bone marrow cells) and expanded until 60-90% confluence for at least two passages (Table 1) from where we calculated the doubling time using the following equation:

$$Td = \frac{t}{\text{Log}_2\left(\frac{N_t}{N_0}\right)}$$

Where Td is the doubling time, N_t is the number of cells at time t and N₀ is the number of cells seeded, from which the number of doublings was calculated based on the time the cells had been in culture.

MSC markers

MSC marker expression was assessed via flow cytometry by staining the cells with anti CD44 (APC, #130-113-893, Miltenyi Biotec), anti CD45 (APC, #130-113-676, Miltenyi Biotec), anti CD73 (APC, #130-097-945, Miltenyi Biotec), anti CD90 (APC, #130-117-534, Miltenyi Biotec), anti CD105 (APC, #130-099-125, Miltenyi Biotec), IgG1 mouse isotype (APC, #130-113-758, Miltenyi Biotec), or IgG2 mouse isotype (APC, #130-113-831, Miltenyi Biotec) according to the manufacturer's instructions. An extra vial of each cell population was used as unstained control. We acquired the data with a FACScalibur (BD Biosciences) flow cytometer and we analysed a minimum of 10^4 events for each marker.

Morphological analysis of pHIV Luc2 ZsGreen transduced MSCs

We cultured MSCs with complete medium overnight. After 16 hours we fixed the cells with paraformaldehyde (4% w/v in PBS, pH 7.4) for 20 minutes at room temperature (RT), then we washed the cells with PBS, permeabilised them with 0.1% (v/v) Triton X-100 in PBS and incubated them with Alexa Fluor 594 Phalloidin (#A12381, ThermoFisher) [165nM] in PBS with 1% (w/v) bovine serum albumin (BSA) for 1 hour at RT. We used 4',6 diamidino-2-phenylindole (DAPI) [143 nM] as a counter stain for the nuclei. We acquired the fluorescence images with a Leica DM2500 microscope coupled to a DFC350 FX camera. Finally, we used ImageJ to perform the data analysis: the shape of the cells was defined manually, based on the Phalloidin staining, and the software was used to calculate the cells' area, perimeter, and circularity.

Bioluminescence imaging of pHIV Luc2 ZsGreen transduced MSCs

We used the IVIS Spectrum system (Perkin Elmer) to characterise the total flux (photons/second) emitted by the Luc2-ZsGreen⁺ MSCs. We plated the cells at increasing densities, from 156 to 2×10^4 cells/well, in optical bottom 96-well plates with black walls (#165305, ThermoFisher) with a technical triplicate. Untransduced cells were used as negative control. We allowed the cells to adhere for 3 hours and then we added D-Luciferin at a final concentration of 5.12 mM with a final volume of 100 μ L per well. We collected the data using the IVIS immediately after the addition of the substrate, acquiring the signal generated using an open filter. The acquired signal was always normalised to radiance

(photons/second/centimeter²/steradian) and analysed using the region of interest (ROI) tool of the IVIS software (Living Image v 4.5.2) to obtain the flux.

Statistical analysis

All values in graphs are represented as mean \pm standard deviation, unless specified differently in the figure legend. The statistical analysis was performed using the GraphPad Prism software. The type of statistical test and the number of replicates included in the analyses are indicated in the figure legends.

Results

Optimisation of a lentiviral transduction protocol

The cells were infected with a lentiviral particle encoding for firefly luciferase (FLuc) and a green fluorescent protein, ZsGreen, whose expression was used to determine the transduction efficiency via flow cytometry. In parallel, an ATP assay was used to investigate the effect of the different protocols on the proliferation of the cells.

UC-MSC transduction efficiency in the presence of Pb was high regardless of the application of the static ($93.8 \pm 4.4\%$) or the centrifuged ($90.4 \pm 4.5\%$) protocol (Figure 1a, fluorescence images of the cells are shown in Supplementary Figure 2). However, the ATP assay of transduced cells in the presence of Pb revealed a strong reduction in the proliferation of the transduced cells, which was only 32.3% ($\pm 8.6\%$) and 44.1% ($\pm 12.5\%$) of the control in static and centrifuged conditions, respectively (data of untransduced UC-MSCs are shown in Supplementary Figure 5a). Cells transduced in the presence of DEAE-dextran resulted in 74.4% ($\pm 3.1\%$) and 53.8% ($\pm 19.4\%$) of positive cells (Figure 1a) in static and centrifuged conditions and were respectively associated with a proliferation reduced to 79.3% ($\pm 17.8\%$) and 74.1% ($\pm 12.2\%$) (Figure 1b). Finally, in the presence of Ps (Figure 1a), 28.2% ($\pm 11.9\%$) and 70.7% ($\pm 16.3\%$) of the cells were positive for ZsGreen in static and centrifuged conditions, respectively. These conditions were associated with a reduction in proliferation to 96.9% ($\pm 10.1\%$) and to 74.0% ($\pm 11.4\%$) (Figure 1b).

Similar results were obtained with BM-MSCs, where the transduction in the presence of Pb was associated with $92.3 \pm 5.3\%$ and $94.1 \pm 3.0\%$ of the cells expressing the ZsGreen protein following the application of the static and the centrifuged protocols, respectively (Figure 1c, fluorescence images of the cells are shown in Supplementary Figure 3). Like for UC-MSCs, the presence of Pb resulted in a reduction in the proliferation of the transduced cells to $72.3 \pm 6.0\%$ of the control in static condition but did not really affect the growth of the cells when the Pb was combined with a centrifugation step ($94.0 \pm 9.5\%$, Figure 1d). Cells transduced in the presence of DEAE-dextran resulted in $80.2\% (\pm 5.1\%)$ and $76.0\% (\pm 11.5\%)$ of positive cells (Figure 1c) in static and centrifuged conditions, respectively. Surprisingly, the presence of the DEAE-dextran seems to have increased the proliferation of the transduced cells to $116.3\% (\pm 10.5\%)$ and to $109.7\% (\pm 2.9\%)$, in static and centrifuged conditions, respectively (Figure 1d). This increase in the proliferation was also observed when the cells were incubated without the lentiviral particles and with the DEAE-dextran only ($120.7 \pm 10.8\%$ in static and $106.0 \pm 3.0\%$ in centrifuged condition, Supplementary Figure 5b). Finally, in the presence of Ps (Figure 1c), $22.5\% (\pm 10.0\%)$ and $75.8\% (\pm 6.7\%)$ of the cells were positive for ZsGreen in static and centrifuged conditions, respectively. The presence of the Ps was also associated with an increase in proliferation of transduced cells both in static ($128 \pm 8.9\%$) and centrifuged conditions ($119.7 \pm 3.5\%$) (Figure 1d).

For A-MSCs, transduction in the presence of Pb in static and centrifuged conditions was associated with only $48.6 \pm 7.5\%$ and $19.2 \pm 3.8\%$ of cells effectively transduced, respectively (Figure 1e, fluorescence images of the cells are shown in Supplementary Figure 4). Cells transduced in the presence of DEAE-dextran resulted in $50.0\% (\pm 9.9\%)$ and $12.3\% (\pm 2.8\%)$ of positive cells (Figure 1e) in static and centrifuged conditions, respectively. The transduction with Ps resulted in only $8.3\% (\pm 2.3\%)$ and $6.3\% (\pm 1.3\%)$ of positive cells (Figure 1e). The ATP assay revealed that the presence of Pb affected the proliferation of the transduced cells following the application of the static protocol ($74.4 \pm 6.8\%$), while none of the other conditions influenced ATP levels (Figure 1f, Supplementary Figure 5c).

A three-way ANOVA analysis performed on the whole dataset of each cell type (Supplementary Figure 5) revealed that the main impact on the viability for BM and A-MSCs is related to the presence of the polycations ($p < 0.0001$ and $p = 0.001$ respectively). On the other hand, the UC-MSCs were found to be susceptible not only to the presence of polycations ($p < 0.001$), but also to the transduction ($p < 0.05$) and to the centrifugation ($p < 0.01$).

Validation of the selected protocol: DEAE-dextran in static condition

The use of DEAE-dextran with overnight incubation in static conditions was selected as the optimal transduction procedure. This is because it resulted in a good transduction efficiency for all the MSCs tested (74.4% for UC-MSCs, 80.2% for BM-MSCs and 50.0% for A-MSCs) without overtly affecting the proliferation/viability of the cells. This protocol was then applied to transduce the three types of MSCs with different lentiviral particles to assess if the transduction efficiency with this protocol is influenced by the lentiviral construct itself.

Supplementary Figure 6 displays representative fluorescence images of all three MSCs transduced with the four lentiviral particles in the absence or presence of DEAE-dextran. Regardless of the lentiviral particle, all the cell types display a low transduction efficiency when transduced without polycations. In contrast, for all four lentiviral particles, the presence of DEAE-Dextran leads to an increase in the number of transduced cells and in the intensity of signal in all three MSCs (Supplementary Figure 6a-c). This increase is particularly noticeable for the pLV-mCherry and for the pHIV-dTomato, while less evident for the pHIV-eGFP and for the pCDH-EF1 α -Luc2-P2A-tdTomato (especially for the BM-MSCs, Supplementary Figure 6b).

The flow cytometry analysis (Figure 2) of these cells revealed that UC-MSCs displayed a statistically significant increase in the fraction of transduced cells when infected in the presence of the DEAE-dextran for all the lentiviral particles (Figure 2a). The proportion of cells positive for the transgene increased from $0.3 \pm 0.18\%$ to $11.7 \pm 4.4\%$ for the pCDH-EF1 α -Luc2-P2A-tdTomato ($p = 0.0005$), from $9.4 \pm 2.8\%$ to $61.8 \pm 11.6\%$ for the pHIV-dTomato ($p < 0.0001$), from $10.5 \pm 2.7\%$ to $58.7 \pm 8.9\%$ for the

pHIV-eGFP ($p < 0.0001$) and from $5.5 \pm 2.2\%$ to $72.9 \pm 9.2\%$ for the pLV-mCherry ($p < 0.0001$, Figure 2a). A two-way ANOVA analysis also revealed that both the presence of DEAE-dextran and the specific lentiviral construct used have a statistically significant impact on the transduction efficiency ($p < 0.01$ and $p < 0.0001$ respectively).

Similar results were obtained with the BM-MSCs (Figure 2b), where the proportion of transduced cells increased from $0.49 \pm 0.46\%$ to $6.3 \pm 1.3\%$ for the pCDH-EF1 α -Luc2-P2A-tdTomato (no statistically significant difference), from $13.7 \pm 0.9\%$ to $57.8 \pm 9.4\%$ for the pHIV-dTomato ($p < 0.0005$), from $13.2 \pm 0.7\%$ to $54.8 \pm 17.6\%$ for the pHIV-eGFP ($p = 0.0005$) and from $11.7 \pm 1.1\%$ to $73.2 \pm 11.2\%$ for the pLV-mCherry ($p < 0.0001$, Figure 2b). A two-way ANOVA analysis confirmed the statistically significant effect of the DEAE-dextran and of the lentiviral construct on the transduction efficiency ($p < 0.05$ and $p = 0.0001$ respectively).

Lastly, the adipose derived cells showed a transduction behaviour comparable to the other two MSCs (Figure 2c), with an increase in the percentage of transduced cells from $0.8 \pm 0.54\%$ to $5.9 \pm 2.1\%$ for the pCDH-EF1 α -Luc2-P2A-tdTomato (no statistically significant difference), from $6.7 \pm 2.6\%$ to $54.0 \pm 10.6\%$ for the pHIV-dTomato ($p < 0.0001$), from $5.6 \pm 2.4\%$ to $38.1 \pm 10.0\%$ for the pHIV-eGFP ($p < 0.0001$) and from $3.2 \pm 1.2\%$ to $54.2 \pm 13.4\%$ for the pLV-mCherry ($p < 0.0001$, Figure 2c). The two-way ANOVA analysis identified a statistically significant impact of the DEAE-dextran and of the type of lentiviral particle on the transduction efficiency ($p < 0.05$ and $p < 0.0001$ respectively).

Characterisation of the MSCs transduced with the DEAE-dextran static protocol

To determine whether the transduction procedure had an impact on the properties of the different MSCs, all cells were transduced with the pHIV-Luc2-ZsGreen construct and sorted to obtain a population approximately 100% positive for the transgenes. Then, transduced and untransduced cells were compared in terms of doubling time, expression levels of MSC markers and morphology (area and circularity of the cells).

Figure 3a shows phase contrast images of control and Luc2-ZsGreen⁺ UC, BM and A-MSCs cells and the respective green fluorescence images (ZsGreen). The morphology of the different transduced cells looks comparable to controls (Figure 3a). The analysis of the cumulative doublings revealed that the transduced cells behaved similarly to the untransduced ones for at least 2 passages (Supplementary Figure 7a-c) and no statistically significant difference in the doubling time was observed between transduced and untransduced cells (Figure 3b and Supplementary Figure 7d).

The flow cytometry analysis of the expression of ZsGreen was performed two passages after sorting (Table 1) to evaluate any loss of the reporter gene following the expansion. All the transduced populations resulted in a positivity level of over 98.5% (Table 2), underlining the stability of the transduction. Furthermore, the flow cytometry analysis of surface markers revealed a similar level of expression of positive markers (CD44, CD73, CD90 and CD105) and the lack of CD45 negative marker (Table 2) in both transduced and untransduced cells. Only MSCs from one A-MSC donor (A-02) resulted in an increase in the percentage of cells positive for CD45, from 9.28% to 75.52% (Table 2).

Figure 3c shows representative fluorescent images of transduced and untransduced cells stained with phalloidin and DAPI. Morphological analysis of the transduced cells revealed no statistically significant difference in the area (Figure 3d) or circularity (Figure 3e) of transduced MSCs compared to untransduced cells. MSCs from one UC-MSC donor (UC-02) displayed an increase in the area (Supplementary Figure 8a), with no impact on the circularity (Supplementary Figure 8b) and MSC from one A-MSC donor (A-03) showed a slight increase in the circularity (Supplementary Figure 8f), with no difference in the area (Supplementary Figure 8e).

To measure the total flux emitted by each donor and confirm that the luciferase gene is also functional, cells were plated at various densities from 156 to 2×10^4 cells/well and subsequently imaged in the presence of the D-Luciferin substrate, at a final concentration of 5.12 mM [5]. A linear regression of the signal measured revealed that the output from the three UC-MSC donor samples was similar (16,958 p/s/cell, 18,682 p/s/cell and 19,498 p/s/cell for UC-01, UC-02 and UC-03, respectively,

Figure 4a) and comparable to the BM-MSCs (18,630 p/s/cell, 15,531 p/s/cell and 18,672 p/s/cell for BM-01, BM-02 and BM-03, respectively, Figure 4b). The output from the A-MSCs revealed one sample displaying higher signal when compared to the other two (20,781 p/s/cell, 30,723 p/s/cell and 22,317 p/s/cell for A-01, A-02 and A-03, respectively, Figure 4c). On average, UC and BM-MSCs displayed very similar signal outputs ($18,379 \pm 1,297$ p/s/c and $17,611 \pm 1,801$ p/s/c, respectively), while A-MSCs resulted in a stronger signal ($24,607 \pm 5,325$ p/s/c), although no statistically significant difference was observed between MSCs from different tissue sources (Figure 4d).

Discussion

The purpose of this study was to identify a lentiviral transduction protocol suitable to infect human umbilical cord, bone marrow and adipose MSCs and generate stable cell populations without affecting their proliferation and properties.

Lentiviral transduction can be used to generate stable cell lines expressing a specific gene of interest [19]. However, the transduction with the LV alone has a low transduction efficiency as shown both in the literature [12, 20] and in this study. There are reports showing that several polycations, like Pb [10, 11], Ps [15, 20] and DEAE-dextran [9], and the combination of these with a spinoculation step [16-18], can increase the transduction efficiency of different cells. Here we aimed at identifying which of these adjuvants are most effective to transduce MSCs from umbilical cord, bone marrow and adipose tissue, and whether spinoculation does indeed have a positive effect on the transduction of these cells.

For that, we tested the impact of different polycations (Pb, Ps and DEAE-dextran) on transduction efficiency and on proliferation alone or in combination with a centrifugation step. The transduction efficiency in the presence of Pb (8 μ g/mL) was high for all the MSCs, with more than 90% of UC and BM cells transduced in both static and centrifuged conditions and with around 50% of adipose cells transduced in static condition. However, proliferation of cells incubated with the Pb in static condition was significantly reduced in comparison to control cells. We observed this effect not only on the cells incubated with the lentiviral particles (32%, 72% and 74% for UC, BM and A-MSCs respectively), but

also on the cells incubated just with the Pb overnight (56%, 69% and 78% for UC, BM and A-MSCs respectively). These results are in line with previous studies [12-14, 20] that reported an association between the presence of polybrene and a reduction in the proliferation of different cell types. Griukova *et al* [14] has previously shown that Pb induces a p38 mitogen-activated protein kinase (MAPK) dependent premature senescence in human endometrial MSCs [14]. They reported enhanced β -galactosidase activity, a marker extensively used for the detection of senescence, and reduction in the expression of CD146, a novel marker for detecting senescence in UC-MSCs [14, 21]. Furthermore, Lin *et al* [13] reported that even when using a potent mitogen, such as fibroblast growth factor 2, it is not possible to overcome the negative effects of Pb [13]. Lin *et al* also reported a significant reduction in proliferation of human bone marrow MSCs with a polybrene concentration of 4 μ g/mL, which is half of what was used here [13].

The p38 MAPK-dependent senescence might explain the reduced proliferation observed here when applying the Pb static condition, but we note that it was possible to reduce the negative effect of the Pb on BM and A-MSCs with the use of the centrifugation protocol, which reduced the exposure to the lentiviral particles to only 2 hours, although the impact on UC-MSCs was still substantial. Furthermore, the number of A-MSCs transduced with Pb following the introduction of the centrifugation step dropped from 48.6% to only 19.2%. For these reasons, we did not further investigate the use of Pb, as we wanted to identify a method suitable for the transduction of all the three cell types. It is important to underline that the use of Pb together with a centrifugation step would be the best option to transduce BM-MSCs, as it is associated with a very high transduction efficiency (> 94%) without impacting on cell viability (94% of control).

As possible alternative protocols we identified: (a) the incubation with Ps coupled with centrifugation, as it was associated with a good transduction efficiency for UC and BM cells (70.7% and 75.8%, respectively) with a reduced impacting on the proliferation of the cells (74% for UC and 119% for BM); or (b) the DEAE-dextran overnight incubation in static condition, as it was associated with a good

transduction efficiency for all the cell types tested (74.4% for UC, 80.2% for BM and 50.0% for A-MSCs) without overtly impacting the proliferation of the cells (77% for UC, 116% for BM and 107% for A-MSCs). Because we were interested in a protocol suitable to transduce all three cell types tested, and because the presence of centrifugation affected the viability of UC-MSCs, we opted for the DEAE-dextran static method. The superiority of the DEAE-dextran to Pb has been previously shown by Denning *et al* [9], but their work was focused on the 293FT and HT-1080 cell lines rather than primary MSCs. It should be noted that some studies have seen further enhancements in transduction efficiency with Ps by increasing its concentration to up to 100 µg/mL [15], although it seems that this is not consistent throughout the literature, with some investigators identifying a plateau at 20 µg/mL [20], the concentration we used here.

A similar analysis where different transduction protocols were tested to optimise the transduction of natural killer (NK) cells was carried out by Malarkannan *et al* [22]. They investigated the transduction efficiency and the impact on cell viability following the lentiviral transduction of primary human and murine NK cells in the presence of either Pb (8 µg/mL), Ps (8 µg/mL) or DEAE-dextran (8 µg/mL) [22]. In this study, they combined a 1h centrifugation step at 1000g and an overnight incubation at 37°C. Interestingly they obtained a strong increase in the transduction efficiency by incubating the cells with DEAE-dextran and almost no transduction with the two other polycations [22].

The data from Denning *et al* [9] and Malarkannan and co-workers [22] suggest that the type of cell and the lentiviral particle itself might influence the outcome of the transduction and one single protocol might not be suitable for all cell types and LV constructs. Because of this, we investigated the DEAE-dextran static protocol for its ability to transduce the MSCs with four different lentiviruses, characterised by differences in their backbone and in the size of the insert. Regardless of the lentiviral construct used, the DEAE-dextran protocol enabled an increase in the transduction efficiency when compared to the cells incubated with the LVs alone. However, we identified a difference in the transduction efficiency of the different cells that was directly related to the type of lentiviral vector

use. This is expected, as several factors can influence not only the transduction efficiency but also the transgene expression level of a specific lentiviral vector. For instance, it has been previously shown that efficiency of transduction decreases with an increase in the size of the construct [23, 24]. In particular, Canté-Barrett *et al* [24] reported that the transduction efficiency of human hematopoietic stem cells decreases significantly when the size of the insert in the LV was close to 6 kb or larger [24]. The results obtained in the present study are in line with those findings, as the plasmid with the largest insert, the pCDH-EF1 α -Luc2-P2A-tdTomato (7046 bp insert size) was associated with a much lower transduction efficiency than the 3 other plasmids tested, which had smaller inserts (5038 bp, 5032 bp and 3938 bp for the pHIV-eGFP, the pHIV-dTomato and the pLV-mCherry, respectively). Furthermore, it was also previously reported that the gene sequence in the insert can affect the lentiviral infection [24]. This can further explain not only what was observed when applying the DEAE-dextran protocol with 4 different LVs, but also the high transduction efficiency reported when infecting the cells with the pHIV-Luc2-ZsGreen LV particles, which has an insert of 6663 bp.

Furthermore, the percentage of transduced cells for each lentiviral particle was significantly correlated with the tissue origin of the MSCs, confirming that the type of cell also has a role in the outcome of the transduction. For example, the percentage of A-MSCs transduced with the pHIV-Luc2-ZsGreen vector was much lower than the percentage of transduced UC or BM-MSCs. Interestingly we saw little variation between donor samples from the same tissue source, indicating that this is not likely to be a major variable when transduction efficiency is considered.

Finally, a critical aspect of this study was the production of transduced MSCs that can be used for further applications without modifying and impairing the properties of the cells. Transduced cells should be as similar as possible to the naïve, non-genetically modified MSCs in order to be representative. We investigated this by assessing their proliferation, expression of MSC markers and morphology. We transduced a total of 9 donor samples (3 per cell type) using the selected protocol (6 μ g/mL DEAE-dextran) and a pHIV_Luc2_ZsGreen LV with an MOI of 5 and we sorted the cells

following the transduction to obtain pure FLuc expressing populations. We showed that all the transduced cells display similar properties to their untransduced counterparts in terms of doubling time, proliferation, and expression of common MSC markers. However, it is interesting to point out that one UC-MSC displayed an area larger than the respective control after transduction, and that one A-MSC was slightly less circular than the respective control. Additionally, one A-MSC has an increased expression of CD45 after transduction, indicating the need to properly characterise the effect of transgene expression on cell properties, as this is no doubt dependent on the transgene.

The 9 donor samples, when genetically modified, exhibited comparable emission of light in the presence of D-Luciferin, underlining the replicability of the methodology. The average flux emitted ranged from 15,531 p/s/cell for one of the bone marrow samples to 30,723 p/s/cell for one of the adipose samples. These values are much higher than those described in a previous study, which reported that transduction of a mouse derived MSC cell line with the pHIV-Luc2-ZsGreen plasmid in the presence of polybrene resulted in 1508 p/s/cell [25], which is around 10 times lower than the values obtained here. Although this lower flux was already sufficient for *in vivo* bioluminescence imaging [25], greater light emissions can yield better imaging sensitivity.

In summary, an overnight incubation with 6 µg/mL DEAE-dextran and a LV with an MOI of 5 can be used universally to enhance the transduction of umbilical cord, bone marrow and adipose derived MSCs with several lentiviral particles. The characterisation of the transduced cells expressing Luc2_ZsGreen displayed no major effects of the transduction on the properties of the cells. Still, as mentioned previously, the efficiency of transduction is strongly correlated with the LV and with the cell type used. Because of this, we suggest that several polycations and protocols should be tested when optimising a transduction process for a specific cell type.

References

1. Fung, M., et al., *Responsible Translation of Stem Cell Research: An Assessment of Clinical Trial Registration and Publications*. Stem Cell Reports, 2017. **8**(5): p. 1190-1201.
2. Amadeo, F., et al., *Mesenchymal stromal cells: what have we learned so far about their therapeutic potential and mechanisms of action?* Emerg Top Life Sci, 2021. **5**(4): p. 549-562.
3. Niess, H., et al., *Genetic engineering of mesenchymal stromal cells for cancer therapy: turning partners in crime into Trojan horses*. Innov Surg Sci, 2016. **1**(1): p. 19-32.
4. Scarfe, L., et al., *Non-invasive imaging reveals conditions that impact distribution and persistence of cells after in vivo administration*. Stem Cell Res Ther, 2018. **9**(1): p. 332.
5. Amadeo, F., et al., *Firefly luciferase offers superior performance to AkaLuc for tracking the fate of administered cell therapies*. Eur J Nucl Med Mol Imaging, 2022. **49**(3): p. 796-808.
6. Damasceno, P.K.F., et al., *Genetic Engineering as a Strategy to Improve the Therapeutic Efficacy of Mesenchymal Stem/Stromal Cells in Regenerative Medicine*. Front Cell Dev Biol, 2020. **8**: p. 737.
7. Pizzato, M., et al., *Initial binding of murine leukemia virus particles to cells does not require specific Env-receptor interaction*. J Virol, 1999. **73**(10): p. 8599-611.
8. Cornetta, K. and W.F. Anderson, *Protamine sulfate as an effective alternative to polybrene in retroviral-mediated gene-transfer: implications for human gene therapy*. J Virol Methods, 1989. **23**(2): p. 187-94.
9. Denning, W., et al., *Optimization of the transductional efficiency of lentiviral vectors: effect of sera and polycations*. Mol Biotechnol, 2013. **53**(3): p. 308-14.
10. Davis, H.E., J.R. Morgan, and M.L. Yarmush, *Polybrene increases retrovirus gene transfer efficiency by enhancing receptor-independent virus adsorption on target cell membranes*. Biophys Chem, 2002. **97**(2-3): p. 159-72.
11. Davis, H.E., et al., *Charged polymers modulate retrovirus transduction via membrane charge neutralization and virus aggregation*. Biophys J, 2004. **86**(2): p. 1234-42.

12. Seitz, B., et al., *Retroviral vector-mediated gene transfer into keratocytes: in vitro effects of polybrene and protamine sulfate*. Graefes Arch Clin Exp Ophthalmol, 1998. **236**(8): p. 602-12.
13. Lin, P., et al., *Polybrene inhibits human mesenchymal stem cell proliferation during lentiviral transduction*. PLoS One, 2011. **6**(8): p. e23891.
14. Griukova, A., et al., *P38 MAPK inhibition prevents polybrene-induced senescence of human mesenchymal stem cells during viral transduction*. PLoS One, 2018. **13**(12): p. e0209606.
15. Lin, P., et al., *Efficient lentiviral transduction of human mesenchymal stem cells that preserves proliferation and differentiation capabilities*. Stem Cells Transl Med, 2012. **1**(12): p. 886-97.
16. O'Doherty, U., W.J. Swiggard, and M.H. Malim, *Human immunodeficiency virus type 1 spinoculation enhances infection through virus binding*. J Virol, 2000. **74**(21): p. 10074-80.
17. Guo, J., et al., *Spinoculation triggers dynamic actin and cofilin activity that facilitates HIV-1 infection of transformed and resting CD4 T cells*. J Virol, 2011. **85**(19): p. 9824-33.
18. Yan, R., et al., *Spinoculation Enhances HBV Infection in NTCP-Reconstituted Hepatocytes*. PLoS One, 2015. **10**(6): p. e0129889.
19. Tandon, N., et al., *Generation of Stable Expression Mammalian Cell Lines Using Lentivirus*. Bio Protoc, 2018. **8**(21).
20. Deryabin, P., et al., *Optimization of lentiviral transduction parameters and its application for CRISPR-based secretome modification of human endometrial mesenchymal stem cells*. Cell Cycle, 2019. **18**(6-7): p. 742-758.
21. Jin, J. H., et al., *Downregulation of Melanoma Cell Adhesion Molecule (MCAM/CD146) Accelerates Cellular Senescence in Human Umbilical Cord Blood-Derived Mesenchymal Stem Cells*. Stem Cells Transl Med, 2016. **5**(4): p. 427-39
22. Nanbakhsh, A., et al., *Dextran Enhances the Lentiviral Transduction Efficiency of Murine and Human Primary NK Cells*. J Vis Exp, 2018(131).
23. Kumar, M., et al., *Systematic determination of the packaging limit of lentiviral vectors*. Hum Gene Ther, 2001. **12**(15): p. 1893-905.

24. Cante-Barrett, K., et al., *Lentiviral gene transfer into human and murine hematopoietic stem cells: size matters*. BMC Res Notes, 2016. **9**: p. 312.
25. Taylor, A., et al., *Multicolour In Vivo Bioluminescence Imaging Using a NanoLuc-Based BRET Reporter in Combination with Firefly Luciferase*. Contrast Media Mol Imaging, 2018. **2018**: p. 2514796.

Figures and supplementary figures




Figure 1 | Static DEAE-dextran is a valid alternative to Pb for the transduction of mesenchymal stromal cells. (a) Percentage of transduced UC cells expressing the ZsGreen as evaluated via flow cytometry. (b) Levels of ATP produced by transduced UC-MSCs after 7 days of culture following the application of the different protocols. (c) Percentage of transduced BM cells expressing ZsGreen as evaluated via flow cytometry. (d) Levels of ATP produced by transduced BM-MSCs after 10 days of culture following the application of the different protocols. (e) Percentage of transduced A-MSCs expressing the ZsGreen as evaluated via flow cytometry. (f) Levels of ATP produced by transduced A-MSCs cells after 7 days of culture following the application of the different protocols. All data are displayed as mean \pm SD with $n = 3$ individual donors for each tissue source. For the ATP assay each donor sample was measured in triplicate and averaged. Data are normalised to the untransduced control group in static condition and expressed as percentage. Charts display only the data from transduced cells. ATP assay data from untransduced cells are displayed in Supplementary Figure 5. Statistical analysis applies only to the ATP assay and was performed on whole dataset as shown in supplementary Figure 5a-c: Three-way ANOVA with Dunnett post-hoc comparison test against the respective untransduced static control; * $p < 0.05$, ** $p < 0.01$, \$ $p < 0.0001$.




Figure 2 | Application of the DEAE-dextran protocol to transduce MSCs with four different lentiviral particles. Percentage of transduced and control UC- (a), BM- (b) and A-MSCs (c) expressing either the tdTomato, the dTomato, the eGFP or the mCherry protein as measured via flow cytometry. Data are displayed as mean \pm SD, $n = 3$ individual donors for each tissue source. Two-way ANOVA analysis with Sidak's multiple comparison post-hoc test; \$ $p \leq 0.0001$ and # $p \leq 0.0005$.




Figure 3 | Luc2-ZsGreen⁺ MSCs display similar properties to untransduced cells. (a) Representative phase contrast and green fluorescence images of UC, BM and A-MSCs after sorting compared to their untransduced counterparts. Scale bar = 200 μm . (b) Average doubling times of transduced and untransduced cells. Data are displayed as mean \pm SD, n = 3 individual donors. Two-way ANOVA analysis with Sidak's multiple comparisons post-hoc test. (c) Fluorescence images of cells 16h after seeding stained with phalloidin (f-actin, red) and DAPI (nuclei, blue) acquired at 100x. Scale bar = 200 μm . (d-e) Average cell area (d) and circularity (e) of transduced and untransduced MSCs. Full dataset shown in Supplementary Figure 8. FLuc = Luc2-ZsGreen⁺ cells. Data are displayed as mean \pm SD, n = 3 individual donors. Two-way ANOVA analysis with Sidak's multiple comparisons post-hoc test. Abbreviations: CU = circular unit (1 = perfect circle, 0 = elongated polygon).







Figure 4 | Light output of Luc2-ZsGreen⁺ MSCs as a function of cell density. Luc2-ZsGreen⁺ expressing MSCs were seeded at a density of 156 to 2×10^4 cells/well and treated with saturating concentration of D-Luciferin (5.12 mM D-Luciferin). **(a-c)** Light output (flux) as a function of cell concentration, with linear regression curves of UC **(a)**, BM **(b)** and A **(c)** MSCs for each individual donor sample. The slope of each curve represents the flux/cell and is shown in the legend of the respective graph. Data are displayed as mean \pm SD from $n = 3$ replicates. Missing values are due to saturation of the signal (A-02) or unexpected dropping with 2×10^4 cells (BM-02). **(d)** Average of the total amount of photon/second emitted by each cell type following transduction. Data obtained by averaging the results from individual donors. Data are displayed as mean \pm SD from $n = 3$. One-way ANOVA with Tukey's multiple comparison post-hoc test.




Supplementary Figure 1 | Lentiviral vector backbones. (a-b) Schematic representation of the lentiviral vectors used to generate Luc2-ZsGreen⁺ cells **(a)** and to perform the validation of the DEAE-Dextran protocol **(b)**.




Supplementary Figure 2 | Effect of polycations and centrifugation on UC-MSC transduction. (a) Representative phase contrast images of untransduced and transduced UC-MSCs in static condition. The images of the transduced cells are coupled with the green fluorescence channel to show the expression of the ZsGreen protein. Scale bar 200 μ m. **(b)** Representative phase contrast images of untransduced and transduced UC-MSCs following the application of the centrifugation protocol. The images of the transduced cells are coupled with the green fluorescence channel to show the expression of the ZsGreen protein. Scale bar 200 μ m. All fluorescence images acquired under the same acquisition conditions.




Supplementary Figure 3 | Effect of polycations and centrifugation on BM-MSCs transduction. (a) Representative phase contrast images of untransduced and transduced bone marrow MSCs in static condition. The images of the transduced cells are coupled with the green fluorescence channel to show the expression of the ZsGreen protein. Scale bar 200 μ m. (b) Representative phase contrast images of untransduced and transduced bone marrow derived MSCs following the application of the centrifugation protocol. The images of the transduced cells are coupled with the green fluorescence channel to show the expression of the ZsGreen protein. Scale bar 200 μ m.




Supplementary Figure 4 | Effect of polycations and centrifugation on A-MSCs transduction. (a) Representative phase contrast images of untransduced and transduced adipose derived MSCs in static condition. The images of the transduced cells are coupled with the green fluorescence channel to show the expression of the ZsGreen protein. Scale bar 200 μm . **(b)** Representative phase contrast images of untransduced and transduced adipose derived MSCs following the application of the centrifugation protocol. The images of the transduced cells are coupled with the green fluorescence channel to show the expression of the ZsGreen protein. Scale bar 200 μm .




Supplementary Figure 5 | ATP assay of transduced and untransduced MSCs. (a-c) Data from UC (a), BM (b) and A-MSCs (c) separated by static or centrifuged protocols. Data are displayed as mean \pm SD from $n = 3$ donors for each tissue source. Each donor sample was measured in triplicate and averaged. Three-way ANOVA with Dunnett post-hoc comparison test against untransduced static control; * $p < 0.05$; ** $p < 0.01$; # $p < 0.0005$; \$ $p < 0.0001$.



Supplementary Figure 6 | The application of the DEAE-dextran static protocol increases the transduction efficiency of all types of MSCs with all the lentiviral particles tested. (a-c) representative fluorescence images of UC- **(a)**, BM- **(b)** and A- **(c)** MSCs transduced with four different lentiviral particles with and without the use of DEAE-dextran as a polycation. Scale bar 200 μ m. Contrast in the images from eGFP cells was enhanced to facilitate the identification of transduced cells.



Supplementary Figure 7 | Luc2-ZsGreen^+ MSCs display similar proliferation to untransduced cells. (a) Cumulative doublings from p7 to p10 (UC-02 and UC-03) and from p7 to p11 (UC-01) for the 3 UC-MSC and respective controls. (b) Cumulative doubling from p5 to p7 (BM-01 and BM-03) and from p7 to p10 (BM-02) of the 3 BM-MSC and respective controls. (c) Cumulative doubling from p7 to p9 (A-02) and p7 to p10 (A-01 and A-03) of the 3 A-MSC samples and respective controls. (d) Average doubling time of transduced and untransduced cells displayed by tissue of origin. FLuc = Luc2-ZsGreen^+ cells. Data are displayed as mean \pm SD, $n \geq 2$ independent experiments. Two-way ANOVA analysis with Sidak's multiple comparisons post-hoc test.



Supplementary Figure 8 | Morphological characterisation of transduced MSCs. (a-b) Violin plot of the area **(a)** and the circularity **(b)** of the transduced and untransduced UC-MSC populations (at least 55 cells from each sample were analysed). **(c-d)** Violin plot of the area **(c)** and the circularity **(d)** of the transduced and untransduced BM-MSC populations (at least 60 cells for each sample were analysed). **(e-f)** Violin plot of the area **(e)** and the circularity **(f)** of the transduced and untransduced A-MSC populations (at least 64 cells for each sample were analysed). For all the graphs, the data were plotted into a grouped graph and cleaned from the outliers using the automated GraphPad tool “remove outliers” using the ROUT method with a $Q = 1\%$. A two-way ANOVA was performed on the cleaned data with a Dunn’s multiple comparison post-hoc test; * $p < 0.05$, **** $p < 0.0001$.

Tables

Table 1: UC, BM and A-MSCs samples and the respective passages at which they were transduced, sorted and used for the experiments.

Cell type	Donor ID	Transduction Passage	Sorting Passage	Proliferation analysis from	Characterisation*
UC-MSCs	UC-01	P5	P6	P7	P8
	UC-02	P5	P6	P7	P8
	UC-03	P5	P6	P7	P8
BM-MSCs	BM-01	P3	P4	P5	P6
	BM-02	P5	P6	P7	P8
	BM-03	P3	P4	P5	P6
A-MSCs	A-01	P5	P6	P7	P8
	A-02	P5	P6	P7	P8
	A-03	P5	P6	P7	P8

*Characterisation includes flow cytometry, morphological and BLI analysis

Table 2: Percentage of Luc2-ZsGreen cells expressing the ZsGreen protein and percentage of untransduced controls and transduced cells positive to CD44, CD45, CD73, CD90 and CD105. FLuc = Luc2-ZsGreen⁺ cells.

Donor ID	Positive cells (ZsGreen)	CD45	CD44	CD73	CD90	CD105
UC-01 Ctr	N/A	1.75%	99.20%	99.88%	99.95%	99.83%
UC-01 FLuc	98.62%	2.68%	99.75%	99.94%	99.95%	99.79%
UC-02 Ctr	N/A	2.96%	99.20%	99.77%	100.00%	99.57%
UC-02 FLuc	99.08%	4.59%	99.65%	99.80%	99.95%	99.49%
UC-03 Ctr	N/A	1.84%	97.37%	99.80%	99.96%	99.95%
UC-03 FLuc	99.36%	5.01%	95.30%	99.74%	99.93%	99.81%
BM-01 Ctr	N/A	6.23%	99.85%	96.84%	100.00%	98.49%
BM-01 FLuc	99.84%	6.96%	99.97%	99.72%	99.99%	99.64%
BM-02 Ctr	N/A	2.17%	99.91%	99.74%	99.83%	97.64%
BM-02 FLuc	99.49%	1.98%	99.96%	98.99%	99.86%	93.58%
BM-03 Ctr	N/A	1.50%	99.91%	99.90%	99.84%	99.04%
BM-03 FLuc	99.86%	1.34%	100.00%	99.88%	99.70%	98.63%
A-01 Ctr	N/A	9.11%	99.81%	99.89%	99.98%	98.82%
A-01 FLuc	99.04%	4.07%	100.00%	99.50%	99.99%	93.79%
A-02 Ctr	N/A	9.28%	99.61%	99.53%	99.99%	98.61%
A-02 FLuc	99.45%	75.52%	99.98%	99.39%	99.98%	99.87%
A-03 Ctr	N/A	12.96%	99.94%	99.84%	100.00%	99.96%
A-03 FLuc	99.40%	11.20%	99.88%	99.65%	99.97%	99.75%