

A formula for the linking number in terms of isometry invariants of straight line segments

Matt Bright, Olga Anosova, Vitaliy Kurlin
Computer Science and Materials Innovation Factory, University of Liverpool
sgmbrigh@liverpool.ac.uk, oanosova@liv.ac.uk, vitaliy.kurlin@gmail.com

Abstract The linking number is usually defined as an isotopy invariant of two non-intersecting closed curves in 3-dimensional space. However, the original definition in 1833 by Gauss in the form of a double integral makes sense for any open disjoint curves considered up to rigid motion. Hence the linking number can be studied as an isometry invariant of rigid structures consisting of straight line segments. For the first time this paper gives a complete proof for an explicit analytic formula for the linking number of two line segments in terms of six isometry invariants, namely the distance and angle between the segments and four coordinates of their endpoints in a natural coordinate system associated with the segments. Motivated by interpenetration of crystalline networks, we discuss potential extensions to infinite periodic structures and review recent advances in isometry classifications of periodic point sets.

1 The Gauss integral for the linking number of disjoint curves

This extended version of the conference paper [8] includes all previously skipped proofs. For any vectors $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathbb{R}^3$, the *triple product* is $(\mathbf{u}, \mathbf{v}, \mathbf{w}) = (\mathbf{u} \times \mathbf{v}) \cdot \mathbf{w}$.

Definition 1 (Gauss integral for the linking number) For piecewise-smooth curves $\gamma_1, \gamma_2 : [0, 1] \rightarrow \mathbb{R}^3$, the *linking number* can be defined as the Gauss integral [15]

$$(1) \quad \text{lk}(\gamma_1, \gamma_2) = \frac{1}{4\pi} \int_0^1 \int_0^1 \frac{(\dot{\gamma}_1(t), \dot{\gamma}_2(s), \gamma_1(t) - \gamma_2(s))}{|\gamma_1(t) - \gamma_2(s)|^3} dt ds,$$

where $\dot{\gamma}_1(t), \dot{\gamma}_2(s)$ are the vector derivatives of the 1-variable functions $\gamma_1(t), \gamma_2(s)$. \square

The formula in Definition 1 gives an integer number for any closed disjoint curves γ_1, γ_2 due to its interpretation as the *degree* of the *Gauss map* $\Gamma(t, s) = \frac{\gamma_1(t) - \gamma_2(s)}{|\gamma_1(t) - \gamma_2(s)|} : S^1 \times S^1 \rightarrow S^2$, i.e. $\deg \Gamma = \frac{\text{area}(\Gamma(S^1 \times S^1))}{\text{area}(S^2)}$, where the area of

the unit sphere is $\text{area}(S^2) = 4\pi$. This integer degree is the *linking number* of the 2-component link $\gamma_1 \sqcup \gamma_2 \subset \mathbb{R}^3$ formed by the two closed curves. Invariance modulo continuous deformation of \mathbb{R}^3 follows easily for closed curves - indeed, the function under the Gauss integral in (1), and hence the integral itself, varies continuously under perturbations of the curves γ_1, γ_2 . This should keep any integer value constant.

For open curves γ_1, γ_2 , the Gauss integral gives a real but not necessarily integral value, which remains invariant under rigid motions or orientation-preserving isometries (see Theorem 2). In \mathbb{R}^3 with the Euclidean metric isometries consist of rotations, translations and reflections. Isometry invariance of the real-valued linking number for open curves has found applications in the study of molecules [1].

Any smooth curve can be well-approximated by a polygonal line, so the computation of the linking number reduces to a sum over pairs of straight line segments L_1, L_2 . In 1976 Banchoff [6] has expressed the linking number $\text{lk}(L_1, L_2)$ in terms of the endpoints of each segment, see details of this and other past work in Section 3.

In 2000 Klenin and Langowski [16] proposed a formula for the linking number $\text{lk}(L_1, L_2)$ of two straight line segments in terms of six isometry invariants of L_1, L_2 , referring to a previous paper [33], which used the formula without any detailed proof. The paper [16] also skipped all details of the invariant-based formula's derivation.

The usefulness of the invariant-based formula can be seen by considering the analogy with the simpler concept of the scalar (dot) product of vectors. The algebraic or *coordinate-based* formula expresses the scalar product of two vectors $\mathbf{u} = (x_1, y_1, z_1)$ and $\mathbf{v} = (x_2, y_2, z_2)$ as $\mathbf{u} \cdot \mathbf{v} = x_1x_2 + y_1y_2 + z_1z_2$, which in turn depend on the coordinates of their endpoints. However, the scalar product for high-dimensional vectors $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$ can also be expressed in terms of only 3 parameters $\mathbf{u} \cdot \mathbf{v} = |\mathbf{u}| \cdot |\mathbf{v}| \cos \angle(\mathbf{u}, \mathbf{v})$. The two lengths $|\mathbf{u}|, |\mathbf{v}|$ and the angle $\angle(\mathbf{u}, \mathbf{v})$ are isometry invariants of the vectors \mathbf{u}, \mathbf{v} . This second geometric or *invariant-based* formula makes it clear that $\mathbf{u} \cdot \mathbf{v}$ is an isometry invariant, while it is harder to show that $\mathbf{u} \cdot \mathbf{v} = x_1x_2 + y_1y_2 + z_1z_2$ is invariant under rotations. It also provides other geometric insights that are hard to extract from the coordinate-based formula - for example, $\mathbf{u} \cdot \mathbf{v}$ oscillates as a cosine wave when the lengths $|\mathbf{u}|, |\mathbf{v}|$ are fixed, but the angle $\angle(\mathbf{u}, \mathbf{v})$ is varying.

In this paper, we provide a detailed proof of the invariant-based formula for the linking number in Theorem 8 and new corollaries in Section 6 formally investigating the asymptotic behaviour of the linking number, which wasn't previously studied.

Our own interest in the asymptotic behaviour is motivated by the *periodic linking number* by Panagiotou [26] as an invariant of crystalline networks [12] that are infinitely periodic in three directions, by calculating the infinite sum of the linking number between one line segment and all translated copies of another such segment.

2 Outline of the invariant-based formula and consequences

Folklore Theorem 2 lists key properties of $\text{lk}(\gamma_1, \gamma_2)$, which will be used later.

Theorem 2 (properties of the linking number) The linking number defined by the Gauss integral in Definition 1 for smooth curves γ_1, γ_2 has the following properties:

- (2a) the linking number is symmetric: $\text{lk}(\gamma_1, \gamma_2) = \text{lk}(\gamma_2, \gamma_1)$;
- (2b) $\text{lk}(\gamma_1, \gamma_2) = 0$ for any curves γ_1, γ_2 that belong to the same plane;
- (2c) $\text{lk}(\gamma_1, \gamma_2)$ is independent of orientation-preserving parameterisations of the open curves γ_1, γ_2 with fixed endpoints;
- (2d) $\text{lk}(-\gamma_1, \gamma_2) = -\text{lk}(\gamma_1, \gamma_2)$, where $-\gamma_1$ has the reversed orientation of γ_1 ;
- (2e) the linking number $\text{lk}(\gamma_1, \gamma_2)$ is invariant under any scaling $\mathbf{v} \rightarrow \lambda \mathbf{v}$ for $\lambda > 0$;
- (2f) $\text{lk}(\gamma_1, \gamma_2)$ is multiplied by $\det M$ under any orthogonal map $\mathbf{v} \mapsto M\mathbf{v}$. \square

Proof (2a) We note that the Euclidean distance is symmetric, and that since the triple product is anti-symmetric and $\gamma_2(s) - \gamma_1(t) = -(\gamma_1(t) - \gamma_2(s))$, the symmetry follows from

$$\begin{aligned} (\dot{\gamma}_2(s), \dot{\gamma}_1(t), \gamma_2(s) - \gamma_1(t)) &= -(\dot{\gamma}_1(t), \dot{\gamma}_2(s), \gamma_2(s) - \gamma_1(t)) \\ &= -(\dot{\gamma}_1(t), \dot{\gamma}_2(s), -(\gamma_1(t) - \gamma_2(s))) \\ &= (\dot{\gamma}_1(t), \dot{\gamma}_2(s), \gamma_1(t) - \gamma_2(s)). \end{aligned}$$

(2b) is obvious from the coplanarity of the normal vectors.

(2c) is simply a consequence of the path-independence of the integrals over s, t .

(2d) follows from $\dot{\gamma}_1(1-t) = -\dot{\gamma}_1(t)$ since the reverse orientation of $\gamma_1(t)$ is $\gamma_1(1-t)$.

(2e) Any scaling $\mathbf{v} \mapsto \lambda \mathbf{v}$ will result in a change of parameterisation $\gamma_i(t) \mapsto \lambda \gamma_i(t)$. Since $\lambda \dot{\gamma}_i(t) = \dot{\lambda \gamma_i(t)}$, the result follows below

$$\begin{aligned} \text{lk}(\lambda \gamma_1(t), \lambda \gamma_2(s)) &= \frac{1}{4\pi} \int_0^1 \int_0^1 \frac{(\lambda \dot{\gamma}_1(t), \lambda \dot{\gamma}_2(s), \lambda(\gamma_1(t) - \gamma_2(s)))}{|\lambda(\gamma_1(t) - \gamma_2(s))|^3} dt ds \\ &= \frac{1}{4\pi} \int_0^1 \int_0^1 \frac{\lambda^3 (\dot{\gamma}_1(t), \dot{\gamma}_2(s), \gamma_1(t) - \gamma_2(s))}{\lambda^3 |\gamma_1(t) - \gamma_2(s)|^3} dt ds \\ &= \text{lk}(\gamma_1(t), \gamma_2(s)), \end{aligned}$$

(2f) For an orthogonal transformation M , we have $M\mathbf{u} \times M\mathbf{v} = (\det M)M(\mathbf{u} \times \mathbf{v})$, while $M\mathbf{u} \cdot M\mathbf{v} = \mathbf{u} \cdot \mathbf{v}$. Therefore $|M\mathbf{v} - M\mathbf{u}| = |\mathbf{v} - \mathbf{u}|$, $(M\mathbf{u}, M\mathbf{v}, M\mathbf{w}) = \det M(\mathbf{u}, \mathbf{v}, \mathbf{w})$ and $\text{lk}(M\gamma_1, M\gamma_2) = (\det M)(\text{lk}(\gamma_1, \gamma_2))$ as expected. \blacksquare

Our main Theorem 8 will prove an analytic formula for the linking number of any line segments L_1, L_2 in terms of 6 isometry invariants of L_1, L_2 , which are introduced in Lemma 5. Simpler Corollary 3 expresses $\text{lk}(L_1, L_2)$ for any simple orthogonal oriented segments L_1, L_2 defined by their lengths $l_1, l_2 > 0$ and initial endpoints O_1, O_2 , respectively, with the Euclidean distance $d(O_1, O_2) = d > 0$, so

that $\mathbf{L}_1, \mathbf{L}_2, \overrightarrow{O_1O_2}$ form a positively oriented orthogonal basis whose signed volume $(\mathbf{L}_1, \mathbf{L}_2, \overrightarrow{O_1O_2}) = l_1 l_2 d$ is the product of the lengths, see the first picture in Fig. 1.

Corollary 3 (linking number for simple orthogonal segments) For any simple orthogonal oriented line segments $L_1, L_2 \subset \mathbb{R}^3$ with lengths l_1, l_2 and a distance d as defined above, the linking number is $\text{lk}(L_1, L_2) = -\frac{1}{4\pi} \arctan\left(\frac{l_1 l_2}{d\sqrt{l_1^2 + l_2^2 + d^2}}\right)$. \square

The above expression is a special case of general formula (8) for $a_1 = a_2 = 0$ and $\alpha = \frac{\pi}{2}$. Both formulae are invariant under the uniform scaling of \mathbb{R}^3 by λ , which agrees with Theorem 2. If $l_1 = l_2 = l$, the linking number in Corollary 3 becomes $\text{lk}(L_1, L_2) = -\frac{1}{4\pi} \arctan \frac{l^2}{d\sqrt{2l^2 + d^2}}$. If $l_1 = l_2 = \frac{d}{2}$, then $\text{lk}(L_1, L_2) = \frac{1}{2\pi} \left(\arcsin \frac{1}{\sqrt{2.5}} - \frac{\pi}{4} \right) \approx -0.016$. If $l_1 = l_2 = d$, then $\text{lk}(L_1, L_2) = -\frac{1}{4\pi} \arctan \frac{1}{\sqrt{3}} = -\frac{1}{24} \approx -0.0417$.

Corollary 3 implies that the linking number is in the range $(-\frac{1}{8}, 0)$ for any simple orthogonal segments with $d > 0$, which wasn't obvious from Definition 1. If L_1, L_2 move away from each other, then $\lim_{d \rightarrow +\infty} \text{lk}(L_1, L_2) = -\frac{1}{4\pi} \arctan 0 = 0$.

Alternatively, if segments with $l_1 = l_2 = l$ become infinitely short, the limit is again zero: $\lim_{l \rightarrow 0} \text{lk}(L_1, L_2) = 0$ for any fixed d . The limit $\lim_{x \rightarrow +\infty} \arctan x = \frac{\pi}{2}$ implies that if segments with $l_1 = l_2 = l$ become infinitely long for a fixed distance d , $\lim_{l \rightarrow +\infty} \text{lk}(L_1, L_2) = -\frac{1}{4\pi} \arctan \frac{l^2}{d\sqrt{2l^2 + d^2}} = -\frac{1}{8}$. If we push the segments L_1, L_2 , which have fixed (possibly different) lengths l_1, l_2 towards each other, the same limit similarly emerges: $\lim_{d \rightarrow 0} \text{lk}(L_1, L_2) = -\frac{1}{8}$. See more general corollaries in section 6.

3 Past results about the Gauss integral for the linking number

The survey [28] reviews the history of the Gauss integral, its use in Maxwell's description of electromagnetic fields [24], and its interpretation as the degree of a map from the torus to the sphere. In classical knot theory $\text{lk}(\gamma_1, \gamma_2)$ is a topological invariant of a link consisting of closed curves $\gamma_1 \sqcup \gamma_2$, whose equivalence relation is ambient isotopy. This relation is too flexible for open curves which can be isotopically unwound, and hence doesn't preserve the Gauss integral for open curves γ_1, γ_2 .

Computing the value of the Gauss integral directly from the parametric equation of two generic curves is only possible by approximation, but this problem is simplified

when we consider straight lines. The first form of the linking number between two straight line segments in terms of their geometry is described by Banchoff [6]. Banchoff considers the projection of segments on to a plane orthogonal to some vector $\xi \in S^2$. The Gauss integral is interpreted as the fraction of the unit sphere covered by those directions of ξ for which the projection will have a crossing.

This interpretation was the foundation of a closed form developed by Arai [5], using van Oosterom and Strackee's closed formula for the solid angle subtended by a tetrahedron given by the origin of a sphere and three points on its surface. An efficient implementation of the solid angle approach to the linking number is discussed in [7].

An alternative calculation for this solid angle is given in [27] as a starting point for calculating further invariants of open entangled curves. This form does not employ geometric invariants, but was used in [16] to claim a formula (without a proof) similar to Theorem 8, which is proved in this paper with more corollaries in section 6.




Fig. 1 Each line segment L_i is in the plane $\{z = (-1)^i \frac{d}{2}\}$, $i = 1, 2$. **Left:** signed distance $d > 0$, the endpoint coordinates $a_1 = 0, b_1 = 1$ and $a_2 = 0, b_2 = 1$, the lengths $l_1 = l_2 = 1$. **Right:** signed distance $d < 0$, the endpoint coordinates $a_1 = -1, b_1 = 1$ and $a_2 = -1, b_2 = 1$, so $l_1 = l_2 = 2$. In both middle pictures $\alpha = \frac{\pi}{2}$ is the angle from $\text{pr}_{xy}(L_1)$ to $\text{pr}_{xy}(L_2)$ with x -axis as the bisector.

4 Six isometry invariants of skew line segments in 3-space

This section introduces six isometry invariants, which uniquely determine positions of any line segments $L_1, L_2 \subset \mathbb{R}^3$ modulo isometries of \mathbb{R}^3 , see Lemma 5.

It suffices to consider only *skew* line segments that do not belong to the same 2-dimensional plane. If L_1, L_2 are in the same plane Π , for example if they are parallel, then $\dot{L}_1(t) \times \dot{L}_2(s)$ is orthogonal to any vector $L_1(t) - L_2(s)$ in the plane Π , hence $\text{lk}(L_1, L_2) = 0$. We denote by $\bar{L}_1, \bar{L}_2 \subset \mathbb{R}^3$ the infinite oriented lines through the given line segments L_1, L_2 , respectively. In a plane with fixed coordinates x, y , all angles are measured anticlockwise from the positive x -axis.

Definition 4 (invariants of line segments) Let $\alpha \in [0, \pi]$ be the angle between oriented line segments $L_1, L_2 \subset \mathbb{R}^3$. Assuming that L_1, L_2 are not parallel, there is a unique pair of parallel planes $\Pi_i, i = 1, 2$, each containing the infinite line \bar{L}_i through the line segment L_i . We choose orthogonal coordinates x, y, z in \mathbb{R}^3 so that

- (4a) the horizontal plane $\{z = 0\}$ is in the middle between Π_1, Π_2 , see Fig. 1;
- (4b) $(0, 0, 0)$ is the intersection of the projections $\text{pr}_{xy}(\bar{L}_1), \text{pr}_{xy}(\bar{L}_2)$ to $\{z = 0\}$;
- (4c) the x -axis bisects the angle α from $\text{pr}_{xy}(\bar{L}_1)$ to $\text{pr}_{xy}(\bar{L}_2)$, the y -axis is chosen so that α is anticlockwise measured from the x -axis to the y -axis in $\{z = 0\}$;
- (4d) the z -axis is chosen so that x, y, z are oriented in the right hand way, then d is the signed distance from Π_1 to Π_2 ; the distance d is negative if the vector $\overrightarrow{O_1 O_2}$ is opposite to the positively oriented z -axis in Fig. 1.

Let a_i, b_i be the coordinates of the initial and final endpoints of the segments L_i in the infinite line \bar{L}_i whose origin is $O_i = \Pi_i \cap (z\text{-axis}) = (0, 0, (-1)^i \frac{d}{2}), i = 1, 2$. \square

The case of segments L_1, L_2 lying in the same plane $\Pi \subset \mathbb{R}^3$ can be formally covered by Definition 4 if we allow the signed distance d from Π_1 to Π_2 to be 0.

Lemma 5 (parameterisation) Any oriented line segments $L_1, L_2 \subset \mathbb{R}^3$ are uniquely determined modulo a rigid motion by their isometry invariants $\alpha \in [0, \pi]$ and $d, a_1, b_1, a_2, b_2 \in \mathbb{R}$ from Definition 4. For $l_i = b_i - a_i, i = 1, 2$, each line segment L_i is

$$(5) \quad L_i(t) = \left((a_i + l_i t) \cos \frac{\alpha}{2}, (-1)^i (a_i + l_i t) \sin \frac{\alpha}{2}, (-1)^i \frac{d}{2} \right), t \in [0, 1].$$

Proof Any line segments $L_1, L_2 \subset \mathbb{R}^3$ that are not in the same plane are contained in distinct parallel planes. For $i = 1, 2$, the plane Π_i is spanned by L_i and the line parallel to L_{3-i} and passing through an endpoint of L_i . Let L'_i be the orthogonal projection of the line segment L_i to the plane Π_{3-i} . The non-parallel lines through the segments L_i and L'_{3-i} in the plane Π_i intersect at a point, say O_i . Then the line segment $O_1 O_2$ is orthogonal to both planes Π_i , hence to both L_i for $i = 1, 2$.

By Theorem 2, to compute $\text{lk}(L_1, L_2)$, one can apply a rigid motion to move the mid-point of the line segment $O_1 O_2$ to the origin $O = (0, 0, 0) \in \mathbb{R}^2$ and make $O_1 O_2$ vertical, i.e. lying within the z -axis. The signed distance d can be defined as the difference between the coordinates of $O_2 = \Pi_2 \cap (z\text{-axis})$ and $O_1 = \Pi_1 \cap (z\text{-axis})$ along the z -axis. Then L_i lies in the horizontal plane $\Pi_i = \{z = (-1)^i \frac{d}{2}\}, i = 1, 2$.

An extra rotation around the z -axis guarantees that the x -axis in the horizontal plane $\Pi = \{z = 0\}$ is the bisector of the angle $\alpha \in [0, \pi]$ from $\text{pr}_{xy}(\bar{L}_1)$ to $\text{pr}_{xy}(\bar{L}_2)$, where $\text{pr}_{xy} : \mathbb{R}^3 \rightarrow \Pi$ is the orthogonal projection. Then the infinite lines \bar{L}_i through L_i have the parametric form $(x, y, z) = (t \cos \frac{\alpha}{2}, (-1)^i t \sin \frac{\alpha}{2}, (-1)^i \frac{d}{2})$ with $s \in \mathbb{R}$.

The point O_i can be considered as the origin of the oriented infinite line \bar{L}_i . Let the line segment L_i have a length $l_i > 0$ and its initial point have the coordinate $a_i \in \mathbb{R}$ in the oriented line \bar{L}_i . Then the final endpoint of L_i has the coordinate $b_i = a_i + l_i$. To cover only the segment L_i , the parameter t should be replaced by $a_i + l_i t, t \in [0, 1]$. \blacksquare

If $t \in \mathbb{R}$ in Lemma 5, the corresponding point $L_i(t)$ moves along the line \bar{L}_i .

Lemma 6 (formulae for invariants) Let $L_1, L_2 \subset \mathbb{R}^3$ be any skewed oriented line segments given by their initial and final endpoints $A_i, B_i \in \mathbb{R}^3$ so that $\mathbf{L}_i = \overrightarrow{A_i B_i}$, $i = 1, 2$. Then the isometry invariants of L_1, L_2 in Lemma 5 are computed as follows:

the lengths $l_i = |\overrightarrow{A_i B_i}|$, the signed distance $d = \frac{[\mathbf{L}_1, \mathbf{L}_2, \overrightarrow{A_1 A_2}]}{|\mathbf{L}_1 \times \mathbf{L}_2|}$, the angle $\alpha = \arccos \frac{\mathbf{L}_1 \cdot \mathbf{L}_2}{l_1 l_2}$, $a_1 = \left(\frac{\mathbf{L}_2}{l_2} \cos \alpha - \frac{\mathbf{L}_1}{l_1} \right) \cdot \frac{\overrightarrow{A_1 A_2}}{\sin^2 \alpha}$, $a_2 = \left(\frac{\mathbf{L}_2}{l_2} - \frac{\mathbf{L}_1}{l_1} \cos \alpha \right) \cdot \frac{\overrightarrow{A_1 A_2}}{\sin^2 \alpha}$, $b_i = a_i + l_i, i = 1, 2$. \square

Proof The vectors along the segments are \mathbf{L}_i , hence the lengths are $l_i = |\mathbf{L}_i| = |\overrightarrow{A_i B_i}|$, $i = 1, 2$. The angle $\alpha \in [0, \pi]$ between $\mathbf{L}_1, \mathbf{L}_2$ can be found from the scalar product $\mathbf{L}_1 \cdot \mathbf{L}_2 = |\mathbf{L}_1| \cdot |\mathbf{L}_2| \cos \alpha$ as $\alpha = \arccos \frac{\mathbf{L}_1 \cdot \mathbf{L}_2}{l_1 l_2}$, because the function $\arccos x : [-1, 1] \rightarrow [0, \pi]$ is bijective. Since the vectors $\mathbf{L}_1, \mathbf{L}_2$ are not proportional to each other, the normalised vector product $\mathbf{e}_3 = \frac{\mathbf{L}_1 \times \mathbf{L}_2}{|\mathbf{L}_1 \times \mathbf{L}_2|}$ is well-defined and orthogonal to both $\mathbf{L}_1, \mathbf{L}_2$. Then $\mathbf{e}_1 = \frac{\mathbf{L}_1}{|\mathbf{L}_1|}$, $\mathbf{e}_2 = \frac{\mathbf{L}_2}{|\mathbf{L}_2|}$ and \mathbf{e}_3 have lengths 1 and form a linear basis of \mathbb{R}^3 , where the last vector is orthogonal to the first two.

Let O be any fixed point of \mathbb{R}^3 , which can be assumed to be the origin $(0, 0, 0)$ in the coordinates of Lemma 5, though its position relative to the vectors $\overrightarrow{A_i B_i}$ is not yet determined. First we express the points $O_i = (0, 0, (-1)^i \frac{d}{2}) \in \bar{L}_i$ from Fig. 3 in terms of given vectors $\overrightarrow{A_i B_i}$. If the initial endpoint A_i has a coordinate a_i in the line \bar{L}_i through the line segment L_i , then $\overrightarrow{O_i A_i} = a_i \mathbf{e}_i$ and

$$\overrightarrow{O_1 O_2} = \overrightarrow{O O_2} - \overrightarrow{O O_1} = (\overrightarrow{O A_2} - \overrightarrow{O_2 A_2}) - (\overrightarrow{O A_1} - \overrightarrow{O_1 A_1}) = \overrightarrow{A_1 A_2} + a_1 \mathbf{e}_1 - a_2 \mathbf{e}_2.$$

By Definition 4, $\overrightarrow{O_1 O_2}$ is orthogonal to the line \bar{L}_i going through the vector $\mathbf{e}_i = \frac{\mathbf{L}_i}{|\mathbf{L}_i|}$ for $i = 1, 2$. Then the product $[\mathbf{e}_1, \mathbf{e}_2, \overrightarrow{O_1 O_2}] = (\mathbf{e}_1 \times \mathbf{e}_2) \cdot \overrightarrow{O_1 O_2}$ equals $|\mathbf{e}_1 \times \mathbf{e}_2| d$, where $\overrightarrow{O_1 O_2}$ is in the z -axis, the signed distance d is the z -coordinate of O_2 minus the z -coordinates O_1 . The triple product $[\mathbf{e}_1, \mathbf{e}_2, \overrightarrow{O_1 O_2}] = (\mathbf{e}_1 \times \mathbf{e}_2) \cdot (\overrightarrow{A_1 A_2} + a_1 \mathbf{e}_1 - a_2 \mathbf{e}_2) = (\mathbf{e}_1 \times \mathbf{e}_2) \cdot \overrightarrow{A_1 A_2}$ doesn't depend on the parameters a_1, a_2 , because $\mathbf{e}_1 \times \mathbf{e}_2$ is orthogonal to both $\mathbf{e}_1, \mathbf{e}_2$. Hence the signed distance is $d = \frac{[\mathbf{e}_1, \mathbf{e}_2, \overrightarrow{A_1 A_2}]}{|\mathbf{e}_1 \times \mathbf{e}_2|} = \frac{[\mathbf{L}_1, \mathbf{L}_2, \overrightarrow{A_1 A_2}]}{|\mathbf{L}_1 \times \mathbf{L}_2|}$, which can be positive or negative, see Fig. 3.

It remains to find the coordinate a_i of the initial endpoint of L_i relative to the origin $O_i \in \bar{L}_i, i = 1, 2$. The vector $\overrightarrow{O_1 O_2} = \overrightarrow{A_1 A_2} + a_1 \mathbf{e}_1 - a_2 \mathbf{e}_2$ is orthogonal to both \mathbf{e}_i if and only if the scalar products vanish: $\overrightarrow{O_1 O_2} \cdot \mathbf{e}_i = 0$. Due to $|\mathbf{e}_1| = 1 = |\mathbf{e}_2|$ and $\mathbf{e}_1 \cdot \mathbf{e}_2 = \cos \alpha$, we get

$$\begin{cases} \mathbf{e}_1 \cdot \overrightarrow{A_1 A_2} + a_1 - a_2(\mathbf{e}_1 \cdot \mathbf{e}_2) = 0, \\ \mathbf{e}_2 \cdot \overrightarrow{A_1 A_2} + a_1(\mathbf{e}_1 \cdot \mathbf{e}_2) - a_2 = 0, \end{cases} \quad \begin{pmatrix} 1 & -\cos \alpha \\ \cos \alpha & -1 \end{pmatrix} \begin{pmatrix} a_1 \\ a_2 \end{pmatrix} = - \begin{pmatrix} \mathbf{e}_1 \cdot \overrightarrow{A_1 A_2} \\ \mathbf{e}_2 \cdot \overrightarrow{A_1 A_2} \end{pmatrix}.$$

The determinant of the 2×2 matrix is $\cos^2 \alpha - 1 = -\sin^2 \alpha \neq 0$, because L_1, L_2 are not parallel. Then $\begin{pmatrix} a_1 \\ a_2 \end{pmatrix} = \frac{1}{\sin^2 \alpha} \begin{pmatrix} -1 & \cos \alpha \\ -\cos \alpha & 1 \end{pmatrix} \begin{pmatrix} \mathbf{e}_1 \cdot \overrightarrow{A_1 A_2} \\ \mathbf{e}_2 \cdot \overrightarrow{A_1 A_2} \end{pmatrix}$. We get the formulae

$$\begin{aligned} a_1 &= \frac{-\mathbf{e}_1 \cdot \overrightarrow{A_1 A_2} + \cos \alpha (\mathbf{e}_2 \cdot \overrightarrow{A_1 A_2})}{\sin^2 \alpha} = \frac{(\mathbf{e}_2 \cos \alpha - \mathbf{e}_1) \cdot \overrightarrow{A_1 A_2}}{\sin^2 \alpha} = \\ &= \left(\frac{\mathbf{L}_2}{l_2} \cos \alpha - \frac{\mathbf{L}_1}{l_1} \right) \cdot \frac{\overrightarrow{A_1 A_2}}{\sin^2 \alpha}, \\ a_2 &= \frac{\cos \alpha (\mathbf{e}_1 \cdot \overrightarrow{A_1 A_2}) - \mathbf{e}_1 \cdot \overrightarrow{A_1 A_2}}{\sin^2 \alpha} = \frac{(\mathbf{e}_2 - \mathbf{e}_1 \cos \alpha) \cdot \overrightarrow{A_1 A_2}}{\sin^2 \alpha} = \\ &= \left(\frac{\mathbf{L}_2}{l_2} - \frac{\mathbf{L}_1}{l_1} \cos \alpha \right) \cdot \frac{\overrightarrow{A_1 A_2}}{\sin^2 \alpha}. \end{aligned}$$

The coordinates of the final endpoints are obtained as $b_i = a_i + l_i$ for $i = 1, 2$. ■

Lemma 7 guarantees that the linking number behaves symmetrically in d , meaning that we may confine any particular analysis to cases where $d > 0$ or $d < 0$.

Lemma 7 (symmetry) Let $L_1, L_2 \subset \mathbb{R}^3$ be parameterised as in Lemma 5. Under the central symmetry $\text{CS} : (x, y, z) \mapsto (-x, -y, -z)$ with respect to the origin $(0, 0, 0) \in \mathbb{R}^3$, the line segments keep their invariants $\alpha, a_1, b_1, a_2, b_2$. The signed distance d and the linking number change their signs: $\text{lk}(\text{CS}(L_1), \text{CS}(L_2)) = -\text{lk}(L_1, L_2)$. □

Proof Under the central symmetry CS , in the notation of Lemma 6 the vectors $\mathbf{L}_1, \mathbf{L}_2, \overrightarrow{A_1 A_2}$ change their signs. Then the formulae for $\alpha, a_1, b_1, a_2, b_2$ gives the same expression, but the triple product $[\mathbf{L}_1, \mathbf{L}_2, \overrightarrow{A_1 A_2}]$ and d change their signs.

Since the central symmetry CS is an orthogonal map M with $\det M = -1$, the new linking number changes its sign as follows: $\text{lk}(\text{CS}(L_1), \text{CS}(L_2)) = \text{lk}(\text{CS}(L_2), \text{CS}(L_1)) = -\text{lk}(L_1, L_2)$, where we also make use of the invariance of the linking number under exchange of the segments from Theorem 2(f). ■

5 Invariant-based formula for the linking number of segments

This section proves main Theorem 8, which expresses the linking number of two line segments in terms of their six isometry invariants from Definition 4. In 2000 Klenin and Langowski claimed a similar but a bit less symmetric formula [16], but gave no proof, which requires substantial lemmas below. For example, one of their six invariants differs from the signed distance d between oriented line segments.

Theorem 8 (invariant-based formula) For any line segments $L_1, L_2 \subset \mathbb{R}^3$ with invariants $\alpha \in (0, \pi)$, $a_1, b_1, a_2, b_2, d \in \mathbb{R}$ from Definition 4, we have $\text{lk}(L_1, L_2) =$

$$\frac{\text{AT}(a_1, b_2; d, \alpha) + \text{AT}(b_1, a_2; d, \alpha) - \text{AT}(a_1, a_2; d, \alpha) - \text{AT}(b_1, b_2; d, \alpha)}{4\pi}, \quad (8)$$

where $\text{AT}(a, b; d, \alpha) = \arctan\left(\frac{ab \sin \alpha + d^2 \cot \alpha}{d\sqrt{a^2 + b^2 - 2ab \cos \alpha + d^2}}\right)$. For $\alpha = 0$ or $\alpha = \pi$, we set $\text{AT}(a, b; d, \alpha) = \text{sign}(d)\frac{\pi}{2}$. We also set $\text{lk}(L_1, L_2) = 0$ when $d = 0$. \square

The expression $a^2 + b^2 - 2ab \cos \alpha$ is the squared third side of the triangle with the first two sides a, b and the angle α between them, hence is always non-negative. Also $a^2 + b^2 - 2ab \cos \alpha = 0$ only when the triangle degenerates for $a = \pm b$ and $\cos \alpha = \pm 1$. For $\alpha = 0$ or $\alpha = \pi$ when L_1, L_2 are parallel, $\text{lk}(L_1, L_2) = 0$ is guaranteed by $\text{AT}(a, b; d, \alpha) = \text{sign}(d)\frac{\pi}{2} = 0$ when $d = 0$ holds in addition to $\alpha = 0$ or $\alpha = \pi$.

The symmetry of the AT function in a, b , i.e. $\text{AT}(a, b; d, \alpha) = \text{AT}(b, a; d, \alpha)$ implies that $\text{lk}(L_1, L_2) = \text{lk}(L_2, L_1)$ by Theorem 8. Since the AT function is odd in d , i.e. $\text{AT}(a, b; -d, \alpha) = -\text{AT}(a, b; d, \alpha)$, Lemma 7 is also respected.

Proof (of Corollary 3) By definition any simple orthogonal line segments L_1, L_2 have the angle $\alpha = \frac{\pi}{2}$ and initial endpoints $a_1 = 0 = a_2$, hence $b_1 = l_1, b_2 = l_2$. Then (8) gives $\text{AT}(0, l_2; d, \frac{\pi}{2}) = 0$, $\text{AT}(l_1, 0; d, \frac{\pi}{2}) = 0$, $\text{AT}(0, 0; d, \frac{\pi}{2}) = 0$. Then

$$\text{lk}(L_1, L_2) = -\frac{1}{4\pi} \text{AT}(l_1, l_2; d, \alpha) = -\frac{1}{4\pi} \arctan\left(\frac{l_1 l_2}{d\sqrt{l_1^2 + l_2^2 + d^2}}\right). \quad \blacksquare$$

Figure 2 shows how the function $\text{AT}(a, b; d, \alpha)$ from Theorem 8 depends on 2 of 4 parameters when others are fixed. For example, if the angle $\alpha = \frac{\pi}{2}$ is fixed, then $\text{AT}(a, b; d, \frac{\pi}{2}) = \arctan\left(\frac{ab}{d\sqrt{a^2 + b^2 + d^2}}\right)$. If also $a = b$, the surface $\text{AT}(a, a; d, \frac{\pi}{2}) = \arctan\left(\frac{a^2}{d\sqrt{2a^2 + d^2}}\right)$ in the first picture of Figure 2 has the horizontal ridge $\text{AT}(0, 0; d, \frac{\pi}{2}) = 0$ and $\lim_{d \rightarrow 0} \text{AT}(a, a; d, \frac{\pi}{2}) = \text{sign}(d)\frac{\pi}{2}$ for $a \neq 0$. If d, α are free, but $a = 0$, then $\text{AT}(0, 0; d, \alpha) = \arctan\left(\frac{d^2 \cot \alpha}{d\sqrt{d^2}}\right) = \text{sign}(d) \arctan(\cot \alpha) = \text{sign}(d)(\frac{\pi}{2} - \alpha)$. Similarly, $\lim_{d \rightarrow \infty} \text{AT}(0, 0; d, \alpha) = \text{sign}(d)(\frac{\pi}{2} - \alpha)$, see the lines $\text{AT} = \frac{\pi}{2} - \alpha$ on the boundaries of the AT surfaces in the middle pictures of Figure 2.

Lemma 9 ($\text{lk}(L_1, L_2)$ is an integral in p, q) In the notations of Definition 4 we have

$$\text{lk}(L_1, L_2) = -\frac{1}{4\pi} \int_{a_1/d}^{b_1/d} \int_{a_2/d}^{b_2/d} \frac{\sin \alpha \, dp \, dq}{(1 + p^2 + q^2 - 2pq \cos \alpha)^{3/2}} \text{ for } d > 0. \quad \square$$




Fig. 2 The graph of $AT(a, b; d, \alpha) = \arctan\left(\frac{ab \sin \alpha + d^2 \cot \alpha}{d\sqrt{a^2 + b^2 - 2ab \cos \alpha + d^2}}\right)$, where 2 of 4 parameters are fixed. **Top left:** $l = b - a = 0$, $\alpha = \frac{\pi}{2}$. **Top right:** $l = d = -1$. **Middle left:** $a = 0$, $d = 1$. **Middle right:** $a = 0$, $l = 1$. **Bottom left:** $a = 1$, $\alpha = \frac{\pi}{2}$. **Bottom right:** $d = -1$, $\alpha = \frac{\pi}{2}$.

Proof Below we assume that $a_1, a_2, l_1, l_2, \alpha$ are given and $t, s \in [0, 1]$.

$$L_1(t) = ((a_1 + l_1 t) \cos \frac{\alpha}{2}, -(a_1 + l_1 t) \sin \alpha, -\frac{d}{2}),$$

$$L_2(s) = ((a_2 + l_2 s) \cos \frac{\alpha}{2}, (a_2 + l_2 s) \sin \alpha, \frac{d}{2}),$$

$$\dot{L}_1(t) = (l_1 \cos \frac{\alpha}{2}, -l_1 \sin \frac{\alpha}{2}, 0),$$

$$\dot{L}_2(s) = (l_2 \cos \frac{\alpha}{2}, l_2 \sin \frac{\alpha}{2}, 0),$$

$$\dot{L}_1(t) \times \dot{L}_2(s) = (0, 0, 2l_1 l_2 \sin \frac{\alpha}{2} \cos \frac{\alpha}{2}) = (0, 0, l_1 l_2 \sin \alpha),$$

$$L_1(t) - L_2(s) = ((a_1 - a_2 + l_1 t - l_2 s) \cos \alpha, -(a_1 + a_2 + l_1 t + l_2 s) \sin \alpha, -d),$$

$$(\dot{L}_1(t), \dot{L}_2(s), L_1(t) - L_2(s)) = -dl_1 l_2 \sin \alpha,$$

$$\begin{aligned}
\text{lk}(L_1, L_2) &= \frac{1}{4\pi} \int_0^1 \int_0^1 \frac{(\dot{L}_1(t), \dot{L}_2(s), L_1(t) - L_2(s))}{|L_1(t) - L_2(s)|^3} dt ds = \\
&= \frac{1}{4\pi} \int_0^1 \int_0^1 \frac{-dl_1 l_2 \sin \alpha dt ds}{(d^2 + (a_1 - a_2 + l_1 t - l_2 s)^2 \cos^2 \frac{\alpha}{2} + (a_1 + a_2 + l_1 t + l_2 s)^2 \sin^2 \frac{\alpha}{2})^{3/2}} = \\
&= \frac{dl_1 l_2 \sin \alpha}{4\pi} \int_0^1 \int_0^1 \frac{dt ds}{(d^2 + (a_1 - a_2 + l_1 t - l_2 s)^2 \cos^2 \frac{\alpha}{2} + (a_1 + a_2 + l_1 t + l_2 s)^2 \sin^2 \frac{\alpha}{2})^{3/2}}
\end{aligned}$$

To simplify the last integral, introduce the variables $p = \frac{a_1 + l_1 t}{d}$ and $q = \frac{a_2 + l_2 s}{d}$.

In the new variables p, q the expression under the power $\frac{3}{2}$ in the denominator becomes

$$\begin{aligned}
&d^2 + (pd - qd)^2 \cos^2 \frac{\alpha}{2} + (pd + qd)^2 \sin^2 \frac{\alpha}{2} = \\
&= d^2 \left(1 + (p^2 - 2pq + q^2) \cos^2 \frac{\alpha}{2} + (p^2 + 2pq + q^2) \sin^2 \frac{\alpha}{2} \right) = d^2 \times \\
&\left(1 + p^2 \left(\cos^2 \frac{\alpha}{2} + \sin^2 \frac{\alpha}{2} \right) + q^2 - 2pq \left(\cos^2 \frac{\alpha}{2} - \sin^2 \frac{\alpha}{2} \right) \right) = d^2 (1 + p^2 + q^2 - 2pq \cos \alpha)
\end{aligned}$$

The old variables are expressed as $t = \frac{pd - a_1}{l_1}$, $s = \frac{qd - a_2}{l_2}$ and have the differentials

$dt = \frac{d}{l_1} dp$, $ds = \frac{d}{l_2} dq$. Since $t, s \in [0, 1]$, the new variables p, q have the ranges $[\frac{a_1}{d}, \frac{b_1}{d}]$ and $[\frac{a_2}{d}, \frac{b_2}{d}]$, respectively. Then we get the required expression:

$$\begin{aligned}
\text{lk}(L_1, L_2) &= -\frac{dl_1 l_2 \sin \alpha}{4\pi} \int_{a_1/d}^{b_1/d} \int_{a_2/d}^{b_2/d} \frac{d^2}{l_1 l_2 d^3 (1 + p^2 + q^2 - 2pq \cos \alpha)^{3/2}} dp dq = \\
&= -\frac{1}{4\pi} \int_{a_1/d}^{b_1/d} \int_{a_2/d}^{b_2/d} \frac{\sin \alpha dp dq}{(1 + p^2 + q^2 - 2pq \cos \alpha)^{3/2}}.
\end{aligned}$$

Due to Lemma 7, the above computations assume that the signed distance $d > 0$. ■

Lemma 10 (the linking number as a single integral) In the notations of Definition 4 we have $\text{lk}(L_1, L_2) = \frac{I(a_2/d) - I(b_2/d)}{4\pi}$, where the function $I(r)$ is defined as

the single integral $I(r) = \int_{a_1/d}^{b_1/d} \frac{\sin \alpha (r - p \cos \alpha) dp}{(1 + p^2 \sin^2 \alpha) \sqrt{1 + p^2 + r^2 - 2pr \cos \alpha}}$ for $d > 0$. □

Proof Complete the square in the expression under power $\frac{3}{2}$ in Lemma 9:

$$1 + p^2 + q^2 - 2pq \cos \alpha = 1 + p^2 \sin^2 \alpha + (q - p \cos \alpha)^2.$$

The substitution $(q - p \cos \alpha)^2 = (1 + p^2 \sin^2 \alpha) \tan^2 \psi$ for the new variable ψ simplifies the sum of squares to $1 + \tan^2 \psi = \frac{1}{\cos^2 \psi}$. Since q varies within $[\frac{a_2}{d}, \frac{b_2}{d}]$, for any fixed $p \in [\frac{a_1}{d}, \frac{b_1}{d}]$, the range $[\psi_0, \psi_1]$ of ψ satisfies $\tan \psi_0 = \frac{\frac{a_2}{d} - p \cos \alpha}{\sqrt{1 + p^2 \sin^2 \alpha}}$ and $\tan \psi_1 = \frac{\frac{b_2}{d} - p \cos \alpha}{\sqrt{1 + p^2 \sin^2 \alpha}}$. Since we treat p, ψ as independent variables, the Jacobian of the substitution $(p, q) \mapsto (p, \psi)$ equals

$$\frac{\partial q}{\partial \psi} = \frac{\partial}{\partial \psi} \left(p \cos \alpha + \tan \psi \sqrt{1 + p^2 \sin^2 \alpha} \right) = \frac{\sqrt{1 + p^2 \sin^2 \alpha}}{\cos^2 \psi}.$$

In the variables p, ψ the expression under the double integral of Lemma 9 becomes

$$\begin{aligned} \frac{\sin \alpha \, dp \, dq}{(1 + p^2 + q^2 - 2pq \cos \alpha)^{3/2}} &= \frac{\sin \alpha \, dp}{((1 + p^2 \sin^2 \alpha) + (1 + p^2 \sin^2 \alpha) \tan^2 \psi)^{3/2}} \frac{\partial q}{\partial \psi} \, d\psi \\ &= \frac{\sin \alpha \, dp}{(1 + p^2 \sin^2 \alpha)^{3/2} (1 + \tan^2 \psi)^{3/2}} \frac{d\psi \sqrt{1 + p^2 \sin^2 \alpha}}{\cos^2 \psi} = \frac{\sin \alpha \, dp \cos \psi \, d\psi}{1 + p^2 \sin^2 \alpha}. \end{aligned}$$

$$\text{lk} = -\frac{1}{4\pi} \int_{a_1/d}^{b_1/d} \frac{\sin \alpha \, dp}{1 + p^2 \sin^2 \alpha} \int_{\psi_0}^{\psi_1} \cos \psi \, d\psi = \frac{1}{4\pi} \int_{a_1/d}^{b_1/d} \frac{\sin \alpha \, dp}{1 + p^2 \sin^2 \alpha} (\sin \psi_0 - \sin \psi_1).$$

We can express the sin functions for the bounds ψ_0, ψ_1 in terms of tan as $\sin \psi_0 = \frac{\tan \psi_0}{\sqrt{1 + \tan^2 \psi_0}}$. Using $\tan \psi_0 = \frac{\frac{a_2}{d} - p \cos \alpha}{\sqrt{1 + p^2 \sin^2 \alpha}}$ obtained above, we get

$$\sqrt{1 + \tan^2 \psi_0} = \sqrt{\frac{(1 + p^2 \sin^2 \alpha) + (\frac{a_2}{d} - p \cos \alpha)^2}{1 + p^2 \sin^2 \alpha}} = \sqrt{\frac{1 + p^2 + (\frac{a_2}{d})^2 - 2\frac{a_2}{d} p \cos \alpha}{1 + p^2 \sin^2 \alpha}}.$$

$$\sin \psi_0 = \frac{\frac{a_2}{d} - p \cos \alpha}{\sqrt{1 + p^2 \sin^2 \alpha}} \sqrt{\frac{1 + p^2 \sin^2 \alpha}{1 + p^2 + (\frac{a_2}{d})^2 - 2\frac{a_2}{d} p \cos \alpha}} = \frac{\frac{a_2}{d} - p \cos \alpha}{\sqrt{1 + p^2 + (\frac{a_2}{d})^2 - 2\frac{a_2}{d} p \cos \alpha}}$$

Then $\sin \psi_1$ has the same expression with a_2 replaced by b_2 . After substituting these expressions in the previous formula for the linking number, we get $\text{lk}(L_1, L_2) =$

$$\begin{aligned} &\frac{1}{4\pi} \int_{a_1/d}^{b_1/d} \frac{\sin \alpha \, dp}{1 + p^2 \sin^2 \alpha} \left(\frac{\frac{a_2}{d} - p \cos \alpha}{\sqrt{1 + p^2 + (\frac{a_2}{d})^2 - 2\frac{a_2}{d} p \cos \alpha}} - \frac{\frac{b_2}{d} - p \cos \alpha}{\sqrt{1 + p^2 + (\frac{b_2}{d})^2 - 2\frac{b_2}{d} p \cos \alpha}} \right) \\ &= \frac{I(a_2/d) - I(b_2/d)}{4\pi} \text{ for } I(r) = \int_{a_1/d}^{b_1/d} \frac{\sin \alpha (r - p \cos \alpha) dp}{(1 + p^2 \sin^2 \alpha) \sqrt{1 + p^2 + r^2 - 2pr \cos \alpha}}. \quad \blacksquare \end{aligned}$$

Lemma 11 ($I(r)$ via \arctan) The integral $I(r)$ in Lemma 10 can be found as

$$\int \frac{\sin \alpha(r - p \cos \alpha) dp}{(1 + p^2 \sin^2 \alpha) \sqrt{1 + p^2 + r^2 - 2pr \cos \alpha}} = \arctan \frac{pr \sin \alpha + \cot \alpha}{\sqrt{1 + p^2 + r^2 - 2pr \cos \alpha}} + C.$$

Proof The easy way is to differentiate $\arctan \omega$ for $\omega = \frac{pr \sin^2 \alpha + \cos \alpha}{\sin \alpha \sqrt{1 + p^2 + r^2 - 2pr \cos \alpha}}$ with respect to the variable p remembering that r, α are fixed parameters. For notational clarity, we use an auxiliary symbol for the expression under the square root:

$$R = 1 + p^2 + r^2 - 2pr \cos \alpha. \text{ Then } \omega = \frac{pr \sin^2 \alpha + \cos \alpha}{\sin \alpha \sqrt{R}} \text{ and}$$

$$\begin{aligned} \frac{d\omega}{dp} &= \frac{1}{R \sin \alpha} \left(r \sin^2 \alpha \sqrt{R} - (rp \sin^2 \alpha + \cos \alpha) \frac{2p - 2r \cos \alpha}{2\sqrt{R}} \right) = \\ &= \frac{1}{R \sqrt{R} \sin \alpha} \left(r \sin^2 \alpha (1 + p^2 + r^2 - 2pr \cos \alpha) - (rp \sin^2 \alpha + \cos \alpha) (p - r \cos \alpha) \right) = \\ &= \frac{rp^2 \sin^2 \alpha + r^3 \sin^2 \alpha - 2pr^2 \cos \alpha \sin^2 \alpha - rp^2 \sin^2 \alpha + pr^2 \cos \alpha \sin^2 \alpha - p \cos \alpha + r}{R \sqrt{R} \sin \alpha} \\ &= \frac{r^3 \sin^2 \alpha - pr^2 \cos \alpha \sin^2 \alpha - p \cos \alpha + r}{R \sqrt{R} \sin \alpha} = \frac{(r - p \cos \alpha)(1 + r^2 \sin^2 \alpha)}{R \sqrt{R} \sin \alpha}. \\ \frac{d}{dp} \arctan \omega &= \frac{1}{1 + \omega^2} \cdot \frac{d\omega}{dp} = \frac{(\sin \alpha \sqrt{R})^2}{(\sin \alpha \sqrt{R})^2 + (pr \sin^2 \alpha + \cos \alpha)^2} \cdot \frac{d\omega}{dp} = \\ &= \frac{R \sin^2 \alpha}{R \sin^2 \alpha + (p^2 r^2 \sin^4 \alpha + 2pr \sin^2 \alpha \cos \alpha + \cos^2 \alpha)} \cdot \frac{(r - p \cos \alpha)(1 + r^2 \sin^2 \alpha)}{R \sqrt{R} \sin \alpha} = \\ &= \frac{\sin \alpha}{\sqrt{R}} \cdot \frac{(r - p \cos \alpha)(1 + r^2 \sin^2 \alpha)}{\sin^2 \alpha (1 + p^2 + r^2 - 2pr \cos \alpha) + (p^2 r^2 \sin^4 \alpha + 2pr \sin^2 \alpha \cos \alpha + \cos^2 \alpha)} = \\ &= \frac{\sin \alpha (r - p \cos \alpha)(1 + r^2 \sin^2 \alpha)}{(1 + p^2 \sin^2 \alpha + r^2 \sin^2 \alpha + p^2 r^2 \sin^4 \alpha) \sqrt{R}} = \frac{\sin \alpha (r - p \cos \alpha)(1 + r^2 \sin^2 \alpha)}{(1 + p^2 \sin^2 \alpha)(1 + r^2 \sin^2 \alpha) \sqrt{R}} = \\ &= \frac{\sin \alpha (r - p \cos \alpha)}{(1 + p^2 \sin^2 \alpha) \sqrt{R}} = \frac{\sin \alpha (r - p \cos \alpha)}{(1 + p^2 \sin^2 \alpha) \sqrt{1 + p^2 + r^2 - 2pq \cos \alpha}}. \end{aligned}$$

Since we got the required expression under the integral $I(r)$, Lemma 11 is proved. ■

Proof (Theorem 8) Consider the right hand side of the equation in Lemma 11 as the 3-variable function $F(p, r; \alpha) = \arctan \left(\frac{pr \sin \alpha + \cot \alpha}{\sqrt{1 + p^2 + r^2 - 2pr \cos \alpha}} \right)$. The function in Lemma 10 is $I(r) = F(b_1/d, r; \alpha) - F(a_1/d, r; \alpha)$. By Lemma 10 $\text{lk}(L_1, L_2) =$

$$\frac{\left(F(b_1/d, a_2/d; \alpha) - F(a_1/d, a_2/d; \alpha)\right) - \left(F(b_1/d, b_2/d; \alpha) - F(a_1/d, b_2/d; \alpha)\right)}{4\pi}.$$

Rewrite a typical function from the numerator above as follows: $F(a/d, b/d; \alpha) =$

$$\arctan \frac{(ab/d^2) \sin \alpha + \cot \alpha}{\sqrt{1 + (a/d)^2 + (b/d)^2 - 2(ab/d^2) \cos \alpha}} = \arctan \frac{ab \sin \alpha + d^2 \cot \alpha}{d\sqrt{a^2 + b^2 - 2ab \cos \alpha + d^2}}.$$

If we denote the last expression as $\text{AT}(a, b; d, \alpha)$, required formula (8) follows.

In Lemma 9, Lemma 10 and above we have used that the signed distance d is positive. By Lemma 7 the signed distance d and $\text{lk}(L_1, L_2)$ simultaneously change their signs under a central symmetry, while all other invariants remain the same. Since $\text{AT}(a, b; -d, \alpha) = -\text{AT}(a, b; d, \alpha)$ due to the arctan function being odd, formula (8) holds for $d < 0$. The formula remains valid even for $d = 0$, when L_1, L_2 are in the same plane. The expected value $\text{lk}(L_1, L_2) = 0$ needs an explicit setting, see the discussion of the linking number discontinuity around $d = 0$ in Corollary 14. ■

6 The asymptotic behaviour of the linking number of segments

This section discusses how the linking number $\text{lk}(L_1, L_2)$ in Theorem 8 behaves with respect to the six parameters of line segments L_1, L_2 . Figure 3 shows how the linking number between two equal line segments varies with different pairs of parameters.

Corollary 12 (bounds of the linking number) For any line segments $L_1, L_2 \subset \mathbb{R}^3$, the linking number $\text{lk}(L_1, L_2)$ is between $\pm \frac{1}{2}$. □

Proof By Theorem 8 $\text{lk}(L_1, L_2)$ is a sum of 4 arctan functions divided by 4π . Since each arctan is strictly between $\pm \frac{\pi}{2}$, the linking number is between $\pm \frac{1}{2}$. ■

Corollary 13 (sign of the linking number) In the notation of Definition 4, we have $\lim_{\alpha \rightarrow 0} \text{lk}(L_1, L_2) = 0 = \lim_{\alpha \rightarrow \pi} \text{lk}(L_1, L_2)$. Any non-parallel L_1, L_2 have $\text{sign}(\text{lk}(L_1, L_2)) = -\text{sign}(d)$. So $\text{lk}(L_1, L_2) = 0$ if and only if $d = 0$ or $\alpha = 0$ or $\alpha = \pi$. □

Proof If $\alpha = 0$ or $\alpha = \pi$, then $\cot \alpha$ is undefined, so Theorem 8 sets $\text{AT}(a, b; d, \alpha) = \text{sign}(d) \frac{\pi}{2}$. Then $\text{lk}(L_1, L_2) = \text{sign}(d) \frac{\pi}{2} (1 + 1 - 1 - 1) = 0$.

Theorem 8 also specifies that $\text{lk}(L_1, L_2) = 0$ for $d = 0$. If $d \neq 0$ and $\alpha \rightarrow 0$ within $[0, \pi]$ while all other parameters remain fixed, then $d^2 \cot \alpha \rightarrow +\infty$. Hence each of the 4 arctan functions in Theorem 8 approaches $\frac{d\pi}{2}$, so $\text{lk}(L_1, L_2) \rightarrow 0$. The same conclusion similarly follows in the case $\alpha \rightarrow \pi$ when $d^2 \cot \alpha \rightarrow -\infty$.




Fig. 3 The linking number $lk(a, a + l; a, a + l; d, \alpha)$ from formula (8), where 2 of 4 parameters are fixed. **Top left:** $l = 1, \alpha = \frac{\pi}{2}$. **Top right:** $l = 1, d = -1$. **Middle left:** $a = 0, d = 1$. **Middle right:** $a = 0, l = 1$. **Bottom left:** $a = 0, \alpha = \frac{\pi}{2}$. **Bottom right:** $d = -1, \alpha = \frac{\pi}{2}$.

If L_1, L_2 are not parallel, the angle α between them belongs to $(0, \pi)$. If $d > 0$, Lemma 9 says that $lk(L_1, L_2) = -\frac{1}{4\pi} \int_{a_1/d}^{b_1/d} \int_{a_2/d}^{b_2/d} \frac{\sin \alpha dp dq}{(1 + p^2 + q^2 - 2pq \cos \alpha)^{3/2}}$. Since the function under the integral is strictly positive, $lk(L_1, L_2) < 0$. By Lemma 7 both $lk(L_1, L_2)$ simultaneously change their signs under a central symmetry. Hence the formula $\text{sign}(lk(L_1, L_2)) = -\text{sign}(d)$ holds for all d including $d = 0$ above. ■

Corollary 14 (lk for $d \rightarrow 0$) If the distance $d \rightarrow 0$ and the curves L_1, L_2 remain disjoint, the expression in formula (8) behaves continuously, so $\lim_{d \rightarrow 0} lk(L_1, L_2) = 0$. If $d \rightarrow 0$ and the interiors of L_1, L_2 intersect each other in the limit case $d = 0$, then $\lim_{d \rightarrow 0} lk(L_1, L_2) = -\frac{\text{sign}(d)}{2}$, where $d \rightarrow 0$ keeps its sign. □

Proof Recall that $\lim_{x \rightarrow \pm\infty} \arctan x = \pm \frac{\pi}{2}$. By Corollary 13 assume that $\alpha \neq 0, \alpha \neq \pi$, so $\alpha \in (0, \pi)$. Then $\sin \alpha > 0, a^2 + b^2 - 2ab \cos \alpha > (a - b)^2 \geq 0$ and

$$\begin{aligned} \lim_{d \rightarrow 0} \text{AT}(a, b; d, \alpha) &= \lim_{d \rightarrow 0} \arctan \left(\frac{ab \sin \alpha + d^2 \cot \alpha}{d \sqrt{a^2 + b^2 - 2ab \cos \alpha + d^2}} \right) = \\ &= \text{sign}(a) \text{sign}(b) \text{sign}(d) \frac{\pi}{2}, \text{ so Theorem 8 gives} \end{aligned}$$

$$\lim_{d \rightarrow 0} \text{lk}(L_1, L_2) = \frac{\text{sign}(d)}{8} (\text{sign}(a_1) - \text{sign}(b_1)) (\text{sign}(b_2) - \text{sign}(a_2)).$$

In the limit case $d = 0$, the line segments $L_1, L_2 \subset \{z = 0\}$ remain disjoint in the same plane if and only if both endpoint coordinates a_i, b_i have the same sign for at least one of $i = 1, 2$, which is equivalent to $\text{sign}(a_i) - \text{sign}(b_i) = 0$, i.e. $\lim_{d \rightarrow 0} \text{lk}(L_1, L_2) = 0$ from the product above. Hence formula (8) is continuous under $d \rightarrow 0$ for any non-crossing segments. Any segments that intersect in the plane $\{z = 0\}$ when $d = 0$ have endpoint coordinates $a_i < 0 < b_i$ for both $i = 1, 2$ and have the limit $\lim_{d \rightarrow 0} \text{lk}(L_1, L_2) = \frac{\text{sign}(d)}{8} (-1 - 1)(1 - (-1)) = -\frac{\text{sign}(d)}{2}$ as required. ■

Corollary 15 (lk for $d \rightarrow \pm\infty$) If the distance $d \rightarrow \pm\infty$, then $\text{lk}(L_1, L_2) \rightarrow 0$. □

Proof If $d \rightarrow \pm\infty$, while other parameters of L_1, L_2 remain fixed, then the function $\text{AT}(a, b; d, \alpha) = \arctan \left(\frac{ab \sin \alpha + d^2 \cot \alpha}{d \sqrt{a^2 + b^2 - 2ab \cos \alpha + d^2}} \right)$ from Theorem 8 has the limit $\arctan(\text{sign}(d) \cot \alpha) = \text{sign}(d) \left(\frac{\pi}{2} - \alpha \right)$. Since the four AT functions in Theorem 8 include the same values of d, α , their limits cancel, so $\text{lk}(L_1, L_2) \rightarrow 0$. ■

Corollary 16 (lk for $a_i, b_i \rightarrow \infty$) If the invariants d, α of line segments $L_1, L_2 \subset \mathbb{R}^3$ remain fixed, but $a_i \rightarrow +\infty$ or $b_i \rightarrow -\infty$ for each $i = 1, 2$, then $\text{lk}(L_1, L_2) \rightarrow 0$. □

Proof If $a_i \rightarrow +\infty$, then $a_i \leq b_i \rightarrow +\infty, i = 1, 2$. If $b_i \rightarrow -\infty$, then $b_i \geq a_i \rightarrow -\infty, i = 1, 2$. Consider the former case $a_i \rightarrow +\infty$, the latter is similar. Since d, α are fixed, $a^2 + b^2 - 2ab \cos \alpha + d^2 \leq (a + b)^2 + d^2 \leq 5b^2$ for large enough b . Since $\arctan(x)$ increases, $\text{AT}(a, b; d, \alpha) \geq \arctan \left(\frac{ab \sin \alpha + d^2 \cot \alpha}{db \sqrt{5}} \right) \rightarrow \text{sign}(d) \frac{\pi}{2}$ as $b \geq a \rightarrow +\infty$. Since the four AT functions in Theorem 8 have the same limit when their first two arguments tend to $+\infty$, these 4 limits cancel, so $\text{lk}(L_1, L_2) \rightarrow 0$. ■

Corollary 17 (lk for $a_i \rightarrow b_i$) If one of segments $L_1, L_2 \subset \mathbb{R}^3$ becomes infinitely short so that its final endpoint tends to the fixed initial endpoint (or vice versa), while all other invariants of L_1, L_2 from Definition 4 remain fixed, then $\text{lk}(L_1, L_2) \rightarrow 0$. □

Proof We show that $\text{lk}(L_1, L_2) = 0$ for $d = 0$. It's enough to consider the case $d \neq 0$. Then $\text{AT}(a, b; d, \alpha) = \arctan \left(\frac{ab \sin \alpha + d^2 \cot \alpha}{d \sqrt{a^2 + b^2 - 2ab \cos \alpha + d^2}} \right)$ is continuous. Let (say

for $i = 1$) $a_1 \rightarrow b_1$, the case $b_1 \rightarrow a_1$ is similar. The continuity of AT implies that $AT(a_1, b_2; d, \alpha) \rightarrow AT(b_1, b_2; d, \alpha)$ and $AT(a_1, a_2; d, \alpha) \rightarrow AT(b_1, a_2; d, \alpha)$. In the limit all terms in Theorem 8 cancel, hence $lk(L_1, L_2) \rightarrow 0$. ■

7 Computations of the linking number for polygonal links

If curves $\gamma_1, \gamma_2 \subset \mathbb{R}^3$ consist of straight line segments, then $lk(\gamma_1, \gamma_2)$ can be computed as the sum of $lk(L_1, L_2)$ over all line segments $L_1 \subset \gamma_1$ and $L_2 \subset \gamma_2$. In [26] there is a complex proof that this sum is convergent for a cubical lattice. The convergence of the periodic linking numbers remains open for arbitrary lattices.




Fig. 4 1st: The Hopf link as two square cycles has $lk = -1$ and vertices with coordinates $L_1 = (-2, 0, -2), (2, 0, -2), (2, 0, 2), (-2, 0, 2)$, $L_2 = (-1, -2, 0), (-1, 2, 0), (1, 2, 0), (1, -2, 0)$
2nd: The Hopf link of triangular cycles $L_1 = (-1, 0, -1), (-1, 0, 1), (1, 0, 0)$ and $L_2 = (0, 0, 0), (2, 1, 0), (2, -1, 0)$ has $lk = +1$. **3rd:** Solomon's link of $L_1 = (-1, 1, 1), (-1, -1, 1), (3, -1, 1), (3, 1, -1), (1, 1, -1), (1, 1, 1)$ and $L_2 = (0, -2, -2), (0, -2, 0), (0, 2, 0), (2, 2, 0), (2, 0, 0), (2, 0, -2), (2, -2, -2), (0, -2, -2)$ has $lk = +2$. **4th:** Whitehead's link of the curves with the vertices $L_1 = (-1, 0, 1), (1, 0, 1), (0, 0, -2), (0, 1, -2), (-1, 1, -2), (-2, -1, 3), (1, -1, 3), (-1, 0, -3), (-1, 2, -3), (-1, 2, 1)$ and $L_2 = (0 - 2 - 1), (0, 2, -1), (0, 2, 2), (0, -2, 2)$ has $lk = 0$.

Figure 4 shows polygonal links whose linking numbers were computed by our Python code implementing formula (8) at https://github.com/MattB-242/Closed_Lk_Form. For all links in Fig 4 formula 8 calculates the linking number between the two components correctly (as equal to -1 and $+1$ respectively in the orientations given in Fig 4), with a computation error of less than 10^{-12} .

The asymptotic linking number introduced by Arnold converges for infinitely long curves [32], while our motivation was a computation of geometric and topology invariants to classify periodic structures such as textiles [11] and crystals [12].

Theorem 8 allows us to compute the *periodic* linking number between a segment J and a growing finite lattice L_n whose unit cell consists of n copies of two oppositely oriented segments orthogonal to J . This periodic linking number is computed for increasing n in a lattice extending periodically in one, two and three directions, see Figure 5. As n increases, the lk function asymptotically approaches an approximate value of 0.30 for 1- and 3-periodic lattice and 0.29 for the 2-periodic lattice.




Fig. 5 **Left:** the line segment $J = (0, 0, -1) + t(0, 0, 2)$ in red and the periodic lattice $L(n^k)$ derived from n copies of the 'unit cell' $L = \{(-1, -1, 0) + t(0, 2, 0), (-1, 1, 0) + s(0, -2, 0)\}$, $t, s \in [0, 1]$, translated in k linearly independent directions for $n \in \mathbb{Z}$. **Right:** the periodic linking number $lk(J, L(n^k))$ is converging fast for $n \rightarrow +\infty$. **Top:** $k = 1$. **Middle:** $k = 2$. **Bottom:** $k = 3$.

The invariant-based formula has allowed us to prove new asymptotic results of the linking number in Corollaries 12-17 of section 6. Since the periodic linking number is a real-valued invariant modulo isometries, it can be used to continuously quantify similarities between periodic crystalline networks [12]. One next possible step is to

use formula (8) to prove asymptotic convergence of the periodic linking number for arbitrary lattices, so that we can show that the limit of the infinite sum is a general invariant that can be used to develop descriptors of crystal structures.

The Milnor invariants generalise the linking number to invariants of links with more than two components. An integral for the three component Milnor invariant [13] may be possible to compute in a closed form similarly to Theorem 8. The interesting open problem is to extend the isometry-based approach to finer invariants of knots.

The Gauss integral in (1) was extended to the infinite Kontsevich integral containing all finite-type or Vassiliev’s invariants of knots [17]. The coefficients of this infinite series were explicitly described [19] as solutions of exponential equations with non-commutative variables x, y in a compressed form modulo commutators of commutators in x, y . The underlying metabelian formula for $\ln(e^x e^y)$ has found an easier proof [20] in the form of a generating series in the variables x, y .

This paper has provided a detailed proof of the analytic formula for the linking number based on six isometry invariants that uniquely determine a relative position of two line segments in \mathbb{R}^3 . Though a similar formula was claimed in [16], no proof was given. Hence this paper fills an important gap in the literature by completing the previously missing proof via highly technical Lemmas 9, 10, 11 in section 5.

We were motivated by detecting inter-penetrations of crystalline networks [12]. Solid crystalline materials (crystals) are periodic structures, which are determined in a rigid form and can be naturally classified up to isometry preserving all inter-atomic distances. The first complete isometry invariant of crystals was found in [3]. The harder problem is to design a continuous metric between crystals. Well-approximated metrics between lattices in any dimension were defined in the first paper [25] in a new area of Geometric Data Science whose periodic case was initiated in [2, 14], see the latest advances for dimension one [4, 22], two [23, 10], and three [18, 9, 31], for finite sets [30, 21], and materials applications [29, 34, 35]. This work was supported by the EPSRC grant “Application-driven Topological Data Analysis” (EP/R018472/1) and by the Royal Academy of Engineering Industrial Fellowship (IF2122/186).

References

1. Rasool Ahmad, Saptarshi Paul, and Sumit Basu. Characterization of entanglements in glassy polymeric ensembles using the gaussian linking number. *Phys. Rev. E*, 101(2):022503, 2020.
2. O Anosova and V Kurlin. Introduction to periodic geometry and topology. *arXiv:2103.02749*, 2021.
3. O Anosova and V Kurlin. An isometry classification of periodic point sets. In *Proceedings of Discrete Geometry and Mathematical Morphology*, 2021.
4. O Anosova and V Kurlin. Density functions of periodic sequences. 2022.
5. Zin Arai. A rigorous numerical algorithm for computing the linking number of links. *Nonlinear Theory and Its Applications*, 4(1):104–110, 2013.
6. T Banchoff. Self-linking numbers of space polygons. *Indiana U. Math. J.*, 25:1171–1188, 1976.
7. Enrico Bertolazzi, Riccardo Ghiloni, and Ruben Specogna. Efficient computation of linking number with certification. *arXiv:1912.13121*, 2019.

8. M Bright, O Anosova, and V Kurlin. A proof of the invariant-based formula for the linking number and its asymptotic behaviour. In *Numerical Geometry, Grid Generation and Scientific Computing*, pages 37–60. Springer, 2021.
9. M Bright, A Cooper, and V Kurlin. Welcome to a continuous world of 3-dimensional lattices. *arXiv:2109.11538*, 2021.
10. M Bright, A Cooper, and V Kurlin. Geographic-style maps for 2-dimensional lattices. *Acta Crystallographica Section A*, to appear.
11. M Bright and V Kurlin. Encoding and topological computation on textile structures. *Computers and Graphics*, 90:51–61, 2020.
12. P Cui, D McMahon, P Spackman, B Alston, M Little, G Day, and A Cooper. Mining predicted crystal structure landscapes with high throughput crystallisation: old molecules, new insights. *Chemical Science*, 10:9988–9997, 2019.
13. D. DeTurck, H. Gluck, R. Komendarczyk, P. Melvin, C. Shonkwiler, and D. Vela-Vick. Pontryagin invariants and integral formulas for milnor’s triple linking number. *arXiv:1101.3374*.
14. H Edelsbrunner, T Heiss, V Kurlin, P Smith, and M Wintraecken. The density fingerprint of a periodic point set. In *Proceedings of SoCG*, pages 32:1–32:16, 2021.
15. Carl Friedrich Gauss. Integral formula for linking number. *Zur mathematischen theorie der electrodynamische wirkungen, Collected Works*, 5:605, 1833.
16. Konstantin Klenin and Jörg Langowski. Computation of writhe in modeling of supercoiled dna. *Biopolymers: Original Research on Biomolecules*, 54(5):307–317, 2000.
17. M. Kontsevich. Vassiliev’s knot invariants. *Adv. Soviet Mathematics*, 16:137 – 150, 1993.
18. V Kurlin. A complete isometry classification of 3-dimensional lattices. *arxiv:2201.10543*.
19. V Kurlin. Compressed Drinfeld associators. *Journal of Algebra*, 292:184–242, 2005.
20. V Kurlin. The Baker-Campbell-Hausdorff formula in the free metabelian lie algebra. *Journal of Lie Theory*, 17(3):525–538, 2007.
21. V Kurlin. Computable complete invariants for finite clouds of unlabeled points. *arxiv:2207.08502*, 2022.
22. V Kurlin. Exactly computable and continuous metrics on isometry classes of finite and 1-periodic sequences. *arXiv:2205.04388*, 2022.
23. V Kurlin. Mathematics of 2-dimensional lattices. *Found. Comp. Mathematics*, to appear.
24. James Clerk Maxwell. A treatise on electricity and magnetism. I, 1873.
25. Marco Mosca and Vitaliy Kurlin. Voronoi-based similarity distances between arbitrary crystal lattices. *Crystal Research and Technology*, 55(5):1900197, 2020.
26. E Panagiotou. The linking number in systems with periodic boundary conditions. *J Computational Physics*, 300:533–573, 2015.
27. E Panagiotou and L Kauffman. Knot polynomials of open and closed curves. *Proc. Royal Society A*. 476: 20200124, 2020.
28. Renzo L Ricca and Bernardo Nipoti. Gauss’ linking number revisited. *Journal of Knot Theory and Its Ramifications*, 20(10):1325–1343, 2011.
29. J Ropers, M Mosca, O Anosova, V Kurlin, and A Cooper. Fast predictions of lattice energies by continuous isometry invariants of crystal structures. In *Proceedings of DAMDID (arXiv:2108.07233)*, 2022.
30. P Smith and V Kurlin. Families of point sets with identical 1d persistence. *arxiv:2202.00577*, 2022.
31. P Smith and V Kurlin. A practical algorithm for degree- k voronoi domains of three-dimensional periodic point sets. In *Proc. Int. Symposium on Visual Computing*, 2022.
32. T Vogel. On the asymptotic linking number. *Proc. Amer. Math. Soc.*, 131:2289–2297, 2003.
33. A.V. Vologodskii, V. V. Anshelevich, A.V. Lukashin, and M. D. Frank-Kamenetskii. Statistical mechanics of supercoils and the torsional stiffness of the dna double helix. *Nature*, 280(5720):294–298, 1974.
34. D Widdowson and V Kurlin. Resolving the data ambiguity for periodic crystals. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2022.
35. Q Zhu et al. Analogy powered by prediction and structural invariants. *J Amer. Chem. Society*, 144:9893–9901, 2022.