

Hopelessness and Burnout in Italian healthcare workers during COVID-19 pandemic: the mediating role of trait Emotional Intelligence

Maria S. Epifanio¹, Sabina La Grutta¹, Marco A. Piombo^{2*}, Martina Riolo¹, Vittoria Spicuzza¹, Marianna Franco¹, Giacomo Mancini³, Leonardo De Pascalis^{2, 4}, Elena Trombini², Federica Andrei²

¹Department of Psychology, Educational Science and Human Movement, University of Palermo, Italy,

²Department of Psychology, University of Bologna, Italy, ³Giovanni Maria Bertin Department of Education Studies, University of Bologna, Italy, ⁴Psychological Sciences, Institute of Health and Life Sciences, University of Liverpool, Liverpool, UK, United Kingdom

Submitted to Journal:
Frontiers in Psychology

Specialty Section:
Health Psychology

Article type:
Original Research Article

Manuscript ID:
1146408

Received on:
17 Jan 2023

Revised on:
30 Mar 2023

Journal website link:
www.frontiersin.org

Conflict of interest statement

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest

Author contribution statement

MSE, SLG, ET, and FA; conceptualization. MSE, SLG, FA and LDP; methodology. MR, VS, MF and GM; data collection and data curation. MAP; data analysis. MSE, MAP, VS, and MR writing the original draft manuscript. MSE, MAP, and FA; review and editing. MSE, SLG, FA, ET; supervision.

Keywords

COVID-19, pandemic, Healthcare workers, burnout, Hopelessness, trait emotional intelligence, TEIQue-SF

Abstract

Word count: 308

Objective: The study aims to assess the impact of COVID-19 on healthcare workers' (HCW) work-related stress during the first wave of the pandemic in Italy. The main objective is to investigate the existence of a positive correlation between hopelessness and burnout, assuming that burnout may be a risk factor for the development of hopelessness, and to analyse the role that TEI and changes in workload could have in this relationship. Furthermore, evaluate any significant differences in burnout and hopelessness levels in the function of some demographic variables such as gender, professional profiles, and different working zones of Italy, to better understand how the diverse diffusion of pandemic had affected Italian HCW.

Methods: An online survey was between April and June, 2020, with 562 responses among nurses (52.1%) and physicians (47.9%). Demographics and changes in workload and work conditions were collected through an ad-hoc questionnaire. The Trait Emotional Intelligence Questionnaire-Short Form (TEIQue-SF;), The Beck Hopelessness Scale (BHS), and the Link Burnout Questionnaire (LBQ) were used to assess Trait Emotional Intelligence (TEI), hopelessness and burnout respectively. Results: correlation analysis showed a significant positive relationship between hopelessness and each burnout dimension. TEI showed negative correlations both with burnout dimensions and hopelessness. Significant differences in burnout and hopelessness levels emerged as a function of some demographic variables such as gender, professional profiles (nurses or physicians), and different working zone of Italy (northern or southern). Results showed that TEI partially mediated the relationship between hopelessness and every burnout dimension, while the interaction of changes in workload was non-significant. **Discussion:** The mediating role of TEI in the burnout-hopelessness relationship partly explains the protective role that individual factors had on HWC mental health. Our findings support the need to integrate considerations on both psychological risk and protective factors into COVID-19 care, including the monitoring of psychological symptoms and social needs especially among healthcare workers.

Contribution to the field

Although much attention has been paid to the effects of the COVID-19 pandemic on burnout and mental health in healthcare workers, little has been carried out so far to understand the underlying specific mediation and moderation mechanisms. To our knowledge, this is the first study to examine the role of trait EI in burnout syndrome in healthcare workers during the COVID-19 pandemic. This study identifies a pathway (i.e., via trait EI) that may help better understand the mechanism by which the extreme consequence of burnout syndrome, such as suicide risk, could be limited. Our findings support the need to integrate considerations on both psychological risk and protective factors into COVID-19 care, including the monitoring of psychological symptoms and social needs, especially among healthcare workers.

Ethics statements

Studies involving animal subjects

Generated Statement: No animal studies are presented in this manuscript.

Studies involving human subjects

Generated Statement: The studies involving human participants were reviewed and approved by Bioethics Committee of the University of Palermo (n. 7/2020). The patients/participants provided their written informed consent to participate in this study.

Inclusion of identifiable human data

Generated Statement: No potentially identifiable human images or data is presented in this study.

In review

Data availability statement

Generated Statement: The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

In review

Hopelessness and Burnout in Italian healthcare workers during COVID-19 pandemic: the mediating role of trait emotional intelligence

Maria Stella Epifanio ^{†1}, Sabina La Grutta*^{†1}, Marco Andrea Piombo* ², Martina Riolo¹, Vittoria Spicuzza¹, Marianna Franco, Giacomo Mancini ³, Leonardo De Pascalis^{2,4}, Elena Trombini^{‡2}, Federica Andrei^{‡2}

¹Department of Psychology, Educational Science and Human Movement, University of Palermo, 90128 Palermo, Italy.

²Department of Psychology “Renzo Canestrari”, Alma Mater Studiorum- University of Bologna, 40127 Bologna, Italy.

³Department of Education Studies “Giovanna Maria Bertin”, Alma Mater Studiorum, University of Bologna, Bologna, Italy.

⁴Psychological Sciences, Institute of Health and Life Sciences, University of Liverpool, Liverpool, UK

[†]These authors share first authorship

[‡] These authors share last authorship

* Correspondence:

Corresponding Author

marcoandrea.piombo2@unibo.it

Keywords: COVID-19, pandemic, healthcare workers, burnout, hopelessness, trait emotional intelligence, TEIQue-SF.

Objective: The study aims to assess the impact of COVID-19 on healthcare workers’ work-related stress during the first wave of the pandemic in Italy. The main objective is to investigate the existence of a positive correlation between hopelessness and burnout, assuming that burnout may be a risk factor for the development of hopelessness, and to analyze the role that trait Emotional Intelligence (TEI) and changes in workload could have in this relationship. Furthermore, evaluate any significant differences in burnout and hopelessness levels in the function of some demographic variables, such as gender, professional profiles, and different working zones of Italy, to better understand how the diverse diffusion of pandemic had affected Italian healthcare workers. **Methods:** An online survey was used to collect data between April and June, 2020, with 562 responses among nurses (52.1%) and physicians (47.9%). Demographics and changes in workload and work conditions were collected through an ad-

hoc questionnaire. The Trait Emotional Intelligence Questionnaire-Short Form (TEIQue-SF;), The Beck Hopelessness Scale (BHS), and the Link Burnout Questionnaire (LBQ) were used to assess Trait Emotional Intelligence (TEI), hopelessness, and burnout, respectively. **Results:** correlation analysis showed a significant positive relationship between hopelessness and each burnout dimension. TEI showed negative correlations both with burnout dimensions and hopelessness. Significant differences in burnout and hopelessness levels emerged as a function of some demographic variables such as gender, professional profiles (nurses or physicians), and different working zone of Italy (northern or southern). Results showed that TEI partially mediated the relationship between hopelessness and every burnout dimension, while the interaction of changes in workload was non- significant. **Discuion:** The mediating role of TEI in the burnout-hopelessness relationship partly explains the protective role that individual factors had on **healthcare workers'** mental health. Our findings support the need to integrate considerations on both psychological risk and protective factors into COVID-19 care, including the monitoring of psychological symptoms and social needs, especially among healthcare workers.

1. INTRODUCTION

Background

The Covid-19 pandemic has been deeply affecting both the general population and the health systems of different countries. Several international studies have shown an increase in mental suffering (Rossi et al., 2020; Zhang et al., 2020; Breslau et al., 2021; Lorant et al., 2021; Daly M., Robinson E., 2022) and the worsening of quality of life (QoL; Epifanio et al., 2020; Van Ballegooijen et al., 2020; White & Van Der Boor, 2020; Horesh et al. 2020; Hansel et al., 2022) especially on healthcare professionals (Bellizzi et al., 2020; Guanche 2020; Semple & Cherrie 2020; Cortés-Álvarez & Vuelvas-Olmos, 2022).

During the first pandemic wave, Italy was among the most affected nations in terms of hospital overload, and its healthcare workforce struggled to face all the challenges related to the pandemic, especially in the northern regions of the country. According to the integrated surveillance report on COVID-19 in Italy¹ in November 2020, 446.875 health workers were infected², and there were 378 deaths among doctors.

¹ <https://portale.fnomceo.it/> ; ² <https://www.epicentro.iss.it/>

In this context, it has become necessary to understand the COVID-19 epidemic health consequences on Italian health professionals at the forefront (Guanche, 2020). Several studies on the impact of COVID-19 on healthcare workers' mental health (Moitra et al., 2021; Saragih et al., 2021; Sun et al., 2021; Dragioti et al., 2022) found a high prevalence of post-traumatic stress disorder, depressive and anxious symptoms, sleep disorders, psychological distress, and burnout syndrome. Stress is caused not only by the continuous contact with the patient's suffering but also by other factors related to the workplace, such as its organization, dynamics, and quality of relations with colleagues and health management (Romano, 2005; Schaufeli, Leiter & Maslach, 2009).

Recurrence of distress situations could lead to a high burnout risk, a particular type of acute and extreme physical and psychological suffering (Maslach, 1982; Maslach & Jackson, 1981; Maslach & Schaufeli, 1993;) resulting from the perception of an imbalance between the number of requests and the resources available, in order to positively respond to such requests. This can lead to energies depletion, with subsequent physical symptoms (frequent headaches, insomnia, drug use), psychological symptoms (guilt, negativism, irritability), behavioural reactions, and absence and delay at work (tendency to avoid telephone contact, changes in attitude, emotional detachment and indifference to patient's problems (Barello et al., 2020; Kang et al., 2020; Lai et al., 2020). Early research on burnout (Maslach, 1993) suggests that it can lead to a deterioration of care or service quality provided by medical professionals. In fact, operator feelings can include tension, irritability, and emotional excitability that could lead to detachment from their users in order to preserve their energies and resources.

In addition, burnout can be characterized by a motivation decline that leads to a loss of idealism, enthusiasm, positivity, and increases in apathy and cynicism toward users (Maslach, 2003). It is clear from this that personal satisfaction and gratification, together with empathy toward their users, are essential to provide effective care. Therefore, healthcare professionals' burnout has negative consequences for staff, patients, and organizations, particularly in the case of frontline workers (Friedberg et al., 2014). For instance, burnout compromises the quality of care due to turnover (Leiter, Harvie & Frizzel, 1998), absenteeism (Davey et al., 2009), and early withdrawal (Linzer et al., 2001). The clinical and economic impact of burnout is such that in 2019 it was included in the 11th revision of the International Classification of Disease (ICD-11, 2019) as an employment phenomenon. According to the definition of ICD-11, burnout is a syndrome resulting from chronic stress in the workplace, not properly managed. It is characterized by a feeling of depletion of energy or exhaustion,

an increase in mental distance and negative or cynical feelings toward work and others, and reduced professional effectiveness.

In their efforts to assist and treat patients with COVID-19, physicians and nurses in particular (Liu et al., 2020), have been exposed to daily events that may lead to the development, or exacerbation, of psychological distress. Moreover, the very long work shifts, the constant fear of contagion, and the frustration caused by the difficulty in providing effective care are all risk factors that may have contributed to the onset of considerable work distress (Lai et al., 2020, Baptista et al., 2021).

Several studies (Morgantini et al., 2020; Franza et al., 2020; Rossi et al., 2020; Martinez-Lopez et al., 2020; Ghahramani et al., 2021; Kimpe et al., 2022; Gualano et al., 2021; Claponea et al., 2022) have shown a high prevalence of burnout syndrome in healthcare workers during COVID-19 pandemic. Zhang and colleagues (2020) found a 56.03% prevalence of anxiety symptoms and burnout in the Intensive Care Unit (ICU) medical staff. Other studies (Baptista et al., 2020; Tuna & Ozdin, 2021; Dinibitun, 2020; Alraxashdeh et al., 2020) show similar results both in family doctors (Baptista et al., 2020) and in frontline physicians (Tuna & Ozdin, 2020).

Regarding nurses, international studies report around an 11% prevalence of burnout worldwide (Woo et al., 2020; Galanis et al., 2021; Zareei et al., 2022). In other words, one-tenth of the world's nurses show symptoms of burnout syndrome. Moreover, among gender-related burnout, data in the literature (Vitale, Galatola & Mea, 2020; Innstrand, Melbye Langball & Falkum, 2011) suggest that females tend to experience higher levels of psychophysical exhaustion probably due to women's double workload and inequality between genders at work.

There are many risk factors for burnout at both individual and institutional levels (Bria, Baban & Dumitrascu, 2012). In a systematic review, Bria et al. (2012) identified some risk factors at the institutional level, such as perceived job control, high workload, organizational justice, social support at work, effort-reward imbalance, perceived burnout complaints among colleagues, and hospital organizational characteristics. Moreover, a recent review by Khosravi, Ghiasi & Ganjali (2022) on healthcare staff's burnout during the COVID-19 pandemic highlights the key role of environmental factors, such as role ambiguity, role conflict, and work overload, in increasing the severity of burnout (Khosravi, Ghiasi & Ganjali, 2022) and report several contributions from the literature on possible strategies for treating and preventing burnout among health care workers, such as interventions focused on increasing the awareness of occupational burnout (Maldonato et al., 2020) or interventions based on the promotion of supportive work culture in order to elevate the health care workers' resilience (Dewey et al., 2020).

Concerning individual risk factors, studies (Demir et al., 2003; Sharma et al., 2007; Sharma et al., 2008) show that healthcare professionals who experienced burnout use more emotion-focused coping such as substance abuse and unhealthy eating habits. In healthcare workers, hope is one of the main coping strategies that influence the ability to interact with stress in life-threatening situations (Franza et al., 2020). Hopelessness, in fact, leads to the absence of determination and catastrophic and destructive thoughts, and it is well documented that physicians take their own lives at rates much higher than the general public, and on average, 400 US physicians die by suicide each year (Bradley et al., 2020). In Italy (Loretto et al., 2020) it seems that there are more than 25 suicide every year among physicians, while data about nurses are not available.

One of the most important risk factor for suicide is certainly hopelessness, a psychological construct (Beck & Weissman 1974) that has been identified as one of the characteristics of depression and has been implicated in a variety of other conditions such as schizophrenia and alcoholism. The few studies that investigated the relationship between hopelessness and burnout (Pompili et al., 2013; Franza et al., 2020; Karagol & Kaya, 2022) found a positive correlation between the two constructs, assuming that burnout may be a risk factor for the development of hopelessness, which could lead to the onset of depression and suicide risk.

In contrast to these risk factors, there are personal dispositions that could be important in preventing burnout and hopelessness, such as Emotional Intelligence (EI). EI is generally defined as a psychological attribute that captures individual differences in how we perceive, communicate, regulate, and understand our own emotions, as well as the emotions of others (Hughes & Evans, 2018). The ability to monitor one's own and others' feelings and emotions and to guide one's thinking and behaviour seems to be a useful competency in dealing with stress work-related (Humpel & Caputi, 2001). In line with other studies (Humpel & Caputi, 2001; Moraes et al., 2015; Swami et al., 2013), Adamson (2014) found that nurses with high EI were less likely to experience burnout than those with low scores. Moreover, other studies found a negative relationship between EI and burnout (Fiorilli et al., 2019; Tesi, 2021; Lopez & Extremera, 2017).

Among the different possible EI formulations, some theories see EI as a factor that promotes general health (Schutte et al., 2007; Zysberg, 2018) and better adaptation (Pellitteri et al., 2022) influencing the way individuals cope with demands and pressures from the environment to be more resilient to challenging situations (Zeidner, Matthews & Roberts, 2006) and uses specific tests to measure emotional abilities. Trait emotional intelligence (TEI), instead, focuses on people's perception of their emotional disposition and is identified as a distinct latent variable that integrates the affective

aspects of personality (Andrei et al., 2014; Andrei et al., 2016). Particularly, the construct of TEI has emerged as an important individual difference variable referring to a constellation of emotional self-perceptions assessed by self-reported questionnaires (Petrides, Pita & Kokkinaki, 2007). Several studies have explored the influence of TEI across the lifespan (Petrides et al., 2016) and its impact on health, underlining that high TEI strongly positively predicts well-being, mental health (e.g., Martins, Ramalho & Morin, 2010), and moderates the responses to stress (Mikolajczak & Luminet, 2008). In this sense, this type of framework and measurement are best suited to the purposes of our research, which assesses how personal disposition (e.g. the tendency towards hopelessness) and situational (the ongoing pandemic and consequent changes in the burden of work) affect work-related stress. To our knowledge, while recent studies examined the role of trait EI in the relationship between several stressful experiences, such as job loss during the COVID-19 Pandemic (Andrei et al., 2022), few studies explore the relationship between trait EI, burnout, and hopelessness in a high-risk sample such as healthcare workers, especially those who were in the frontline during COVID-19 Pandemic.

The present study

The study aims to assess the impact of COVID-19 on work-related stress and risks for healthcare workers' mental health during the first wave of the pandemic in Italy. The main objective was to investigate the existence of a positive correlation between hopelessness and burnout, assuming that burnout may be a risk factor for the development of hopelessness, and to analyze the role that trait EI and changes in workload due to the pandemic situation could have in this relationship. Another aim of this study was to evaluate the differences in burnout and hopelessness levels taking into account gender, professional profiles (nurses or physicians), and different working zone of Italy (northern or southern).

We hypothesized that: (1) healthcare workers who work in a COVID-19 ward and northern regions of Italy have higher levels of burnout and hopelessness than others due to the different diffusion that the pandemic had in the first wave; (2) nurses have higher levels of burnout and hopelessness than physicians due to their first-responder role; (3) females have a higher level of psychophysical exhaustion burnout dimension, as confirmed by previous literature, due to women's double workload and inequality between men and women in work contexts; (4) the existence of a positive correlation between hopelessness and burnout, assuming that burnout may be a risk factor for the development of hopelessness; (5) trait EI (TEI) will mediate the relationship between burnout and hopelessness, while changes in workload due to the pandemic situation could be a moderator factor of this relationship; (6) higher workload could cause more burnout, consequentially increasing the feeling of hopelessness.

2. MATERIALS AND METHODS

Procedures

An online cross-sectional survey was performed with Qualtrics® (Qualtrics, Provo, UT, USA) Survey Platform. This data collection strategy was chosen as it allowed us to reach as many voluntary participants as possible in a phase of forced social distancing. The survey started after 7 weeks of quarantine in Italy (25 April 2020) and was performed for about 6 weeks, until the end of lockdown measures (2 June 2020). The sample was recruited via a snowball sampling strategy. A link to Qualtrics questionnaires were sent via e-mail, social networks (Facebook and WhatsApp), and official working platforms (website of the University of Palermo, Italy).

A brief presentation informed the participants about the aims of the study, and electronic informed consent, assuring maximum confidentiality in the handling and analysis of the responses, was requested from each participant before starting the investigation. The survey took approximately 30 min to complete. Participation was voluntary and free of charge. To guarantee anonymity, no personal data, which could allow the identification of participants, were collected.

Participants could withdraw from the study at any time without providing any justification, and the data were not saved. Only the questionnaire data with a complete set of answers by respondents were considered. The study was conducted in accordance with the Declaration of Helsinki and was approved by the Bioethics Committee of the University of Palermo (n. 7/2020).

Participants

Italian healthcare workers who were working in public hospital wards COVID-19 and NO-COVID-19 and outpatient clinics at the time of the first wave pandemic (April-July 2020) were eligible for participation to data collection. The recruited sample size through the online survey included 774 Italian adults, with an attrition rate of approximately 27%. Our final sample comprised 562 respondents, physicians ($n = 269$) and nurses ($n = 293$) Demographic characteristics of the study sample are presented in Table 1.

Measures

Demographics and changes in workload information Questionnaire

An ad hoc questionnaire was created to collect demographic data (such as gender, age, region of residence in Italy, job position) and work information (i.e., changes in employment status, workload, years of professional experience, public hospital, COVID-19 and NO-COVID-19 wards and public out-patient clinic).

The Trait Emotional Intelligence Questionnaire–Short Form (TEIQue-SF)

The TEIQue-SF (Petrides, 2009; Di Fabio & Palazzeschi, 2011) was used to assess trait EI. Consist of 30 items rated on 7-point Likert scale, provides a global trait EI score and scores at four factors (i.e., wellbeing, self-control, emotionality and sociability), which higher scores corresponds to high levels of trait EI. Reliability for the present study was good (Cronbach's $\alpha = 0.90$).

The Beck Hopelessness Scale (BHS)

The BHS (Beck, Weissman, Lester & Trexler, 1974; Pompili et al. 2009) was used to measure feeling of hopelessness, BHS includes 20 true/false items about the future (9 positives and 11 negatives), that can be summed up to give a total score ranging from 0 to 20. Higher scores reflect increased hopelessness. In this study, this questionnaire showed good reliability (Cronbach's $\alpha = 0.89$).

The Link Burnout Questionnaire (LBQ)

The LBQ (Santinello, Verzelletti & Altoè, 2006) was used to measure professional burnout, LBQ includes 24 items rated on a 6-point Likert scale. A total score of LBQ allows measuring 4 dimensions of professional burnout: psychophysical exhaustion (dimension referring to the self-assessment of one's own psychophysical resources); deterioration of relations with clients (dimension referring to the quality of those relations); job ineffectiveness (dimension referring to the self-assessment of one's own professional competences); disappointment (dimension of existential expectations). For each dimension cut-off scores change differently for physicians and nurses: psychophysical exhaustion (low: 6-13 physicians, 6-12 nurses; average: 14-20 physicians, 13-19 nurses; moderate: 21-26 physicians, 20-25 nurses; high: 27-36 physicians, 26-36 nurses); deterioration of relations with clients (low: 6-9 physicians, 6-8 nurses; average: 10-13 physicians, 9-13 nurses; moderate: 14-21 physicians,

14-20 nurses; high: 22-36 physicians, 21-36 nurses); job ineffectiveness (low: 6-10 physicians, 6-10 nurses; average: 11-17 physicians, 11-17 nurses; moderate: 18-23 physicians, 18-24 nurses; high: 24-36 physicians, 25-36 nurses); disappointment (low: 6-9 physicians, 6-9 nurses; average: 10-16 physicians, 10-16 nurses; moderate: 17-24 physicians, 17-25 nurses; high: 25-36 physicians, 26-36 nurses). Reliability for the present study was good (Cronbach's $\alpha = 0.91$).

Data analysis

Statistical analyses were performed using programs available in the Statistical Package for Social Sciences (SPSS for Windows release 25.0). Descriptive statistics were utilized to describe the data (frequencies, percentages, mean and standard deviation). Pearson's correlations were used to investigate associations among variables. Change in workload was coded as a dummy variable: -1= less workload; 0= same workload; 1= more workload than pre-pandemic period. Moreover, also gender (0= females; 1=males), region (0=south and 1=north) and COVID-19ward (0=No; 1=Yes) were coded as dummy variables. Multivariate measures of ANOVA were used to investigate differences between groups. Finally, to explain more in-depth the relationship between burnout and hopelessness and to analyze the effect that changes in workload and TEI have on this relationship, some mediation- and moderation models were performed with the computational tool for SPSS, PROCESS (Model 5). In particular, every burnout dimension was included in the model as a predictor, while hopelessness was included as a dependent variable. Moreover, changes in workload were included as a moderator factor of this relationship, while TEI was included as a mediator (Figure 1).

3. RESULTS

Descriptive statistics

The final sample comprised a total of 562 healthcare workers (72.2% females), 52.15% nurses, and 47.9% physicians, collected mainly from the northern (78.3%) and southern (21.7%) regions of Italy and from hospitals (75.1%) and outpatient clinics (24.9%). Respondents were mostly middle (age 35 - 54; 51.6%) and older adults (age ≥ 55 ; 28.3%), while the group of young adults was smaller (age

≤ 34 ; 20.1%). Among years of professional experience, most of them had more than 20 years of experience (46.3%), while ≤ 10 and 11-20 range were respectively 28.8% and 24.9%.

Most participants worked in COVID-19 wards at the time of the survey (66.5%), while only 33.1% were in the other wards. Concerning the change in workload, 52% of participants experienced a greater workload, while 33.1% were the same and only 15.7% had a lower workload. Finally, regarding COVID-19 contagion, only 7.3% of participants had personally contracted COVID-19 while 17.8% had relatives who contracted the virus and, among them, 20% died.

Gender, regions, and working wards comparisons

Table 1 presents results from multivariate ANOVAs. Results showed significant gender differences in two burnout dimensions, with females scoring higher in psychophysical exhaustion ($F = 10.51$; $p < .001$) and job ineffectiveness ($F = 7.23$; $p < .01$) than males so as in workload changes ($F = 5.13$; $p < .05$). Contrary, males score higher than females in TEI ($F = 4.55$; $p < .05$). As regard regions of Italy, healthcare workers from North reported significantly higher workload changes ($F = 19.87$; $p < .05$) and in two burnout dimensions: psychophysical exhaustion ($F = 7.89$; $p < .01$) and disappointment ($F = 4.37$; $p < .05$). Finally, regarding professional profiles, nurses reported significantly higher workload changes than physicians ($F = 15.97$; $p < .001$;) so as those who work in COVID wards ($F = 25.71$; $p < .001$). Finally, no differences between groups were found in Hopelessness (all $ps > .05$).

Correlational Analyses

Table 2 presents Pearson's correlation coefficients of study variables. Specifically, hopelessness showed positive correlations with every burnout dimension (psychophysical exhaustion $r = .57$, $p < .001$; deterioration of relations with clients $r = .35$, $p < .01$; job ineffectiveness $r = .46$, $p < .01$; Disappointment $r = .63$, $p < .01$) and negative correlations with TEI ($r = -.59$, $p < .01$); while TEI showed negative correlation with every burnout dimensions (psychophysical exhaustion $r = -.55$, $p < .001$; deterioration of relations with clients $r = -.42$, $p < .01$; job ineffectiveness $r = -.55$, $p < .01$; disappointment $r = -.56$, $p < .01$). Finally, Change in Workload showed a significant relationship only with two burnout dimensions: psychophysical exhaustion ($r = .17$ $p < .01$) and deterioration of relation with Clients ($r = .13$, $p < .01$).

Differences in Burnout levels and hopelessness

Table 3 presents data on burnout dimension levels. Most participants obtained an average score in the dimension of psychophysical exhaustion (41.8%), while 20.6% obtained high, 29% moderate and 8.5% low scores, respectively. Regarding the dimension deterioration of the relationship with customers, 48.6% of the participants obtained a moderate score, while 20.6% high, 29.7% average and only the 1.1% low scores. Levels of the dimension labelled as work ineffectiveness, scores were high for 5% of participants, moderate for 16.9%, medium for 66.5%, and low for 11.6%. Disappointment subscale scores were high for 13.2% of participants, moderate for 26.9%, medium for 50.7%, and low for 9.6%. Furthermore, women showed higher levels than men in two dimensions of burnout: psychophysical exhaustion ($\chi^2 = 17.59$; $p < .01$) and work ineffectiveness ($\chi^2 = 29.04$; $p < .01$). Nurses, on the other hand, showed higher levels than physicians in the three dimensions of burnout (psychophysical exhaustion: $\chi^2 = 8.340$; $p < .05$; deterioration of the relationship with clients: $\chi^2 = 10.87$; $p < .05$; work Ineffectiveness: $\chi^2 = 9.67$; $p < .05$). Regarding region differences, participants from North Italy showed more high or moderate levels than participants from South in three burnout dimensions (psychophysical exhaustion ($\chi^2 = 8.340$; $p < .05$; deterioration of the relationship with clients: $\chi^2 = 10.87$; $p < .05$; disappointment: $\chi^2 = 13.50$; $p < .05$). Finally, regarding comparisons working with COVID-19 patients or not, participants who worked in COVID-19 wards showed more high or moderate scores only in job ineffectiveness ($\chi^2 = 8.00$; $p < .05$). Contrary, participants who worked in COVID-19 wards showed more low hopelessness levels than who worked in other wards ($\chi^2 = 11.07$; $p < .05$; see Table 3).

Mediation by trait EI and moderation by changes in workload

Regarding the mediation-moderation hypothesis, results showed that the trait EI partially mediated the relationship between every burnout dimension, which were included in the model as predictors, and hopelessness, which instead was the outcome variable (psychophysical exhaustion-BHS: $B = -2.80$, $p < .01$, 95% CI= [-3.31, -2.29]; deterioration of relations with clients – BHS: $B = -3.77$, $p < .01$, 95% CI= [-3.31, -2.29]; job ineffectiveness – BHS: $B = 3.41$, $p < .01$, 95% CI= [-3.96, -2.87]; disappointment – BHS: $B = 2.42$, $p < .01$, 95% CI= [-2.49, -1.92]). However, the interaction between each predictor and changes in workload was non-significant (all $ps > .05$). Finally, all these full models accounted from 37 to 49% of the variance in hopelessness ($R^2 = .44$, $p < .001$; $R^2 = .37$, $p < .001$; $R^2 = .38$, $p < .001$; $R^2 = .49$, $p < .001$; see Table 4).

4. DISCUSSION

The study aimed to assess the impact of the COVID-19 pandemic on the work-related stress of Italian healthcare workers involved in facing the first COVID-19 pandemic wave that, in Europe, mainly affected Italy. Our results show that northern workers have higher levels than southern workers in two burnout dimensions: psychophysical exhaustion and disappointment. This could be attributed to the increased spread of the virus in northern Italy during the first wave of the COVID-19 pandemic, with consequent greater pressure on hospitals. Furthermore, healthcare workers from the North reported significantly higher workload changes. Consistent with our findings, Swami and colleagues (2013) found a strong correlation between the physical fatigue dimension and changes in workload, also due to, as Karagall and colleagues (2022) found healthcare workers declared not having enough time for themselves or their own family obtained higher average scores in all dimensions of burnout and hopelessness. The authors suggest that this is due to working in departments with an extensive workload and highlight the importance to “have time for oneself or one's own family” for mental health, as confirmed by previous studies (Tiete et al., 2021; Marcia-Rodriguez et al., 2021; Hacimusalar, Kahve & Yasar, 2020).

Regarding those who worked in COVID-19 wards, we found significant differences from those who worked in other wards only in workload changes and in the job effectiveness burnout dimension. Moreover, in contrast with our hypothesis, we found lower levels of hopelessness than those who worked in usual wards. In this regard, the literature reports some interesting data: a study conducted by Wu and colleagues (2020) showed that, compared with medical staff who worked on their usual wards for uninfected patients, medical staff worked on the COVID-19 front-line ward had a lower frequency of burnout, in line with another study conducted by Dinibitum and colleagues (2020). According to Wu and colleagues (2020), a possible explanation could be that those at the front-line wards have experienced a greater closeness to the key decision-makers and were allowed to access to more timely and accurate information. On the contrary, results of other studies showing worse mental health outcomes among frontline healthcare workers during COVID-19 (Lai et al., 2020; Lu et al., 2020; Wu Y. et al., 2020).

However, the job ineffectiveness dimension was high, and this result could be explained by the fact that physicians who were in the first line in the emergency had access to their resources to face the emergency, experiencing acute stress, but facing an unknown virus and a new disease, without standardized, has probably destabilized their sense of professional self-efficacy. These findings suggest that healthcare workers' mental health outcomes seem to be not directly associated with the fact of

working on COVID-19 front-line wards and that further studies are needed to investigate possible other variables involved.

With respect to the differences between professional profiles, nurses showed higher workload changes than physicians and, in line with our hypothesis, they scored higher in three further dimensions assessed by the LBQ: psychophysical exhaustion, deterioration with clients, and work effectiveness, as confirmed by previous studies (Lai et al., 2020; Tiete et al., 2021; Bisesti et al., 2021). These findings could be possible because nurses had more frequent contacts with patients affected by COVID-19 and were thus exposed to higher risk (Bisesti et al., 2021). Moreover, these findings also align with those reported in a previous study on the SARS epidemic (Wong et al., 2005).

Regarding gender differences, our results show that females scored higher than males only in two burnout dimensions: psychophysical exhaustion and job ineffectiveness, partially in line with previous studies in which the female gender was associated with higher levels of emotional exhaustion than the male gender (Richards et al., 2021; Vitale, 2022). However, it is important to note that the gender-burnout relationship is still unclear. For example, in their qualitative review of the burnout literature, Maslach et al. (2001) observed a tendency for women to score higher on emotional exhaustion than men, whereas men tend to score higher on depersonalization than women, as confirmed by a 2010 meta-analysis (Purvanova and Muros, 2010) that highlight the fact that burnout is experienced differently by men and women, in line with gender role theory (Eagly, 1987; Eagly & Wood, 1982) which predicts that women tend to express feelings of emotional and physical fatigue (e.g., emotional exhaustion) due to the fact that they learn to display their emotions, whereas men tend to “shut off and withdraw under stress” (i.e., depersonalization) because they learn to conceal their emotions. In addition, as a possible gender roles outcome, Karagall, and colleagues (2022) indicate that women's increasing domestic workload and their less opportunity to distinguish between home and work requests, compared to men, may have led women to experience greater burnout (Karagall et al., 2022). On the other hand, males score higher than females in TEI, in line with Petrides and Furnham (2000) and Bar-On, Brown, Kirkcaldy, & Thomé (2000).

Therefore, it's interesting to note that the findings in the literature are controversial. In this regard, a review by Shazad and Bagum (2012) on gender differences in TEI reported a difference between mean scores of males and females concerning EI, with males having higher levels of TEI than females. According to the authors, a possible explanation could be that men perceive themselves as more emotionally intelligent, which could affect the results due to the self-report nature of the TEIQue. On the other hand, a gender role issue related to the social context, which defines gender roles

differently, could explain the possible effects on female's emotional development (Duckelt and Raffalli, 1989; Sandhu & Mehrotra, 1999), especially if we consider the sociability subdomain in which male score slightly higher than females (Bar-On, Brown, Kirkcaldy, & Thomé; 2000; Petrides & Furhman, 2000).

Regarding correlation analysis, we found that hopelessness and each burnout dimension are positively related, as confirmed by previous literature (Pompili et al., 2006; Franza et al., 2020; Akova et al., 2022; Karagoll et al., 2022) and negatively related with trait EI. Despite many studies on burnout in healthcare workers, few studies have investigated the relationship between burnout and hopelessness during the COVID-19 pandemic (Franza et al., 2020; Karagoll., 2022; Akova et al., 2022). However, assuming that burnout may be a risk factor for the development of hopelessness and that both conditions could represent an increased risk of committing suicide, we believe that this relationship requires greater attention in order to implement psychological interventions in care setting (i.e. hospital) in the future, not only for patients and their families but also to provide support to health care workers and prevent the extreme consequences of burnout.

The exhaustion stage is the most severe stage of burnout and is characterized by symptoms such as chronic sadness or depression, chronic physical disorders (i.e. chronic stomach or bowel and chronic headaches), chronic mental and physical fatigue, the desire to withdraw from society, family, and friends, and also recurrent suicidal ideation (Pompili et al., 2006). In healthcare workers, hope is among the main coping strategies, able to influence people's ability to deal with stress related to life-threatening situations (Jones-Schenk, 2020), however “emotional exhaustion” in physicians and nurses due to providing care for a large number of patients, excessive workload and not receiving adequate financial gratifications, could promote hopelessness feelings (Karagoll et al., 2022).

Moreover, our results showed that trait EI could act as a protective factor against burnout levels and hopelessness, as confirmed by previous research that highlights the positive role of EI over burnout levels (Vlachou et al., 2016; Nastasa & Farcas, 2015; Humpel & Caputi, 2001). According to the literature, EI is associated with lower rates of burnout and higher rates of job satisfaction, self-compassion, and communication skills (De Moraes et al., 2015). Results also showed that trait EI partially mediated the relationship between every burnout dimension and hopelessness. However, full mediation was not obtained as further risk factors, including alienation and/or resources, such as self-efficacy and social support, could play a significant role in the buffering process and could potentially contribute to mediating the association between burnout and hopelessness.

The role of EI as a protective factor against burnout in a nurse's sample during the COVID-19 pandemic has already been observed by Soto-Rubio et al. (2020). The authors highlight the importance

of nurses' abilities to perceive and regulate their emotions while being aware of and empathizing with others' emotions (Soto-Rubio et al., 2020). Several studies conducted during the COVID-19 pandemic took into account EI as a protective factor against psychosocial risks such as burnout and psychosomatic complaints and a favourable effect on job satisfaction (Shariatpanahi et al., 2022; Lisigurski et al., 2012; Soto-Rubio et al., 2020; Zeb et al.; 2020; Richards et al., 2021), but to our knowledge, no study on the relationship between trait EI, burnout, and hopelessness has been conducted yet.

Finally, our results in the mediation and moderation model partially confirmed our hypothesis: TEI partially mediates the relationship between burnout and hopelessness, and this could reduce the risk of the onset of psychopathology. Being flexible and willing to adapt to new conditions, knowing how to recognize, regulate, express, and communicate emotions, being successful, self-confident, talented in networking, and capable of coping with pressure and regulate stress act as a buffer to burnout consequences, reducing the risk of developing negative cognitive patterns in which the negative expectation towards the future prevails. Hopelessness is, in fact, characterized by the tendency to attribute negative life events to internal, stable, generalizable factors that could lead to suicidal ideation. For these reasons, there seems to be a clear need for psychological interventions aimed at improving the emotional awareness of healthcare workers. Working on these abilities, people who work in a high-risk context could better manage emergencies and stress. The COVID-19 pandemic, in its dramatic nature, has highlighted the importance of the role of the psychologist within hospitals. Interventions such as Balint group (Huang et al., 2020) and mental health services (Sultana et al., 2020) have already proven to be effective ways to relieve burnout and improve the quality of work life for healthcare workers.

In Italy, during the COVID-19 pandemic, the National Council of the Order of Psychologists (CNOP) sought to provide emergency and structural responses to the growing need for psychological support for healthcare workers, such as the creation of task forces of psychologists in healthcare facilities³.

However, in this study, the mediation of TEI in the burnout-hopelessness relationship partially explains the protective role of individual factors against burnout, even in emergencies such as the first COVID-19 pandemic wave.

³ <https://www.psy.it/covid-19-e-gestione-dello-stress-al-via-liniziativa-per-il-supperto-psicologico-del-personale-sanitario/>

Regarding the possible moderator role of changes in workload in the burnout-hopelessness relationship, our analysis doesn't support this hypothesis. However, this finding highlights how individual factors have more influence on suicide risk, even in high-risk individuals such as healthcare workers with burnout.

Individuals with high levels of hopelessness believe that nothing will turn out in their favor, that they will never be successful in life, that their important goals will not be achieved, and that their problems will not be solved, regardless of the actual situation.

LIMITATIONS

Our study suffers from some limitations. First of all, the cross-sectional design does not allow us to grasp the effect of the changes in the variables over time. Moreover, the use of self-reports only may be associated with common method bias. Moreover, even though recruitment procedures (i.e., snowball sampling method through social media, emails, and the university's website) allowed us to reach as many voluntary participants as possible during the lockdown, the sample composition could be biased due to the fact that online recruitment procedures tend to select individuals who are more active on both the internet and social media platforms. Finally, some other variables (personal and/or situational) that should influence the results could be excluded from the analysis and not controlled.

CONCLUSION

Although much attention has been paid to the effects of the COVID-19 pandemic on burnout and mental health in healthcare workers, little has been carried out so far to understand the underlying specific mediation and moderation mechanisms. To our knowledge, this is the first study to examine the role of trait EI in burnout syndrome in healthcare workers during the COVID-19 pandemic. The model we tested could more effectively grasp the complexity of the conditions through which a significant specific work-related stress can affect healthcare workers' mental health. The model, in fact, included protective (TEI) and risk (burnout) psychological factors that could have influenced psychopathological risks such as hopelessness and suicide risk. Our findings indicated that trait EI partially mediated the relationship between burnout and hopelessness; these effects were gender- and age-range-independent. This study identifies a pathway (i.e., via trait EI) that may help better understand the mechanism by which the extreme consequence of burnout syndrome, such as suicide

risk, could be limited. Our findings support the need to integrate considerations on both psychological risk and protective factors into COVID-19 care, including the monitoring of psychological symptoms and social needs, especially among healthcare workers. Therefore, we believe that health promotion policies must take into account both contextual and personal variables, in order to plan in advance intervention for the high-risk group.

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

ETHICS STATEMENT

The studies involving human participants were reviewed and approved by the Bioethics Committee of the University of Palermo (n. 7/2020). The participants provided their written informed consent to participate in this study.

ACKNOWLEDGEMENTS

The authors thank all physicians and nurses who participated in this study. Thanks to healthcare workers on the whole for their efforts and dedication to guarantee assistance to the general population in a dramatic moment such as the first wave of the COVID-19 pandemic.

AUTHOR CONTRIBUTIONS

MSE, SLG, ET, and FA; conceptualization. MSE, SLG, FA and LDP; methodology. MR, VS, MF and GM; data collection and data curation. MAP; data analysis. MSE, MAP, VS, and MR writing the original draft manuscript. MSE, MAP, and FA; review and editing. MSE, SLG, FA, ET; supervision. All authors contributed to the article and approved the submitted version.

FUNDING

This research receives no external funding.

CONFLICT OF INTEREST

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

REFERENCES

- Adamson, E. (2014). Caring behaviour of nurses in Malaysia is influenced by spiritual and emotional intelligence, psychological ownership and burnout. *Evid. Based Nurs.*, 17(4), 121-121. doi.org/10.1136/eb-2013-101704.
- Alrawashdeh, H. M., Al-Tammemi, A. B., Alzawahreh, M. K., Al-Tamimi, A., Elkholy, M., Al Sarireh, F., Abusamak, M., Elehamer, N. M., Malkawi, A., Al-Dolat, W., Abu-Ismael, L., Al-Far, A., & Ghoul, I. (2021). Occupational Burnout and job satisfaction among physicians in times of covid-19 crisis: A convergent parallel mixed-method study. *BMC Public Health*, 21(1). doi.org/10.1186/s12889-021-10897-4
- Andrei, F.; Mancini, G.; Baldaro, B.; Trombini, E.; Agnoli, S. (2014). A systematic review on the predictive utility of the Trait Emotional Intelligence Questionnaire (TEIQue). *Appl. Psychol. Bull.* 62, 2–29.
- Andrei, F.; Siegling, A.; Aloe, A.M.; Baldaro, B.; Petrides, K.V. (2015). The Incremental Validity of the Trait Emotional Intelligence Questionnaire (TEIQue): A Systematic Review and Meta-Analysis. *J. Pers. Assess*, 98, 261–276. doi.org/10.1080/00223891.2015.1084630.
- Andrei, F.; Mancini, G.; Agostini, F.; Epifanio, M.S.; Piombo, M.A.; Riolo, M.; Spicuzza, V.; Neri, E.; Lo Baido, R.; La Grutta, S. and Trombini E. (2022). Quality of Life and Job Loss during the COVID-19 Pandemic: Mediation by Hopelessness and Moderation by Trait Emotional Intelligence. *Int. J. Environ. Res. Public Health*, 19, 2756. <https://doi.org/10.3390/ijerph19052756>
- Baptista, S., Teixeira, A., Castro, L., Cunha, M., Serrão, C., Rodrigues, A., & Duarte, I. (2021). Physician Burnout in Primary Care during the COVID-19 Pandemic: A Cross-Sectional Study in Portugal. *J. Prim. Care & Community Health*, 12, 21501327211008437. doi.org/10.1177/21501327211008437
- Barello, S., Palamenghi, L., & Graffigna, G. (2020). Burnout and somatic symptoms among frontline healthcare professionals at the peak of the Italian COVID-19 pandemic. *Psychiatry Res.*, 290, 113129. doi.org/10.1016/j.psychres.2020.113129
- Bar-On, R., Brown, J. M., Kirkcaldy, B. D., & Thomé, E. P., (2000). Emotional expression and implication for occupational stress; an implication for the Emotional Quotient Inventory (EQ-i). *Pers. Individ. Differ.*, 28, 1107-1118. [doi.org/10.1016/S0191-8869\(99\)00160-9](https://doi.org/10.1016/S0191-8869(99)00160-9)

Beck, A. T., Weissman, A., Lester, D., & Trexler, L. (1974). The measurement of pessimism: the hopelessness scale. *J. Consult Clin Psychol*, 42(6), 861. doi.org/10.1037/h0037562

Bellizzi, S., Fiamma, M., Arru, L., Farina, G., & Manca, A. (2020). COVID-19: The daunting experience of healthcare workers in Sardinia, Italy. *Infect. Control Hosp Epidemiol.*, 41(9), 1118-1119. doi.org/10.1017/ice.2020.149

Bisesti, A., Mallardo, A., Gambazza, S., Binda, F., Galazzi, A., Pazzaglia, S., Laquintana, D. (2021). Facing COVID-19 Pandemic in a Tertiary Hospital in Milan: Prevalence of Burnout in Nursing Staff Working in Sub-Intensive Care Units. *Int. J. Environ. Res. Public Health*, 18, 6684. doi.org/10.3390/ijerph18136684

Bradley, M., & Chahar, P. (2020). Burnout of healthcare providers during COVID-19. *Clevel. Clin. J. Med.* Jul. doi: 10.3949/ccjm.87a.ccc051.

Breslau, L., Finucane, M.L., Locker, A.R., Baird, M.D., Roth, E.A., Collins. R.L. (2021). A longitudinal study of psychological distress in the United States before and during the COVID-19 pandemic. *Preventive Medicine*, Volume 143, 106362, ISSN 0091-7435. doi.org/10.1016/j.ypmed.2020.106362.

Bria, M., Baban, A., & Dumitrascu, D. L. (2012). Systematic review of burnout risk factors among European healthcare professionals. *Cogn. Brain, Behav. An Interdiscip. J.*, 16(3), 423-452.

Claponea, R. M., Pop, L. M., Iorga, M., & Iurcov, R. (2022). Symptoms of Burnout Syndrome among Physicians during the Outbreak of COVID-19 Pandemic—A Systematic Literature Review. *Healthcare*, Vol. 10, No. 6, p. 979. MDPI. doi.org/10.3390/healthcare10060979

Cortés-Álvarez, N., & Vuelas-Olmos, C. (2022). COVID 19: Psychological Effects and Associated Factors in Mexican Nurses. *Disaster Med Public Health Prep*, 16(4), 1377-1383. [doi:10.1017/dmp.2020.495](https://doi.org/10.1017/dmp.2020.495)

Daly, M., Robinson, E. (2022). Psychological distress associated with the second COVID-19 wave: Prospective evidence from the UK Household Longitudinal Study. *J.Affect Disord.*, Vol.310, Pag 274-278, ISSN 0165-0327. doi.org/10.1016/j.jad.2022.05.025.

Davey, M. M., Cummings, G., NEWBURN-COOK, C. V., & Lo, E. A. (2009). Predictors of nurse absenteeism in hospitals: a systematic review. *J. Nurs. Manag.*, 17(3), 312-330. doi.org/10.1111/j.1365-2834.2008.00958.x

De Moraes, A. G., Wilson, M. E., Gajic, O., & Benzo, R. (2015). A101 EFFECTIVE COMMUNICATION STRATEGIES FOR IMPROVEMENT OF PATIENT CARE: The Impact Of Emotional Intelligence On Burnout, Job Satisfaction, Self-Compassion, And Communication Skills In Critical Care Trainees. *Am. J. Respir. Crit. Care Med.*, 191, 1.

Demir, A., Ulusoy, M., & Ulusoy, M. F. (2003). Investigation of factors influencing burnout levels in the professional and private lives of nurses. *Int. J. Nurs. stud.*, 40(8), 807-827. [doi.org/10.1016/S0020-7489\(03\)00077-4](https://doi.org/10.1016/S0020-7489(03)00077-4)

Dewey C, Hingle S, Goelz E, et al. Supporting clinicians during the COVID-19 pandemic. *Ann Intern Med* 2020; 172:752–753. <https://doi.org/10.7326/M20-1033>

Dinibutun, S. R. (2020). Factors associated with burnout among physicians: an evaluation during a period of COVID-19 pandemic. *J. Healthc. Leadersh.*, Sep 15;12:85-94. doi: 10.2147/JHL.S270440.

Dragioti, E., Tsartsalis, D., Mentis, M., Mantzoukas, S., & Gouva, M. (2022). Impact of the COVID-19 pandemic on the mental health of hospital staff: An umbrella review of 44 meta-analyses. *Inter. J. Nurs Stud.*, 104272. doi.org/10.1016/j.ijnurstu.2022.104272.

Duckett, E., Raffaelli, M. & Richards, M.H. (1989). “Taking care”: Maintaining the self and the home in early adolescence. *J Youth Adolescence* 18, 549–565. doi.org/10.1007/BF02139073

Eagly, A. H. (1987). *Sex Differences in Social Behavior: A Social-role Interpretation*. Lawrence Erlbaum Associates.

Eagly, A. H., & Wood, W. (1982). Inferred sex differences in status as a determinant of gender stereotypes about social influence. *Journal of personality and social psychology*, 43(5), 915. doi.org/10.1037/0022-3514.43.5.915

Epifanio, M. S., Andrei, F., Mancini, G., Agostini, F., Piombo, M. A., Spicuzza, V., ... & La Grutta, S. (2021). The impact of COVID-19 pandemic and lockdown measures on quality of life among Italian general population. *J. Clinic. Med.*, 10(2), 289. doi.org/10.3390/jcm10020289.

Fiorilli, C., Benevene, P., De Stasio, S., Buonomo, I., Romano, L., Pepe, A., et al. (2019). Teachers’ burnout: the role of trait emotional intelligence and social support. *Front. Psychol.* . 10:2743. doi.org/10.1016/j.tate.2015.06.001

Franza, F., Basta, R., Pellegrino, F., Solomita, B., & Fasano, V. (2020). The Role of Fatigue of Compassion, Burnout and Hopelessness in Healthcare: Experience in the Time of COVID-19 Outbreak. *Psychiatr. Danub.*, 32(Suppl 1), 10-14. from <https://pubmed.ncbi.nlm.nih.gov/32890354/>

Friedberg, M. W., Chen, P. G., Van Busum, K. R., Aunon, F., Pham, C., Caloyeras, J., ... & Tutty, M. (2014). Factors affecting physician professional satisfaction and their implications for patient care, health systems, and health policy. *Rand Health Q.*, 3(4). PMID: 28083306; PMCID: PMC5051918.

Galanis, P., Vraha, I., Fragkou, D., Bilali, A., & Kaitelidou, D. (2021). Nurses' burnout and associated risk factors during the COVID-19 pandemic: A systematic review and meta-analysis. *J. Adv. Nurs.*, 77(8), 3286-3302. doi.org/10.1111/jan.14839

Ghahramani, S., Lankarani, K. B., Yousefi, M., Heydari, K., Shahabi, S., & Azmand, S. (2021). A systematic review and meta-analysis of burnout among healthcare workers during COVID-19. *Front psychiatry*, 12. doi: [10.3389/fpsyt.2021.758849](https://doi.org/10.3389/fpsyt.2021.758849)

Gualano, M. R., Sinigaglia, T., Lo Moro, G., Rousset, S., Cremona, A., Bert, F., & Siliquini, R. (2021). The burden of burnout among healthcare professionals of intensive care units and

emergency departments during the COVID-19 pandemic: a systematic review. *Int. J. Environ. Res. Public Health*, 18(15), 8172. doi.org/10.3390/ijerph18158172

Guanche Garcell, H. (2020). COVID-19. Un reto para los profesionales de la salud. *Rev haban cienc méd* [online]. vol.19, n.2, e3284. Epub 22-Abr-2020. ISSN 1729-519X

Hacimusalar, Y., Kahve, AC., Yasar, AB., Aydin, MS., (2020). Anxiety and hopelessness levels in COVID-19 pandemic: a comparative study of healthcare professionals and other community samples in Turkey. *J Psychiatr Res.* 2020 Oct;129:181– 8. doi.org/10.1016/j.jpsychires.2020.07.024

Hansel, T.C., Saltzman, L.Y., Melton, P.A. *et al.* (2022). COVID-19 behavioral health and quality of life. *Sci Rep* 12, 961 . doi.org/10.1038/s41598-022-05042-z

Horesh, D., Kapel Lev-Ari, R., & Hasson-Ohayon, I. (2020). Risk factors for psychological distress during the COVID-19 pandemic in Israel: Loneliness, age, gender, and health status play an important role. *Br. J. Health Psychol.*, 25(4), 925-933. doi.org/10.1111/bjhp.12455

<https://portale.fnomceo.it/>

Huang, H., Zhang, H., Xie, Y., Wang, S. B., Cui, H., Li, L., ... & Geng, Q. (2020). Effect of balint group training on burnout and quality of work life among intensive care nurses: a randomized controlled trial. *Neurology, Psychiatry and Brain Research*, 35, 16-21. doi.org/10.1016/j.npbr.2019.12.002

Hughes, D. J., & Evans, T. R. (2018). Putting ‘emotional intelligences’ in their place: Introducing the integrated model of affect-related individual differences. *Front. Psychol.*, 9, 2155. doi.org/10.3389/fpsyg.2018.02155

Humpel, N., & Caputi, P. (2001). Exploring the relationship between work stress, years of experience and emotional competency using a sample of Australian mental health nurses. *J. of Psychiatr. Ment. Health Nurs.*, 8(5), 399-403. doi.org/10.1046/j.1365-2850.2001.00409

Innstrand, S.T., Langballe, E.M., Falkum, E. *et al.* Exploring within- and between-gender differences in burnout: 8 different occupational groups. *Int Arch Occup Environ Health* 84, 813–824 (2011). doi.org/10.1007/s00420-011-0667-y

Jones-Schenk, J. (2020). Hope as a generative force: Lifting our gaze to the future. *J Contin Educ Nurs*, 51(5), 203-204. doi.org/10.3928/00220124-20200415-03

Kang, L., Ma, S., Chen, M., Yang, J., Wang, Y., Li, R., ... & Liu, Z. (2020). Impact on mental health and perceptions of psychological care among medical and nursing staff in Wuhan during the 2019 novel coronavirus disease outbreak: A cross-sectional study. *Brain Behav. Immun.*, 87, 11-17. doi.org/10.1016/j.bbi.2020.03.028

Karagol, A., Zulal Torenli Kaya, Z.T., (2022). Healthcare workers’ burn-out, hopelessness, fear of COVID-19 and perceived social support levels. *Eur. Psychiatry* 36 (2022) 200– 206. doi.org/10.1016/j.ejpsy.2022.01.001

Kimpe, V., Sabe, M., & Sentissi El Idrissi, O. (2022). No increase in burnout in health care workers during the initial COVID-19 outbreak: Systematic review and meta-analysis. *World J. Meta-Anal.*, 10(4), 206-219. doi: [10.13105/wjma.v10.i4.206](https://doi.org/10.13105/wjma.v10.i4.206)

Khosravi, M., Ghiasi, Z., & Ganjali, A. (2022). A narrative review of research on healthcare staff's burnout during the COVID-19 pandemic. *Proceedings of Singapore Healthcare*, 31, 20101058211040575. doi.org/10.1177/20101058211040575

Lai, J., Ma, S., Wang, Y., Cai, Z., Hu, J., Wei, N., ... & Hu, S. (2020). Factors associated with mental health outcomes among health care workers exposed to coronavirus disease 2019. *JAMA network open*, 3(3), e203976-e203976. doi:10.1001/jamanetworkopen.2020.3976

Leiter, M. P., Harvie, P., & Frizzell, C. (1998). The correspondence of patient satisfaction and nurse burnout. *Soc. Sci. Med.*, 47(10), 1611-1617. [doi.org/10.1016/S0277-9536\(98\)00207-X](https://doi.org/10.1016/S0277-9536(98)00207-X)

Linzer, M., Visser, M. R., Oort, F. J., Smets, E. M., McMurray, J. E., & De Haes, H. C. (2001). Predicting and preventing physician burnout: results from the United States and the Netherlands. *Am. J. Med.*, 111(2), 170-175. [doi.org/10.1016/S0002-9343\(01\)00814-2](https://doi.org/10.1016/S0002-9343(01)00814-2)

Lisigurski, M. Z., Shaikh, U., & Toston, B. (2021). Physicians' Emotional Intelligence: Improving Performance While Reducing Burnout. *HCA Healthc. J. Med.*, 2(1), 4. doi.org/10.36518/2689-0216.1074

Liu, Y., Long, Y., Cheng, Y., Guo, Q., Yang, L., Lin, Y., ... & Du, L. (2020). Psychological impact of the COVID-19 outbreak on nurses in China: a nationwide survey during the outbreak. *Front. Psychiatry*, 11, 598712. doi.org/10.3389/fpsyt.2020.598712

Lorant, V., Smith, P., Van den Broeck, K. *et al.* Psychological distress associated with the COVID-19 pandemic and suppression measures during the first wave in Belgium. *BMC Psychiatry* 21, 112 (2021). doi.org/10.1186/s12888-021-03109-1

Loretto, L., Dessanti, A., Nivoli, A., & Bellizzi, S. (2020). The COVID-19 Pandemic in Italy and the World: to be or not to be? That is the real problem. *Health security*, 18(6), 499-501. doi.org/10.1089/hs.2020.0076

Maldonato NM, Bottone M, Chiodi A, et al. A mental health first aid service in an Italian university public hospital during the coronavirus disease 2019 outbreak. *Sustainability* 2020; 12:4244. <https://doi.org/10.3390/su12104244>

Marcia-Rodríguez, C., Alejandro de Ona, A., Martín-Iglesias, D., Barrera-Lopez, L., Perez-Sanz, MT., Moreno-Díaz, J., et al., (2021). Burn-out syndrome in Spanish internists during the COVID-19 outbreak and associated factors: a cross-sectional survey. *BMJ Open*. Feb;11(2):e042966. doi.org/10.1136/bmjopen-2020-042966

Martínez-López, J. Á., Lázaro-Pérez, C., Gómez-Galán, J., & Fernández-Martínez, M. D. M. (2020). Psychological impact of COVID-19 emergency on health professionals: Burnout incidence at the most critical period in Spain. *J. Clinic. Med.*, 9(9), 3029. doi.org/10.3390/jcm9093029

Martins, A.; Ramalho, N.; Morin, E. (2010). A comprehensive meta-analysis of the relationship between Emotional Intelligence and health. *Pers. Individ. Differ.*, 49, 554–564. doi.org/10.1016/j.paid.2010.05.029

Maslach, C. (2003). *Burnout: the cost of caring*. Cambridge, MA: ISHK.

Maslach, C., & Jackson, S. E. (1981). The measurement of experienced burnout. *J. Organ. Behav.*, 2(2), 99-113. doi.org/10.1002/job.4030020205

Maslach, C., & Schaufeli, W. B. (1993). Historical and conceptual development of burnout. Professional burnout: Recent developments in theory and research, 12, 1-16.

Mérida-López, S., & Extremera, N. (2017). Emotional intelligence and teacher burnout: A systematic review. *Int. J. Educ. Res.*, 85, 121-130. doi.org/10.1016/j.ijer.2017.07.006

Mikolajczak, M.; Luminet, O. (2008). Trait emotional intelligence and the cognitive appraisal of stressful events: An exploratory study. *Pers. Individ. Differ.*, 44, 1445–1453. doi.org/10.1016/j.paid.2007.12.012

Moitra, M., Rahman, M., Collins, P. Y., Gohar, F., Weaver, M., Kinuthia, J., ... & Kumar, M. (2021). Mental health consequences for healthcare workers during the COVID-19 pandemic: a scoping review to draw lessons for LMICs. *Front. Psychiatry*, 12, 602614. [doi:10.3389/fpsyt.2021.602614](https://doi.org/10.3389/fpsyt.2021.602614)

Morgantini, L. A., Naha, U., Wang, H., Francavilla, S., Acar, Ö., Flores, J. M., ... & Weine, S. M. (2020). Factors contributing to healthcare professional burnout during the COVID-19 pandemic: a rapid turnaround global survey. *PloS one*, 15(9), e0238217. doi.org/10.1371/journal.pone.0238217

Năstasă, L. E., & Fărcaș, A. D. (2015). The effect of emotional intelligence on burnout in healthcare professionals. *Procedia Soc Behav Sci.*, 187, 78-82. doi.org/10.1016/j.sbspro.2015.03.015

Pellitteri, J. (2002). The relationship between emotional intelligence and ego defense mechanisms. *J. Psychol. Interdiscip. Appl.*, 136, 182–194. doi.org/10.1080/00223980209604149

Petrides, K. V. (2009). Psychometric properties of the Trait Emotional Intelligence Questionnaire. In C. Stough, D. H. Saklofske, and J. D. Parker, *Advances in the assessment of emotional intelligence*. New York: Springer. doi: 10.1007/978-0-387-88370-0_5

Petrides, K.V.; Mikolajczak, M.; Mavroveli, S.; Sanchez-Ruiz, M.-J.; Furnham, A.; Pérez-González, J.-C. (2016). Developments in Trait Emotional Intelligence Research. *Emot. Rev.* 8, 335–341. doi: 10.1177/1754073916650493

Petrides, K.V.; Pita, R.; Kokkinaki, F. (2007). The location of trait emotional intelligence in personality factor space. *Br. J. Psychol.* 98, 273–289. doi.org/10.1348/000712606X120618

Ping, W., Zheng, J., Niu, X., Guo, C., Zhang, J., Yang, H., & Shi, Y. (2020). Evaluation of health-related quality of life using EQ-5D in China during the COVID-19 pandemic. *PloS one*, 15(6), e0234850. doi.org/10.1371/journal.pone.0234850

Pompili, M., Innamorati, M., Gonda, X., Serafini, G., Sarno, S., Erbuto, D., ... & Girardi, P. (2013). Affective temperaments and hopelessness as predictors of health and social functioning in mood disorder patients: a prospective follow-up study. *J. Affect. Disord.*, 150(2), 216-222. doi.org/10.1016/j.jad.2013.03.026

Purvanova, R. K., & Muros, J. P. (2010). Gender differences in burnout: A meta-analysis. *J. Vocat. Behav.* 77(2), 168-185. doi.org/10.1016/j.jvb.2010.04.006

Richards, A. E., Curley, K. L., Zhang, N., Bendok, B. R., Zimmerman, R. S., Patel, N. P., ... & Neal, M. T. (2021). Burnout and Emotional Intelligence in Neurosurgical Advanced Practice Providers Across the United States: A Cross-Sectional Analysis. *World Neurosurg.*, 155, e335-e344. doi.org/10.1016/j.wneu.2021.08.066

Romano, G. A. (2005). La sindrome del burnout nelle helping professions nel paradigma biopsicosociale. *ACP-Rivista di Studi Rogersiani*, 1-14.

Rossi, R., Socci, V., Talevi, D., Mensi, S., Niolu, C., Pacitti, F., ... & Di Lorenzo, G. (2020). COVID-19 pandemic and lockdown measures impact on mental health among the general population in Italy. *Front. in psychiatry*, 11, 790. doi.org/10.3389/fpsyt.2020.00790

Saragih, I. D., Tonapa, S. I., Saragih, I. S., Advani, S., Batubara, S. O., Suarilah, I., & Lin, C. J. (2021). Global prevalence of mental health problems among healthcare workers during the Covid-19 pandemic: a systematic review and meta-analysis. *Int. J. Nurs. Stud.*, 121, 104002. doi.org/10.1016/j.ijnurstu.2021.104002

Schaufeli, W. B., Leiter, M. P., & Maslach, C. (2009). Burnout: 35 years of research and practice. *Career Dev. Int.* Vol. 14 No. 3, pp. 204-220. doi.org/10.1108/13620430910966406

Schenk, J., (2020). Hope as a Generative Force: Lifting Our Gaze to the Future. *J Contin Educ Nurs*; 51:203-204. doi.org/10.3928/00220124-20200415-03

Schutte, N.S.; MalouffJ, M.; Thorsteinsson, E.; Bhullar, N.; Rooke, S.E. (2007). A meta-analytic investigation of the relationship between emotional intelligence and health. *Pers. Individ. Differ.* , 42, 921–933. doi.org/10.1016/j.paid.2006.09.003

Semple, S., Cherrie, J. (2020). COVID-19: Protecting Worker Health. *Ann. Work Expo. Health*, 64, 461–464. doi.org/10.1093/annweh/wxaa033

Shahzad, S., & Bagum, N. (2012). Gender differences in trait emotional intelligence: A comparative study. *Business Review*, 7(2), 106-112.

Shariatpanahi, G., Asadabadi, M., Rahmani, A., Effatpanah, M., & Esslami, G. G. (2022). The Impact of Emotional Intelligence on Burnout Aspects in Medical Students: Iranian Research. *Educ. Res. Int*, 2022. doi.org/10.1155/2022/5745124

Sharma, A., Sharp, D. M., Walker, L. G., & Monson, J. R. (2008). Stress and burnout among colorectal surgeons and colorectal nurse specialists working in the National Health Service. *Colorectal Disease*, 10(4), 397-406.

Sharma, A., Sharp, D. M., Walker, L. G., & Monson, J. R. T. (2008). Stress and burnout in colorectal and vascular surgical consultants working in the UK National Health Service. *Psycho-Oncology: J. of the Psychological, Social and Behavioral Dimensions of Cancer*, 17(6), 570-576. doi.org/10.1111/j.1463-1318.2007.01338.x

Soto-Rubio, A., Giménez-Espert, M.d.C., Prado-Gascó, V. (2020). Effect of Emotional Intelligence and Psychosocial Risks on Burnout, Job Satisfaction, and Nurses' Health during the COVID-19 Pandemic. *Int. J. Environ. Res. Public Health*, 17, 7998. doi.org/10.3390/ijerph17217998

Sultana, A., Sharma, R., Hossain, M. M., Bhattacharya, S., & Purohit, N. (2020). Burnout among healthcare providers during COVID-19: Challenges and evidence-based interventions. *Indian J Med Ethics*, 5(4), 308-11. doi.org/10.31235/osf.io/4hxda

Sun, P., Wang, M., Song, T., Wu, Y., Luo, J., Chen, L., & Yan, L. (2021). The psychological impact of COVID-19 pandemic on health care workers: a systematic review and meta-analysis. *Front. Psych*, 12, 626547. doi: 10.3389/fpsyg.2021.626547

Swami, M. K., Mathur, D. M., & Pushp, B. K. (2013). Emotional intelligence, perceived stress and burnout among resident doctors: an assessment of the relationship. *Nat. Med. J. India*, 26(4), 210-213. PMID: 24758443.

Tesi, A., (2021). A Dual Path Model of Work-Related Well-Being in Healthcare and Social Work Settings: The Interweaving Between Trait Emotional Intelligence, End-User Job Demands, Coworkers Related Job Resources, Burnout, and Work Engagement. *Front. Psychol*. 12:660035. doi: 10.3389/fpsyg.2021.660035.

Tiete, J., Guatteri, M., Lachaux, A., Matossian, A., Hougardy, J. M., Loas, G., & Rotsaert, M. (2021). Mental health outcomes in healthcare workers in COVID-19 and non-COVID-19 care units: a cross-sectional survey in Belgium. *Front. Psychol.*, 11, 612241. doi.org/10.3389/fpsyg.2020.612241

Tuna, T., & Özdin, S. (2021). Levels and Predictors of Anxiety, Depression, and Burnout Syndrome in Physicians During the COVID-19 Pandemic. *Int J Ment Health Addiction* 19, 2470–2483. doi.org/10.1007/s11469-021-00505-2

Van Ballegooijen, H., Goossens, L., Bruin, R. H., Michels, R., & Krol, M. (2021). Concerns, quality of life, access to care and productivity of the general population during the first 8 weeks of the coronavirus lockdown in Belgium and the Netherlands. *BMC Health Serv Res* 21, 227 (2021). doi.org/10.1186/s12913-021-06240-7

Vitale, E., Galatola, V., & Mea, R. (2020). Exploring within and between gender differences in burnout levels in Italian nurses engaged in the Covid-19 health emergency: a cohort observational study. *Minerva Psichiatr.*, 162-170.

Vlachou, E. M., Damigos, D., Lyrakos, G., Chanopoulos, K., Kosmidis, G., & Karavis, M. (2016). The relationship between burnout syndrome and emotional intelligence in healthcare professionals. *Health Sci. J.*, 10(5), 0-0. doi: 10.4172/1791-809X.1000100502

Wang, W., Song, W., Xia, Z., He, Y., Tang, L., Hou, J., & Lei, S. (2020). Sleep disturbance and psychological profiles of medical staff and non-medical staff during the early outbreak of COVID-19 in Hubei Province, China. *Front. Psychiatry*, 11, 733. doi.org/10.3389/fpsyt.2020.00733

White, R. G., & Van Der Boor, C. (2020). Impact of the COVID-19 pandemic and initial period of lockdown on the mental health and well-being of adults in the UK. *BJPsych Open*, 6(5), E90. doi:10.1192/bjo.2020.79

Wong TW., Yau JKY., Chan CLW., Kwong RSY., Ho SMY., Lau CC., et al. (2005). The psychological impact of severe acute respiratory syndrome outbreak on healthcare workers in emergency departments and how they cope. *Eur J Emerg Med.* 12:13–8.

Woo T., Ho R., Tang A., Tam W. (2020). Global prevalence of burnout symptoms among nurses: a systematic review and meta-analysis. *J. Psychiatr. Res.* 123, 9–20. doi.org/10.1016/j.jpsychires.2019.12.015

Wu, Y., Wang, J., Luo, C., Hu, S., Lin, X., Anderson, A. E., ... & Qian, Y. (2020). A comparison of burnout frequency among oncology physicians and nurses working on the frontline and usual wards during the COVID-19 epidemic in Wuhan, China. *J. Pain Symptom Manag.* 60(1), e60-e65. doi.org/10.1016/j.jpainsymman.2020.04.008

Zareei, M., Tabanejad, Z., Oskouie, F., Ebadi, A., & Mesri, M. (2022). Job burnout among nurses during COVID-19 pandemic: A systematic review. *J Educ Health Promot.* , 11. doi: 10.4103/jehp.jehp_797_21.

Zeb S, Akbar A, Gul A, Haider SA, Poulouva P, Yasmin F. Work-Family Conflict, Emotional Intelligence, and General Self-Efficacy Among Medical Practitioners During the COVID-19 Pandemic. *Psychol Res Behav Manag.* 2021 Nov 15;14:1867-1876. doi: 10.2147/PRBM.S333070.

Zeidner, M.; Matthews, G.; Roberts, R.D. (2006). Emotional intelligence, coping with stress, and adaptation. In *Emotional Intelligence in Everyday Life*; Psychology Press: New York, NY, USA; pp. 100–125.

Zhang, H., Ye, Z., Tang, L., Zou, P., Du, C., Shao, J., ... & Mu, S. Y. (2020). Anxiety symptoms and burnout among Chinese medical staff of intensive care unit: the moderating effect of social support. *BMC Psychiatry* 20, 197 (2020). doi.org/10.1186/s12888-020-02603-2

Zysberg, L. Emotional Intelligence and Health Outcomes. (2018). *Psychology* , 09, 2471–2481. doi.org/10.4236/psych.2018.911142

TABLE 1: Mean \pm standard deviation for main study variables divided by gender, region, professional profile and COVID-19 ward.

	Gender		Region		Professional profiles		Covid Wards	
	Female	Male	North	South	Nurses	Physicians	Yes	No
PHE	20.77 \pm 6.50	18.77 \pm 6.63**	20.60 \pm 6.35	18.66 \pm 7.23**	20.08 \pm 6.72	20.35 \pm 6.45	20.40 \pm 6.55	19.82 \pm 6.66
DET	16.94 \pm 5.20	16.28 \pm 4.8	16.90 \pm 4.83	16.22 \pm 5.97	17.00 \pm 5.24	16.49 \pm 4.92	16.96 \pm 5.03	16.35 \pm 5.21
JIN	14.62 \pm 4.90	13.37 \pm 5.05**	14.37 \pm 4.76	13.88 \pm 5.67	14.08 \pm 4.83	14.48 \pm 5.11	14.29 \pm 5.080	14.23 \pm 4.75
DIS	15.82 \pm 7.37	15.37 \pm 7.09	16.02 \pm 7.13*	14.42 \pm 7.763	15.92 \pm 7.52	15.45 \pm 7.04	15.63 \pm 7.42	15.82 \pm 7.05
WOR	.41 \pm .71	.26 \pm .79*	.43 \pm .711	0.14 \pm .79**	0.49 \pm .67	0.24 \pm .78**	0.48 \pm .69	0.15 \pm .77***
TEI	4.96 \pm .70	5.11 \pm .76*	4.97 \pm .70	5.12 \pm .78*	4.92 \pm .71	5.09 \pm .72**	4.98 \pm .72	5.04 \pm .71
HPS	6.93 \pm 4.98	6.80 \pm 5.08	7.00 \pm 4.92	6.53 \pm 5.29	6.61 \pm 4.95	7.21 \pm 5.05	6.99 \pm 5.02	6.70 \pm 4.991

Note: PHE= Psychophysical Exhaustion, DET= Deterioration of relations with clients, JIN=Job Ineffectiveness, DIS=Disappointment, WOR=Workload, TEI=trait Emotional Intelligence, HPS=Hopelessness.

*p<.05, **p<.01,***p<.001

TABLE 2: Correlational analysis

	2	3	4	5	6	7	8	9
1.Gender	.13**	-.09*	-.13**	-.05	-.11**	-.02	.09*	-.01
2.Region		-.16**	-.11**	-.03	-.02	-.08*	.09*	-.03
3.Workload			.17**	.13**	.04	.02	-.01	.07
4.PE				.55**	.62**	.71**	-.55**	.57**
5.DET					.50**	.52**	-.42**	.35**
6.JIN						.59**	-.55**	.46**
7.DIS							-.56**	.63**
8.TEI								-.59**
9. HPS								

Note: PE= Psychophysical Exhaustion, DET= Deterioration of relations with clients, JIN=Job Ineffectiveness, DIS=Disappointment, TEI=trait Emotional Intelligence, HPS=Hopelessness

*p<0.05, **p<0.01

TABLE 3: Frequencies of Burnout levels, Hopelessness and change in Workload

		Region		Gender		Professional profiles		COVID-19 Wards	
		North	South	Female	Male	Nurses	Phys.	Yes	No
PE	Average	41.6%	42.5%	39.7%	47.4%	43.7%	39.8%	41.9%	41.8%

	High	20.7%	20.4%	22.1%	16.7%	23.5%	17.5%	19.9%	21.0%
	Low	5.9%	19.4%	6.2%	14.7%	6.1%	11.2%	11.8%	6.9%
	Moderate	31.8%	17.7%	32.0%	21.2%	26.6%	31.6%	26.3%	30.3%
Chi Squared Test	χ^2	22.00**		17.60**		8.34*		4.27	
DET	Average	27.7%	36.9%	28.1%	34.0%	27.3%	32.3%	33.9%	27.7%
	High	20.2%	22.1%	21.9%	17.3%	25.3%	15.6%	18.8%	21.5%
	Low	0.7%	2.5%	1.2%	0.6%	1.7%	0.4%	1.1%	1.1%
	Moderate	51.4%	38.5%	48.8%	48.1%	45.7%	51.7%	46.2%	49.7%
Chi Squared Test	χ^2	8.93*		2.86		10.87*		2.38	
JIN	Average	70.7%	51.6%	70.2%	57.1%	71.7%	61.0%	62.4%	68.6%
	High	5.0%	4.9%	5.7%	3.2%	4.1%	5.9%	2.7%	6.1%
	Low	8.4%	23.0%	7.1%	23.1%	8.2%	15.2%	13.4%	10.6%
	Moderate	15.9%	20.5%	17.0%	16.7%	16.0%	17.8%	21.5%	14.6%
Chi Squared Test	χ^2	29.86**		29.04***		9.68*		7.99*	
DIS	Average	50.7%	50.8%	51.0%	50.0%	49.8%	51.7%	48.4%	51.9%
	High	12.5%	15.6%	13.8%	11.5%	13.3%	13.0%	12.4%	13.6%
	Low	7.5%	17.2%	8.6%	12.2%	9.9%	9.3%	9.1%	9.8%
	Moderate	29.3%	16.4%	26.6%	26.3%	27.0%	26.0%	30.1%	24.7%
Chi Squared Test	χ^2	16.08**		1.95		.20		1.85	
HPS	High	10.2%	9.8%	10.1%	10.3%	9.2%	11.2%	7.5%	11.4%

	Low	35.7%	27.9%	34.5%	32.7%	34.1%	33.8%	26.9%	37.5%
	Moderate	23.6%	21.3%	22.7%	24.4%	21.5%	24.9%	28.5%	20.5%
	Normal	30.5%	41.0%	32.8%	32.7%	35.2%	30.1%	37.1%	30.6%
Chi Squared Test	χ^2	5.19		0.25		2.31		11.07*	
WOR	Less	13.0%	25.6%	13.3%	21.9%	10.2%	21.6%	23.7%	11.7%
	Same	30.7%	34.7%	32.0%	30.3%	30.7%	32.5%	37.6%	28.5%
	More	56.3%	39.7%	54.7%	47.7%	59.0%	45.9%	38.7%	59.7%
Chi Squared Test	χ^2	15.20**		6.45*		16.32**		24.95***	

Note: PE= Psychophysical Exhaustion, DET= Deterioration of relations with clients, JIN=Job Ineffectiveness, DIS=Disappointment, WOR=Workload, HPS=Hopelessness

*p<.05, **p<.01,*** p<.001

TABLE 4: Testing the mediation and moderation effect of trait Emotional Intelligence and change in workload on the relationship between Burnout dimensions and Hopelessness.

Model	Predictors	B	SE	t	95%CI
Model 1	Psychophysical Ex.	.25***	.31	8.05	[.19; .31]
	TEI	-2.80***	.26	10.63	[-3.31; -2.28]
	Workload	-.58	.68	-.84	[-1.92; .76]
	Workload x Psychophysical Ex.	.03	.03	.35	[-.03; -.09]
	R ²	.44			
	F	110.31***			

Model 2	Det. relations with clients	.10	.03	2.64	[.02; .18]
	TEI	-3.78***	.25	-14.73	[-4.29; -3.28]
	Workload	-.05	.78	-.06	[-1.59; 1.49]
	Workload x Det. relations with clients	.02	.04	.47	[-.06; .11]
	R ²	.37			
	F	82.09***			
Model 3	Job ineffectiveness	.18***	.04	4.24	[.09; .27]
	TEI	-3.41***	.27	-12.35	[-3.96; -2.87]
	Workload	.38	.66	.57	[-.93; 1.69]
	Workload x Job ineffectiveness	.00	.04	-.31	[-.08; .08]
	R ²	.38			
	F	86.64***			
Model 4	Disappointment	.30***	.02	11.02	[11.10; 17.16]
	TEI	-2.41***	.25	-9.59	[-2.91; -1.92]
	Workload	.36	.48	.73	[-.59; 1.31]
	Workload x Disappointment	-.00	.02	-.02	[-.05; .05]
	R ²	.49			
	F	135.31***			

Note. Change in workload was dummy coded such that -1= less workload, 0= no changes in workload, 1= more workload. TEI= Trait emotional intelligence. *p<0.05, **p<0.01, *p<0.001**

Figure 1. Mediation and moderation model of association between all Burnout dimensions, trait Emotional Intelligence, changes in workload and Hopelessness.

