

Appendix III: Descriptive statistics for the samples used in the study

Variable	H3	H4	H8	R5	R9	R10	U2	UMProduct
African Pygmies M (<i>n</i> =8)								
Median	36.7	51.72	50.5	19.6	37	28.14	28.735	151.3
Mean	37.43	51.94	53.19	19.72	37.7	28.69	29.3	161.17
SD	2.592	3.626	6.186	1.593	4.148	1.777	1.461	39.631
Variance	6.72	13.145	38.272	2.537	17.21	3.157	2.135	1570.663
Coefficient of Variation	6.92	6.98	11.63	8.08	11.004	6.19	4.99	24.59
Southeast Asian Negrito's M (<i>n</i> =15)								
Median	39.76	53.05	60.5	19.55	40	30.1	31.69	155.3
Mean	38.89	52.37	58.8	19.48	40.2	29.47	30.92	158.33
SD	2.824	4.984	5.249	1.67	4.538	2.855	2.457	24.354
Variance	7.977	24.839	27.548	2.79	20.59	8.154	6.035	593.119
Coefficient of Variation	7.262	9.52	8.93	8.575	11.288	9.69	7.945	15.382
South African Bantu M (<i>n</i> =23)								
Median	44.36	60.97	66	22.1	43.8	33.87	36.52	233.7
Mean	44.62	61.05	66.7	22.46	44.3	33.69	36.31	227.31
SD	2.77	3.171	5.207	1.705	3.243	1.993	2.711	31.497
Variance	7.676	10.056	27.113	2.908	10.52	3.974	7.352	992.077
Coefficient of Variation	6.21	5.194	7.806	7.593	7.321	5.917	7.467	13.856

Appendix III: Descriptive statistics for the samples used in the study

Variable	H3	H4	H8	R5	R9	R10	U2	UMProduct
Medieval Hungarians M ($n=22$).								
Median	46.28	63.08	68.25	21.815	45.75	34.91	35.12	241.4
Mean	46.24	63.28	67.7	22.19	45.8	34.82	34.83	235.24
SD	3.378	3.251	4.591	2.303	4.134	3.001	3.619	35.238
Variance	11.412	10.57	21.081	5.305	17.089	9.007	13.095	1241.75
Coefficient of Variation	7.305	5.138	6.782	10.379	9.025	8.619	10.39	14.98
Australian Aborigines M ($n=20$)								
Median	41.57	58.4	64.25	21.535	43.5	32.77	33.77	195.55
Mean	41.8	58.83	65.24	21.46	44.03	32.56	34.1	198.69
SD	1.44	2.36	5.05	1.71	3.49	1.62	1.85	35.97
Variance	2.09	5.59	25.48	2.93	12.16	2.62	3.41	1294.23
Coefficient of Variation	3.445	4.011	7.741	7.968	7.926	4.975	5.425	18.103
Southwest Amerindians M ($n=22$)								
Median	43.785	59.705	62.25	20.68	40.25	31.85	33.86	178.9
Mean	43.63	59.7	62.2	21.03	40.24	32.65	33.82	175.41
SD	2.24	3.24	5.46	1.59	2.99	2.21	1.82	30.74
Variance	5.025	10.47	29.8	2.53	8.96	4.9	3.32	945.18
Coefficient of Variation	5.134	5.427	8.778	7.561	7.43	6.769	5.381	17.525

Appendix III: Descriptive statistics for the samples used in the study

Variable	H3	H4	H8	R5	R9	R10	U2	UMProduct
Libben Amerindians M ($n=15$)								
Median	44.48	60.52	64.7	21.71	37.5	32.18	36	188.1
Mean	44.48	61.41	64.7	21.6	38	32.5	36.19	186.85
SD	2.72	3.74	3.99	0.88	2.17	1.81	1.63	22.49
Variance	7.4	13.98	16	0.777	4.7	3.29	2.67	505.7
Coefficient of Variation	6.115	6.09	6.167	4.074	5.71	5.57	4.504	12.036
Recent Human "Macro-Sample" M ($n=88$).								
Median	41.85	56.795	62.25	20.72	42.5	31.585	34.01	186.11
Mean	41.84	56.986	61.88	20.74	42.07	31.46	33.61	184.01
SD	4.183	5.071	6.928	1.954	4.304	3.073	3.618	41.667
Variance	17.501	25.721	47.998	3.818	18.527	9.447	13.09	1736.106
Coefficient of Variation	9.999	8.99	11.196	9.42	10.23	9.77	10.764	22.643
<i>Pan pansicus</i> M ($n=10$)								
Median	38.78	56.655	64	21.945	41.5	35.615	30.955	175.1
Mean	38.11	56.28	63	21.58	41.2	35.28	29.65	168.87
SD	1.57	3	5	1.42	2.98	2	6.17	32.98
Variance	2.45	9.01	21	2.03	8.9	4.02	38.11	1087.8
Coefficient of Variation	4.12	5.33	7.936	6.58	7.233	5.669	20.81	19.53

Appendix III: Descriptive statistics for the samples used in the study

Variable	H3	H4	H8	R5	R9	R10	U2	UMProduct
<i>Pan troglodytes</i> M (n=19)								
Median	43.03	64.21	76	25.55	50	38.51	35.67	242.5
Mean	43.21	64.46	76.1	35.67	51	38.42	35.82	239.55
SD	2.64	3.02	5.01	1.61	4	2.19	2.79	36.4
Variance	6.95	9.11	25.1	2.59	14	4.81	7.77	1325.31
Coefficient of Variation	6.11	4.685	6.583	4.513	7.843	5.7	7.789	15.195
<i>Gorilla gorilla</i> M (n=24)								
Median	66.29	103.32	112.125	34.36	66.9	52.385	50.525	523.15
Mean	65.64	102.1	109.15	33.91	65.28	51.97	50.48	519.79
SD	4.25	7.93	10.33	2.89	5.74	3.41	3.442	75.46
Variance	18.09	62.9	106.64	8.36	32.96	11.61	11.85	5694.3
Coefficient of Variation	6.475	7.767	9.464	8.522	8.793	6.561	6.818	14.517

Appendix III: Descriptive statistics for the samples used in the study

Variable	F6	F16	F17	T3	DTProduct	T13
African Pygmies M (<i>n</i> =8)						
Median	69.5	37.65	65.84	62.32	1368	71.75
Mean	70.5	37.99	65.48	61.83	1383	71.9
SD	7.111	1.907	2.77	2.964	186.574	4.606
Variance	50.571	3.636	7.673	8.785	34809.714	21.218
Coefficient of Variation	10.088	5.02	4.23	4.794	13.49	6.406
Southeast Asian Negrito's M (<i>n</i> =15)						
Median	78	40.24	69.99	63.77	1537	75
Mean	76.5	39.79	69.38	64.08	1499.33	73.9
SD	4.959	3.161	5.055	4.785	210.177	5.852
Variance	24.588	9.993	25.559	22.898	44174.352	34.245
Coefficient of Variation	6.482	7.944	7.286	7.467	14.018	7.919
South African Bantu M (<i>n</i> =23)						
Median	88	44.8	78.73	73.69	1920	87.5
Mean	88.18	45.46	79.19	74.01	1915.26	85.7
SD	8.059	3.452	4.515	3.631	220.109	6.038
Variance	64.95	11.914	20.387	13.183	48447.838	36.454
Coefficient of Variation	9.139	7.593	5.702	4.906	11.492	7.045

Appendix III: Descriptive statistics for the samples used in the study

Variable	F6	F16	F17	T3	DTProduct	T13
Medieval Hungarians M ($n=22$).						
Median	88.5	46.92	80.76	73.85	2038	81.25
Mean	88.2	47.21	80.41	73.54	2050.82	81.2
SD	7.305	3.323	4.858	4.278	275.635	7.573
Variance	53.366	11.044	23.597	18.304	75974.632	57.351
Coefficient of Variation	8.282	7.039	6.041	5.818	13.44	9.326
Australian Aborigines M ($n=20$)						
Median	84.625	42.92	75.355	69.71	3304.5	82.5
Mean	84.6	42.62	75.37	70.55	3414.45	82.71
SD	5.54	2.37	3.4	3.8	321.32	7.32
Variance	30.66	5.61	11.54	14.44	103249.1	53.63
Coefficient of Variation	6.548	5.561	4.511	5.386	9.41	8.85
Southwest Amerindians M ($n=22$)						
Median	83.5	43.445	78.58	72.08	1893.5	83.25
Mean	83.17	43.63	78.47	71.99	1889.59	84.4
SD	6.36	2.19	3.59	3.62	195.43	5.44
Variance	40.44	4.79	12.9	13.09	38195.2	29.6
Coefficient of Variation	7.647	5.02	4.575	5.028	10.342	6.445

Appendix III: Descriptive statistics for the samples used in the study

Variable	F6	F16	F17	T3	DTProduct	T13
Libben Amerindians M (<i>n</i> =15)						
Median	85	45.5	79.72	75.5	1820	84
Mean	84	45.5	79.72	75.39	1834.92	83.2
SD	4.67	2.37	4.07	3.26	184.54	4.67
Variance	21.9	5.62	16.53	10.6	34054.64	21.8
Coefficient of Variation	5.559	5.209	5.105	4.324	10.057	5.613
Recent Human "Macro-Sample" M (<i>n</i> =88).						
Median	82.15	42.17	74.375	68.57	1715.935	78.75
Mean	81.039	42.45	73.918	68.21	1692.65	78.21
SD	7.243	3.993	6.649	6.923	362.027	7.138
Variance	52.257	15.942	44.21	47.933	131063.635	50.953
Coefficient of Variation	8.937	9.405	8.995	10.15	21.388	9.127
<i>Pan pansicus</i> M (<i>n</i> =10)						
Median	68	30.295	57.475	53.945	877.265	58.5
Mean	68	30.62	57.86	54.73	890.29	58.4
SD	5	1.74	2.9	3.34	107.67	5.78
Variance	28	3.022	8.41	11.15	11592.77	33.4
Coefficient of Variation	7.353	5.682	5.01	6.103	12.094	9.897

Appendix III: Descriptive statistics for the samples used in the study

Variable	F6	F16	F17	T3	DTProduct	T13
<i>Pan troglodytes</i> M (n=19)						
Median	79.5	34.15	63.46	60.61	1117	66
Mean	79.9	34.51	64.61	60.15	1118.54	67
SD	3.5	1.91	3.42	4.58	123.75	6.5
Variance	12.2	3.67	11.72	20.97	15313.18	42
Coefficient of Variation	4.38	5.535	5.293	7.615	11.063	9.701
<i>Gorilla gorilla</i> M (n=24)						
Median	116.3	51.14	97.675	90.8	2395	89.75
Mean	114.9	50.6	96.8	89.63	2337.37	87.5
SD	9.22	3.36	6.35	5.81	282.8	7.43
Variance	84.97	11.28	40.27	33.81	79972.94	55.2
Coefficient of Variation	8.024	6.64	6.56	6.482	12.099	8.491

Appendix III: Descriptive statistics for the samples used in the study

Variable	H3	H4	H8	R5	R9	R10	U2	UMProduct
African Pygmies F (<i>n</i> =6)								
Median	32.88	47.575	50.5	18.13	35.5	26.55	27.62	117.25
Mean	33.7	48.59	50.58	17.85	35.8	26.98	28.19	130.47
SD	2.42	4.57	4.042	1.565	2.043	2.62	1.363	30.372
Variance	5.858	20.881	16.342	2.45	4.175	6.866	1.859	922.467
Coefficient of Variation	7.181	9.404	7.992	8.769	5.707	9.712	4.836	23.279
Southeast Asian Negrito's F (<i>n</i> =5)								
Median	33.83	44.87	49	16.76	33	25.29	28.64	98.58
Mean	34.42	46.46	51.8	16.48	34.5	26.34	28.43	117.66
SD	1.633	4.45	5.307	1.21	3.9	2.124	2.787	36.214
Variance	2.666	19.801	28.163	1.465	15.208	4.513	7.768	1311.468
Coefficient of Variation	4.744	9.578	10.245	7.345	11.304	8.065	9.803	30.779
South African Bantu F (<i>n</i> =28)								
Median	39.79	56.46	61	20.155	40	31.34	34.37	173.25
Mean	40.06	56.45	61.5	20.24	40.1	30.79	33.91	178.58
SD	2.3	3.016	6.317	1.253	2.972	3.297	2.371	34.566
Variance	5.288	9.069	39.902	1.57	8.834	10.868	5.622	1194.812
Coefficient of Variation	5.74	5.343	10.271	6.19	7.412	10.707	6.992	19.356

Appendix III: Descriptive statistics for the samples used in the study

Variable	H3	H4	H8	R5	R9	R10	U2	UMProduct
Medieval Hungarians F ($n=22$).								
Median	39.695	54.46	59	19.36	40.75	30.21	31.945	164
Mean	40.17	54.61	58.7	19.03	40.2	30.32	31.16	163.48
SD	1.816	2.488	3.427	2.159	2.585	1.466	2.954	18.591
Variance	3.297	6.192	11.748	4.663	6.683	2.15	8.729	345.63
Coefficient of Variation	4.52	4.557	5.839	11.347	6.431	4.836	9.482	11.372
Australian Aborigines F ($n=8$)								
Median	36.91	50.78	55	18.525	38.5	30.075	30.81	156.1
Mean	36.18	51.33	54.81	18.77	37.56	29.99	30.36	145.51
SD	2.002	3.48	3.21	1.4	3.33	1.43	2.33	20.52
Variance	4.01	12.08	10.28	1.95	11.11	2.036	5.45	420.96
Coefficient of Variation	5.533	6.78	5.856	7.459	8.866	4.768	7.674	14.102
Southwest Amerindians F ($n=10$)								
Median	38.025	53.74	56	18.66	37	29.35	31.77	157.6
Mean	38.47	53.53	58.1	18.85	36.78	29.35	31.72	155.43
SD	2.27	2.95	4.52	0.85	2.43	1.9	1.42	21.95
Variance	5.16	8.72	20.5	0.72	5.92	3.617	2.01	481.65
Coefficient of Variation	5.901	5.511	7.78	4.509	6.607	6.473	4.477	14.122

Appendix III: Descriptive statistics for the samples used in the study

Variable	H3	H4	H8	R5	R9	R10	U2	UMProduct
Libben Amerindians F ($n=19$)								
Median	39.47	54.81	59	19.82	35	30.1	31.18	159.63
Mean	39.23	55.22	59.4	19.5	35.8	30.17	31.57	161.06
SD	1.79	2.67	3.1	1.16	1.72	1.61	1.23	16.72
Variance	3.19	7.13	9.63	1.36	2.97	2.58	1.52	279.61
Coefficient of Variation	4.563	4.835	5.219	5.949	4.804	5.336	3.896	10.381
Recent Human "Macro-Sample" F ($n=33$).								
Median	36.36	52.08	54.75	18.07	37	28.28	30.38	142.54
Mean	36.82	51.346	55.488	17.845	37.143	28.582	30.34	147.63
SD	3.666	4.613	6.918	2.266	3.954	2.979	3.141	38.659
Variance	13.442	21.277	47.86	5.135	15.632	8.873	9.867	1494.547
Coefficient of Variation	9.956	8.983	12.467	12.698	10.645	10.421	10.353	26.186
<i>Pan pansicus</i> F ($n=8$)								
Median	37.425	53.72	64	20.855	40.5	33.815	30.165	177.55
Mean	36.47	52.64	60	20.04	39.1	33.4	29.94	166.5
SD	4.28	4.25	8	2.88	3.89	2.13	2.69	28.87
Variance	18.31	18.06	63	8.28	15.1	4.52	7.26	833.48
Coefficient of Variation	11.736	8.074	13.333	14.371	9.949	6.377	8.985	17.339

Appendix III: Descriptive statistics for the samples used in the study

Variable	H3	H4	H8	R5	R9	R10	U2	UMProduct
<i>Pan troglodytes</i> F (n=25)								
Median	40.66	60.55	72.5	23.98	47	34.89	32.89	201.7
Mean	40.34	59.92	73	23.98	48	34.57	32.59	215.92
SD	2.18	4.05	5.14	1.47	5	2.38	2.42	51.44
Variance	4.76	16.45	26.5	2.17	21	5.68	5.86	2646.13
Coefficient of Variation	5.404	6.759	7.041	6.13	10.417	6.884	7.425	23.824
<i>Gorilla gorilla</i> F (n=25)								
Median	51.44	81.06	87.5	26.76	52	40.97	39.11	306.7
Mean	51.5	79.46	87.19	26.83	51.43	40.87	39.81	319.37
SD	2.73	4.75	6.97	1.37	4.01	3.2	2.45	69.61
Variance	7.46	22.59	48.65	1.88	16.09	10.23	5.99	4845.01
Coefficient of Variation	5.301	5.978	7.994	5.106	7.797	7.83	6.154	21.796

Appendix III: Descriptive statistics for the samples used in the study

Variable	F6	F16	F17	T3	DTProduct	T13
African Pygmies F (<i>n</i> =6)						
Median	70.25	34.595	61.715	57.44	1076.5	66
Mean	69.33	35.65	62.49	57.61	1156.5	66.7
SD	4.936	2.418	3.441	4.572	162.051	5.373
Variance	24.367	5.849	11.843	20.902	26260.7	28.867
Coefficient of Variation	7.12	6.784	5.507	7.936	14.012	8.055
Southeast Asian Negrito's F (<i>n</i> =5)						
Median	67	35.6	59.24	56.67	1144	63
Mean	69	35.33	61.48	56.46	1186.1	63.9
SD	5.964	2.553	4.844	5.287	280.7	4.45
Variance	35.575	6.363	23.461	27.957	78792.55	19.8
Coefficient of Variation	8.644	7.14	7.878	9.365	23.666	6.963
South African Bantu F (<i>n</i> =28)						
Median	80	40.68	72.255	67.475	1634	79.75
Mean	81.31	40.5	72.3	67.71	1639.43	78.55
SD	5.098	1.656	3.35	3.229	172.83	6.35
Variance	25.994	2.744	11.226	10.429	29870.99	40.321
Coefficient of Variation	6.27	4.09	4.634	4.769	10.542	8.084

Appendix III: Descriptive statistics for the samples used in the study

Variable	F6	F16	F17	T3	DTProduct	T13
Medieval Hungarians F (<i>n</i> =22).						
Median	77	40.74	71.255	66.05	1636.5	70
Mean	76.9	40.98	70.9	65.72	1631.41	70.7
SD	3.686	1.669	2.398	2.413	130.526	4.275
Variance	13.586	2.786	5.751	5.823	17037.015	18.227
Coefficient of Variation	4.793	4.073	3.382	3.672	8	6.047
Australian Aborigines F (<i>n</i> =8)						
Median	75.5	37.155	67.85	63.135	2724.5	76.5
Mean	76.23	37.71	66.49	62.33	2676.62	77.13
SD	4.08	2.33	4.35	3.77	227.75	4.02
Variance	16.63	5.42	18.9	14.18	51871.7	16.13
Coefficient of Variation	5.352	6.179	6.542	6.048	8.509	5.212
Southwest Amerindians F (<i>n</i> =10)						
Median	76.75	38.475	69.815	65.395	1538	73.75
Mean	75.6	38.62	69.85	64.36	1533.8	73.7
SD	5.17	1.97	3.15	2.66	139.51	4.81
Variance	26.78	3.86	9.92	7.05	19461.11	23.1
Coefficient of Variation	6.839	5.101	4.51	4.133	9.096	6.526

Appendix III: Descriptive statistics for the samples used in the study

Variable	F6	F16	F17	T3	DTProduct	T13
Libben Amerindians F (<i>n</i> =19)						
Median	76	41.5	72.12	69	1537	72.5
Mean	77	41.61	71.8	68.6	1545.97	73.2
SD	3.23	1.55	2.47	1.98	108.38	2.96
Variance	10.5	2.39	6.1	3.94	11747.01	8.78
Coefficient of Variation	4.195	3.725	3.44	2.886	7.01	4.044
Recent Human "Macro-Sample" F (<i>n</i> =33).						
Median	73.25	38.11	67.26	62.465	1437.56	68.37
Mean	75.198	38.02	67.32	61.89	1387.82	69.931
SD	9.782	3.593	6.003	5.375	315.485	7.856
Variance	95.692	12.91	36.039	28.891	99530.796	61.711
Coefficient of Variation	13.008	9.45	8.918	8.684	22.732	11.233
<i>Pan pansicus</i> F (<i>n</i> =8)						
Median	67	30.175	57.175	53.24	845.5	56.9
Mean	65	29.89	55.27	51.71	823.72	54.1
SD	8	3.24	5.17	5.07	118.57	5.96
Variance	57	10.48	26.73	25.7	14058.4	435.5
Coefficient of Variation	12.308	10.84	9.35	9.805	14.394	11.017

Appendix III: Descriptive statistics for the samples used in the study

Variable	F6	F16	F17	T3	DTProduct	T13
<i>Pan troglodytes</i> F (n=25)						
Median	75	32.33	59.55	55.99	1012	62
Mean	76	32.37	59.18	55.97	1008.07	63
SD	6.34	2.28	3.12	2.91	130.18	6.1
Variance	40.2	5.19	9.76	8.48	16947.26	37
Coefficient of Variation	8342	7.043	5.272	5.199	12.914	9.682
<i>Gorilla gorilla</i> F (n=25)						
Median	92	41.92	76.95	71.99	1465	68.5
Mean	94.64	41.69	77.37	72.58	1476.84	68.2
SD	7.26	2.44	3.82	4.2	151.88	4.96
Variance	53.7	5.94	14.61	17.61	23067.06	24.6
Coefficient of Variation	7.671	5.853	4.937	5.787	10.284	7.273

Appendix III: Descriptive statistics for the samples used in the study

Variable	Acetabulum	Biacetabular	<i>W</i>	<i>Wb</i>	<i>M</i>	<i>J</i>	<i>Dw</i>	<i>Dm</i>	<i>Dw/Dm</i>
African Pygmies M (<i>n</i> =10)									
Median	44.28	116.22	278.52	232.01	348.7	618.8	69.63	58.26	121.17
Mean	44.46	115.49	278.48	231.98	346.4	624.9	69.42	58.35	120.79
SD	1.001	5.412	47.02	39.17	48.78	90.82	2.866	4.875	7.588
Variance	1.001	29.296	2210.911	1534.081	2379.886	8249.159	8.212	23.765	57.852
Coefficient of Variation	2.251	4.686	16.884	16.885	14.082	14.533	4.128	8.355	6.282
Southeast Asian Negrito's M (<i>n</i> =19)									
Median	49.2	128.615	362.3	301.8	419.8	770.7	77.055	64.85	117.275
Mean	48.18	125.17	385.2	320.8	445.1	789.2	75.13	64.44	118.19
SD	4.559	7.609	55.87	46.56	59.83	149.66	4.878	4.647	6.326
Variance	20.782	57.905	3121.652	2167.778	3579.994	22398.989	23.792	21.598	40.021
Coefficient of Variation	9.462	6.079	14.504	14.514	13.441	18.963	6.493	7.211	5.352
South African Bantu M (<i>n</i> =30)									
Median	53.52	135.2	458.52	382.34	525.32	989.34	81.79	73.2	111.87
Mean	53.95	135.2	458.5	382.3	525.32	989.34	81.79	73.2	111.87
SD	2.849	8.644	60.87	51.87	85.62	145.68	4.605	5.16	8.061
Variance	8.119	74.717	3704.58	2690.113	7330.497	21222.774	21.208	26.628	64.992
Coefficient of Variation	5.281	6.393	13.276	13.568	16.299	14.725	5.63	7.049	7.206

Appendix III: Descriptive statistics for the samples used in the study

Variable	Acetabulum	Biacetabular	<i>W</i>	<i>Wb</i>	<i>M</i>	<i>J</i>	<i>Dw</i>	<i>Dm</i>	<i>Dw/Dm</i>
Medieval Hungarians M (<i>n</i> =26).									
Median	56.38	151.25	572.81	477.15	668.62	1245.73	90.555	76.27	120.74
Mean	56.36	150.6	576.4	480.14	682.41	1258.81	90.15	75.68	119.56
SD	3.355	4.851	82.56	68.77	88.12	168.741	2.8	4.655	5.868
Variance	11.259	23.531	6816.552	4730.004	7765.48	28473.60516	7.843	21.671	34.434
Coefficient of Variation	5.953	3.221	14.323	14.323	12.913	0.134048029	3.106	6.151	4.908
Caucasus Tartars M (<i>n</i> =15)									
Median	54.77	151.99	570.9	475.6	758.75	1329.5	89.3	71.69	126.485
Mean	53.99	149.19	560.4	466.9	746.6	1312	88.61	70.55	126.95
SD	4.126	11.565	62.39	51.97	134.2	189.973	6.666	6.112	8.341
Variance	17.022	133.76	3893.044	2700.554	18017.346	36089.6	44.434	37.359	69.581
Coefficient of Variation	7.642	7.752	11.133	11.131	17.975	14.479	7.523	8.663	6.57
Australian Aborigines M (<i>n</i> =21)									
Median	51.03	133.42	458.25	381.72	529.6	992.15	80.18	68.17	116.61
Mean	51.27	132.28	464.35	386.8	534.3	998.65	79.61	68.26	115.52
SD	2.085	5.736	70.717	58.908	77.94	146.742	3.128	4.417	4.991
Variance	4.327	32.907	5000.938	3470.21	6074.389	21533.315	9.786	19.507	24.907
Coefficient of Variation	4.067	4.336	15.229	15.229	14.587	14.694	3.929	6.471	4.32

Appendix III: Descriptive statistics for the samples used in the study

Variable	Acetabulum	Biacetabular	<i>W</i>	<i>Wb</i>	<i>M</i>	<i>J</i>	<i>Dw</i>	<i>Dm</i>	<i>Dw/Dm</i>
Southwest Amerindians M (<i>n</i> =25)									
Median	51.06	144.31	582.69	485.38	697.93	1280.63	85.57	71.98	119.55
Mean	51.06	144.31	582.69	485.38	697.93	1280.63	85.57	71.98	119.55
SD	2.798	10.084	66.178	55.126	81.002	143.542	5.598	4.824	6.006
Variance	7.829	101.687	4379.539	3038.905	6561.357	20604.16	31.34	23.266	36.079
Coefficient of Variation	5.479	6.988	11.357	11.357	11.606	11.209	6.542	6.702	5.024
Libben Amerindians M (<i>n</i> =19)									
Median	52.31	143.71	584.13	486.58	666.19	1257.17	85.46	73.51	117.01
Mean	52.03	143.47	583.66	486.19	681.47	1265.14	85.33	72.92	116.92
SD	2.122	6.876	73.584	61.296	94.687	164.576	3.996	4.591	6.267
Variance	4.502	47.286	5414.41	3757.18	8965.57	27085.16	15.966	21.076	39.274
Coefficient of Variation	4.078	4.793	12.607	12.607	13.894	13.001	4.683	6.296	5.36
Recent Human "Macro-Sample" M (<i>n</i> =97).									
Median	52.92	133	473.6	394.5	562.65	1040.33	80.5	67.02	120.55
Mean	52.73	133.08	465.2	387.5	566.71	1029.5	80.88	67.12	122.09
SD	4.92	18.33	107.1	89.19	145.4	248.95	8.072	7.405	10.218
Variance	24.206	336.018	11463.778	7954.854	21142.348	61973.99	65.166	54.841	104.41
Coefficient of Variation	9.33	13.774	23.022	23.017	2.566	24.182	9.98	11.032	8.369

Appendix III: Descriptive statistics for the samples used in the study

Variable	Acetabulum	Biacetabular	<i>W</i>	<i>Wb</i>	<i>M</i>	<i>J</i>	<i>Dw</i>	<i>Dm</i>	<i>Dw/Dm</i>
African Pygmies F (<i>n</i> =7)									
Median	42.18	122.33	316	263.22	408.8	732.5	72.54	53.43	129.34
Mean	42.79	120.9	311.03	259.08	403.2	714.2	71.69	54.52	129.44
SD	2.173	6.797	34.197	28.49	55.63	88.27	3.748	5.238	7.24
Variance	4.72	46.207	1169.44	811.49	3094.49	7791.891	14.045	27.433	52.433
Coefficient of Variation	5.078	5.622	10.995	10.996	13.797	12.359	5.228	9.607	5.593
Southeast Asian Negrito's F (<i>n</i> =6)									
Median	42.34	128.29	375.8	313	466.9	842.7	74.6	57.685	130.67
Mean	43.92	130.23	367.7	306.3	486.7	854.5	76.65	57.64	133
SD	3.79	8.826	75.18	62.61	93.11	165.77	5.283	5.002	11.031
Variance	14.364	77.904	5652.013	3920.248	8669.423	27478.845	27.913	25.021	121.692
Coefficient of Variation	8.629	6.777	20.446	20.441	19.131	19.399	6.892	8.678	8.294
South African Bantu F (<i>n</i> =30)									
Median	49.55	137.23	454.2	378.3	581.07	1048.45	81.835	65.51	124.89
Mean	49.3	137.99	461.5	384.4	580.58	1042.4	82.13	65.65	125.45
SD	2.273	9.137	44.56	37.12	79.06	120.76	4.882	3.76	8.359
Variance	5.168	83.484	1985.886	1377.93	6251.24	14582.01	23.834	14.134	69.886
Coefficient of Variation	4.61	6.621	9.655	9.656	13.617	11.585	5.944	5.723	6.663

Appendix III: Descriptive statistics for the samples used in the study

Variable	Acetabulum	Biacetabular	<i>W</i>	<i>Wb</i>	<i>M</i>	<i>J</i>	<i>Dw</i>	<i>Dm</i>	<i>Dw/Dm</i>
Medieval Hungarians F (<i>n</i> =24).									
Median	48.55	147.33	512.15	426.62	628.78	1130.92	85.62	66.775	129.27
Mean	48.88	147.77	507.05	422.38	646.83	1153.89	86.74	68.35	127.46
SD	3.057	8.271	63.08	52.55	93.65	153.385	4.522	3.798	7.617
Variance	9.347	68.41	3979.475	2761.19	8769.881	23526.811	20.452	14.426	58.019
Coefficient of Variation	6.254	5.597	12.44	12.441	14.478	13.293	5.213	5.557	5.976
Australian Aborigines F (<i>n</i> =8)									
Median	45.79	130.55	422.48	351.93	533.3	955.78	76.96	61.03	128.46
Mean	45.5	130.05	433.1	360.77	572.6	1005.68	76.98	59.63	130.89
SD	1.022	6.389	60.279	50.214	95.1	153.306	3.241	4.499	6.88
Variance	1.045	40.829	3633.536	2521.411	9043.53	23502.63	10.504	20.243	47.34
Coefficient of Variation	2.246	4.913	13.918	13.918	16.608	15.244	4.21	7.545	5.256
Southwest Amerindians F (<i>n</i> =18)									
Median	45.03	140.41	473.88	394.74	631.715	1123.28	81.66	63.88	128.79
Mean	45.07	139.68	482.27	401.73	625.04	1107.31	81.66	63	129.81
SD	2.612	7.721	34.872	29.047	61.139	91.226	4.021	3.082	7.48
Variance	6.822	59.615	1216.045	843.752	3738.014	8322.18	16.166	9.497	55.954
Coefficient of Variation	5.795	5.528	7.231	7.23	9.782	8.238	4.924	4.892	5.762

Appendix III: Descriptive statistics for the samples used in the study

Variable	Acetabulum	Biacetabular	<i>W</i>	<i>Wb</i>	<i>M</i>	<i>J</i>	<i>Dw</i>	<i>Dm</i>	<i>Dw/Dm</i>
Libben Amerindians F (<i>n</i> =21)									
Median	48.45	137.3	509.99	424.82	642.9	1144.22	81.72	65.06	124.63
Mean	48.44	137.62	516.98	430.65	645.84	1162.82	81.53	66.16	124.76
SD	1.901	9.205	49.76	41.44	84.94	128.688	4.784	4.987	9.849
Variance	3.614	84.726	2476.064	1717.64	7215.057	16560.555	22.889	24.872	97.018
Coefficient of Variation	3.924	6.689	9.625	9.623	13.152	11.066	5.868	7.538	7.894
Recent Human "Macro-Sample" F (<i>n</i> =41).									
Median	48.46	128.29	392.9	327.25	510.72	922.17	76.3	61.04	131.14
Mean	48.21	132.46	395.7	329.6	507.46	894.13	78.71	60.64	131.92
SD	3.865	11.758	83.67	69.69	96.022	171.284	6.732	7.086	12.476
Variance	14.939	138.249	7001.213	4857.34	9220.213	29338.038	45.326	50.216	155.648
Coefficient of Variation	8.017	8.876	21.145	21.144	18.922	19.156	8.553	11.685	9.457

Appendix III: Descriptive statistics for the samples used in the study

Variable	Total Load	<i>RP(FemHead)</i>	Stature	Bi-iliac/Stature	Bicristal	FMArea	FMML	FMAP
African Pygmies M (<i>n</i> =10)								
Median	71050	56.91	136.4	14.36	21.23	412.15	18.69	17.21
Mean	71041.67	55.5	139.52	14.23	20.99	397.33	18.46	16.79
SD	11995.25	7.643	6.891	0.529	0.909	69.346	1.488	1.578
Variance	143886093.8	58.424	47.491	0.28	0.827	4808.867	2.215	2.491
Coefficient of Variation	16.885	13.771	4.939	3.721	4.333	17.453	8.063	0.094
Southeast Asian Negrito's M (<i>n</i> =19)								
Median	92425	72.22	150.34	14.17	22.45	460.2	20.31	17.71
Mean	98047.73	67.72	149.57	14.53	22.94	459.83	20.38	17.69
SD	13934.91	8.477	7.894	0.966	1.446	56.412	1.616	1.411
Variance	194181681.8	71.864	62.337	0.934	2.091	3182.273	2.61	1.992
Coefficient of Variation	14.212	12.518	5.278	6.65	6.303	12.268	7.927	7.978
South African Bantu M (<i>n</i> =30)								
Median	116967.39	59.67	165.1	13.49	23.9	639.77	23.67	21.12
Mean	116967.39	59.67	165.1	13.5	23.89	639.77	23.67	21.12
SD	15525.99	6.848	5.179	1.203	1.674	114.03	2.962	1.785
Variance	241056502	46.902	26.826	1.448	2.802	13002.751	8.772	3.188
Coefficient of Variation	13.274	11.476	3.137	8.913	7.007	17.823	12.512	8.545

Appendix III: Descriptive statistics for the samples used in the study

Variable	Total Load	<i>RP(FemHead)</i>	Stature	Bi-iliac/Stature	Bicristal	FMArea	FMML	FMAP
Medieval Hungarians M (<i>n</i> =26).								
Median	146125	72.04	166.13	16.16	27.14	637.1	21.96	22.1
Mean	147041.67	72.52	166.5	15.76	27.48	630.98	22.3	21.32
SD	21061.983	8.249	6.966	1.422	1.486	97.163	1.88	4.323
Variance	443607145.8	68.044	48.521	2.024	2.209	9440.588	3.534	18.685
Coefficient of Variation	14.324	11.375	4.183	9.026	5.408	15.399	8.43	20.275
Caucasus Tartars M (<i>n</i> =15)								
Median	145637.5	67.365	164.95	15.37	27.65	615.85	22.485	21.29
Mean	142966.67	70.78	165.15	15.52	27.38	622.2	22.62	21.6
SD	15920.26	7.917	5.015	1.283	2.025	56.596	1.237	1.59
Variance	253454810.6	62.677	25.153	1.645	4.099	3203.097	1.53	2.529
Coefficient of Variation	11.136	11.185	3.037	8.264	7.395	9.096	5.469	7.363
Australian Aborigines M (<i>n</i> =21)								
Median	114750	71.74	165.19	14.16	24.345	571.45	22.55	19.675
Mean	107913.46	68.53	165.64	14.07	24.42	578.61	22.79	19.86
SD	31520.723	12.193	8.872	1.115	1.25	88.07	1.894	1.727
Variance	993556001.6	148.689	78.714	1.244	1.563	7756.356	3.589	2.982
Coefficient of Variation	29.209	17.792	5.356	7.928	5.119	15.221	8.313	8.696

Appendix III: Descriptive statistics for the samples used in the study

Variable	Total Load	<i>RP(FemHead)</i>	Stature	Bi-iliac/Stature	Bicristal	FMArea	FMML	FMAP
Southwest Amerindians M (<i>n</i> =25)								
Median	148646.59	86.72	172.18	14.91	26.99	555.09	22.17	18.78
Mean	148646	86.72	172.18	14.91	26.99	555.09	22.17	18.78
SD	16882.167	11.478	4.676	1.141	1.736	88.22	2.164	3.62
Variance	285006208.1	131.741	21.867	1.302	3.015	7782.85	4.685	13.106
Coefficient of Variation	11.357	13.236	2.716	7.654	6.433	15.893	9.763	19.277
Libben Amerindians M (<i>n</i> =19)								
Median	149012.5	82.44	167.82	15.4	26.79	554.65	20.03	22.18
Mean	148337.5	81.71	169.39	15.37	26.85	569.67	20.24	22.08
SD	19292.236	11.735	7.34	1.305	1.731	57.309	1.192	1.485
Variance	372190386	137.722	53.882	1.702	2.996	3284.361	1.42	2.206
Coefficient of Variation	13.056	14.362	4.333	8.489	6.447	10.06	5.888	6.726
Recent Human "Macro-Sample" M (<i>n</i> =97).								
Median	119225	62.23	159.965	14.345	24.86	546.75	21.735	19.615
Mean	118012.36	62.62	158.36	14.54	24.69	540.44	21.44	19.61
SD	27841.96	8.797	12.439	1.54	2.562	108.244	2.294	2.255
Variance	775174969.1	77.385	154.73	2.731	6.563	11716.683	5.261	5.085
Coefficient of Variation	23.592	14.048	7.855	10.589	10.376	20.029	10.698	11.499

Appendix III: Descriptive statistics for the samples used in the study

Variable	Total Load	<i>RP(FemHead)</i>	Stature	Bi-iliac/Stature	Bicristal	FMArea	FMML	FMAP
African Pygmies F (<i>n</i> =7)								
Median	82912.5	71.27	132.8	14.98	21.91	394	17.65	16.28
Mean	83595.83	72.37	134.86	15.12	21.62	369.89	17.51	16.45
SD	10305.79	6.843	9.555	1.416	1.111	60.179	1.772	1.305
Variance	106209354.2	46.826	91.306	2.006	1.235	3621.475	3.14	1.704
Coefficient of Variation	12.328	9.455	7.085	9.368	5.141	16.269	10.119	7.935
Southeast Asian Negrito's F (<i>n</i> =6)								
Median	95875	80.08	140.99	16.42	22.64	390.6	17.64	16.245
Mean	93790	77.86	139.79	16.24	23.75	338.27	18.16	16.78
SD	19179.37	7.384	8.156	1.573	2.214	56.074	1.521	1.947
Variance	367848312.5	54.524	66.516	2.473	4.9	3144.335	2.312	3.792
Coefficient of Variation	20.449	9.484	5.834	9.684	9.32	16.577	8.374	11.605
South African Bantu F (<i>n</i> =30)								
Median	115875	75.64	155.61	14.41	24.02	525.1	21.12	19.105
Mean	117723.91	75.6	156.65	14.21	23.97	542.54	21.72	19.59
SD	11364.79	7.746	4.567	0.893	1.269	75.293	1.846	1.492
Variance	129158663.5	60	20.854	0.797	1.61	5669.001	3.408	2.227
Coefficient of Variation	9.654	10.246	2.921	6.282	5.293	13.878	8.499	7.618

Appendix III: Descriptive statistics for the samples used in the study

Variable	Total Load	<i>RP(FemHead)</i>	Stature	Bi-iliac/Stature	Bicristal	FMArea	FMML	FMAP
Medieval Hungarians F (<i>n</i> =24).								
Median	130650	80.89	160.11	15.21	26.63	484.15	19.67	19.215
Mean	129350	82.06	158.93	15.41	26.57	484.18	19.58	19.41
SD	16093.134	9.542	5.938	1.353	1.426	53.417	1.995	0.899
Variance	258988977.3	91.056	35.26	1.83	2.035	2853.434	3.981	0.809
Coefficient of Variation	12.441	11.628	3.736	8.778	0.537	11.032	10.19	4.634
Australian Aborigines F (<i>n</i> =8)								
Median	107775	84.26	153.93	15.09	24.1	436.05	20.33	17.17
Mean	110483.33	86.02	153.43	14.96	23.95	459.65	20.44	17.63
SD	15376.896	15.259	7.85	1.6	1.413	50.629	1.304	1.19905841
Variance	236448916.7	232.834	61.624	2.562	1.998	2563.288	1.691	1.437741071
Coefficient of Variation	13.918	17.739	5.116	10.699	5.9	11.015	6.362	0.068012389
Southwest Amerindians F (<i>n</i> =18)								
Median	120887.5	92.35	155.78	15.7	25.79	483.3	19.73	18.72
Mean	123028.13	93.08	155.01	15.64	25.29	464.05	19.74	18.45
SD	8895.705	4.617	2.982	1.137	1.66	50.811	1.777	1.166
Variance	79133572.92	21.319	8.895	1.293	2.755	2588.893	3.156	1.359
Coefficient of Variation	7.231	4.96	1.924	7.269	6.563	10.964	9	6.318

Appendix III: Descriptive statistics for the samples used in the study

Variable	Total Load	<i>RP(FemHead)</i>	Stature	Bi-iliac/Stature	Bicristal	FMArea	FMML	FMAP
Libben Amerindians F (<i>n</i> =21)								
Median	130100	82.41	157.07	16.47	26.22	468.3	18.85	19.63
Mean	131881.61	84.87	156.72	16.4	26.48	476	18.99	19.67
SD	12693.472	7.478	5.374	0.964	1.101	45.785	0.771	1.474
Variance	161124238.1	55.919	28.876	0.929	1.212	2096.3	0.594	2.173
Coefficient of Variation	9.625	8.811	3.429	5.879	4.158	9.619	4.058	7.494
Recent Human "Macro-Sample" F (<i>n</i> =41).								
Median	100225	67.47	144.82	14.485	22.39	428.65	18.915	17.535
Mean	100930.36	68.44	146.06	15.04	23.36	446.28	19.25	17.73
SD	21345.16	9.127	10.678	1.857	2.456	120.594	2.591	1.607
Variance	455616081.3	83.302	114.016	3.449	6.034	14542.978	6.716	2.584
Coefficient of Variation	21.148	13.336	7.31	12.349	10.516	27.022	13.463	9.066

Appendix III: Descriptive statistics for the samples used in the study

Variable	STArea	STML	STAP	Iliac Ht.	Iliac Br.	Pubic Lth.	AP Pelvis	Sacrum	Iliac (Theta)
African Pygmies M (<i>n</i> =10)									
Median	433.1	16.755	19.495	81.7	122.82	73.85	121.69	94.49	29.98
Mean	423.17	17.02	19.45	82.01	124.5	75.65	119.7	93.19	29.94
SD	63.019	1.382	1.756	4.959	6.579	5.031	5.422	5.313	1.526
Variance	3971.398	1.909	3.085	24.588	43.288	25.314	29.404	28.229	2.329
Coefficient of Variation	14.892	8.118	9.031	6.046	5.285	6.651	4.53	5.701	5.097
Southeast Asian Negrito's M (<i>n</i> =19)									
Median	487.8	18.21	21.53	87.56	135.78	81.43	120.67	101.16	27.96
Mean	509.35	18.38	21.71	86.65	134.38	80.51	123.23	100.14	27.63
SD	64.545	1.144	1.757	8.021	10.085	3.549	12.721	9.112	3.968
Variance	4166.025	1.308	3.086	64.343	101.712	12.594	161.816	83.022	15.742
Coefficient of Variation	12.672	6.223	8.092	9.257	7.505	4.408	10.333	9.099	14.36
South African Bantu M (<i>n</i> =30)									
Median	703.04	21.33	25.75	94.23	145.25	84.82	129.68	104.4	28.74
Mean	703.04	21.33	25.75	94.23	145.25	84.82	129.68	104.39	28.74
SD	128.819	2.335	2.48	6.718	9.293	4.518	10.83	6.409	2.762
Variance	16594.365	5.454	6.15	45.127	86.358	20.411	117.295	41.07	7.629
Coefficient of Variation	18.323	10.949	9.631	7.139	6.398	5.326	8.351	6.139	9.61

Appendix III: Descriptive statistics for the samples used in the study

Variable	STArea	STML	STAP	Iliac Ht.	Iliac Br.	Pubic Lth.	AP Pelvis	Sacrum	Iliac (Theta)
Medieval Hungarians M (<i>n</i> =26).									
Median	728.3	21.17	26.45	105.38	159.83	96.17	136.875	116.62	30.37
Mean	730.07	21.52	26.63	106.01	159.86	95.19	137.51	117.21	30.38
SD	87.958	1.634	2.238	7.063	8.155	5.26	4.664	8.404	2.734
Variance	7736.589	2.67	5.008	49.89	66.51	27.666	21.753	70.621	7.477
Coefficient of Variation	12.048	7.593	8.403	6.663	5.101	5.526	3.392	7.17	9.001
Caucasus Tartars M (<i>n</i> =15)									
Median	706.7	21.88	25.83	100.38	157.94	90.755	133.35	116.46	31.92
Mean	738.3	22.28	26.04	101.86	157.99	92.69	131.87	116.68	30.98
SD	84.89	1.602	2.484	5.852	11.122	6.462	5.682	6.882	2.155
Variance	7206.357	2.567	6.171	34.246	123.711	41.756	32.287	47.369	4.643
Coefficient of Variation	11.498	7.192	9.54	5.745	7.04	6.971	4.309	5.899	6.955
Australian Aborigines M (<i>n</i> =21)									
Median	615.25	19.99	24.01	89.96	145.74	83.08	125.55	103.17	30.705
Mean	638.22	20.45	24.39	91.69	145.12	81.83	126.79	102.8	31.41
SD	104.381	1.888	2.103	6.974	7.368	5.884	6.103	4.485	2.366
Variance	10895.332	3.563	4.424	48.642	54.288	34.623	37.242	20.116	5.6
Coefficient of Variation	16.355	9.231	8.623	7.606	5.077	7.191	4.813	4.363	7.534

Appendix III: Descriptive statistics for the samples used in the study

Variable	STArea	STML	STAP	Iliac Ht.	Iliac Br.	Pubic Lth.	AP Pelvis	Sacrum	Iliac (Theta)
Southwest Amerindians M (<i>n</i> =25)									
Median	620.85	19.79	24.52	98.13	151.24	92.61	130.37	112.23	32.6
Mean	620.85	19.79	24.52	98.13	151.24	92.61	130.37	112.23	32.6
SD	94.728	1.703	1.965	5.089	9.515	4.573	8.819	6.376	2.306
Variance	8973.5	2.901	3.86	25.903	90.546	20.909	77.712	40.648	5.316
Coefficient of Variation	15.26	8.607	8.013	5.186	6.292	4.937	6.764	5.681	7.072
Libben Amerindians M (<i>n</i> =19)									
Median	672.6	20.47	25.92	103.93	153.24	93.59	135.07	118.83	30.54
Mean	676.6	20.31	26.1	103.82	153.6	94.03	135.51	120.25	30.95
SD	72.643	1.423	1.612	2.847	4.109	5.094	9.678	5.125	3.416
Variance	5277.053	2.026	2.598	8.106	16.888	25.951	93.657	26.262	11.67
Coefficient of Variation	10.736	7.009	6.176	2.742	2.675	5.418	7.142	4.262	11.037
Recent Human "Macro-Sample" M (<i>n</i> =97).									
Median	596.65	19.905	23.355	93.77	145	86.79	128.36	105.9	30.31
Mean	609.13	19.95	23.7	94.38	144.78	86.62	128.33	107.51	30.65
SD	139.696	2.402	2.965	9.586	14.333	9.21	10.05	13.023	3.084
Variance	19515.06	5.769	8.793	91.891	205.452	84.831	101.013	169.614	9.514
Coefficient of Variation	22.93	12.039	12.512	10.157	9.9	10.633	7.832	12.114	10.063

Appendix III: Descriptive statistics for the samples used in the study

Variable	STArea	STML	STAP	Iliac Ht.	Iliac Br.	Pubic Lth.	AP Pelvis	Sacrum	Iliac (Theta)
African Pygmies F (<i>n</i> =7)									
Median	408.5	16.07	19.84	83.6	127.42	79.29	126.87	89.15	30.625
Mean	390.91	15.77	18.73	81.38	128.45	79.91	125.08	92.8	30.34
SD	87.423	2.346	2.745	4.736	3.906	4.499	7.255	6.858	2.518
Variance	7642.828	5.504	7.538	22.433	15.254	20.244	52.642	47.03	6.341
Coefficient of Variation	22.364	14.876	14.658	5.82	3.04	5.63	5.801	7.39	8.299
Southeast Asian Negrito's F (<i>n</i> =6)									
Median	405.5	16.445	18.725	84.62	128.51	79.75	125.38	100.2	28.28
Mean	424.55	16.63	19.96	82.25	132.23	82.91	127.59	100.07	29.73
SD	76.426	1.335	2.413	4.68	10.624	7.413	9.287	4.243	3.177
Variance	5840.971	1.783	5.823	21.903	112.878	54.955	86.25	18	10.091
Coefficient of Variation	18.002	8.03	12.089	5.69	8.035	8.941	7.279	4.24	10.685
South African Bantu F (<i>n</i> =30)									
Median	575.9	19.3	23.4	89.34	144.865	87.4	138.75	108.325	29.12
Mean	585.15	19.67	23.3	89.99	145.47	88.3	133.79	108.58	29.46
SD	79.819	1.788	1.567	5.812	7.947	5.35	22.787	7.652	2.631
Variance	6371.023	3.198	2.456	33.777	63.162	28.627	519.26	58.551	6.925
Coefficient of Variation	13.64	9.091	6.727	6.458	5.463	6.059	17.032	7.047	8.933

Appendix III: Descriptive statistics for the samples used in the study

Variable	STArea	STML	STAP	Iliac Ht.	Iliac Br.	Pubic Lth.	AP Pelvis	Sacrum	Iliac (Theta)
Medieval Hungarians F (<i>n</i> =24).									
Median	579.25	19.02	23.83	97.845	151.55	93.89	132.42	116.38	31.3
Mean	578.99	19.05	23.79	96.92	151.65	94.8	132.93	116.95	31.35
SD	84.06	1.268	2.478	6.573	9.572	5.056	8.109	6.57	3.099
Variance	7066.093	1.607	6.141	43.204	91.626	25.564	65.755	43.172	9.607
Coefficient of Variation	14.518	6.655	10.416	6.782	6.312	5.333	6.1	5.618	9.887
Australian Aborigines F (<i>n</i> =8)									
Median	517.2	18.59	21.79	86.6	139.3	80.44	118.73	101.55	31.72
Mean	505.49	18.23	21.73	87.3	139.13	80.04	119.75	104.04	31.99
SD	55.4551669	0.886	1.641	4.002	7.201	3.916	6.402	8.027	3.346
Variance	3075.275536	0.786	2.692	16.016	51.858	15.333	40.984	64.434	11.194
Coefficient of Variation	10.97	4.862	7.55	4.584	5.176	4.892	5.346	7.715	10.459
Southwest Amerindians F (<i>n</i> =18)									
Median	515.6	18.65	22.23	91.97	138.62	89.09	127.78	112.79	32.64
Mean	528.19	18.82	22.03	91.38	140.1	89.63	128.68	114.08	32.43
SD	58.385	1.449	1.559	5.699	7.496	4.671	6.394	9.342	2.696
Variance	3408.77	2.099	2.431	32.481	56.198	21.823	40.888	87.273	7.217
Coefficient of Variation	11.054	7.698	7.078	6.237	5.351	5.212	4.969	8.189	8.315

Appendix III: Descriptive statistics for the samples used in the study

Variable	STArea	STML	STAP	Iliac Ht.	Iliac Br.	Pubic Lth.	AP Pelvis	Sacrum	Iliac (Theta)
Libben Amerindians F ($n=21$)									
Median	565.5	18.46	24.72	94.19	148.12	93.75	127.51	117.525	32.94
Mean	579.83	18.31	24.8	95.84	148.04	94.34	126.53	118.38	33.53
SD	62.621	1.164	1.742	4.925	8.377	3.062	9.385	4.487	2.453
Variance	3921.414	1.354	3.035	24.254	70.183	9.375	88.089	20.132	6.018
Coefficient of Variation	10.799	6.356	7.024	5.139	5.659	3.246	7.418	3.79	7.317
Recent Human "Macro-Sample" F ($n=41$).									
Median	500.5	17.69	21.84	86.25	133.02	83.31	127.05	104.025	29.965
Mean	498.23	17.84	21.6	88.49	137.65	85.86	130.6	104.77	29.72
SD	116.619	2.263	2.929	8.556	12.198	8.179	14.541	10.449	4.24
Variance	13599.926	5.12	8.512	73.202	148.805	66.892	211.429	109.187	17.977
Coefficient of Variation	23.407	12.683	13.562	9.669	8.862	9.526	11.134	9.973	14.266