

MALLIAVIN DIFFERENTIABILITY OF SOLUTIONS OF HYPERBOLIC STOCHASTIC PARTIAL DIFFERENTIAL EQUATIONS WITH IRREGULAR DRIFTS

ANTOINE-MARIE BOGSO AND OLIVIER MENOUKEU PAMEN

ABSTRACT. We prove path-by-path uniqueness of solutions to hyperbolic stochastic partial differential equations when the drift coefficient is the difference of two componentwise monotone Borel measurable functions of spatial linear growth. The Yamada-Watanabe principle for SDEs driven by Brownian sheet then allows to derive strong uniqueness for such equation and thus extending the results in [Bogso, Dieye and Menoukeu Pamen, *Elect. J. Probab.*, 27:1-26, 2022] and [Nualart and Tindel, *Potential Anal.*, 7(3):661–680, 1997]. Assuming that the drift is globally bounded, we show that the unique strong solution is Malliavin differentiable. The case of a spatial linear growth drift coefficient is also studied.

1. INTRODUCTION

The existence, uniqueness and Malliavin differentiability of strong solutions of SDEs on the plane with smooth coefficients have been obtained in several settings of varying generality. However there are not many results when the coefficients of the such equation are singular. The purpose of the present paper is two-fold: first we obtain the existence and uniqueness of strong solution of the following integral form equation

$$(1.1) \quad X_{s,t} = \xi + \int_0^t \int_0^s b(s_1, t_1, X_{s_1, t_1}) ds_1 dt_1 + W_{s,t}, \text{ for } (s, t) \in \mathbb{R}_+^2,$$

when W is a d -dimensional Brownian sheet and the drift b is the difference of two componentwise monotone functions and of spatial linear growth. We address this problem by using the Yamada-Watanabe argument for SDEs driven by Brownian sheet derived in [37] (see also [47], [45, Remark 2]), that is, we combine weak existence and pathwise uniqueness to obtain the existence of a unique strong solution. More particularly, we replace the pathwise uniqueness by a stronger notion of uniqueness, namely, the path-by-path uniqueness introduced in [14] (see also [19]) in the case of SDEs driven by one-parameter Brownian motion. This notion was introduced in [5] for the two parameter process, as follows:

Definition 1.1. *Let \mathcal{V} (resp. $\partial\mathcal{V}$) be the space of \mathbb{R}^d -valued continuous functions on $[0, T]^2$ (resp. $\{0\} \times [0, T] \cup ([0, T] \times \{0\})$) for some $T > 0$. We say that the path-by-path uniqueness of solutions to (1.1) holds when there exists a full \mathbb{P} -measure set $\Omega_0 \subset \Omega$ such that for all $\omega \in \Omega_0$ the following statement is true: there exists at most one function $y \in \mathcal{V}$ which satisfies*

$$\int_0^T \int_0^T |b(s, t, y_{s,t})| ds dt < \infty, \quad \partial y = x, \text{ for some } x \in \partial\mathcal{V}$$

and

$$(1.2) \quad y_{s,t} = x + \int_0^s \int_0^t b(s_1, t_1, y_{s_1, t_1}) ds_1 dt_1 + W_{s,t}(\omega), \quad \forall (s, t) \in [0, T]^2.$$

Date: November 14, 2023.

2020 Mathematics Subject Classification. Primary 60H07, 60H50, 60H17; Secondary 60H15 .

Key words and phrases. Brownian sheet, SDEs on the plane, path by path uniqueness, Malliavin derivative.

The project on which this publication is based has been carried out with funding provided by the Alexander von Humboldt Foundation, under the programme financed by the German Federal Ministry of Education and Research entitled German Research Chair No 01DG15010.

The study of path-by-path uniqueness is motivated by the problem of regularisation by noise of random ordinary (or partial) differential equations (ODEs or PDEs). In the case of SDEs driven by Brownian motion, path-by-path uniqueness of equation (1.1) was proved in [14] assuming that the drift is bounded and measurable, and the diffusion is constant. This result was extended to the non-constant diffusion in [15] using rough path analysis. There has now been several generalisation of this result. The authors in [4] proved a Sobolev regularity of solutions to the linear stochastic transport and continuity equations with drift in critical L^p spaces. Such a result does not hold for the corresponding deterministic equations. In [7], the authors analysed the regularisation by noise for a non-Lipschitz stochastic heat equation and proved path-by-path uniqueness for any initial condition in a certain class of a set of probability one. In [2], the path-by-path uniqueness for transport equations driven by the fractional Brownian motion of Hurst index $H < 1/2$ with bounded and integrable vector-fields is investigated. In [10; 21] the authors solved the regularisation by noise problem from the point of view of additive perturbations. In particular, the work [10] considered generic perturbations without any specific probabilistic setting whereas authors in [1] construct a new Gaussian noise of fractional nature and proved that it has a strong regularising effect on a large class of ODEs. More recently, the regularisation by noise problem for ODEs with vector fields given by Schwartz distributions in the setting of non-linear Young type of integrals was studied in [23]. It was also proved that if one perturbs such an equation by adding an infinitely regularising path, then it has a unique solution. Let us also mention the recent work [27] in which the authors looked at multidimensional SDEs with distributional drift driven by symmetric α -stable Lévy processes for $\alpha \in (1, 2]$. In all of the above mentioned works, the driving noise considered are one parameter processes.

Our method to prove path-by-path uniqueness follows as in [5]. We show the path-by-path uniqueness on $\Gamma_0 = [0, 1]^2$. More precisely, we consider the integral equation

$$(1.3) \quad X_{s,t} = \xi + \int_0^t \int_0^s b(s_1, t_1, X_{s_1, t_1}) ds_1 dt_1 + W_{s,t} \text{ for } (s, t) \in \Gamma_0,$$

where the drift is of spatial linear growth. We denote by \mathcal{V}_0^1 the space of continuous \mathbb{R}^d -valued functions on Γ_0 which vanish on $\{0\} \times [0, 1] \cup [0, 1] \times \{0\}$. It is shown in [5, Section 1] (see also [14, Section 1]) that path-by-path uniqueness of solutions to (1.3) holds if and only if, with probability one, there is no nontrivial solution $u \in \mathcal{V}_0^1$ of

$$(1.4) \quad u(s, t) = \int_0^s \int_0^t \{b(s_1, t_1, W_{s_1, t_1} + u(s_1, t_1)) - b(s_1, t_1, W_{s_1, t_1})\} ds_1 dt_1, \text{ for } (s, t) \in \Gamma_0.$$

This is the statement of Theorem 2.10 which is extended to unbounded drifts in Theorem 2.9. The proof of Theorem 2.10 relies on some estimates for an averaging operator along the sheet (see Lemma 2.3). This result plays a key role in the proof of a Gronwall type lemma (see Lemma 2.15) which enables us to prove path-by-path uniqueness of solutions to (1.3). The latter combined with the weak existence yield the existence of a unique strong solution. A crucial idea to obtain the Gronwall type inequality is to take advantage of the fact that the set of dyadic numbers is dense in $[-1, 1]^d$. Note that Lemma 2.6 involves real numbers $x \in [-1, 1]^d$ and not functions $u : [0, 1]^2 \rightarrow [-1, 1]^d$. In order to apply Lemma 2.6 in the proof of Lemma 2.15 for a drift that is the difference of two componentwise monotone functions, one needs to first rewrite (1.4) in each dyadic square and carefully replace the function u by the maximum of either its positive part or its negative part. Note that when the drift coefficient is componentwise nondecreasing, similar result can be found in [5; 36].

Secondly, in this paper, we prove Malliavin smoothness of the unique solution to the SDE (1.3). When the coefficients are smooth, the authors in [34; 35] showed existence, uniqueness, Malliavin differentiability and smoothness of density of solutions to SDEs on the plane. Here, assuming that the drift is the difference of two componentwise nondecreasing functions, we show that the solution is Malliavin differentiable. In the one parameter case, the Malliavin differentiability of solutions to SDEs with bounded and measurable coefficients was studied in [31] under an additional commutativity assumption. The later assumption was removed in [30]. It is worth mentioning that in the above work, the Malliavin smoothness of the unique solutions to the SDEs with rough coefficients and driven by Brownian motion is obtained as a byproduct of the method used to study existence and uniqueness. This technique was introduced in [39] and has now been extensively utilised; see for example the work [22] for the case of singular SDEs driven by Lévy noise, [29] for the case of random coefficients and [2; 1] for

the case of singular SDEs driven by fractional noise. In order to prove the Malliavin differentiability of the solution to the SDE (1.3), we take advantage of Gaussian white noise theory and a local time-space integration formula provided in [6, Proposition 3.1] to show that the sequence of approximating sequence of solutions converges strongly in $L^2(\Omega, \mathbb{R}^d)$ to the solution of the SDE (compare with [30]) and we use a compactness criteria given in [32, Lemma 1.2.3] to conclude. An essential step in showing this is to obtain good enough estimates for the Malliavin derivative of the approximating sequence. This task is not trivial and requires the use of the Wendroff inequality (Theorem B.1) which plays a crucial role in the proof.

Equation (1.1) can also be written in a differential form as the following hyperbolic stochastic partial differential equation

$$(1.5) \quad \begin{cases} \frac{\partial^2 X_{s,t}}{\partial s \partial t} = b(s, t, X_{s,t}) + \dot{W}_{s,t}, & (s, t) \in \Gamma, \\ \partial X = \xi, \end{cases}$$

where ∂X is the restriction of X to the boundary $\partial\Gamma = \{0\} \times \mathbb{R}_+ \cup \mathbb{R}_+ \times \{0\}$ of $\Gamma := \mathbb{R}_+^2$, $b : \Gamma \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is Borel measurable, $\dot{W} = (\dot{W}^{(1)}, \dots, \dot{W}^{(d)})$ is a d -dimensional white noise of the Brownian sheet on Γ given on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and $(s, t) \mapsto \xi_{s,t}(\omega)$ is continuous on $\partial\Gamma$ for all $\omega \in \Omega$. Recall that a d -dimensional white noise on Γ is a mean-zero Gaussian process $\dot{W} = (\dot{W}^{(1)}, \dots, \dot{W}^{(d)})$ indexed by the Borel field $\mathcal{B}(\Gamma)$ on Γ with covariance functions

$$\mathbb{E}[\dot{W}^{(i)}(A)\dot{W}^{(j)}(B)] = \delta_{i,j}|A \cap B|, \quad \forall A, B \in \mathcal{B}(\Gamma)$$

where $|\cdot|$ denotes the Lebesgue measure on Γ and $\delta_{i,j} = 1$ if $i = j$ and $\delta_{i,j} = 0$ otherwise. The process $W = (W_{s,t} := \dot{W}([0, s] \times [0, t]), (s, t) \in \Gamma)$ is mean-zero Gaussian process with covariance functions

$$\mathbb{E}[W_{s,t}^{(i)}W_{s',t'}^{(j)}] = \delta_{i,j}(s \wedge s')(t \wedge t'), \quad \forall (s, t), (s', t') \in \Gamma.$$

By the Kolmogorov continuity theorem, there exists a continuous version of W , still denoted by W , which is a d -dimensional Brownian sheet. We consider a nondecreasing and right-continuous family $\mathbb{F} = (\mathcal{F}_{s,t})$ of sub- σ -algebras of \mathcal{F} each of which contains all negligible sets in $(\Omega, \mathcal{F}, \mathbb{P})$ such that W and ξ are \mathbb{F} -adapted, that is $W_{s,t}$ (respectively, $\xi_{s,t}$) is $\mathcal{F}_{s,t}$ -measurable for every $(s, t) \in \Gamma$ (respectively, $(s, t) \in \partial\Gamma$). We refer the reader to Khoshnevisan [26] for a complete analysis on multi-parameter processes and their applications.

Equation (1.5) is a particular case of the quasilinear stochastic hyperbolic differential equation

$$(1.6) \quad \begin{cases} \frac{\partial^2 X_{s,t}}{\partial s \partial t} = b(s, t, X_{s,t}) + a(s, t, X_{s,t})\dot{W}_{s,t}, & (s, t) \in \Gamma \\ \partial X = \xi, \end{cases}$$

where $a : \mathbb{R}_+^2 \times \mathbb{R}^d \rightarrow \mathbb{R}^d \times \mathbb{R}^d$ is a Borel measurable matrix function. A formal $\frac{\pi}{4}$ rotation transforms (1.6) into the following nonlinear stochastic wave equation

$$(1.7) \quad \frac{\partial^2 Y_{\rho,\theta}}{\partial \rho^2} - \frac{\partial^2 Y_{\rho,\theta}}{\partial \theta^2} = \tilde{b}(\rho, \theta, Y_{\rho,\theta}) + \tilde{a}(\rho, \theta, Y_{\rho,\theta})\dot{W}_{\rho,\theta}, \quad (\rho, \theta) \in \tilde{\Gamma},$$

with the Goursat-Darboux type boundary condition $\partial Y = \tilde{\xi}$, where $\tilde{\Gamma} = \{(\rho, \theta) : \theta \geq 0 \text{ and } |\rho| \leq \theta\}$, \dot{W} is a d -dimensional white noise of the Brownian sheet on $\tilde{\Gamma}$, $\tilde{b}(\rho, \theta, y) = b(\frac{\theta+\rho}{\sqrt{2}}, \frac{\theta-\rho}{\sqrt{2}}, y)$ (the same applies to \tilde{a}), $Y_{\rho,\theta} = X_{\frac{\theta+\rho}{\sqrt{2}}, \frac{\theta-\rho}{\sqrt{2}}}$, $\tilde{\xi}_{\theta,\theta} = \xi_{\sqrt{2}\theta, 0}$ and $\tilde{\xi}_{-\theta,\theta} = \xi_{0, \sqrt{2}\theta}$. The $\frac{\pi}{4}$ rotation has been used by Carmona and Nualart [9] (see also [18, Section 0] and [41, Section 1]) to prove existence and uniqueness of solution to (1.7) under a different boundary condition when \tilde{a} and \tilde{b} are time-homogeneous.

Equation (1.5) can also be seen as a noisy analogue of the so-called Darboux problem given by

$$(1.8) \quad \frac{\partial^2 y}{\partial s \partial t} = b\left(s, t, y, \frac{\partial y}{\partial s}, \frac{\partial y}{\partial t}\right) \quad \text{for } (s, t) \in [0, T] \times [0, T],$$

with the initial conditions

$$(1.9) \quad y(0, t) = \sigma(t) \quad \text{on } [0, T] \quad \text{and} \quad y(s, 0) = \tau(s) \quad \text{on } [0, T],$$

where σ and τ are absolutely continuous on $[0, T]$. Using Caratheodory's theory of differential equations, Deimling [16] proved an existence theorem for the system (1.8)-(1.9) when b is Borel measurable in the first two variables and bounded and continuous in the last three variables. Hence the results obtained here can also be seen as a generalisation to the stochastic setting of the above mentioned one.

The remainder of the paper is organised as follows: In Section 2, we provide a path-by-path uniqueness result for (1.3) when the drift b is of linear growth. In Section 3, we study the Malliavin differentiability of the strong solution to (1.3). We show that this solution is Malliavin differentiable for uniformly bounded drifts and when the drift b is of linear growth, we obtain Malliavin differentiability of the solution only for sufficiently small time parameters.

2. EXISTENCE AND UNIQUENESS RESULTS

In this section, we show that the SDE (1.3) has a unique strong solution. Our approach is based on the Yamada-Watanabe principle introduced in [37]. As pointed out earlier, instead of showing the weak existence and pathwise uniqueness, we show weak existence and path by path uniqueness (which implies pathwise uniqueness as shown in [10]). The following preliminary results that have been obtained by applying a local time-space integration formula for Brownian sheets (see [6] for more information) are needed to show path-by-path uniqueness.

2.1. Preliminary results. Let $f : [0, 1]^2 \times \mathbb{R}^d \rightarrow \mathbb{R}$ be a continuous function such that for any $(s, t) \in [0, 1]^2$, $f(s, t, \cdot)$ is differentiable and for any $i \in \{1, \dots, d\}$, the partial derivative $\partial_{x_i} f$ is continuous. We also know from [6, Proposition 3.1] that for a d -dimensional Brownian sheet $(W_{s,t} := (W_{s,t}^{(1)}, \dots, W_{s,t}^{(d)}); s \geq 0, t \geq 0)$ defined on a filtered probability space and for any $(s, t) \in [0, 1]^2$ and any $i \in \{1, \dots, d\}$, we have

$$(2.1) \quad \begin{aligned} & \int_0^s \int_0^t \partial_{x_i} f(s_1, t_1, W_{s_1, t_1}) dt_1 ds_1 \\ &= - \int_0^s \int_0^t f(s_1, t_1, W_{s_1, t_1}) \frac{d_{t_1} W_{s_1, t_1}^{(i)}}{s_1} ds_1 - \int_0^s \int_{1-t}^1 f(s_1, 1-t_1, \widehat{W}_{s_1, t_1}) \frac{d_{t_1} B_{s_1, t_1}^{(i)}}{s_1} ds_1 \\ & \quad + \int_0^s \int_{1-t}^1 f(s_1, 1-t_1, \widehat{W}_{s_1, t_1}) \frac{\widehat{W}_{s_1, t_1}^{(i)}}{s_1(1-t_1)} dt_1 ds_1, \end{aligned}$$

where $\widehat{W}^{(i)} = (\widehat{W}_{s_1, t_1}^{(i)} := W_{s_1, 1-t_1}^{(i)}; 0 \leq s_1, t_1 \leq 1)$ and $B^{(i)} := (B_{s_1, t_1}^{(i)}; 0 \leq s_1, t_1 \leq 1)$ is a standard Brownian sheet with respect to the filtration of $\widehat{W}^{(i)}$, independent of $(W_{s,1}^{(i)}; s \geq 0)$. Here “ $d_{t_1} W_{s_1, t_1}^{(i)}$ ”, resp. “ $d_{t_1} B_{s_1, t_1}^{(i)}$ ” denotes the stochastic line integral with respect to the Brownian motion $(W_{s_1, t_1}^{(i)}; 0 \leq t_1 \leq 1)$, resp. $(B_{s_1, t_1}^{(i)}; 0 \leq t_1 \leq 1)$ for s_1 fixed.

The following result will be extensively used and can be found in [5].

Proposition 2.1. *Let $W := (W_{s,t}^{(1)}, \dots, W_{s,t}^{(d)}; (s, t) \in [0, 1]^2)$ be a \mathbb{R}^d -valued Brownian sheet defined on a filtered probability space $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$, where $\mathbb{F} = (\mathcal{F}_{s,t}; s, t \in [0, 1])$. Let $b \in \mathcal{C}([0, 1]^2, \mathcal{C}^1(\mathbb{R}^d))$, $\|b\|_\infty \leq 1$. Let $(a, a', \varepsilon, \varepsilon') \in [0, 1]^4$. Then there exist positive constants α and C (independent of $\nabla_y b$, a , a' , ε and ε') such that*

$$(2.2) \quad \mathbb{E} \left[\exp \left(\alpha \varepsilon' \varepsilon \left| \int_0^1 \int_0^1 \nabla_y b \left(s, t, \widetilde{W}_{s,t}^{\varepsilon, \varepsilon'} \right) dt ds \right|^2 \right) \right] \leq C.$$

Here $\nabla_y b$ denotes the gradient of b with respect to the third variable, $|\cdot|$ is the usual norm on \mathbb{R}^d and the \mathbb{R}^d -valued two-parameter Gaussian process $\widetilde{W}^{\varepsilon, \varepsilon'} := \left(\widetilde{W}_{s,t}^{(\varepsilon, \varepsilon', 1)}, \dots, \widetilde{W}_{s,t}^{(\varepsilon, \varepsilon', d)}; (s, t) \in [0, 1]^2 \right)$ is given by

$$\widetilde{W}_{s,t}^{(\varepsilon, \varepsilon', i)} = W_{a'+\varepsilon's, a+\varepsilon t}^{(i)} - W_{a', a+\varepsilon t}^{(i)} - W_{a'+\varepsilon's, a}^{(i)} + W_{a', a}^{(i)} \quad \text{for all } i \in \{1, \dots, d\}.$$

For every $0 \leq a < \gamma \leq 1$, $0 \leq a' < \gamma' \leq 1$ and for $(x, y) \in \mathbb{R}^d$ let us define the function ϱ by:

$$\varrho(x, y) = \int_{a'}^{\gamma'} \int_a^\gamma \left\{ b(s, t, W_{s,t} + x) - b(s, t, W_{s,t} + y) \right\} dt ds.$$

Here is a direct consequence of the previous estimation.

Corollary 2.2. *Let $b : [0, 1]^2 \times \mathbb{R}^d \rightarrow \mathbb{R}$ be a bounded and Borel measurable function such that $\|b\|_\infty \leq 1$. Let α , C and $\widetilde{W}^{\varepsilon, \varepsilon'}$ be defined as in Proposition 2.1. Then the following two bounds are valid:*

(1) *For every $(x, y) \in \mathbb{R}^{2d}$, $x \neq y$ and every $(\varepsilon, \varepsilon') \in [0, 1]^2$, we have*

$$(2.3) \quad \mathbb{E} \left[\exp \left(\frac{\alpha \varepsilon' \varepsilon}{|x - y|^2} \left| \int_0^1 \int_0^1 \left\{ b(s, t, \widetilde{W}_{s,t}^{\varepsilon', \varepsilon} + x) - b(s, t, \widetilde{W}_{s,t}^{\varepsilon', \varepsilon} + y) \right\} dt ds \right|^2 \right) \right] \leq C.$$

(2) *For any $(x, y) \in \mathbb{R}^2$ and any $\eta > 0$, we have*

$$(2.4) \quad \mathbb{P} \left(|\varrho(x, y)| \geq \eta \sqrt{(\gamma - a)(\gamma' - a')} |x - y| \right) \leq C e^{-\alpha \eta^2}.$$

For any positive integer n , we divide $[0, 1]$ into 2^n intervals $I_{nk} =]k2^{-n}, (k+1)2^{-n}]$. We define the random real valued function $\varrho_{nkk'}$ on $[-1, 1]^{2d}$ by

$$\varrho_{nkk'}(x, y) := \int_{I_{nk'}} \int_{I_{nk}} \left\{ b(s, t, W_{s,t} + x) - b(s, t, W_{s,t} + y) \right\} dt ds.$$

The next two lemmas provide an estimate for $\varrho_{nkk'}(x, y)$ and $\varrho_{nkk'}(0, x)$ for every dyadic numbers $x, y \in [-1, 1]^d$. Their proofs can be found in [5, Section 5].

Lemma 2.3. *Suppose $b : [0, 1]^2 \times \mathbb{R}^d \rightarrow \mathbb{R}$ is a Borel measurable function such that $|b(s, t, x)| \leq 1$ everywhere on $[0, 1]^2 \times \mathbb{R}^d$. Then there exists a subset Ω_1 of Ω with $\mathbb{P}(\Omega_1) = 1$ such that for all $\omega \in \Omega_1$,*

$$|\varrho_{nkk'}(x, y)(\omega)| \leq C_1(\omega) 2^{-n} \left[\sqrt{n} + \left(\log^+ \frac{1}{|x - y|} \right)^{1/2} \right] |x - y| \quad \text{on } \Omega_1$$

for all dyadic numbers $x, y \in [-1, 1]^d$ and all choices of integers n, k, k' with $n \geq 1$, $0 \leq k, k' \leq 2^n - 1$, where $\log^+ z = \max\{0, \log z\}$ for $z \in (0, \infty)$ and $C_1(\omega)$ is a positive random constant that does not depend on x, y, n, k and k' .

Lemma 2.4. *Suppose b is as in Lemma 2.3. Then there exists a subset Ω_2 of Ω with $\mathbb{P}(\Omega_2) = 1$ such that for all $\omega \in \Omega_2$, for any choice of n, k, k' , and any choice of a dyadic number $x \in [-1, 1]^d$*

$$(2.5) \quad |\varrho_{nkk'}(0, x)(\omega)| \leq C_2(\omega) \sqrt{n} 2^{-n} \left(|x| + 2^{-4n} \right),$$

where $C_2(\omega)$ is a positive random constant that does not depend on x, n, k and k' .

Observe that the above two results require only the drift to be bounded and Borel measurable. Assuming in addition b is nondecreasing, the next two results state that Lemmas 2.3 and 2.4 can be extended to any $x, y \in [-1, 1]^d$ (not only dyadic). The proof of Lemma 2.6 is omitted since it is similar to that of Lemma 2.5.

Lemma 2.5. *Suppose b , Ω_1 and C_1 are as in Lemma 2.3. Suppose in addition that b is componentwise nondecreasing. Then for all $\omega \in \Omega_1$,*

$$|\varrho_{nkk'}(x, y)(\omega)| \leq C_1(\omega)2^{-n} \left[\sqrt{n} + \left(\log^+ \frac{1}{|x-y|} \right)^{1/2} \right] |x-y| \text{ on } \Omega_1$$

for all $x, y \in [-1, 1]^d$ and all choices of integers n, k, k' with $n \geq 1, 0 \leq k, k' \leq 2^n - 1$.

Proof. Fix $\omega \in \Omega_1, x, y \in [-1, 1]^d, n \geq 1$ and $0 \leq k, k' \leq 2^n - 1$. Suppose without loss of generality that $\varrho_{nkk'}(x, y)(\omega) > 0$. For every $i \in \{1, \dots, d\}$ and $\ell \in \mathbb{N}$, define $y_{i,\ell}^- = 2^{-\ell} \lfloor 2^\ell y_i \rfloor, x_{i,\ell}^+ = 1 - 2^{-\ell} \lfloor 2^\ell (1 - x_i) \rfloor, y_\ell^- = (y_{1,\ell}^-, \dots, y_{d,\ell}^-)$ and $x_\ell^+ = (x_{1,\ell}^+, \dots, x_{d,\ell}^+)$. Observe that x_ℓ^+ (respectively y_ℓ^-) is a componentwise non-increasing (respectively non-decreasing) sequence of dyadic vectors that converges to x (respectively y). Hence, as $b(s, t, W_{s,t}(\omega) + x) \leq b(s, t, W_{s,t}(\omega) + x_\ell^+)$ and $b(s, t, W_{s,t}(\omega) + y_\ell^-) \leq b(s, t, W_{s,t}(\omega) + y)$, it follows from Lemma 2.3 that

$$\begin{aligned} |\varrho_{nkk'}(x, y)(\omega)| &= \varrho_{nkk'}(x, y)(\omega) = \int_{I_{nk'}} \int_{I_{nk}} \{b(s, t, W_{s,t}(\omega) + x) - b(s, t, W_{s,t}(\omega) + y)\} dt ds \\ &\leq \int_{I_{nk'}} \int_{I_{nk}} \{b(s, t, W_{s,t}(\omega) + x_\ell^+) - b(s, t, W_{s,t}(\omega) + y_\ell^-)\} dt ds \\ &\leq C_1(\omega)2^{-n} \left[\sqrt{n} + \left(\log^+ \frac{1}{|x_\ell^+ - y_\ell^-|} \right)^{1/2} \right] |x_\ell^+ - y_\ell^-|. \end{aligned}$$

Then, letting ℓ tends to ∞ , we obtain the result. \square

Lemma 2.6. *Suppose b , Ω_2 and C_2 are as in Lemma 2.4. Suppose in addition that b is componentwise nondecreasing. Then for all $\omega \in \Omega_2$*

$$|\varrho_{nkk'}(0, x)(\omega)| \leq C_2(\omega) \sqrt{n} 2^{-n} \left(|x| + 2^{-4n} \right) \text{ on } \Omega_2$$

for all $x \in [-1, 1]^d$ and all choices of integers n, k, k' with $n \geq 1, 0 \leq k, k' \leq 2^n - 1$.

2.2. Main results and proofs. In this section, we prove the path-by-path uniqueness of the solution to (1.3). We use this result to derive the existence and uniqueness of a strong solution to (1.3). We assume the following conditions on the drift. We endow \mathbb{R}^d with the partial order “ \preceq ” defined by

$$x \preceq y \text{ when } x_i \leq y_i \text{ for all } i \in \{1, \dots, d\}.$$

Hypothesis 2.7.

- (1) $b : [0, 1]^2 \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is Borel measurable and admits the decomposition $b = \hat{b} - \check{b}$, where $\hat{b}(s, t, \cdot)$ and $\check{b}(s, t, \cdot)$ are componentwise nondecreasing functions, that is each component \hat{b}_i and $\check{b}_i, 1 \leq i \leq d$ is componentwise nondecreasing. Precisely, for every $x, y \in \mathbb{R}^d$,

$$x \preceq y \Rightarrow \hat{b}_i(s, t, x) \leq \hat{b}_i(s, t, y) \text{ and } \check{b}_i(s, t, x) \leq \check{b}_i(s, t, y).$$

- (2) b is of linear growth uniformly on (s, t) ; precisely, there exists a positive constant M such that

$$|b(s, t, x)| \leq M(1 + |x|), \quad \forall (s, t, x) \in [0, 1]^2 \times \mathbb{R}^d.$$

The main results of this section are the following :

Theorem 2.8. *Suppose b satisfies Hypothesis 2.7. Then the SDE (1.3) admits a unique strong solution.*

The above result constitutes an extension to those in [5; 36] by allowing the drift b to be the difference of two monotone functions. It is proved by using the Yamada-Watanabe principle. However, instead of showing the pathwise uniqueness we show the following path-by-path uniqueness property.

Theorem 2.9. *Suppose b satisfies Hypothesis 2.7. Then for almost every Brownian sheet path W , there exists a unique continuous function $X : [0, 1]^2 \rightarrow \mathbb{R}^d$ satisfying (1.3).*

The proof of Theorem 2.9 is omitted since it follows the same lines as that of [5, Theorem 3.2]. It follows from both Gronwall inequality on the plane and the next result.

Theorem 2.10. *Suppose b is as in Theorem 2.9. Suppose in addition that b is uniformly bounded. Then for almost every Brownian sheet path W , there exists a unique continuous function $X : [0, 1]^2 \rightarrow \mathbb{R}^d$ satisfying (1.3).*

Proof of Theorem 2.8. It follows from the conditions of the theorem that, (1.3) has a weak solution. In addition, since path-by-path uniqueness implies pathwise uniqueness (see e.g. [4, Page 9, Section 1.8.4] where the result is provided in the one-parameter case. This may be extend easily to the two-parameter case.), the result follows from the Yamada-Watanabe type principle for SDEs driven by Brownian sheets (see e.g. Nualart and Yeh [37]). \square

Corollary 2.11. *Suppose that b is as in Theorem 2.9. Then for almost every Brownian sheet path W , there exists a unique continuous function $X : [0, 1]^2 \rightarrow \mathbb{R}^d$ satisfying (1.5).*

Corollary 2.12. *Suppose that b is as in Theorem 2.9. Then for almost every Brownian sheet path W , there exists a unique continuous function $X : [0, 1]^2 \rightarrow \mathbb{R}^d$ satisfying the stochastic wave equation (1.7) when a is the identity matrix.*

Below we provide a non trivial example of functions satisfying hypothesis of Theorem 2.9. This comes from the Jordan decomposition of real-valued functions of bounded variation on \mathbb{R} (see e.g. [20][Theorem 3.27, b.]).

Example 2.13. *Let g_1, \dots, g_d be real-valued functions of bounded variation on \mathbb{R} and let h_1, \dots, h_d be the functions defined on \mathbb{R} by $h_i(z) = |z|g_i(z)$ for all $z \in \mathbb{R}$ and i . It follows from Jordan decomposition that $g_i = \hat{g}_i - \check{g}_i$, where \hat{g}_i, \check{g}_i are two bounded nondecreasing functions on \mathbb{R} . It also holds that $h_i = \hat{h}_i - \check{h}_i$, where \hat{h}_i, \check{h}_i are two nondecreasing functions of linear growth on \mathbb{R} . This follows from the fact that $z \mapsto |z|$ is the difference of two non-decreasing functions and $z \mapsto |z|(\hat{g}_i(z) - \check{g}_i(0))$ (resp. $z \mapsto |z|(\check{g}_i(z) - \hat{g}_i(0))$) is non-decreasing on \mathbb{R} . Then*

1. the function $b := (b_1, \dots, b_d) : [0, 1]^2 \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ defined by

$$b_i(s, t, x_1, \dots, x_d) = g_i\left(\kappa_i(s, t) + \sum_{\ell=1}^d x_\ell\right), \quad \text{for all } (s, t, x_1, \dots, x_d) \text{ and } i$$

satisfies Hypothesis 2.7 for any Borel measurable functions $\kappa_1, \dots, \kappa_d : [0, 1]^2 \rightarrow \mathbb{R}$,

2. the function $\tilde{b} := (\tilde{b}_1, \dots, \tilde{b}_d) : [0, 1]^2 \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ defined by

$$\tilde{b}_i(s, t, x_1, \dots, x_d) = h_i\left(\zeta_i(s, t) + \sum_{\ell=1}^d x_\ell\right), \quad \text{for all } (s, t, x_1, \dots, x_d) \text{ and } i$$

satisfies Hypothesis 2.7 for any bounded Borel measurable functions $\zeta_1, \dots, \zeta_d : [0, 1]^2 \rightarrow \mathbb{R}$.

2.3. Proof of Theorem 2.10. In this subsection, we prove Theorem 2.10. As already pointed out in the introduction, this is equivalent to showing that for almost all Brownian sheet path, the unique continuous solution u to (1.4) is zero. More precisely, Theorem 2.10 is equivalent to:

Theorem 2.14. *Let $W := (W_{s,t}, (s, t) \in [0, 1]^2)$ be a d -dimensional Brownian sheet defined on a filtered probability space $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$, where $\mathbb{F} = \{\mathcal{F}_{s,t}\}_{s,t \in [0,1]}$. Let $b : [0, 1]^2 \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ be a Borel measurable function such that for every $i \in \{1, \dots, d\}$, $b_i(s, t, \cdot) = \hat{b}_i(s, t, \cdot) - \check{b}_i(s, t, \cdot)$, where \hat{b}_i, \check{b}_i are bounded and componentwise nondecreasing in x for all (s, t) . Then there exists $\Omega_1 \subset \Omega$ with $\mathbb{P}(\Omega_1) = 1$ such that for any $\omega \in \Omega_1$, $u = 0$ is the unique continuous solution of the integral equation*

$$(2.6) \quad u(s, t) = \int_0^t \int_0^s \{b(s_1, t_1, W_{s_1, t_1}(\omega) + u(s_1, t_1)) - b(s_1, t_1, W_{s_1, t_1}(\omega))\} ds_1 dt_1, \quad \forall (s, t) \in [0, 1]^2.$$

The proof of Theorem 2.14 relies on the following Gronwall type result.

Lemma 2.15. *Suppose conditions of Theorem 2.14 are valid. Then there exists $\Omega_1 \subset \Omega$ with $\mathbb{P}(\Omega_1) = 1$ and a positive random constant C_2 such that for any $\omega \in \Omega_1$, any sufficiently large positive integer n , any $(k, k') \in \{0, 1, 2, \dots, 2^n\}^2$, any $\beta(n) \in [2^{-4^{3n/4}}, 2^{-4^{2n/3}}]$, and any solution u of the integral equation*

$$(2.7) \quad \begin{aligned} & u(s, t) - u(s, 0) - u(0, t) + u(0, 0) \\ &= \int_0^t \int_0^s \left\{ b(s_1, t_1, W_{s_1, t_1}(\omega) + u(s_1, t_1)) - b(s_1, t_1, W_{s_1, t_1}(\omega)) \right\} ds_1 dt_1, \quad \forall (s, t) \in [0, 1]^2 \end{aligned}$$

satisfying

$$(2.8) \quad \max_{1 \leq i \leq d} \max\{|u_i|(s, 0), |u_i|(0, t)\} \leq \beta(n), \quad \forall (s, t) \in [0, 1]^2,$$

we have

$$(2.9) \quad \max_{1 \leq i \leq d} \max\{\bar{u}_{n,i}(k, k'), \underline{u}_{n,i}(k, k')\} \leq 3^{k+k'-1} \left(1 + 3C_2(\omega)\sqrt{dn}2^{-n}\right)^{k+k'} \beta(n),$$

where $\bar{u}_n = (\bar{u}_{n,1}, \dots, \bar{u}_{n,d})$, $\underline{u}_n = (\underline{u}_{n,1}, \dots, \underline{u}_{n,d})$ and for every $i \in \{1, \dots, d\}$,

$$\bar{u}_{n,i}(k, k') = \sup_{(s,t) \in I_{n,k-1} \times I_{n,k'-1}} \max\{0, u_i(s, t)\} \text{ and } \underline{u}_{n,i}(k, k') = \sup_{(s,t) \in I_{n,k-1} \times I_{n,k'-1}} \max\{0, -u_i(s, t)\}.$$

Proof. Suppose without loss of generality that $\|\hat{b}_i\|_\infty \leq 1$ and $\|\check{b}_i\|_\infty \leq 1$ for every $i \in \{1, \dots, d\}$. By Lemma 2.6, there exists a subset $\Omega_2 \subset \Omega$ with $\mathbb{P}(\Omega_2) = 1$ such that for all $\omega \in \Omega_2$,

$$(2.10) \quad \max_{1 \leq i \leq d} \left| \hat{\varrho}_{nkk'}^{(i)}(0, x)(\omega) \right| \leq C_2(\omega)\sqrt{n}2^{-n} (|x| + \beta(n)) \text{ on } \Omega_2$$

and

$$(2.11) \quad \max_{1 \leq i \leq d} \left| \check{\varrho}_{nkk'}^{(i)}(0, x)(\omega) \right| \leq C_2(\omega)\sqrt{n}2^{-n} (|x| + \beta(n)) \text{ on } \Omega_2$$

for all integers n, k, k' with $n \geq 1, 0 \leq k, k' \leq 2^n - 1$ and all $x \in [-1, 1]^d$, where

$$\hat{\varrho}_{nkk'}^{(i)}(0, x)(\omega) = \int_{k2^{-n}}^{(k+1)2^{-n}} \int_{k'2^{-n}}^{(k'+1)2^{-n}} \left\{ \hat{b}_i(s_1, t_1, W_{s_1, t_1}(\omega) + x) - \hat{b}_i(s_1, t_1, W_{s_1, t_1}(\omega)) \right\} ds_1 dt_1$$

and

$$\check{\varrho}_{nkk'}^{(i)}(0, x)(\omega) = \int_{k2^{-n}}^{(k+1)2^{-n}} \int_{k'2^{-n}}^{(k'+1)2^{-n}} \left\{ \check{b}_i(s_1, t_1, W_{s_1, t_1}(\omega) + x) - \check{b}_i(s_1, t_1, W_{s_1, t_1}(\omega)) \right\} ds_1 dt_1.$$

For any $\omega \in \Omega_2$, we choose $n \in \mathbb{N}^*$ such that $C_2(\omega)\sqrt{dn}2^{-n} \leq 1/6$ and split the set $[0, 1] \times [0, 1]$ onto 4^n squares $I_{nk} \times I_{nk'}$. We set $u = (u_1, \dots, u_d)$, with $u^+ = (u_1^+, \dots, u_d^+)$, $u^- = (u_1^-, \dots, u_d^-)$, $u_i^+ = \max\{0, u_i\}$ and $u_i^- = \max\{0, -u_i\}$ for every $i \in \{1, \dots, d\}$. Since $\hat{b}_i(s_1, t_1, \cdot)$ and $\check{b}_i(s_1, t_1, \cdot)$ are nondecreasing, we deduce from

(2.7) that for all $i \in \{1, \dots, d\}$ and all $(s, t) \in I_{nk} \times I_{nk'}$,

$$\begin{aligned}
& u_i(s, t) - u_i(s, k'2^{-n}) - u_i(k2^{-n}, t) + u_i(2^{-n}(k, k')) \\
&= \int_{k2^{-n}}^s \int_{k'2^{-n}}^t \left\{ \hat{b}_i(s_1, t_1, W_{s_1, t_1}(\omega) + u(s_1, t_1)) - \hat{b}_i(s_1, t_1, W_{s_1, t_1}(\omega)) \right\} ds_1 dt_1 \\
&\quad - \int_{k2^{-n}}^s \int_{k'2^{-n}}^t \left\{ \check{b}_i(s_1, t_1, W_{s_1, t_1}(\omega) + u(s_1, t_1)) - \check{b}_i(s_1, t_1, W_{s_1, t_1}(\omega)) \right\} ds_1 dt_1 \\
&\leq \int_{k2^{-n}}^s \int_{k'2^{-n}}^t \left\{ \hat{b}_i(s_1, t_1, W_{s_1, t_1}(\omega) + u^+(s_1, t_1)) - \hat{b}_i(s_1, t_1, W_{s_1, t_1}(\omega)) \right\} ds_1 dt_1 \\
&\quad - \int_{k2^{-n}}^s \int_{k'2^{-n}}^t \left\{ \check{b}_i(s_1, t_1, W_{s_1, t_1}(\omega) - u^-(s_1, t_1)) - \check{b}_i(s_1, t_1, W_{s_1, t_1}(\omega)) \right\} ds_1 dt_1.
\end{aligned}$$

Then, using the fact that $\max\{0, x + y\} \leq \max\{0, x\} + \max\{0, y\}$, we have

$$\begin{aligned}
u_i^+(s, t) &\leq \max\{0, u_i(s, k'2^{-n}) + u_i(k2^{-n}, t) - u_i(2^{-n}(k, k'))\} \\
&\quad + \int_{k2^{-n}}^s \int_{k'2^{-n}}^t \left\{ \hat{b}_i(s_1, t_1, W_{s_1, t_1}(\omega) + u^+(s_1, t_1)) - \hat{b}_i(s_1, t_1, W_{s_1, t_1}(\omega)) \right\} ds_1 dt_1 \\
&\quad - \int_{k2^{-n}}^s \int_{k'2^{-n}}^t \left\{ \check{b}_i(s_1, t_1, W_{s_1, t_1}(\omega) - u^-(s_1, t_1)) - \check{b}_i(s_1, t_1, W_{s_1, t_1}(\omega)) \right\} ds_1 dt_1.
\end{aligned}$$

As a consequence,

$$\begin{aligned}
(2.12) \quad u_i^+(s, t) &\leq u_i^+(s, k'2^{-n}) + u_i^+(k2^{-n}, t) + u_i^-(2^{-n}(k, k')) + \hat{\varrho}_{nkk'}^{(i)}(0, \bar{u}_n(k+1, k'+1))(\omega) \\
&\quad - \check{\varrho}_{nkk'}^{(i)}(0, -\underline{u}_n(k+1, k'+1))(\omega)
\end{aligned}$$

for all $(s, t) \in I_{nk} \times I_{nk'}$. Similarly, we can show that

$$\begin{aligned}
(2.13) \quad u_i^-(s, t) &\leq u_i^-(s, k'2^{-n}) + u_i^-(k2^{-n}, t) + u_i^+(2^{-n}(k, k')) - \hat{\varrho}_{nkk'}^{(i)}(0, -\underline{u}_n(k+1, k'+1))(\omega) \\
&\quad + \check{\varrho}_{nkk'}^{(i)}(0, \bar{u}_n(k+1, k'+1))(\omega)
\end{aligned}$$

for all $(s, t) \in I_{nk} \times I_{nk'}$. For any $k, k' \in \{1, 2, \dots, 2^n\}$ and $i \in \{1, \dots, d\}$, we define $\hat{u}_{n,i}(k, k') = \max\{\bar{u}_{n,i}(k, k'), \underline{u}_{n,i}(k, k')\}$. We deduce from Inequalities (2.10)-(2.13) that

$$\begin{aligned}
\bar{u}_{n,i}(k+1, k'+1) &\leq \max_{1 \leq j \leq d} \hat{u}_{n,j}(k, k'+1) + \max_{1 \leq j \leq d} \hat{u}_{n,j}(k+1, k') + \max_{1 \leq j \leq d} \hat{u}_{n,j}(k, k') \\
&\quad + 2C_2(\omega)\sqrt{n}2^{-n} \left(\sqrt{d} \max_{1 \leq j \leq d} \hat{u}_{n,j}(k+1, k'+1) + \beta(n) \right)
\end{aligned}$$

and

$$\begin{aligned}
\underline{u}_{n,i}(k+1, k'+1) &\leq \max_{1 \leq j \leq d} \hat{u}_{n,j}(k, k'+1) + \max_{1 \leq j \leq d} \hat{u}_{n,j}(k+1, k') + \max_{1 \leq j \leq d} \hat{u}_{n,j}(k, k') \\
&\quad + 2C_2(\omega)\sqrt{n}2^{-n} \left(\sqrt{d} \max_{1 \leq j \leq d} \hat{u}_{n,j}(k+1, k'+1) + \beta(n) \right).
\end{aligned}$$

Then

$$\begin{aligned}
\max_{1 \leq i \leq d} \hat{u}_{n,i}(k+1, k'+1) &\leq \max_{1 \leq i \leq d} \hat{u}_{n,i}(k, k'+1) + \max_{1 \leq i \leq d} \hat{u}_{n,i}(k+1, k') + \max_{1 \leq i \leq d} \hat{u}_{n,i}(k, k') \\
&\quad + 2C_2(\omega)\sqrt{n}2^{-n} \left(\sqrt{d} \max_{1 \leq i \leq d} \hat{u}_{n,i}(k+1, k'+1) + \beta(n) \right).
\end{aligned}$$

Since $C_2(\omega)\sqrt{dn}2^{-n} \leq 1/6$, we have $(1 - 2C_2(\omega)\sqrt{dn}2^{-n})^{-1} \leq (1 + 3C_2(\omega)\sqrt{dn}2^{-n})$ and the above inequality implies

$$\begin{aligned} \max_{1 \leq i \leq d} \widehat{u}_{n,i}(k+1, k'+1) &\leq (1 + 3C_2(\omega)\sqrt{dn}2^{-n}) \left(\max_{1 \leq i \leq d} \widehat{u}_{n,i}(k, k'+1) + \max_{1 \leq i \leq d} \widehat{u}_{n,i}(k+1, k') + \max_{1 \leq i \leq d} \widehat{u}_{n,i}(k, k') \right) \\ &\quad + 2C_2(\omega)\sqrt{n}2^{-n}\beta(n). \end{aligned}$$

The desired result then follows by induction on k and k' as in the proof of Lemma 3.9 in [5]. \square

We now turn to the proof of Theorem 2.14.

Proof of Theorem 2.14. Choose Ω_1 , ω , n and $\beta(n)$ as in Lemma 2.15. Let u be a solution of (2.6). We have $\max\{|u|(s, 0), |u|(0, t)\} = 0 \leq \beta(n)$ for all $(s, t) \in [0, 1]^2$. Moreover, we deduce from (2.9) that

$$(2.14) \quad \sup_{k, k' \in \{0, 1, 2, \dots, 2^n\}} \max_{1 \leq i \leq d} \max\{\bar{u}_{n,i}(k, k'), \underline{u}_{n,i}(k, k')\} \leq 2^{2^{n+2}} \beta(n)$$

for all n satisfying $C_2(\omega)\sqrt{dn}2^{-n} \leq 1/9$. Since the right hand side of (2.14) converges to 0 as n goes to ∞ , then, for all (s, t) , we have $u(s, t) = 0$ on Ω_1 . \square

3. MALLIAVIN REGULARITY

In this section we study the Malliavin regularity of the strong solution to (1.3).

3.1. Basic facts on Malliavin calculus and compactness criterion on the plane. We first recall some basic facts on Malliavin calculus for Wiener functionals on the plane which can be found in [34, Section 2] (see also [35, Section 1]). Let $(\Omega, \mathcal{F}, \mathbb{P})$ be the canonical space associated to the d -dimensional Brownian sheet, that is Ω is the space of all continuous functions $\omega : \Gamma \rightarrow \mathbb{R}^d$ which vanish on the axes, \mathbb{P} is the Wiener measure and \mathcal{F} is the completion of the Borel σ -algebra of Ω with respect to \mathbb{P} . Let $(\mathcal{F}_{s,t}, (s, t) \in \Gamma)$ denote the nondecreasing family of σ -algebras where $\mathcal{F}_{s,t}$ is generated by the functions $(s_1, t_1) \mapsto \omega(s_1 \wedge s, t_1 \wedge t)$, $(s_1, t_1) \in \Gamma$, $\omega \in \Omega$ and the null sets of \mathcal{F} . Consider the following subset H of Ω :

$$H = \left\{ \omega \in \Omega : \text{there exists } \dot{\omega} \in L^2(\Gamma, \mathbb{R}^d) \text{ such that } \omega(s, t) = \int_0^s \int_0^t \dot{\omega}(s_1, t_1) dt_1 ds_1, \text{ for any } (s, t) \in \Gamma \right\}.$$

Endowed with the inner product

$$\langle \omega_1, \omega_2 \rangle_H = \sum_{i=1}^d \int_{\Gamma} \dot{\omega}_1^{(i)}(s_1, t_1) \dot{\omega}_2^{(i)}(s_1, t_1) ds_1 dt_1,$$

the set H is a Hilbert space. We call Wiener functional any measurable function defined on the Wiener space $(\Omega, \mathcal{F}, \mathbb{P})$. A Wiener functional $F : \Omega \rightarrow \mathbb{R}$ is said to be smooth if there exists some integer $n \geq 1$ and an infinitely differentiable function f on \mathbb{R}^n such that

- (i) f and all its derivatives have at most polynomial growth order,
- (ii) $F(\omega) = f(\omega(s_1, t_1), \dots, \omega(s_n, t_n))$ for some $(s_1, t_1), \dots, (s_n, t_n) \in \Gamma$.

Every smooth functional F is Fréchet-differentiable and the Fréchet-derivative of F along any vector $h \in H$ is given by

$$DF(h) = \sum_{j=1}^d \sum_{i=1}^n \frac{\partial f}{\partial x_i^{(j)}}(\omega(s_1, t_1), \dots, \omega(s_n, t_n)) h^{(j)}(s_i, t_i) = \sum_{j=1}^d \int_{\Gamma} \xi_j(q, r) \dot{h}^{(j)}(q, r) dq dr,$$

where

$$\xi_j(q, r) = \sum_{i=1}^n \frac{\partial f}{\partial x_i^{(j)}}(\omega(s_1, t_1), \dots, \omega(s_n, t_n)) \mathbf{1}_{[0, s_i] \times [0, t_i]}(q, r).$$

Let $\mathbb{D}_{2,1}$ denote the closed hull of the family of smooth functionals with respect to the norm

$$\|F\|_{2,1}^2 = \|F\|_{L^2(\Omega)}^2 + \|DF\|_{L^2(\Omega;H)}^2.$$

Now we present a useful characterization of relatively compact subsets in the space $L^2(\Omega, \mathbb{R}^d)$. Let us recall the following compactness criterion provided in [11, Theorem 1].

Theorem 3.1. *Let A be a self-adjoint compact operator on H . Then, for any $c > 0$, the set*

$$\mathcal{G} = \{G \in \mathbb{D}_{2,1} : \|G\|_{L^2(\Omega)} + \|A^{-1}DG\|_{L^2(\Omega;H)} \leq c\}$$

is relatively compact in $L^2(\Omega, \mathbb{R}^d)$.

In order to apply the above result, we consider the fractional Sobolev space:

$$\mathbb{G}_\beta^{2,p}(U; \mathbb{R}) := \left\{ g \in L^2(U, \mathbb{R}) : \int_U \int_U \frac{|g(u) - g(u')|^2}{|u - u'|^{p+2\beta}} du du' < \infty \right\},$$

where U is a domain of \mathbb{R}^p , $p \geq 1$ and the norm is given by

$$\|g\|_{\mathbb{G}_\beta^{2,p}(U; \mathbb{R})} := \|g\|_{L^2(U; \mathbb{R})} + \left(\int_U \int_U \frac{|g(u) - g(u')|^2}{|u - u'|^{p+2\beta}} du du' \right)^{1/2}.$$

We need the next compact embedding result from [38, Lemma 10] (see also [17, Theorem 7.1]).

Lemma 3.2. *Let $p \geq 1$, $U \subset \mathbb{R}^p$ be a Lipschitz bounded open set and \mathcal{J} be a bounded subset of $L^2(U; \mathbb{R})$. Suppose that*

$$\sup_{g \in \mathcal{J}} \int_U \int_U \frac{|g(u) - g(u')|^2}{|u - u'|^{p+2\beta}} du du' < \infty$$

for some $\beta \in (0, 1)$. Then \mathcal{J} is relatively compact in $L^2(U; \mathbb{R})$.

As a consequence of Theorem 3.1 and Lemma 3.2, we have the following compactness criterion for subsets in the space $L^2(\Omega, \mathbb{R}^d)$.

Corollary 3.3. *Denote by \mathcal{F}_∞^W the σ -algebra generated by the d -dimensional Brownian sheet $W = (W^{(1)}, \dots, W^{(d)})$. Let $(X^{(n)}, n \in \mathbb{N})$ be a sequence of $(\mathcal{F}_\infty^W, \mathcal{B}(\mathbb{R}^d))$ -measurable random variables and let $D_{s,t}$ be the Malliavin derivative associated with the random vector $W_{s,t} = (W_{s,t}^{(1)}, \dots, W_{s,t}^{(d)})$. Suppose*

$$(3.1) \quad \sup_{n \in \mathbb{N}} \|X^{(n)}\|_{L^2(\Omega, \mathbb{R}^d)} < \infty \quad \text{and} \quad \sup_{n \in \mathbb{N}} \|D_{\cdot, \cdot} X^{(n)}\|_{L^2(\Omega \times [0,1]^2, \mathbb{R}^{d \times d})} < \infty$$

as well as

$$(3.2) \quad \sup_{n \in \mathbb{N}} \int_0^1 \int_0^1 \int_0^1 \int_0^1 \frac{\mathbb{E} [\|D_{s,t} X^{(n)} - D_{s',t'} X^{(n)}\|^2]}{(|s - s'| + |t - t'|)^{2+2\beta}} ds ds' dt dt' < \infty$$

for some $\beta \in (0, 1)$. Then $(X^{(n)}, n \in \mathbb{N})$ is relatively compact in $L^2(\Omega, \mathbb{R}^d)$.

Proof. The proof is inspired from [22, Section 5]. We consider the symmetric form \mathcal{L} on $L^2((0, 1)^2, \mathbb{R}^d)$ defined as

$$\mathcal{L}(f, g) = \int_0^1 \int_0^1 f(s, t) \cdot g(s, t) ds dt + \int_0^1 \int_0^1 \int_0^1 \int_0^1 \frac{(f(s, t) - f(s', t')) \cdot (g(s, t) - g(s', t'))}{(|s - s'| + |t - t'|)^{2+2\beta}} ds ds' dt dt'$$

for functions f, g in the dense domain $\mathcal{D}(\mathcal{L}) \subset L^2((0, 1)^2, \mathbb{R}^d)$ and a fixed $\beta \in (0, 1)$, where

$$\mathcal{D}(\mathcal{L}) = \left\{ g : \|g\|_{L^2((0,1)^2, \mathbb{R}^d)}^2 + \int_0^1 \int_0^1 \int_0^1 \int_0^1 \frac{|g(s, t) - g(s', t')|^2}{(|s - s'| + |t - t'|)^{2+2\beta}} ds ds' dt dt' < \infty \right\}.$$

Then \mathcal{L} is a positive symmetric closed form and, by Kato's first representation theorem (see e.g. [24]), one can find a positive self-adjoint operator $T_{\mathcal{L}}$ such that

$$\mathcal{L}(f, g) = \langle f, T_{\mathcal{L}} g \rangle_{L^2((0,1)^2, \mathbb{R}^d)}$$

for all $g \in \mathcal{D}(T_{\mathcal{L}})$ and $f \in \mathcal{D}(\mathcal{L})$. Further, one may observe that the form \mathcal{L} is bounded from below by a positive number. Indeed,

$$(3.3) \quad \mathcal{L}(g, g) \geq \|g\|_{L^2((0,1)^2, \mathbb{R}^d)}$$

for all $g \in \mathcal{D}(\mathcal{L})$. Hence, we have that $\mathcal{D}(\mathcal{L}) = \mathcal{D}(T_{\mathcal{L}}^{1/2})$.

Now, define the operator A as $A = T_{\mathcal{L}}^{1/2}$. It follows from Lemma 1 in [28, Section 1] (see also [22, Lemma 9]) and Lemma 3.2 applied to $p = 2$, $U = (0, 1)^2$ that A has a discrete spectrum and a compact inverse A^{-1} . Then, using (3.1)-(3.3), the operator A and the sequence $(X^{(n)}, n \in \mathbb{N})$ satisfy the assumptions of Theorem 3.1. \square

3.2. Malliavin differentiability for bounded drifts. In this subsection, we assume in addition the drift b is bounded. The main result of this subsection which significantly generalised those in [35] when the diffusion is constant is the following

Theorem 3.4. *The strong solution $\{X_{s,t}^{\xi}, s, t \in [0, 1]\}$ of the SDE (1.3) is Malliavin differentiable.*

The proof of this theorem is done in two steps:

Step 1: We use standard approximation procedure to approximate the drift coefficient $b = \hat{b} - \check{b}$ by a sequence of functions

$$b_n := \hat{b}_n - \check{b}_n, n \geq 1$$

such that $\hat{b}_n = (\hat{b}_{1,n}, \dots, \hat{b}_{d,n})$, $\check{b}_n = (\check{b}_{1,n}, \dots, \check{b}_{d,n})$, $(\hat{b}_{j,n})_{n \geq 1}$ and $(\check{b}_{j,n})_{n \geq 1}$ are smooth and componentwise non-decreasing functions with $\sup_n \|\hat{b}_{j,n}\|_{\infty} \leq \|\hat{b}_j\|_{\infty} < \infty$ and $\sup_n \|\check{b}_{j,n}\|_{\infty} \leq \|\check{b}_j\|_{\infty} < \infty$. In addition, $(\hat{b}_n)_{n \geq 1}$ (respectively, $(\check{b}_n)_{n \geq 1}$) converges to \hat{b} (respectively, \check{b}) in $(s, t, x) \in [0, 1]^2 \times \mathbb{R}^d$ $ds \times dt \times dx$ -a.e. We know that for such smooth drift coefficients, the corresponding SDEs have a unique strong solution denoted by $X^{\xi, n}$. We then show that for each $s, t \in [0, 1]$, the sequence $(X_{s,t}^{\xi, n})_{n \geq 1}$ is relatively compact in $L^2(\Omega, \mathbb{P}; \mathbb{R}^d)$.

Step 2: We show that the sequence of solutions $(X^{\xi, n})_{n \geq 1}$ converges strongly in $L^2(\Omega, \mathbb{P}; \mathbb{R}^d)$.

From **Step 1** and **Step 2**, the result will follow by application of a compactness criteria [32, Lemma 1.2.3].

The next result corresponds to a $L^2(\Omega)$ compactness criteria. It is analogous to the result derived in [30] for the case of Brownian motion.

Theorem 3.5. *For every $(s, t) \in \Gamma_0$, the sequence $(X_{s,t}^{\xi, n})_{n \geq 1}$ is relatively compact in $L^2(\Omega, \mathbb{P}; \mathbb{R}^d)$.*

The proof of the above theorem uses Corollary 3.3, which in our case is reduced to proving:

Lemma 3.6. *There exists $C_1 > 0$ such that for every $(s, t) \in \Gamma_0$, the sequence $(X_{s,t}^{\xi, n})_{n \geq 1}$ satisfies*

$$(3.4) \quad \sup_{n \geq 1} \|X_{s,t}^{\xi, n}\|_{L^2(\Omega, \mathbb{P}; \mathbb{R}^d)}^2 \leq C_1$$

and

$$(3.5) \quad \sup_{n \geq 1} \sup_{\substack{0 \leq r \leq s \\ 0 \leq u \leq t}} \mathbb{E} \left[\|D_{r,u} X_{s,t}^{\xi, n}\|^2 \right] \leq C_1.$$

Moreover for all $0 \leq r', r \leq s$, $0 \leq u', u \leq t$,

$$(3.6) \quad \mathbb{E} \left[\|D_{r,u} X_{s,t}^{\xi, n} - D_{r',u'} X_{s,t}^{\xi, n}\|^2 \right] \leq C_1(|r - r'| + |u - u'|),$$

where $\|\cdot\|$ denotes the max norm.

Remark 3.7. *If (3.6) is satisfied, then for any $\beta \in (0, 1/2)$, (3.2) holds. Indeed, since, for any s, s', t, t' , one has*

$$|s - s'|^{1/2} |t - t'|^{1/2} \leq \frac{1}{2}(|s - s'| + |t - t'|),$$

then for any $n \in \mathbb{N}$,

$$\begin{aligned} & \int_0^1 \int_0^1 \int_0^1 \int_0^1 \frac{\mathbb{E} [\|D_{s,t} X^{(n)} - D_{s',t'} X^{(n)}\|^2]}{(|s-s'|+|t-t'|)^{2+2\beta}} ds ds' dt dt' \\ & \leq C_1 \int_0^1 \int_0^1 \int_0^1 \int_0^1 \frac{ds ds' dt dt'}{(|s-s'|+|t-t'|)^{1+2\beta}} \leq \frac{C_1}{2} \left(\int_0^1 \int_0^1 \frac{ds ds'}{|s-s'|^{1/2+\beta}} \right) \left(\int_0^1 \int_0^1 \frac{dt dt'}{|t-t'|^{1/2+\beta}} \right) < \infty. \end{aligned}$$

To prove the result above, we need some preliminary estimates.

Lemma 3.8. *There exists a non-decreasing function $\tilde{C}_1 : \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that for any $0 < r < s \leq 1$, any $0 < u < t \leq 1$, any $k \in \mathbb{R}_+$ and any $i, j \in \{1, \dots, d\}$,*

$$(3.7) \quad \mathbb{E} \left[\exp \left(\frac{k}{\delta(r,s)\delta(u,t)} \int_r^s \int_u^t \partial_i \hat{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) dt_1 ds_1 \right) \right] \leq \tilde{C}_1(k, \|\hat{b}_{j,n}\|_\infty)$$

and

$$(3.8) \quad \mathbb{E} \left[\exp \left(\frac{k}{\delta(r,s)\delta(u,t)} \int_r^s \int_u^t \partial_i \check{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) dt_1 ds_1 \right) \right] \leq \tilde{C}_1(k, \|\check{b}_{j,n}\|_\infty),$$

where $\delta(r,s) = \sqrt{s-r}$ and $\delta(u,t) = \sqrt{t-u}$.

Proof. We only prove (3.7) and the proof of (3.8) follows analogously. Using integration with respect to local time formula (see e.g. [6, Corollary 2.3]), we have

$$(3.9) \quad \begin{aligned} & \int_r^s \int_u^t \partial_i \hat{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) dt_1 ds_1 \\ & = - \int_r^s \int_u^t b_{j,n}(s_1, t_1, W_{s_1, t_1}) \frac{d_{t_1} W_{s_1, t_1}^{(i)}}{s_1} ds_1 + \int_r^s \int_{1-t}^{1-u} b_{j,n}(s_1, 1-t_1, W_{s_1, 1-t_1}) \frac{d_{t_1} B_{s_1, t_1}^{(i)}}{s_1} ds_1 \\ & \quad - \int_r^s \int_{1-t}^{1-u} b_{j,n}(s_1, 1-t_1, W_{s_1, 1-t_1}) \frac{W_{s_1, 1-t_1}^{(i)}}{s_1(1-t_1)} dt_1 ds_1, \end{aligned}$$

where $(B_{s,t}^{(i)}, (s,t) \in [0,1]^2)$ is the Brownian sheet given by the following representation provided by Dalang and Walsh [13]

$$W_{s,1-t}^{(i)} = W_{s,1}^{(i)} + B_{s,t}^{(i)} - \int_0^t \frac{W_{s,1-u}^{(i)}}{1-u} du.$$

Since the function $x \mapsto e^{3x}$ is convex,

$$(3.10) \quad \begin{aligned} & \mathbb{E} \left[\exp \left(\frac{k}{\delta(r,s)\delta(u,t)} \left| \int_r^s \int_u^t \partial_i \hat{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) dt_1 ds_1 \right| \right) \right] \\ & \leq \frac{1}{3} \left\{ \mathbb{E} \left[\exp \left(\frac{3k}{\delta(r,s)\delta(u,t)} \left| \int_r^s \int_u^t \hat{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) \frac{d_{t_1} W_{s_1, t_1}^{(i)}}{s_1} ds_1 \right| \right) \right] \right. \\ & \quad + \mathbb{E} \left[\exp \left(\frac{3k}{\delta(r,s)\delta(u,t)} \left| \int_r^s \int_{1-t}^{1-u} \hat{b}_{j,n}(s_1, t_1, W_{s_1, 1-t_1}) \frac{d_{t_1} B_{s_1, t_1}^{(i)}}{s_1} ds_1 \right| \right) \right] \\ & \quad \left. + \mathbb{E} \left[\exp \left(\frac{3k}{\delta(r,s)\delta(u,t)} \left| \int_r^s \int_{1-t}^{1-u} \hat{b}_{j,n}(s_1, t_1, W_{s_1, 1-t_1}) \frac{W_{s_1, 1-t_1}^{(i)}}{s_1(1-t_1)} dt_1 ds_1 \right| \right) \right] \right\} = \frac{1}{3} (I_1 + I_2 + I_3). \end{aligned}$$

By Jensen inequality

$$\begin{aligned}
I_1 &= \mathbb{E} \left[\exp \left(\frac{3k}{\delta(r,s)\delta(u,t)} \left| \int_r^s \int_u^t \hat{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) \frac{d_{t_1} W_{s_1, t_1}^{(i)}}{s_1} ds_1 \right| \right) \right] \\
&\leq \int_r^s \mathbb{E} \left[\exp \left(\frac{6k(\sqrt{s} - \sqrt{r})}{\delta(r,s)\delta(u,t)} \left| \int_u^t \hat{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) \frac{d_{t_1} W_{s_1, t_1}^{(i)}}{\sqrt{s_1}} \right| \right) \right] \frac{ds_1}{2\sqrt{s_1}(\sqrt{s} - \sqrt{r})} \\
&\leq \int_r^s \mathbb{E} \left[\exp \left(\frac{6k}{\delta(u,t)} \left| \int_u^t \hat{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) \frac{d_{t_1} W_{s_1, t_1}^{(i)}}{\sqrt{s_1}} \right| \right) \right] \frac{ds_1}{2\sqrt{s_1}(\sqrt{s} - \sqrt{r})}.
\end{aligned}$$

Since for every $s_1 \in [r, s]$,

$$(Y_{s_1, t_1} := \int_u^{t_1} \hat{b}_{j,n}(s_1, t_2, W_{s_1, t_2}) \frac{d_{t_2} W_{s_1, t_2}^{(i)}}{\sqrt{s_1}}, u \leq t_1 \leq t)$$

is a square integrable martingale, it follows from the Barlow-Yor inequality (see e.g. [3, Proposition 4.2] and [8, Theorem A in Appendix]) that there exists a positive constant c_1 such that for any positive integer m and any $s_1 \in [r, s]$,

$$\begin{aligned}
\mathbb{E}[|Y_{s_1, t}|^m] &\leq \mathbb{E} \left[\sup_{u \leq t_1 \leq t} |Y_{s_1, t_1}|^m \right] \leq c_1^m \sqrt{m}^m \mathbb{E}[\langle Y_{s_1, \cdot} \rangle_t^{m/2}] = c_1^m \sqrt{m}^m \mathbb{E} \left[\left(\int_u^t \hat{b}_{j,n}^2(s_1, t_1, W_{s_1, t_1}) dt_1 \right)^{m/2} \right] \\
&\leq c_1^m \sqrt{m}^m \delta(u, t)^m \|\hat{b}_{j,n}\|_\infty^m.
\end{aligned}$$

Hence, using the following exponential expansion formula,

$$\exp \left(\frac{6k}{\delta(u, t)} |Y_{s_1, t}| \right) = \sum_{m=0}^{\infty} \frac{6^m k^m |Y_{s_1, t}|^m}{\delta(u, t)^m m!},$$

we obtain

$$\begin{aligned}
I_1 &\leq \int_r^s \mathbb{E} \left[\exp \left(\frac{6k}{\delta(u, t)} \left| \int_u^t \hat{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) \frac{d_{t_1} W_{s_1, t_1}}{\sqrt{s_1}} \right| \right) \right] \frac{ds_1}{2\sqrt{s_1}(\sqrt{s} - \sqrt{r})} \\
&= \int_r^s \mathbb{E} \left[\exp \left(\frac{6k}{\delta(u, t)} |Y_{s_1, t}| \right) \right] \frac{ds_1}{2\sqrt{s_1}(\sqrt{s} - \sqrt{r})} = \int_r^s \sum_{m=0}^{\infty} \frac{6^m k^m \mathbb{E}[|Y_{s_1, t}|^m]}{\delta(u, t)^m m!} \frac{ds_1}{2\sqrt{s_1}(\sqrt{s} - \sqrt{r})} \\
&\leq \sum_{m=0}^{\infty} \frac{6^m k^m c_1^m \sqrt{m}^m \|\hat{b}_{j,n}\|_\infty^m}{m!} := \tilde{C}_{1,1}(k, \|\hat{b}_{j,n}\|_\infty),
\end{aligned}$$

which is finite by ratio test. Similarly, we also have

$$I_2 \leq \tilde{C}_{1,1}(k, \|\hat{b}_{j,n}\|_\infty).$$

To estimate I_3 we apply one more Jensen inequality and obtain

$$\begin{aligned}
I_3 &= \mathbb{E} \left[\exp \left(\frac{3k}{\delta(r, s)\delta(u, t)} \left| \int_r^s \int_{1-t}^{1-u} \hat{b}_{j,n}(s_1, t_1, W_{s_1, 1-t_1}) \frac{W_{s_1, 1-t_1}^{(i)}}{s_1(1-t_1)} dt_1 ds_1 \right| \right) \right] \\
&\leq \int_r^s \int_{1-t}^{1-u} \mathbb{E} \left[\exp \left(\frac{12k(\sqrt{s} - \sqrt{r})(\sqrt{1-u} - \sqrt{1-t})}{\delta(r, s)\delta(u, t)} \left| \hat{b}_{j,n}(s_1, t_1, W_{s_1, 1-t_1}) \right| \left| \frac{W_{s_1, 1-t_1}^{(i)}}{\sqrt{s_1}\sqrt{1-t_1}} \right| \right) \right] \\
&\quad \times \frac{dt_1}{2\sqrt{1-t_1}(\sqrt{1-u} - \sqrt{1-t})} \frac{ds_1}{2\sqrt{s_1}(\sqrt{s} - \sqrt{r})} \\
&\leq \int_r^s \int_{1-t}^{1-u} \mathbb{E} \left[\exp \left(12k \|\hat{b}_{j,n}\|_\infty \left| \frac{W_{s_1, 1-t_1}^{(i)}}{\sqrt{s_1}\sqrt{1-t_1}} \right| \right) \right] \frac{dt_1}{2\sqrt{1-t_1}(\sqrt{1-u} - \sqrt{1-t})} \frac{ds_1}{2\sqrt{s_1}(\sqrt{s} - \sqrt{r})} \\
&\leq 2 \exp \left(72k^2 \|\hat{b}_{j,n}\|_\infty^2 \right) := \tilde{C}_{1,2}(k, \|\hat{b}_{j,n}\|_\infty),
\end{aligned}$$

since $\left| \frac{W_{s_1, 1-t_1}^{(i)}}{\sqrt{s_1}\sqrt{1-t_1}} \right|$ is a mean-zero Gaussian random variable of variance 1.

The proof is completed by choosing $\tilde{C}_1 = 2\tilde{C}_{1,1} + \tilde{C}_{1,2}$. \square

Proof of Lemma 3.6. Without loss of generality, we suppose $\xi = 0$. Further we set $X_{s,t}^n := X_{s,t}^{0,n}$. We start with the proof of (3.5). Suppose for every $j \in \{1, \dots, d\}$, $\hat{b}_{j,n}$ (respectively, $\check{b}_{j,n}$) is componentwise nondecreasing and continuously differentiable with bounded derivatives. Then for any $(r, u) \in [0, 1]^2$ with $(0, 0) \prec (r, u) \prec (s, t)$,

$$(3.11) \quad D_{r,u} X_{s,t}^n = I_d + \int_r^s \int_u^t \nabla b_n(s_1, t_1, X_{s_1, t_1}^n) D_{r,u} X_{s_1, t_1}^n dt_1 ds_1,$$

where I_d is the identity matrix, $\nabla b_n = (\partial_i b_{j,n})_{1 \leq i, j \leq d}$, $\partial_i b_{j,n}$ being the partial derivative of $b_j(s, t, \cdot)$ with respect to x_i , and $D_{r,u} X_{s,t}^n = (D_{r,u}^i X_{s,t}^{n,j})_{1 \leq i, j \leq d}$. Since $\partial_i \hat{b}_{j,n}$ and $\partial_i \check{b}_{j,n}$ are nonnegative, we have

$$\|D_{r,u} X_{s,t}^n\| \leq 1 + \int_r^s \int_u^t \sum_{i,j=1}^d \left\{ \partial_i \hat{b}_{j,n}(s_1, t_1, X_{s_1, t_1}^n) + \partial_i \check{b}_{j,n}(s_1, t_1, X_{s_1, t_1}^n) \right\} \|D_{r,u} X_{s_1, t_1}^n\| dt_1 ds_1.$$

Therefore (see e.g. Theorem B.1),

$$(3.12) \quad \|D_{r,u} X_{s,t}^n\| \leq \exp \left(\int_r^s \int_u^t \sum_{i,j=1}^d \left\{ \partial_i \hat{b}_{j,n}(s_1, t_1, X_{s_1, t_1}^n) + \partial_i \check{b}_{j,n}(s_1, t_1, X_{s_1, t_1}^n) \right\} dt_1 ds_1 \right).$$

Squaring both sides of the inequality, taking the expectation and using the Girsanov theorem (see e.g. [12, Theorem 3.5] and [33, Proposition 1.6]) and the Cauchy-Schwarz inequality, we obtain

$$\begin{aligned}
&\mathbb{E}[\|D_{r,u} X_{s,t}^n\|^2] \\
&\leq \mathbb{E} \left[\exp \left(2 \int_r^s \int_u^t \sum_{i,j=1}^d \left\{ \partial_i \hat{b}_{j,n}(s_1, t_1, X_{s_1, t_1}^n) + \partial_i \check{b}_{j,n}(s_1, t_1, X_{s_1, t_1}^n) \right\} dt_1 ds_1 \right) \right] \\
&= \mathbb{E} \left[\mathcal{E} \left(\int_0^1 \int_0^1 b_n(s_1, t_1, W_{s_1, t_1}) \cdot dW_{s_1, t_1} \right) \exp \left(2 \int_r^s \int_u^t \sum_{i,j=1}^d \left\{ \partial_i \hat{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) + \partial_i \check{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) \right\} dt_1 ds_1 \right) \right] \\
&\leq C_0 \mathbb{E} \left[\exp \left(4 \int_r^s \int_u^t \sum_{i,j=1}^d \left\{ \partial_i \hat{b}_{j,n}(s_1, t_1, X_{s_1, t_1}^n) + \partial_i \check{b}_{j,n}(s_1, t_1, X_{s_1, t_1}^n) \right\} dt_1 ds_1 \right) \right]^{\frac{1}{2}} \\
&= C_0 \mathbb{E} \left[\prod_{i,j=1}^d \exp \left(4 \int_r^s \int_u^t \partial_i \hat{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) dt_1 ds_1 \right) \exp \left(4 \int_r^s \int_u^t \partial_i \check{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) dt_1 ds_1 \right) \right]^{\frac{1}{2}},
\end{aligned}$$

where

$$(3.13) \quad C_0 := \sup_{n \geq 1} \mathbb{E} \left[\mathcal{E} \left(\int_0^1 \int_0^1 b_n(s_1, t_1, W_{s_1, t_1}) \cdot dW_{s_1, t_1} \right)^2 \right]^{\frac{1}{2}}$$

is finite (see Lemma A.1 and Remark A.2 in the Appendix), with

$$\mathcal{E} \left(\int_0^1 \int_0^1 b_n(s_1, t_1, W_{s_1, t_1}) \cdot dW_{s_1, t_1} \right) = \exp \left(\int_0^1 \int_0^1 b_n(s_1, t_1, W_{s_1, t_1}) \cdot dW_{s_1, t_1} - \frac{1}{2} \int_0^1 \int_0^1 |b_n(s_1, t_1, W_{s_1, t_1})|^2 \cdot ds_1 dt_1 \right).$$

Hence, by Hölder inequality, we have

$$\begin{aligned} & \mathbb{E}[\|D_{r,u} X_{s,t}^n\|^2] \\ & \leq C_0 \prod_{i,j=1}^d \mathbb{E} \left[\exp \left(8d^2 \int_r^s \int_u^t \partial_i \hat{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) dt_1 ds_1 \right) \right]^{\frac{1}{4d^2}} \mathbb{E} \left[\exp \left(8d^2 \int_r^s \int_u^t \partial_i \check{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) dt_1 ds_1 \right) \right]^{\frac{1}{4d^2}}. \end{aligned}$$

Then, by Lemma 3.8, we obtain

$$\mathbb{E}[\|D_{r,u} X_{s,t}^n\|^2] \leq C_0 \times \tilde{C}_1 \left(8d^2, \max_{1 \leq j \leq d} \{ \|\hat{b}_{j,n}\|_\infty + \|\check{b}_{j,n}\|_\infty \} \right) \leq C_0 \times \tilde{C}_1 \left(8d^2, \|\hat{b}\|_\infty + \|\check{b}\|_\infty \right),$$

which means that the Malliavin derivative of X^n is bounded in $L^2(\Omega, \mathbb{P}; \mathbb{R}^d)$.

Next we prove (3.6). We deduce from (3.11) that for all $0 \leq r' \leq r \leq s \leq 1$, $0 \leq u' \leq u \leq t \leq 1$,

$$\begin{aligned} & D_{r,u} X_{s,t}^n - D_{r',u'} X_{s,t}^n \\ & = \int_r^s \int_u^t \nabla b_n(s_1, t_1, X_{s_1, t_1}^n) D_{r,u} X_{s_1, t_1}^n dt_1 ds_1 - \int_{r'}^s \int_{u'}^t \nabla b_n(s_1, t_1, X_{s_1, t_1}^n) D_{r',u'} X_{s_1, t_1}^n dt_1 ds_1 \\ & = \int_r^s \int_u^t \nabla b_n(s_1, t_1, X_{s_1, t_1}^n) \left(D_{r,u} X_{s_1, t_1}^n - D_{r',u'} X_{s_1, t_1}^n \right) dt_1 ds_1 - \int_{r'}^s \int_{u'}^u \nabla b_n(s_1, t_1, X_{s_1, t_1}^n) D_{r',u'} X_{s_1, t_1}^n dt_1 ds_1 \\ & \quad - \int_{r'}^r \int_u^t \nabla b_n(s_1, t_1, X_{s_1, t_1}^n) D_{r',u'} X_{s_1, t_1}^n dt_1 ds_1. \end{aligned}$$

Taking the norm on both sides and using the fact that $b_n = \hat{b}_n - \check{b}_n$, with \hat{b}_n, \check{b}_n nondecreasing, gives

$$\begin{aligned} & \left\| D_{r,u} X_{s,t}^n - D_{r',u'} X_{s,t}^n \right\| \\ & \leq \int_r^s \int_u^t \sum_{i,j=1}^d \left\{ \partial_i \hat{b}_{j,n}(s_1, t_1, X_{s_1, t_1}^n) + \partial_i \check{b}_{j,n}(s_1, t_1, X_{s_1, t_1}^n) \right\} \left\| D_{r,u} X_{s_1, t_1}^n - D_{r',u'} X_{s_1, t_1}^n \right\| dt_1 ds_1 \\ & \quad + \int_{r'}^s \int_{u'}^u \sum_{i,j=1}^d \left\{ \partial_i \hat{b}_{j,n}(s_1, t_1, X_{s_1, t_1}^n) + \partial_i \check{b}_{j,n}(s_1, t_1, X_{s_1, t_1}^n) \right\} \left\| D_{r',u'} X_{s_1, t_1}^n \right\| dt_1 ds_1 \\ & \quad + \int_{r'}^r \int_u^t \sum_{i,j=1}^d \left\{ \partial_i \hat{b}_{j,n}(s_1, t_1, X_{s_1, t_1}^n) + \partial_i \check{b}_{j,n}(s_1, t_1, X_{s_1, t_1}^n) \right\} \left\| D_{r',u'} X_{s_1, t_1}^n \right\| dt_1 ds_1. \end{aligned}$$

Applying Theorem B.1, we obtain

$$\begin{aligned}
(3.14) \quad & \left\| D_{r,u} X_{s,t}^n - D_{r',u'} X_{s,t}^n \right\| \\
& \leq \left(\int_{r'}^s \int_{u'}^u \sum_{i,j=1}^d \left\{ \partial_i \hat{b}_{j,n}(s_1, t_1, X_{s_1, t_1}^n) + \partial_i \check{b}_{j,n}(s_1, t_1, X_{s_1, t_1}^n) \right\} \left\| D_{r',u'} X_{s_1, t_1}^n \right\| dt_1 ds_1 \right. \\
& \quad \left. + \int_{r'}^r \int_u^t \sum_{i,j=1}^d \left\{ \partial_i \hat{b}_{j,n}(s_1, t_1, X_{s_1, t_1}^n) + \partial_i \check{b}_{j,n}(s_1, t_1, X_{s_1, t_1}^n) \right\} \left\| D_{r',u'} X_{s_1, t_1}^n \right\| dt_1 ds_1 \right) \\
& \quad \times \exp \left(\int_r^s \int_u^t \sum_{i,j=1}^d \left\{ \partial_i \hat{b}_{j,n}(s_1, t_1, X_{s_1, t_1}^n) + \partial_i \check{b}'_{j,n}(s_1, t_1, X_{s_1, t_1}^n) \right\} dt_1 ds_1 \right)
\end{aligned}$$

Since $\partial_i \hat{b}_{j,n}$ and $\partial_i \check{b}_{j,n}$ are nonnegative, it follows from (3.12) that

$$\left\| D_{r',u'} X_{s_1, t_1}^n \right\| \leq \exp \left(\int_{r'}^s \int_{u'}^t \sum_{i,j=1}^d \left\{ \partial_i \hat{b}_{j,n}(s_2, t_2, X_{s_2, t_2}^n) + \partial_i \check{b}_{j,n}(s_2, t_2, X_{s_2, t_2}^n) \right\} dt_2 ds_2 \right).$$

Hence, we deduce from (3.14) that it holds:

$$\begin{aligned}
\left\| D_{r,u} X_{s,t}^n - D_{r',u'} X_{s,t}^n \right\| & \leq \left(\int_{r'}^s \int_{u'}^u \sum_{i,j=1}^d \left\{ \partial_i \hat{b}_{j,n}(s_1, t_1, X_{s_1, t_1}^n) + \partial_i \check{b}_{j,n}(s_1, t_1, X_{s_1, t_1}^n) \right\} dt_1 ds_1 \right. \\
& \quad \left. + \int_{r'}^r \int_u^t \sum_{i,j=1}^d \left\{ \partial_i \hat{b}_{j,n}(s_1, t_1, X_{s_1, t_1}^n) + \partial_i \check{b}_{j,n}(s_1, t_1, X_{s_1, t_1}^n) \right\} dt_1 ds_1 \right) \\
& \quad \times \exp \left(2 \int_{r'}^s \int_{u'}^t \sum_{i,j=1}^d \left\{ \partial_i \hat{b}_{j,n}(s_1, t_1, X_{s_1, t_1}^n) + \partial_i \check{b}_{j,n}(s_1, t_1, X_{s_1, t_1}^n) \right\} dt_1 ds_1 \right).
\end{aligned}$$

Squaring both sides of the inequality, taking the expectation and using Cauchy-Schwarz inequality and Girsanov theorem give

$$\begin{aligned}
& \mathbb{E} \left[\left\| D_{r,u} X_{s,t}^n - D_{r',u'} X_{s,t}^n \right\|^2 \right] \\
& \leq 4^7 C_0 \left(\sum_{i,j=1}^d \left\{ \mathbb{E} \left[\left(\int_{r'}^s \int_{u'}^u \partial_i \hat{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) dt_1 ds_1 \right)^8 \right] + \mathbb{E} \left[\left(\int_{r'}^s \int_{u'}^u \partial_i \check{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) dt_1 ds_1 \right)^8 \right] \right. \right. \\
& \quad \left. \left. + \mathbb{E} \left[\left(\int_{r'}^r \int_u^t \partial_i \hat{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) dt_1 ds_1 \right)^8 \right] + \mathbb{E} \left[\left(\int_{r'}^r \int_u^t \partial_i \check{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) dt_1 ds_1 \right)^8 \right] \right\} \right)^{\frac{1}{4}} \\
& \quad \times \mathbb{E} \left[\exp \left(16 \int_{r'}^s \int_{u'}^t \sum_{i,j=1}^d \left\{ \partial_i \hat{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) + \partial_i \check{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) \right\} dt_1 ds_1 \right) \right]^{\frac{1}{4}} \\
& = 4^7 C_0 (J_1 + J_2 + J_3 + J_4)^{1/4} \times J_5,
\end{aligned}$$

where C_0 is given by (3.13). It follows from Hölder inequality and the estimates (3.7)-(3.8) that

$$\begin{aligned} J_5 &= \mathbb{E} \left[\exp \left(16 \int_{r'}^s \int_{u'}^t \sum_{i,j=1}^d \left\{ \partial_i \hat{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) + \partial_i \check{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) \right\} dt_1 ds_1 \right) \right]^{\frac{1}{4}} \\ &\leq \prod_{i,j=1}^d \mathbb{E} \left[\exp \left(32d^2 \int_r^s \int_u^t \partial_i \hat{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) dt_1 ds_1 \right) \right]^{\frac{1}{8d^2}} \mathbb{E} \left[\exp \left(32d^2 \int_r^s \int_u^t \partial_i \check{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) dt_1 ds_1 \right) \right]^{\frac{1}{8d^2}} \\ &\leq \tilde{C}_1 (32d^2, \|\hat{b}\|_\infty + \|\check{b}\|_\infty)^{\frac{1}{4}}. \end{aligned}$$

Moreover, using the inequality $x^8 \leq 8!e^x$ ($x \in \mathbb{R}$), we have

$$\begin{aligned} &\mathbb{E} \left[\left(\int_{r'}^s \int_{u'}^u \partial_i \hat{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) dt_1 ds_1 \right)^8 \right] + \mathbb{E} \left[\left(\int_{r'}^s \int_{u'}^u \partial_i \check{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) dt_1 ds_1 \right)^8 \right] \\ &\leq 8! \delta(r', s)^8 \delta(u', u)^8 \mathbb{E} \left[\exp \left(\frac{1}{\delta(r', s) \delta(u', u)} \left| \int_{r'}^s \int_{u'}^u \partial_i \hat{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) dt_1 ds_1 \right| \right) \right] \\ &\quad + 8! \delta(r', s)^8 \delta(u', u)^8 \mathbb{E} \left[\exp \left(\frac{1}{\delta(r', s) \delta(u', u)} \left| \int_{r'}^s \int_{u'}^u \partial_i \check{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) dt_1 ds_1 \right| \right) \right] \\ &\leq 2(8!) \tilde{C}_1 (1, \|\hat{b}\|_\infty + \|\check{b}\|_\infty) \delta(u', u)^4 \end{aligned}$$

and

$$\begin{aligned} &\mathbb{E} \left[\left(\int_{r'}^r \int_u^t \partial_i \hat{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) dt_1 ds_1 \right)^8 \right] + \mathbb{E} \left[\left(\int_{r'}^r \int_u^t \partial_i \check{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) dt_1 ds_1 \right)^8 \right] \\ &\leq 8! \delta(r', r)^8 \delta(u, t)^8 \mathbb{E} \left[\exp \left(\frac{1}{\delta(r', r) \delta(u, t)} \left| \int_{r'}^r \int_u^t \partial_i \hat{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) dt_1 ds_1 \right| \right) \right] \\ &\quad + 8! \delta(r', r)^8 \delta(u, t)^8 \mathbb{E} \left[\exp \left(\frac{1}{\delta(r', r) \delta(u, t)} \left| \int_{r'}^r \int_u^t \partial_i \check{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) dt_1 ds_1 \right| \right) \right] \\ &\leq 2(8!) \tilde{C}_1 (1, \|\hat{b}\|_\infty + \|\check{b}\|_\infty) \delta(r, r')^8. \end{aligned}$$

As a consequence,

$$J_1 + J_2 + J_3 + J_4 \leq 4(8!) \tilde{C}_1 (1, \|\hat{b}\|_\infty + \|\check{b}\|_\infty) (|r - r'|^4 + |u - u'|^4).$$

Therefore

$$\mathbb{E} \left[\left\| D_{r,u} X_{s,t}^n - D_{r',u'} X_{s,t}^n \right\|^2 \right] \leq 4^8 (8!) C_0 \times \tilde{C}_1 (32d^2, \|\hat{b}\|_\infty + \|\check{b}\|_\infty)^{1/2} (|r - r'| + |u - u'|).$$

Finally, by the Girsanov theorem and the Cauchy-Schwarz inequality, we have

$$\begin{aligned} \sup_{n \geq 1} \|X_{s,t}^n\|_{L^2(\Omega, \mathbb{P}; \mathbb{R}^d)}^2 &= \sup_{n \geq 1} \mathbb{E} [|X_{s,t}^n|^2] = \sup_{n \geq 1} \mathbb{E} \left[\mathcal{E} \left(\int_0^1 \int_0^1 b_n(s_1, t_1, W_{s_1, t_1}) \cdot dW_{s_1, t_1} \right) \middle| W_{s,t} \right]^2 \\ &\leq \sup_{n \geq 1} \mathbb{E} \left[\mathcal{E} \left(\int_0^1 \int_0^1 b_n(s_1, t_1, W_{s_1, t_1}) \cdot dW_{s_1, t_1} \right)^2 \right]^{\frac{1}{2}} \mathbb{E} [|W_{s,t}|^4]^{\frac{1}{2}} \leq C_0 \sqrt{3d}, \end{aligned}$$

where C_0 is given by (3.13).

The proof is completed by taking $C_1 = C_0 \max\{\sqrt{3d}, 4^8(8!) \tilde{C}_1(32d^2, \|\hat{b}\|_\infty + \|\check{b}\|_\infty)\}$. \square

For $q \geq 1$, let us consider the following space $L^q(\mathbb{R}^d; \mathbf{p}(x)dx)$ defined by

$$(3.15) \quad L^q(\mathbb{R}^d; \mathbf{p}(x)dx) = \left\{ h : \mathbb{R}^d \rightarrow \mathbb{R}^d \text{ measurable and such that } \int_{\mathbb{R}^d} |h(x)|^q \mathbf{p}(x) dx < \infty \right\},$$

where the weight function $\mathbf{p}(x)$ is defined by

$$\mathbf{p}(x) = e^{-\frac{|x|^2}{2st}}, \quad x \in \mathbb{R}^d.$$

Theorem 3.9. *Let b_n be defined as before and let $(X_{s,t}^{\xi,n})_{n \geq 1}$ be the sequence of corresponding strong solutions to the SDE (1.3). Then for any fixed $s, t \in [0, 1]$, $(X_{s,t}^{\xi,n})_{n \geq 1}$ converges strongly in $L^2(\Omega, \mathbb{P}; \mathbb{R}^d)$ to $X_{s,t}^{\xi}$.*

In order to prove the above theorem we need the subsequent result.

Lemma 3.10. *Let $(X_{s,t}^{\xi,n})_{n \geq 1}$ be the sequence of corresponding strong solutions as given before. Then for every $s, t \in [0, 1]$ and function $h \in L^4(\mathbb{R}^d; \mathbf{p}(x)dx)$, it holds that the sequence $(h(X_{s,t}^{\xi,n}))_{n \geq 1}$ converges weakly in $L^2(\Omega, \mathbb{P}; \mathbb{R}^d)$ to $h(X_{s,t}^{\xi})$.*

Proof of Theorem 3.9. Using Theorem 3.5, we know that for each s, t , there exists a subsequence $(X_{s,t}^{\xi,n_k})_{k \geq 1}$ that converges strongly in $L^2(\Omega, \mathbb{P}; \mathbb{R}^d)$. Set $h(x) = x$, $x \in \mathbb{R}^d$ and use Lemma 3.10 to obtain that $(X_{s,t}^{\xi,n_k})_{k \geq 1}$ converges weakly to $X_{s,t}^{\xi}$ in $L^2(\Omega, \mathbb{P}; \mathbb{R}^d)$. Thanks to the uniqueness of the limit, there exists a subsequence n_k such that $(X_{s,t}^{\xi,n_k})_{k \geq 1}$ converges strongly to $X_{s,t}^{\xi}$ in $L^2(\Omega, \mathbb{P}; \mathbb{R}^d)$. The convergence then holds for the entire sequence by uniqueness of the limit. To see this, suppose by contradiction that for some s, t , there exist $\epsilon > 0$ and a subsequence $n_l, l \geq 0$ such that

$$\|X_{s,t}^{\xi,n_l} - X_{s,t}^{\xi}\|_{L^2(\Omega, \mathbb{P}; \mathbb{R}^d)} \geq \epsilon.$$

We also know from the compactness criteria that there exists a further subsequence $n_m, m \geq 0$ of $n_l, l \geq 0$ such that

$$X_{s,t}^{\xi,n_m} \rightarrow \tilde{X}_{s,t} \text{ in } L^2(\Omega, \mathbb{P}; \mathbb{R}^d) \text{ as } m \rightarrow \infty.$$

However, $(X_{s,t}^{\xi,n_k})_{k \geq 1} \rightarrow X_{s,t}^{\xi}$ as $k \rightarrow \infty$ weakly in $L^2(\Omega, \mathbb{P}; \mathbb{R}^d)$, and hence by the uniqueness of the limit, we obtain

$$\tilde{X}_{s,t} = X_{s,t}^{\xi}.$$

Since

$$\|X_{s,t}^{\xi,n_m} - X_{s,t}^{\xi}\|_{L^2(\Omega, \mathbb{P}; \mathbb{R}^d)} \geq \epsilon,$$

we obtain a contradiction. \square

Proof of Theorem 3.4. We know from Theorem 3.9 that $(X_{s,t}^{\xi,n})_{n \geq 1}$ converges strongly in $L^2(\Omega, \mathbb{P}; \mathbb{R}^d)$ to $X_{s,t}^{\xi}$ and from (3.5) in Lemma 3.6 that $(D_{r,u} X_{s,t}^{\xi,n})_{n \geq 1}$ is bounded in the $L^2([0, 1]^2 \times \Omega, ds \times dt \times \mathbb{P}; \mathbb{R}^{d \times d})$ -norm uniformly in n . Therefore, using [32, Lemma 1.2.3], we also have that the limit $X_{s,t}^{\xi}$ is Malliavin differentiable. \square

Proof of Lemma 3.10. Let us first noticing that the space

$$\left\{ \mathcal{E} \left(\int_0^1 \int_0^1 \frac{\partial^2 \varphi_{s_1, t_1}}{\partial s_1 \partial t_1} \cdot dW_{s_1, t_1} \right) : \varphi \in C_{2,b}([0, 1]^2, \mathbb{R}^d) \right\}$$

is a dense subspace of $L^2(\Omega, \mathbb{P}; \mathbb{R}^d)$. Here $C_{2,b}([0, 1]^2, \mathbb{R}^d)$ is the space of bounded vector functions φ such that each component φ^i has a second partial derivative $\frac{\partial^2 \varphi_{s_1, t_1}^i}{\partial s_1 \partial t_1}$ of bounded variation with values in \mathbb{R} . Hence, it suffices to show that for every i ,

$$\mathbb{E} \left[h_i(X_{s,t}^{\xi,n}) \mathcal{E} \left(\int_0^1 \int_0^1 \frac{\partial^2 \varphi_{s_1, t_1}}{\partial s_1 \partial t_1} \cdot dW_{s_1, t_1} \right) \right] \longrightarrow \mathbb{E} \left[h_i(X_{s,t}^{\xi}) \mathcal{E} \left(\int_0^1 \int_0^1 \frac{\partial^2 \varphi_{s_1, t_1}}{\partial s_1 \partial t_1} \cdot dW_{s_1, t_1} \right) \right] \text{ as } n \rightarrow \infty.$$

Since Ω is a Wiener space, then, as in [25, proof Lemma 2] or [46, proof of Theorem 2], one can show a multidimensional analog of the Cameron-Martin translation theorem. Precisely for every $g : \mathbb{R}^d \rightarrow \mathbb{R}$ measurable, one has

$$(3.16) \quad \mathbb{E} \left[g(X_{s,t}^{\xi}) \mathcal{E} \left(\int_0^1 \int_0^1 \frac{\partial^2 \varphi_{s_1, t_1}}{\partial s_1 \partial t_1} \cdot dW_{s_1, t_1} \right) \right] = \int_{\Omega} g(X_{s,t}^{\xi}(\omega + \varphi)) d\mathbb{P}(\omega).$$

Let $\varphi \in C_{2,b}([0, 1]^2, \mathbb{R}^d)$. For every n , the process $\tilde{X}^{\xi,n}$ given by $\tilde{X}_{s,t}^{\xi,n}(\omega) := X_{s,t}^{\xi,n}(\omega + \varphi)$ solves the SDE

$$(3.17) \quad d\tilde{X}_{s,t}^{\xi,n} = \left(b_n(t, \tilde{X}_{s,t}^{\xi,n}) + \frac{\partial^2 \varphi_{s,t}}{\partial s \partial t} \right) ds dt + dW_{s,t}.$$

Since X^ξ is also the solution to the SDE it holds that $\tilde{X}^\xi(\omega) := X^\xi(\omega + \varphi)$ satisfies

$$(3.18) \quad d\tilde{X}_{s,t}^\xi = \left(b(t, \tilde{X}_{s,t}^\xi) + \frac{\partial^2 \varphi_{s,t}}{\partial s \partial t} \right) ds dt + dW_{s,t}, \quad \mathbb{P}\text{-a.s.}$$

Applying (3.16) and the Girsanov theorem, we have

$$(3.19) \quad \begin{aligned} & \mathbb{E} \left[h_i(X_{s,t}^{\xi,n}) \mathcal{E} \left(\int_0^1 \int_0^1 \frac{\partial^2 \varphi_{s_1,t_1}}{\partial s_1 \partial t_1} \cdot dW_{s_1,t_1} \right) - h_i(X_{s,t}^\xi) \mathcal{E} \left(\int_0^1 \int_0^1 \frac{\partial^2 \varphi_{s_1,t_1}}{\partial s_1 \partial t_1} \cdot dW_{s_1,t_1} \right) \right] \\ &= \mathbb{E} \left[\left(h_i(X_{s,t}^{\xi,n}) - h_i(X_{s,t}^\xi) \right) \mathcal{E} \left(\int_0^1 \int_0^1 \frac{\partial^2 \varphi_{s_1,t_1}}{\partial s_1 \partial t_1} \cdot dW_{s_1,t_1} \right) \right] \\ &= \mathbb{E} \left[h_i(\xi + W_{s,t}) \left\{ \mathcal{E} \left(\int_0^1 \int_0^1 \left\{ b_n(s_1, t_1, \xi + W_{s_1,t_1}) + \frac{\partial^2 \varphi_{s_1,t_1}}{\partial s_1 \partial t_1} \right\} \cdot dW_{s_1,t_1} \right) \right. \right. \\ & \quad \left. \left. - \mathcal{E} \left(\int_0^1 \int_0^1 \left\{ b(s_1, t_1, \xi + W_{s_1,t_1}) + \frac{\partial^2 \varphi_{s_1,t_1}}{\partial s_1 \partial t_1} \right\} \cdot dW_{s_1,t_1} \right) \right\} \right]. \end{aligned}$$

Using the fact that $|e^a - e^b| \leq |e^a + e^b||a - b|$, the Hölder inequality and Burkholder-Davis-Gundy inequality, we get

$$(3.20) \quad \begin{aligned} & \mathbb{E} \left[h_i(X_{s,t}^{\xi,n}) \mathcal{E} \left(\int_0^1 \int_0^1 \frac{\partial^2 \varphi_{s_1,t_1}}{\partial s_1 \partial t_1} \cdot dW_{s_1,t_1} \right) - h_i(X_{s,t}^\xi) \mathcal{E} \left(\int_0^1 \int_0^1 \frac{\partial^2 \varphi_{s_1,t_1}}{\partial s_1 \partial t_1} \cdot dW_{s_1,t_1} \right) \right] \\ & \leq C \mathbb{E} \left[h_i(x + W_{s,t})^2 \right]^{\frac{1}{2}} \mathbb{E} \left[\left(\mathcal{E} \left(\int_0^1 \int_0^1 \left\{ b_n(s_1, t_1, \xi + W_{s_1,t_1}) + \frac{\partial^2 \varphi_{s_1,t_1}}{\partial s_1 \partial t_1} \right\} \cdot dW_{s_1,t_1} \right) \right. \right. \\ & \quad \left. \left. + \mathcal{E} \left(\int_0^1 \int_0^1 \left\{ b(s_1, t_1, \xi + W_{s_1,t_1}) + \frac{\partial^2 \varphi_{s_1,t_1}}{\partial s_1 \partial t_1} \right\} \cdot dW_{s_1,t_1} \right) \right)^4 \right]^{\frac{1}{4}} \\ & \quad \times \left\{ \mathbb{E} \left[\left(\int_0^1 \int_0^1 \left(b_n(s_1, t_1, \xi + W_{s_1,t_1}) - b(s_1, t_1, \xi + W_{s_1,t_1}) \right) \cdot dW_{s_1,t_1} \right)^4 \right] \right\} \\ & \quad + \mathbb{E} \left[\left(\frac{1}{2} \int_0^1 \int_0^1 \left(b_n(s_1, t_1, \xi + W_{s_1,t_1}) + \frac{\partial^2 \varphi_{s_1,t_1}}{\partial s_1 \partial t_1} \right)^2 - \left(b(s_1, t_1, \xi + W_{s_1,t_1}) + \frac{\partial^2 \varphi_{s_1,t_1}}{\partial s_1 \partial t_1} \right)^2 ds_1 dt_1 \right)^4 \right]^{\frac{1}{4}} \\ & \leq C \mathbb{E} \left[h_i(x + W_{s,t})^2 \right]^{\frac{1}{2}} \mathbb{E} \left[\left(\mathcal{E} \left(\int_0^1 \int_0^1 \left\{ b_n(s_1, t_1, \xi + W_{s_1,t_1}) + \frac{\partial^2 \varphi_{s_1,t_1}}{\partial s_1 \partial t_1} \right\} \cdot dW_{s_1,t_1} \right) \right. \right. \\ & \quad \left. \left. + \mathcal{E} \left(\int_0^1 \int_0^1 \left\{ b(s_1, t_1, \xi + W_{s_1,t_1}) + \frac{\partial^2 \varphi_{s_1,t_1}}{\partial s_1 \partial t_1} \right\} \cdot dW_{s_1,t_1} \right) \right)^4 \right]^{\frac{1}{4}} \\ & \quad \times \left\{ \int_0^1 \int_0^1 \mathbb{E} \left[\left| b_n(r, \xi + W_{s_1,t_1}) - b(s_1, t_1, \xi + W_{s_1,t_1}) \right|^4 \right] ds_1 dt_1 \right. \\ & \quad \left. + \frac{1}{16} \int_0^1 \int_0^1 \mathbb{E} \left[\left| \left(b_n(s_1, t_1, \xi + W_{s_1,t_1}) + \frac{\partial^2 \varphi_{s_1,t_1}}{\partial s_1 \partial t_1} \right)^2 - \left(b(s_1, t_1, \xi + W_{s_1,t_1}) + \frac{\partial^2 \varphi_{s_1,t_1}}{\partial s_1 \partial t_1} \right)^2 \right|^4 \right] ds_1 dt_1 \right]^{\frac{1}{4}} \\ & = I_1 \times I_{2,n} \times (I_{3,n} + I_{4,n}). \end{aligned}$$

I_1 is finite since $h \in L^4(\mathbb{R}; \mathbf{p}(x)dx)$. Next observe that

$$\begin{aligned} & \mathcal{E} \left(\int_0^1 \int_0^1 \left\{ b_n(s_1, t_1, \xi + W_{s_1, t_1}) + \frac{\partial^2 \varphi_{s_1, t_1}}{\partial s_1 \partial t_1} \right\} \cdot dW_{s_1, t_1} \right) \\ &= \mathcal{E} \left(\int_0^1 \int_0^1 b_n(s_1, t_1, \xi + W_{s_1, t_1}) \cdot dW_{s_1, t_1} \right) \mathcal{E} \left(\int_0^1 \int_0^1 \frac{\partial^2 \varphi_{s_1, t_1}}{\partial s_1 \partial t_1} \cdot dW_{s_1, t_1} \right) \\ & \quad \times \exp \left(\int_0^1 \int_0^1 b_n(s_1, t_1, \xi + W_{s_1, t_1}) \frac{\partial^2 \varphi_{s_1, t_1}}{\partial s_1 \partial t_1} ds_1 dt_1 \right). \end{aligned}$$

Using the boundedness of $\frac{\partial^2 \varphi_{s_1, t_1}}{\partial s_1 \partial t_1}$ and the uniform boundedness of b_n , it follows that $I_{2,n}$ is bounded. Using the dominated convergence theorem, we get that $I_{3,n}$ and $I_{4,n}$ converge to 0 as n goes to infinity. Let us for example consider the term $I_{3,n}$. Using the density of the Brownian sheet for every $q \geq 1$ it holds:

$$\begin{aligned} \mathbb{E} \left[\left| b_n(s, t, \xi + W_{s, t}) - b(s, t, \xi + W_{s, t}) \right|^q \right] &= \frac{1}{\sqrt{2\pi st}} \int_{\mathbb{R}} |b_n(s, t, \xi + z) - b(s, t, \xi + z)|^q e^{-\frac{|z|^2}{2st}} dz \\ &= \frac{1}{\sqrt{2\pi st}} \int_{\mathbb{R}} |b_n(s, t, z) - b(s, t, z)|^q e^{-\frac{|z-\xi|^2}{2st}} dz \\ &= \frac{1}{\sqrt{2\pi st}} \int_{\mathbb{R}} |b_n(s, t, z) - b(s, t, z)|^q e^{-\frac{|z-2\xi|^2}{4st}} e^{-\frac{|z|^2}{4st}} e^{\frac{|\xi|^2}{4st}} dz \\ &\leq \frac{e^{\frac{|\xi|^2}{4st}}}{\sqrt{2\pi st}} \int_{\mathbb{R}} |b_n(s, t, z) - b(s, t, z)|^q e^{-\frac{|z|^2}{4st}} dz. \end{aligned}$$

Thus the result follows by the dominated convergence theorem. \square

3.3. Malliavin regularity under linear growth condition. As in the previous section, we approximate the drift coefficient $b = \hat{b} - \check{b}$ by a sequence of functions $b_n := \hat{b}_n - \check{b}_n, n \geq 1$, where $\hat{b}_n = (\hat{b}_{1,n}, \dots, \hat{b}_{d,n})$, $\check{b}_n = (\check{b}_{1,n}, \dots, \check{b}_{d,n})$, $(\hat{b}_{j,n})_{n \geq 1}$ and $(\check{b}_{j,n})_{n \geq 1}$ are smooth, componentwise non-decreasing and bounded functions satisfying:

- There exists $\tilde{M} > 0$ such that $\|\hat{b}_{j,n}(s, t, x)\| \leq \tilde{M}(1 + |x|)$ and $\|\check{b}_{j,n}(s, t, x)\| \leq \tilde{M}(1 + |x|)$ for all $n \geq 1$ and $(s, t, x) \in \Gamma_0 \times \mathbb{R}^d$,
- $(\hat{b}_n)_{n \geq 1}$ (respectively, $(\check{b}_n)_{n \geq 1}$) converges to \hat{b} (respectively, \check{b}) in $(s, t, x) \in \Gamma_0 \times \mathbb{R}^d$ $ds \times dt \times dx$ -a.e.

One verifies that for such smooth drift coefficients, the corresponding SDEs have a unique strong solution denoted by $X^{\xi, n}$. We show that for s, t small enough, the sequence $(X_{s, t}^{\xi, n})_{n \geq 1}$ is relatively compact in $L^2(\Omega, \mathbb{P}; \mathbb{R}^d)$. The proofs of the next two results are similar to that of Lemmas 3.8 and 3.6 and are found in Appendix.

Lemma 3.11. *There exist $\tilde{C}_2 > 0$ and $\zeta > 0$ such that, for any $i, j \in \{1, \dots, d\}$, any $0 < r < s \leq 1$, any $0 < u < t \leq 1$ and any $k \in \mathbb{R}_+$,*

$$(3.21) \quad \mathbb{E} \left[\exp \left(\frac{\zeta}{\delta(r, s)\delta(u, t)} \int_r^s \int_u^t \partial_i \hat{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) dt_1 ds_1 \right) \right] \leq \tilde{C}_2,$$

and

$$(3.22) \quad \mathbb{E} \left[\exp \left(\frac{\zeta}{\delta(r, s)\delta(u, t)} \int_r^s \int_u^t \partial_i \check{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) dt_1 ds_1 \right) \right] \leq \tilde{C}_2,$$

where $\delta(r, s) = \sqrt{s-r}$ and $\delta(u, t) = \sqrt{t-u}$.

Lemma 3.12. *There exist $C_2 > 0$ and $\tau \in (0, 1)$ such that for every $(s, t) \in [0, \tau]$, the sequence $(X_{s, t}^{\xi, n})_{n \geq 1}$ satisfies*

$$(3.23) \quad \sup_{n \geq 1} \|X_{s, t}^n\|_{L^2(\Omega, \mathbb{P}; \mathbb{R}^d)}^2 \leq C_2$$

and

$$(3.24) \quad \sup_{n \geq 1} \sup_{\substack{0 \leq r \leq s \\ 0 \leq u \leq t}} \mathbb{E} [\|D_{r,u} X_{s,t}^n\|^2] \leq C_2.$$

Moreover, for all $0 \leq r', r \leq s \leq \tau$, $0 \leq u', u \leq t \leq \tau$,

$$(3.25) \quad \mathbb{E} [\|D_{r,u} X_{s,t}^n - D_{r',u'} X_{s,t}^n\|^2] \leq C_2(|r - r'| + |u - u'|).$$

Here is the main result of this section which is a consequence of Lemma 3.12 and the compactness criterion provided in Corollary 3.3.

Theorem 3.13. *There exist $\tau \in (0, 1)$ such that for any the strong solution $\{X_{s,t}^\xi, s, t \in [0, \tau]\}$ of the SDE (1.3) is Malliavin differentiable.*

Proof. As in the proof of Lemma 3.10, we show that $(X_{s,t}^{\xi,n})_{n \geq 1}$ converges weakly in $L^2(\Omega, \mathbb{P}; \mathbb{R}^d)$ to $X_{s,t}^\xi$ for every $(s, t) \in [0, \tau]^2$. Hence, using Lemma 3.12 and the compactness criterion provided in Corollary 3.3, we deduce that $(X_{s,t}^{\xi,n})_{n \geq 1}$ converges strongly in $L^2(\Omega, \mathbb{P}; \mathbb{R}^d)$ to $X_{s,t}^\xi$. Then, since $(D_{r,u} X_{s,t}^{\xi,n})_{n \geq 1}$ is bounded in the $L^2([0, 1]^2 \times \Omega, ds \times dt \times \mathbb{P}; \mathbb{R}^{d \times d})$ -norm uniformly in n (see (3.24)), it follows from [32, Lemma 1.2.3] that the limit $X_{s,t}^\xi$ is also Malliavin differentiable. \square

APPENDIX A. PROOFS OF LEMMAS 3.11 AND 3.12

In this section we provide the proofs of Lemmas 3.11 and 3.12. Let us start with a useful estimate.

Lemma A.1. *There exist $\tau_1 \in (0, 1)$ such that*

$$(A.1) \quad \sup_{n \geq 1} \mathbb{E} \left[\mathcal{E} \left(\int_0^{\tau_1} \int_0^{\tau_1} b_n(s, t, W_{s,t}) \cdot dW_{s,t} \right)^2 \right] < \infty.$$

Proof. By Cauchy-Schwarz inequality, we have

$$\begin{aligned} & \mathbb{E} \left[\mathcal{E} \left(\int_0^{\tau_1} \int_0^{\tau_1} b_n(s, t, W_{s,t}) \cdot dW_{s,t} \right)^2 \right] \\ &= \mathbb{E} \left[\exp \left(2 \int_0^{\tau_1} \int_0^{\tau_1} b_n(s, t, W_{s,t}) \cdot dW_{s,t} - \int_0^{\tau_1} \int_0^{\tau_1} |b_n(s, t, W_{s,t})|^2 ds dt \right) \right] \\ &= \mathbb{E} \left[\exp \left(2 \int_0^{\tau_1} \int_0^{\tau_1} b_n(s, t, W_{s,t}) \cdot dW_{s,t} - 4 \int_0^{\tau_1} \int_0^{\tau_1} |b_n(s, t, W_{s,t})|^2 ds dt + 3 \int_0^{\tau_1} \int_0^{\tau_1} |b_n(s, t, W_{s,t})|^2 ds dt \right) \right] \\ &\leq \mathbb{E} \left[\mathcal{E} \left(4 \int_0^{\tau_1} \int_0^{\tau_1} b_n(s, t, W_{s,t}) \cdot dW_{s,t} \right) \right]^{\frac{1}{2}} \mathbb{E} \left[\exp \left(6 \int_0^{\tau_1} \int_0^{\tau_1} |b_n(s, t, W_{s,t})|^2 ds dt \right) \right]^{\frac{1}{2}}. \end{aligned}$$

Since b_n is bounded, we have (see e.g. [33, Proposition 1.6])

$$\mathbb{E} \left[\mathcal{E} \left(4 \int_0^{\tau_1} \int_0^{\tau_1} b_n(s, t, W_{s,t}) \cdot dW_{s,t} \right) \right] = 1, \quad \forall n \geq 1, \tau_1 > 0.$$

Moreover, by Jensen inequality,

$$\begin{aligned} \mathbb{E} \left[\exp \left(6 \int_0^{\tau_1} \int_0^{\tau_1} |b_n(s, t, W_{s,t})|^2 ds dt \right) \right] &\leq \frac{1}{\tau_1^2} \int_0^{\tau_1} \int_0^{\tau_1} \mathbb{E} \left[\exp \left(6\tau_1^2 |b_n(s, t, W_{s,t})|^2 \right) \right] ds dt \\ &\leq \frac{1}{\tau_1^2} \int_0^{\tau_1} \int_0^{\tau_1} \mathbb{E} \left[\exp \left(24\tau_1^2 \tilde{M} d (1 + |W_{s,t}|^2) \right) \right] ds dt \\ &= \frac{\exp \left(24\tau_1^2 \tilde{M} d \right)}{\tau_1^2} \int_0^{\tau_1} \int_0^{\tau_1} \mathbb{E} \left[\exp \left(24\tau_1^2 \tilde{M} d st |G|^2 \right) \right] ds dt \\ &\leq \exp \left(24\tau_1^2 \tilde{M} d \right) \mathbb{E} \left[\exp \left(24\tau_1^4 \tilde{M} d |G|^2 \right) \right], \end{aligned}$$

where $G = (G_1, \dots, G_d)$ is a mean-zero random vector with identity covariance matrix.

The proof is completed since $\mathbb{E} \left[\exp(12\tau_1^4 \tilde{M}d|G|^2) \right] < \infty$ for τ_1 small enough. \square

Remark A.2. *When the drift b is bounded, the functions b_n , $n \geq 1$ are uniformly bounded and, as a consequence,*

$$C_0 = \sup_{n \geq 1} \mathbb{E} \left[\mathcal{E} \left(\int_0^1 \int_0^1 b_n(s, t, W_{s,t}) \cdot dW_{s,t} \right)^2 \right]^{\frac{1}{2}} < \infty.$$

Proof of Lemma 3.11. We only prove (3.21) since the proof of (3.22) follows the same lines. We deduce from the local time-space integration formula (2.1) that

$$\begin{aligned} & \mathbb{E} \left[\exp \left(\frac{\zeta}{\delta(r, s)\delta(u, t)} \left| \int_r^s \int_u^t \partial_i \hat{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) dt_1 ds_1 \right| \right) \right] \\ (A.2) \quad & \leq \frac{1}{3} \left\{ \mathbb{E} \left[\exp \left(\frac{3\zeta}{\delta(r, s)\delta(u, t)} \left| \int_r^s \int_u^t \hat{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) \frac{d_{t_1} W_{s_1, t_1}^{(i)}}{s_1} ds_1 \right| \right) \right] \right. \\ & + \mathbb{E} \left[\exp \left(\frac{3\zeta}{\delta(r, s)\delta(u, t)} \left| \int_r^s \int_{1-t}^{1-u} \hat{b}_{j,n}(s_1, t_1, W_{s_1, 1-t_1}) \frac{d_{t_1} B_{s_1, t_1}^{(i)}}{s_1} ds_1 \right| \right) \right] \\ & \left. + \mathbb{E} \left[\exp \left(\frac{3\zeta}{\delta(r, s)\delta(u, t)} \left| \int_r^s \int_{1-t}^{1-u} \hat{b}_{j,n}(s_1, t_1, W_{s_1, 1-t_1}) \frac{W_{s_1, 1-t_1}^{(i)}}{s_1(1-t_1)} dt_1 ds_1 \right| \right) \right] \right\} = \frac{1}{3} (I_1 + I_2 + I_3). \end{aligned}$$

By Jensen inequality,

$$\begin{aligned} I_1 &= \mathbb{E} \left[\exp \left(\frac{3\zeta}{\delta(r, s)\delta(u, t)} \left| \int_r^s \int_u^t \hat{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) \frac{d_{t_1} W_{s_1, t_1}^{(i)}}{s_1} ds_1 \right| \right) \right] \\ &\leq \int_r^s \mathbb{E} \left[\exp \left(\frac{6\zeta(\sqrt{s} - \sqrt{r})}{\delta(r, s)\delta(u, t)} \left| \int_u^t \hat{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) \frac{d_{t_1} W_{s_1, t_1}^{(i)}}{\sqrt{s_1}} \right| \right) \right] \frac{ds_1}{2\sqrt{s_1}(\sqrt{s} - \sqrt{r})} \\ &\leq \int_r^s \mathbb{E} \left[\exp \left(\frac{6\zeta}{\delta(u, t)} \left| \int_u^t \hat{b}_{j,n}(s_1, t_1, W_{s_1, t_1}) \frac{d_{t_1} W_{s_1, t_1}^{(i)}}{\sqrt{s_1}} \right| \right) \right] \frac{ds_1}{2\sqrt{s_1}(\sqrt{s} - \sqrt{r})} \end{aligned}$$

Since, for every $s_1 \in [r, s]$,

$$\left(Y_{s_1, t_1} := \int_u^{t_1} \hat{b}_{j,n}(s_1, t_2, W_{s_1, t_2}) \frac{d_{t_2} W_{s_1, t_2}^{(i)}}{\sqrt{s_1}}, u \leq t_1 \leq t \right)$$

is a square integrable martingale and similar reasoning as before gives

$$\mathbb{E}[|Y_{s_1, t}|^m] \leq 2^m c_1^{2m} \tilde{M}^m m^m \delta(u, t)^m.$$

From this and the exponential expansion formula, we get

$$I_1 \leq \sum_{m=0}^{\infty} \frac{2^{4m} \zeta^m c_1^{2m} m^m \tilde{M}^m}{m!} := \tilde{C}_{1,1},$$

which is finite if $\zeta < e/2^4 c_1^2 \tilde{M}$ (by ratio test). Similarly, we also have

$$I_2 \leq \tilde{C}_{1,1}.$$

To estimate I_3 we apply Jensen inequality again and we obtain

$$\begin{aligned}
I_3 &= \mathbb{E} \left[\exp \left(\frac{3\zeta}{\delta(r,s)\delta(u,t)} \left| \int_r^s \int_{1-t}^{1-u} \hat{b}_{j,n}(s_1, t_1, W_{s_1, 1-t_1}) \frac{W_{s_1, 1-t_1}^{(i)}}{s_1(1-t_1)} dt_1 ds_1 \right| \right) \right] \\
&\leq \int_r^s \int_{1-t}^{1-u} \mathbb{E} \left[\exp \left(\frac{12\zeta(\sqrt{s}-\sqrt{r})(\sqrt{1-u}-\sqrt{1-t})}{\delta(r,s)\delta(u,t)} \left| \hat{b}_{j,n}(s_1, t_1, W_{s_1, 1-t_1}) \right| \frac{W_{s_1, 1-t_1}^{(i)}}{\sqrt{s_1}\sqrt{1-t_1}} \right) \right] \\
&\quad \times \frac{dt_1}{2\sqrt{1-t_1}(\sqrt{1-u}-\sqrt{1-t})} \frac{ds_1}{2\sqrt{s_1}(\sqrt{s}-\sqrt{r})} \\
&\leq \int_r^s \int_{1-t}^{1-u} \mathbb{E} \left[\exp \left\{ 24\tilde{M}\zeta \left(1 + \frac{|W_{s_1, 1-t_1}^{(i)}|^2}{s_1(1-t_1)} \right) \right\} \right] \frac{dt_1}{2\sqrt{1-t_1}(\sqrt{1-u}-\sqrt{1-t})} \frac{ds_1}{2\sqrt{s_1}(\sqrt{s}-\sqrt{r})} := \tilde{C}_{1,2},
\end{aligned}$$

with $\tilde{C}_{1,2}$ finite provided that $24\tilde{M}\zeta < 1/2$.

The proof is completed by choosing $\zeta = 1/2^6 c_1^2 \tilde{M}$ and $\tilde{C}_1 = (2\tilde{C}_{1,1} + \tilde{C}_{1,2})/3$. \square

Proof of Lemma 3.12. Fix $\tau \in (0, \min\{\tau_1, \zeta/32d^2\})$, where τ_1 is the constant in Lemma A.1 and ζ is the constant in Lemma 3.11. We deduce from Theorem B.1 and the linear growth condition on the drift b_n that $\mathbb{E}[\|X_{s,t}^n\|^2] \leq C_{2,1}$ for all $(s, t) \in [0, \tau]^2$ and $n \geq 1$, where $C_{2,1}$ does not depend of (s, t) and n . Let $0 \leq r' \leq r \leq s \leq \tau$ and $0 \leq u' \leq u \leq t \leq \tau$. Since

$$\tau \leq \tau_1 \text{ and } \delta(r', s)\delta(u', t) \leq \tau \leq \frac{\zeta}{32d^2},$$

then, using similar computations as in the proof of Lemma 3.6, one can deduce from Lemma A.1, Girsanov theorem and Hölder inequality that

$$\sup_{n \geq 1} \sup_{\substack{0 \leq r \leq s \\ 0 \leq u \leq t}} \mathbb{E} \left[\|D_{r,u} X_{s,t}^n\|^2 \right] := C_{2,2} < \infty$$

and

$$\mathbb{E} \left[\|D_{r,u} X_{s,t}^n - D_{r',u'} X_{s,t}^n\|^2 \right] \leq C_{2,3} (|r - r'| + |u - u'|)$$

for some positive constant \tilde{C}_2 independent of n . The proof is completed by taking $C_2 = \max\{C_{2,1}, C_{2,2}, C_{2,3}\}$. \square

Remark A.3. *It is worth noting that if the drift is in addition the difference of two convex or concave functions, then the solution to the equation (1.3) is twice Malliavin differentiable. Indeed, $b = \hat{b} - \check{b}$ is Lipschitz with the second order weak derivatives of \hat{b}, \check{b} positive or negative.*

APPENDIX B. A GRONWALL TYPE INEQUALITY FOR FUNCTIONS OF TWO VARIABLES

The next result which is originally due to Wendroff, extends Gronwall inequality to functions of two variables (see e.g. [40, Theorem 5.1.1]).

Theorem B.1. *Let $g(s, t)$, $a(s, t)$, $k(s, t)$ be non-negative continuous functions for all $s \geq s_0$, $t \geq t_0$, and let $a(s, t)$ be non-decreasing in each of the variables for all $s \geq s_0$, $t \geq t_0$. Suppose that for all $s \geq s_0$, $t \geq t_0$,*

$$g(s, t) \leq a(s, t) + \int_{s_0}^s \int_{t_0}^t k(s_1, t_1) g(s_1, t_1) dt_1 ds_1.$$

Then for all $s \geq s_0$, $t \geq t_0$,

$$g(s, t) \leq a(s, t) \exp \left(\int_{s_0}^s \int_{t_0}^t k(s_1, t_1) dt_1 ds_1 \right).$$

REFERENCES

- [1] O. Amine, David Baños, and Frank Proske. C^∞ regularization by noise of singular ODE's. *arXiv preprint arXiv:1710.05760*, 2017.
- [2] O. Amine, A.-R. Mansouri, and F. Proske. Well-posedness of the deterministic transport equation with singular velocity field perturbed along fractional brownian paths. *arXiv preprint arXiv:2003.06200*, 2020.
- [3] M. T. Barlow and M. Yor. Semi-martingale inequalities via the garsia-rodemich-rumsey lemma, and applications to local times. *Journal of Functional Analysis*, 49(2):198–229, 1982.
- [4] L. Beck, F. Flandoli, M. Gubinelli, and M. Maurelli. Stochastic ODEs and stochastic linear PDEs with critical drift: regularity, duality and uniqueness. *Elect. J. Probab.*, 24, 2019.
- [5] A.-M. Bogso, M. Dieye, and O. Menoukeu Pamen. Path-by-path uniqueness of multidimensional sde's on the plane with nondecreasing coefficients. *Elect. J. Probab.*, 27: 1-26. DOI: 10.1214/22-EJP844, 2022.
- [6] A.-M. Bogso, M. Dieye, and O. Menoukeu Pamen. Stochastic integration with respect to local time of the brownian sheet and regularising properties of Brownian sheet paths. *Bernoulli (accepted)*, 2022.
- [7] O. Butkovsky and L. Mytnik. Regularization by noise and flows of solutions for a stochastic heat equation. *Annals of Probability*, 47:169–212, 2019.
- [8] E. Carlen and P. Kree. Lp estimates on iterated stochastic integrals. *The Annals of Probability*, pages 354–368, 1991.
- [9] R. Carmona and D. Nualart. Random non-linear wave equations: smoothness of the solutions. *Probability Theory and Related Fields*, 79(4):469–508, 1988.
- [10] R. Catellier and M. Gubinelli. Averaging along irregular curves and regularisation of ODEs. *Stochastic Process. Appl.*, 126:2323–2366, 2016.
- [11] G. D. Da Prato, P. Malliavin, and D. Nualart. Compact families of wiener functionals. *Comptes rendus de l'Académie des sciences. Série 1, Mathématique*, 315(12):1287–1291, 1992.
- [12] R. C. Dalang and C. Mueller. Multiple points of the brownian sheet in critical dimensions. *The Annals of Probability*, 43(4):1577–1593, 2015.
- [13] R. C. Dalang and J. B. Walsh. Time-reversal in hyperbolic S.P.D.E.'s. *The Annals of Probability*, 30(1):213–252, 2002.
- [14] A. M. Davie. Uniqueness of solutions of stochastic differential equations. *International Mathematics Research Notices*, Vol. 2007, 2007.
- [15] A. M. Davie. Individual path uniqueness of solutions of stochastic differential equations. In *Stochastic analysis 2010*, pages 213–225. Springer, 2011.
- [16] K. Deimling. A Carathéodory theory for systems of integral equations. *Annali di Matematica Pura ed Applicata*, 86(1):217–260, 1970.
- [17] E. Di Nezza, G. Palatucci, and E. Valdinoci. Hitchhiker's guide to the fractional sobolev spaces. *Bulletin des sciences mathématiques*, 136(5):521–573, 2012.
- [18] M. Farré and D. Nualart. Nonlinear stochastic integral equations in the plane. *Stochastic processes and their applications*, 46(2):219–239, 1993.
- [19] F. Flandoli. *Random Perturbation of PDE's and Fluid Dynamic Models: École d'été de Probabilités de Saint-Flour XL 2010*, volume 2015. Springer Science & Business Media, 2010.
- [20] Folland, G. B. (1999). *Real analysis: modern techniques and their applications*, volume 40. John Wiley & Sons.
- [21] L. Galeati and M. Gubinelli. Noiseless regularisation by noise. *Revista Matemática Iberoamericana*, 38(2):433–502, 2021.
- [22] S. Haadem and F. Proske. On the construction and malliavin differentiability of solutions of levy noise driven sde's with singular coefficients. *Journal of functional analysis*, 266(8):5321–5359, 2014.
- [23] F. A. Harang and N. Perkowski. C^∞ regularization of ODEs perturbed by noise. *Stochastics and Dynamics*, 21(08):2140010, 2021.
- [24] Tosio Kato. *Perturbation theory for linear operators*, volume 132. Springer Science & Business Media, 2013.

- [25] T. Kitagawa. Analysis of variance applied to function spaces. *Memoirs of the Faculty of Sciences, Kyusyu University*, VI(1):41–53, 1951.
- [26] D. Khoshnevisan. *Multiparameter Processes: an introduction to random fields*. Springer Science & Business Media, 2002.
- [27] H. Kremp and N. Perkowski. Multidimensional SDE with distributional drift and Lévy noise. *Bernoulli*, 28(3):1757–1783, 2022.
- [28] R. T. Lewis. Singular elliptic operators of second order with purely discrete spectra. *Transactions of the American Mathematical Society*, 271(2):653–666, 1982.
- [29] O. Menoukeu-Pamen and L. Tangpi. Strong solutions of some one-dimensional SDEs with random and unbounded drifts. *SIAM J. Appl. Math.*, 51:4105–4141, 2019.
- [30] Olivier Menoukeu-Pamen, Thilo Meyer-Brandis, Torstein Nilssen, Frank Proske, and Tusheng Zhang. A variational approach to the construction and Malliavin differentiability of strong solutions of SDE’s. *Mathematische Annalen*, 357(2):761–799, 2013.
- [31] T. Meyer-Brandis and F. Proske. Construction of strong solutions of SDE’s via Malliavin calculus. *Journal of Functional Analysis*, 258(11):3922–3953, 2010.
- [32] D. Nualart. *The Malliavin calculus and related topics*, volume 1995. Springer, 2006.
- [33] D. Nualart and E. Pardoux. Markov field properties of solutions of white noise driven quasi-linear parabolic pdes. *Stochastics: An International Journal of Probability and Stochastic Processes*, 48(1-2):17–44, 1994.
- [34] D. Nualart and M. Sanz. Malliavin calculus for two-parameter wiener functionals. *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete*, 70(4):573–590, 1985.
- [35] D. Nualart and M. Sanz. Stochastic differential equations on the plane: smoothness of the solution. *Journal of multivariate analysis*, 31(1):1–29, 1989.
- [36] D. Nualart and S. Tindel. Quasilinear stochastic hyperbolic differential equations with nondecreasing coefficient. *Potential Analysis*, 7(3):661–680, 1997.
- [37] D. Nualart and J. Yeh. Existence and uniqueness of a strong solution to stochastic differential equations in the plane with stochastic boundary process. *Journal of Multivariate Analysis*, 28(1):149–171, 1989.
- [38] G. Palatucci, O. Savin, and E. Valdinoci. Local and global minimizers for a variational energy involving a fractional norm. *Annali di matematica pura ed applicata*, 192(4):673–718, 2013.
- [39] F. Proske. Stochastic differential equations—some new ideas. *Stochastic: An international Journal of Probability and Stochastic Processes*, 79:563–600, 2007.
- [40] Y. Qin. *Integral and Discrete Inequalities and Their Applications*. Springer, 2016.
- [41] L. Quer-Sardanyons and S. Tindel. The 1-d stochastic wave equation driven by a fractional brownian sheet. *Stochastic processes and their applications*, 117(10):1448–1472, 2007.
- [42] D. L. Rasmussen. Gronwall’s inequality for functions of two independent variables. *Journal of Mathematical Analysis and Applications*, 55(2):407–417, 1976.
- [43] A. V. Shaposhnikov. Some remarks on Davie’s uniqueness theorem. *Proceedings of the Edinburgh Mathematical Society*, 59(4):1019–1035, 2016.
- [44] D. R. Snow. Gronwall’s inequality for systems of partial differential equations in two independent variables. *Proceedings of the American Mathematical Society*, 33(1):46–54, 1972.
- [45] C. Tudor. On the two parameter Itô equations. *Publications mathématiques et informatique de Rennes*, (1):1–17, 1983.
- [46] J. Yeh. Cameron-Martin translation theorems in the Wiener space of functions of two variables. *Transactions of the American Mathematical Society*, 107(3):409–420, 1963.
- [47] J. Yeh. Uniqueness of strong solutions to stochastic differential equations in the plane with deterministic boundary process. *Pacific journal of mathematics*, 128(2):391–400, 1987.

UNIVERSITY OF YAOUNDE I, FACULTY OF SCIENCES, DEPARTMENT OF MATHEMATICS, P.O. Box 812, YAOUNDE, CAMEROON, AND
AFRICAN INSTITUTE FOR MATHEMATICAL SCIENCES GHANA, P.O. Box LGDTD 20046, SUMMERHILL ESTATES, EAST LEGON HILLS,
SANTOE, ACCRA

Email address: antoine.bogso@facsciences-uy1.cm, antoine@aims.edu.gh

INSTITUTE FOR FINANCIAL AND ACTUARIAL MATHEMATICS (IFAM), DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF
LIVERPOOL, LIVERPOOL L69 7ZL, UK, AND AIMS GHANA

Email address: menoukeu@liverpool.ac.uk