

**A transdiagnostic investigation of ‘theory of mind’ and ‘jumping to conclusions’ in patients with persecutory delusions.**

Running Head: Transdiagnostic processes in persecutory delusions.

Word count: 2923.

Rhiannon Corcoran, <sup>1a</sup>

Georgina Rowse, <sup>4</sup>

Rosie Moore, <sup>3</sup>

Nigel Blackwood, <sup>3</sup>

Peter Kinderman, <sup>5</sup>

Robert Howard, <sup>2</sup>

Sinead Cummings, <sup>2</sup>

Richard P. Bentall, <sup>2</sup>

<sup>a</sup> to whom correspondence should be addressed.

<sup>1</sup> Division of Psychiatry, School of Community Health Sciences, University of Nottingham, A Floor, South Block, Queens Medical Centre, Nottingham NG7 2UH.

<sup>2</sup> School of Psychological Sciences, University of Manchester, Oxford Road, Manchester M13 9PL, United Kingdom.

<sup>3</sup> Section of Old Age Psychiatry, Institute of Psychiatry, Kings College London, DeCrespigny Park, London SE5

<sup>4</sup> Department of Clinical Psychology, University of Sheffield

<sup>5</sup> Department of Clinical Psychology, University of Liverpool

## **Abstract.**

Background: A tendency to make hasty decisions on probabilistic reasoning tasks and a difficulty attributing mental states to others are key cognitive features of persecutory delusions (PD) in the context of schizophrenia. This study examines whether these same psychological anomalies characterise PD when they present in the context of psychotic depression.

Method: Performance on measures of probabilistic reasoning and theory of mind were examined in 5 subgroups differing in diagnostic category and current illness status.

Results: The tendency to draw hasty decisions in probabilistic settings and poor theory of mind (ToM) tested using story format feature in PD irrespective of diagnosis.

Furthermore, performance on the ToM story task correlated with the degree of distress caused by and preoccupation with the current PD in the currently deluded groups. By contrast, performance on the non-verbal ToM task appears to be more sensitive to diagnosis as patients with schizophrenia spectrum disorders perform worse on this task than those with depression irrespective of the presence of PD.

Conclusions: The psychological anomalies associated with PD examined here are transdiagnostic but different measures of ToM may be more or less sensitive to indices of severity of the PD, diagnosis and trait- or state- related cognitive effects.

## **Introduction.**

Psychopathological research conducted over the past 2 decades has undermined the view that delusions are un-understandable (e.g. Bentall, et al. 2001; Blackwood, et al., 2001). In particular there are two key reasoning anomalies that have been found in adults with delusions. First, people with PD have theory of mind (ToM) difficulties in that they have trouble accurately inferring other people's beliefs and intentions (e.g. Corcoran et al., 1995; Corcoran and Frith, 2003; Harrington et al., 2005). Secondly, deluded individuals have a tendency to jump to conclusions (JTC) on the basis of insufficient information when reasoning in probabilistic contexts (e.g. Huq et al., 1988; Garety et al., 1991; Dudley et al., 1997). This characteristic style of decision-making seems to apply to delusions irrespective of content and, as Freeman (2007) notes, there has been little research exploring the JTC bias in relation to PD specifically. The balance of evidence suggests that the JTC finding is robust and has been found to relate particularly to delusional conviction (e.g. Garety and Freeman, 1999; Peters and Garety, 2006; Garety et al. 2005). This contrasts with the findings on ToM reasoning which are not so well replicated but which are theoretically compelling in their account of the psychological origins of PD in particular (for review see Harrington et al., 2005). While it is clear that neither of these reasoning anomalies can comprehensively account for the generation and maintenance of PD (see for example Bentall et al., in press) it seems likely that they both contribute to paranoid ideation.

The vast majority of the studies conducted to date on these psychological variables have focussed on schizophrenia as a diagnostic category. However, the context within which psychological models of delusional processes have grown over the last 15 years or so

has been one which emphasises the symptoms of psychosis over the diagnosis of a psychotic illness. This has been driven by compelling theoretical reasons which question the utility of methods relying on data gathered from groups of patients for whom the diagnosis is often the only shared characteristic. Instead, psychologists have argued that the best way to advance understanding of psychosis is to take a symptom-based approach. Indeed, it is clear that just such an approach has led to significant advances in our understanding of the psychological basis of psychotic symptoms (see for example Frith, 1992; Bentall 2003). Of course, a strict symptom-based approach if correct, implies that the psychological bases of a psychotic symptom should be the same irrespective of the diagnostic context in which it is seen and PD are found in several other diagnostic groups including affective disorder. The study reported below is the first to investigate whether the drawing of hasty decisions in probabilistic settings and poor theory of mind are seen in PD across diagnostic categories.

## **Method.**

### Participants.

115 patients in four groups were recruited from in-patient and out-patient clinics in South London and the North West of England. Informed consent was obtained using procedures agreed by a Multi-Centre Research Ethics Committee. The *currently paranoid group* consisted of 26 male and 13 female patients with a DSM-IV diagnosis of schizophrenia, schizoaffective disorder or delusional disorder who were experiencing PD as assessed by the endorsement of the Schedules for Clinical Assessment in Neuropsychiatry (SCAN; World Health Organization, 1997) item 19.012 (Does anyone seem to be trying to harm you (trying to poison or kill you?)) at interview, examination

of case notes and endorsement of item 4 ('Do you ever feel as if you are being persecuted in some way?') on the Peters et al. Delusions Inventory (Peters et al., 1999). They had a mean age of 33.95 years (SD = 8.38), a mean IQ assessed by the Wechsler Abbreviated Scale of Intelligence (WASI, Wechsler, 1999) of 92.18 (SD = 10.74). Their mean age of illness onset was 23.00 years (SD = 7.33), and they were in receipt of a mean antipsychotic dosage of 417.13 mg chlorpromazine (CPZ) equivalent (SD = 510.51). The mean number of other positive symptoms present in this group was 0.85 (SD=0.90).

The *remitted paranoid group* consisted of 18 male and 11 female schizophrenia spectrum patients who had experienced PD in the past as evidenced by endorsement of SCAN item 19.012 for a representative illness period, examination of case notes and endorsement of item 4 on the PDI for the representative period identified by the SCAN. However, participants in this group indicated that they did not currently feel paranoid and that they had not suffered from PD for at least the past fortnight. The current status of these patients was confirmed by case note examination and / or by clinical staff. Patients in this group did report the presence of other positive symptoms, notably auditory hallucinations. Their mean age was 34.66 years (SD = 10.35); their mean IQ was 98.38 (SD = 13.17); their mean age of illness onset was 21.82 years (SD = 5.81) and they were in receipt of a mean of 227.78 mg CPZ equivalent of antipsychotic medication (SD = 261.14). The mean number of other positive symptoms present in this group was 0.72 (SD=0.96).

A *paranoid depressed group* consisted of 11 male and 9 female patients with a primary DSM-IV diagnosis of major depression, currently experiencing persecutory

delusions as indicated by the SCAN. They had a mean age of 36.00 years (SD = 10.11), a mean IQ of 98.50 (SD = 15.75), a mean age of illness onset of 23.93 years (SD = 11.63) and were in receipt of a mean of 17.50 mg CPZ equivalent of antipsychotic medication (SD = 37.36). The mean number of other positive symptoms present in this group was 1.05 (SD= 0.99).

No significant differences existed in terms of the number of other positive symptoms assessed to be present by the SCAN between the three psychotic patient groups. (i.e. depersonalisation, hallucinations, perceptual experiences;  $F(2,85)= 0.706$ , ns). The presence of clear negative signs of psychosis (i.e. underactivity, slow speech, poverty of speech or blunted affect as assessed by the SCAN) or overt formal thought disorder were exclusion criteria for the three psychotic patient groups.

A *non-psychotic depressed group* consisted of 9 male and 18 female patients with a diagnosis of major depression and not experiencing PD. Their mean age was 48.37 years (SD = 10.97), their mean IQ was 108.22 (SD = 12.99), their mean age of onset was 29.11 years (SD = 14.45), and they were in receipt of a mean dose of 5.55 mg CPZ equivalent of antipsychotic medication (SD = 28.87).

A *healthy adult group* of 14 males and 19 females, recruited by advertisement, had a mean age of 39.03 years (SD = 13.96), and a mean IQ of 108.64 (SD = 14.48). The initial screening questions from the SCAN were used to determine that they were symptom free.

### Measures.

A version of the probabilistic reasoning task developed by Phillips and Edwards (1966), commonly known as the 'beads in a jar task' was used to investigate the tendency to 'jump to conclusions' when considering neutral stimuli. This task was

originally used to investigate probabilistic reasoning in deluded patients by Huq et al. (1988) and has enjoyed widespread use since (eg. Garety *et al.*, 2005) The proportion of the two coloured beads in the jars in this version was 60:40 with the majority colour being yellow in one jar and black in the other. The participants were shown the order in which beads were drawn from one of the two jars until they felt confident that they knew which of the jars the beads were being drawn from. The number of beads requested before reaching a decision and the jar chosen were recorded. Each participant was shown 3 sequences of black and yellow beads which had been randomly determined prior to commencement of the study and which were held constant across participants.

A social version of the 'beads in a jar' task developed by Dudley et al. (1997) was used as a second measure of the tendency to jump to conclusions when considering more salient information. In this task participants were presented with 2 surveys containing 100 comments about a person. They were told that survey A contained 60 negative and 40 positive comments, while survey B contained 60 positive and 40 negative comments. Participants could see as many comments as they needed before deciding which of the 2 surveys they came from. The number of comments requested before reaching a decision and the survey chosen were recorded. The same 3 random sequences used as for the 'beads' task were used for each participant.

Six theory of mind stories developed by Snowden, Thompson, Corcoran, Gibbons and Doubleday (unpublished) were used to assess understanding of false belief and deception to second order level and to provide measures of narrative memory and non-mental state inference capacity. These stories have recently been used by Moore et al. (2006) in a study examining PD in people with very late onset schizophrenia-like psychosis and are similar in format to those previously used by Frith and Corcoran (1996) except that they develop from first (when the contents of a single person's mind

must be inferred) to second order ToM (where the contents of two people's minds must be inferred) within each story. The stories were read out to participants accompanied by a set of cartoon pictures depicting events in the story. At set points during the course of the story participants were asked questions to assess theory of mind, narrative memory and non-mental state inference skills which examined the ability to draw inferences from the text beyond what was actually stated but which did not involve an inference about the contents of the character's minds.

Finally, Langdon and colleagues' (1997) false belief picture sequencing task was used as a non-verbal measure of theory of mind. Following two practice sequences, participants were presented with 4 stories involving a person acting on a false belief. The cards were presented in a fixed random order and participants were instructed to put the cards into the right order so that they would tell a story. The standard scoring system devised by Langdon et al. (1997) was used.

#### Analysis Strategy.

In the first instance analysis of variance with planned contrasts was used to test for group differences on the four reasoning tasks. The first contrast compared groups with current PD to those without. Contrast 2 compared the performance of patients with PD in the context of schizophrenia spectrum to those with PD in the context of depression. Contrast 3 compared the performance of the remitted paranoid patients to that of the non-psychotic depressed patients and the healthy adults and the final contrast compared the performance of the non-psychotic depressed patients to the healthy adults. It was expected that contrast one would produce a statistically significant result while contrasts 2-4 would prove to be non-significant. Preliminary analyses prior to ANOVA showed that the beads in the jar task and the ToM stories task were associated with

between group heterogeneity of variance (Levene's statistic beads (4,140) = 2.403,  $p = .053$ ; Levene's statistic ToM stories (4, 139) = 3.014,  $p = .02$ ). Therefore, the Brown-Forsythe robust test for the equality of means (B-F) will be reported for the 4 experimental tasks as it is recommended for use with ANOVA in preference to the F statistic when variances are unequal between groups and where the data is skewed as is the case with all 4 tasks (Clinch and Keselman, 1988).

Preliminary analyses of the data indicated that the assumptions with respect to the use of linear regression were not violated with plots of the residuals indicating that these were normally distributed for all 4 measures and the Cook's distance statistics indicating that outliers were not exerting an undue influence on regression equations. Collinearity statistics (tolerance and VIF) were all acceptable. Therefore a series of linear regressions using the enter method are reported which investigate possibly confounding factors such as between group differences in IQ, age and gender. For these analyses three dummy variables were created to enable clinical status to be considered as predictor variables. These were whether or not each case was currently paranoid (paranoid schizophrenia + paranoid depressed versus the other groups), whether or not each case had ever been paranoid (paranoid schizophrenia+ paranoid depressed+ remitted paranoid versus the other groups) and whether or not each case was currently depressed (paranoid depressed + depressed versus the other groups). Other predictor variables were maleness, age, and IQ . Additionally, for the theory of mind stories, the percent correct on the narrative memory questions and the percent correct on the non-mental state inference questions were also included.

Finally separate correlation analyses were conducted to examine the relationship between performance on the tasks, antipsychotic medication dosage (expressed as

chlorpromazine equivalents), and the conviction, distress and preoccupation scales of the PDI (each measured on 1-5 scales) in the groups for whom these variables are relevant.

The performance of each group on the measures and their background demographics can be seen in Table One.

## **Results.**

### **Jumping to Conclusions.**

Performance on the two measures of probabilistic reasoning correlated significantly in these participants ( $r=.73$ ,  $p<.001$ ).

The ANOVA findings relating to performance on the beads in the jar task were consistent with the hypothesis with a significant overall effect for group ( $B-F(4,97.9) = 4.436$ ,  $p<.005$ ). The planned contrasts indicated that contrast 1 was significant at the  $p<.001$  level indicating that the groups with current PD tended to draw conclusions more hastily on this task compared to the other groups. As expected contrasts 2-4 were non-significant for this task. The regression analyses for the beads task was significant ( $r^2 = .16$  ;  $F(6,137) = 4.27$ ,  $p<.001$ ). The significant predictors of performance on this task were age ( $\beta = -.19$ ;  $t=-2.22$ ;  $p<.03$ ) and currently paranoid status ( $\beta = -.35$ ;  $t=-3.1$ ;  $p<.003$ ).

The results of the social probabilistic task were consistent with the beads task but not as compelling. There was an overall trend indicating between group differences using ANOVA ( $B-F(4,103.39) = 1.99$ ,  $p=.1$ ) Mean scores clearly indicated that those with current PD were doing less well on this task and this was supported by a significant result for planned contrast 1 ( $p<.01$ ). As expected contrasts 2-4 were non-significant. The regression analyses for this task was significant ( $r^2 = .13$  ;  $F(6,136) = 3.49$ ;  $p<.005$ )

with IQ significantly predicting performance ( $\beta = .24$ ;  $t=2.6$ ;  $p<.02$ ) and current paranoid status just failing to reach significance at  $p<.05$  ( $\beta = -.20$ ;  $t=-1.77$ ;  $p<.08$ ).

### Theory of Mind.

Performance on the two measures of theory of mind correlated significantly in these participants ( $r=.41$ ,  $p<.001$ ).

The ANOVA findings relating to performance on theory of mind stories task were consistent with the hypothesis with a significant overall effect for group (B-F(4,127.96) = 4.9,  $p<.002$ ). The planned contrasts indicated that contrast 1 was significant at the  $p<.001$  level indicating that the groups with current PD tended to score lower on this task than the other groups. As predicted contrasts 2-4 were non-significant. The regression analyses for this task was significant ( $r^2 = .35$  ; F(8,134) = 9.01,  $p<.001$ ). The significant predictors of performance were IQ ( $\beta = -.28$ ;  $t=3.32$ ;  $p<.002$ ), current paranoid status ( $\beta = -.25$ ;  $t=-2.49$ ;  $p<.02$ ) and performance on the narrative memory questions within the task ( $\beta = .37$ ;  $t=4.42$ ;  $p<.001$ ).

The results of the false belief picture sequencing task were rather different to those of the stories task. While for this task the overall ANOVA was again significant (B-F(4,132.68) = 2.92,  $p<.025$ ), this was accounted for by contrast 3 ( $p<.02$ ) indicating that the remitted paranoid patients performed worse on this task than the depressed patients and the healthy adults. Contrasts 1,2 and 4 were all non-significant for this task. The regression analyses for this task was also significant ( $r^2 = .25$  ; F(6,140) = 7.83;  $p<.001$ ) with age ( $\beta = -.23$ ;  $t=-2.82$ ;  $p<.01$ ), IQ ( $\beta = .39$ ;  $t = 4.71$ ;  $p<.001$ ) predicting performance on this task. Having a history of paranoid delusions just failed to significantly predict performance on this task ( $\beta = -.214$ ;  $t=-1.90$ ;  $p=.059$ ) and no other independent variables proved to be significant predictors.

### Correlation of task performance with medication dosage, conviction, preoccupation and distress as measured by the PDI.

A summary of these findings can be found in Table Two.

Medication dosage expressed as chlorpromazine equivalents was only considered for the groups with current psychosis or a history of psychosis as the other groups were not taking antipsychotic medication. No significant correlations were found between performance on any of the four experimental tasks and medication dosage.

The conviction, preoccupation and distress scales of the PDI were considered in the groups who were currently expressing PD. As shown in Table Two, there were no significant correlations between any of these PDI severity scales and performance on the beads task, the social version of the beads task or the picture sequencing task. Modest but significant correlations were found between performance on the ToM stories and the preoccupation and distress sub-scales but not the conviction subscale.

### **Discussion.**

These results replicate previous findings showing that people with PD in the context of schizophrenia spectrum disorders tend to make hasty decisions on probabilistic reasoning tasks (Garety and Freeman, 1999) and have difficulty inferring the beliefs and intentions of others (eg. Frith and Corcoran, 1996). This study is however the first to demonstrate that both of these cognitive tendencies are transdiagnostic features of persecutory delusions. The second contrast of the planned comparisons analyses consistently failed to demonstrate any differences in performance across the tasks between the patients with PD in the context of schizophrenia spectrum disorders and those with PD in the context of affective disorder. Thus, the finding

indicates that the same level of difficulty on these tasks exists in patients with affective disorder as exists in patients with schizophrenia spectrum disorders. Recently, Moore et al. (2006) have shown that theory of mind difficulties but not the tendency to jump to conclusions was a feature of PD in the context of very late onset schizophrenia-like psychosis. The authors accounted for the null finding on the JTC task by arguing that any difference was over-ridden by general age-related changes in higher order cognition as their comparison groups were matched for age with their psychosis group.

The regression analyses demonstrated that performance on these measures is influenced by demographic variables (in particular age) and cognitive ability. However, the specific transdiagnostic associations with PD survived the effects of these confounding variables. For 3 of the 4 tests examined (the 2 probabilistic reasoning tasks and the theory of mind stories), abnormal performance was associated with current delusional state indicating a clear state-related cognitive dysfunction associated with PD irrespective of diagnosis. While it is notable that the performance on the JTC task did not correlate with the delusional conviction component of the PDI as has been previously reported (e.g. Garety *et al.*, 2006), the fact that the performance on the ToM stories task correlated significantly with the PDI subscales of preoccupation and distress supports the state-related nature of performance on this verbal ToM task. However, for the false belief picture sequencing task there was some evidence to support a trait effect both in the ANOVA where the only significant contrast showed that the remitted paranoid patients performed this task worse than the non-psychotic depressed and the healthy adults. Marginal support for this finding was seen in the regression analyses where having a history of PD just failed to reach significance at the 0.05 level as a predictor of performance on this task. The idea that the nature of the ToM dysfunction

in schizophrenia can be distinguished in terms of state and trait requires further exploration as it could resolve inconsistent findings in the earlier literature exploring this matter (see Harrington et al.,2005). It is possible that the non-verbal nature of this task makes it less sensitive to any further deleterious effects of acute episodes of PD. Furthermore, there is evidence in the data that the false belief picture sequencing task may be more sensitive to diagnosis than to the presence of PD per se. Examination of the data provide in Table One shows that on this task the remitted paranoid and the currently paranoid patients with a diagnosis of schizophrenia are both, in fact, performing poorly and at the same level while the performance of the two depressed groups is also comparable and somewhat better than that seen in the context of schizophrenia. It may be that this task is sensitive to diagnosis because of its strong sequencing and problem-solving components which uncover more general cognitive difficulties associated with the diagnosis of schizophrenia. Had we included the logical sequencing stories that Langdon et al. devised as part of the Picture Sequencing task, we could have explored this issue further. Unfortunately constraints of time and the consideration of patient's tolerance to the battery dictated the exclusion of this measure in this study.

The fact that the probabilistic reasoning task findings were clearly state related in this study with no evidence in support of trait effects on this task does not support other research where authors have reported that hasty decisions are also taken by participants who are vulnerable to psychosis (Van Dael et al., 2006, Freeman , 2007). It is unclear why these findings differ but it may reflect the different methodologies used.

## **Conclusions.**

This study demonstrates that the same underlying reasoning difficulties that are associated with PD in the context of schizophrenia spectrum disorders are also associated with PD in the context of depression. While it is clear that the tendency to JTC is related to the presence of current delusions (i.e. a state effect), there is evidence in this study to suggest that false belief picture sequencing may be more sensitive to diagnosis and thus may be indicative of a trait - related cognitive features of paranoid schizophrenia. Thus it seems that the generic theory of mind skill may be a cognitive marker of paranoid schizophrenia which undergoes further decline during acute episodes. It is possible that verbal ToM tasks may be more sensitive to this state related decline than non-verbal ToM tasks. As the diagnostic groups were well categorised in this study and since no differences existed for the presence of other signs and symptoms of psychosis, these transdiagnostic findings relating to PD are compelling.

#### Declaration of Interest.

The work reported here was undertaken with the financial support of an award provided by the Wellcome Trust given to Dr. Corcoran as the PI and the other authors as co-applicants (Grant number: GR064134MA). There are no conflicts of interest associated with this research.

## **References.**

**Bentall, R.P.** (2003) *Madness Explained. Psychosis and Human Nature*. Penguin Books: London.

**Bentall, R.P., Corcoran, R., Howard, R., Blackwood, N. & Kinderman, P.** (2001). Persecutory delusions: A review and theoretical integration. *Clinical Psychology Review* **21**, 1143-1192.

**Bentall, R.P., Corcoran, R., Rowse, G., Moore, R., Cummins, S., Blackwood, N., Howard and Kinderman, P.** Paranoid delusions in schizophrenia and depression: The transdiagnostic role of expectations of negative events and negative self-esteem. *Journal of Nervous and Mental Disease*, in press.

**Blackwood, N.J., Howard, R.J. , Bentall, R.P. & Murray, R.M.** (2001). Cognitive Neuropsychiatric Models of Persecutory Delusions. *American Journal of Psychiatry* **158**, 527-539.

**Clinch, J.J. & Keselman, H.J.** (1988) Parametric alternatives to the analysis of variance. *Journal of Educational Statistics*, **7**, 207-214

**Corcoran, R. & Frith, C.D.** (2003). Autobiographical memory and theory of mind: evidence of a relationship in schizophrenia. *Psychological Medicine* **33**, 897-905.

**Corcoran, R., Frith, C.D. & Mercer, G.** (1995). Schizophrenia, symptomatology and social inference: investigating 'theory of mind' in people with schizophrenia. *Schizophrenia Research* **17**, 5–13

**Dudley, R.E.J., John, C.H., Young, A.W. & Over, D.E.** (1997). Normal and abnormal reasoning in people with delusions. *British Journal of Clinical Psychology* **36**, 243-258.

**Freeman, D.** (2007) Suspicious minds: The psychology of persecutory delusions. *Clinical Psychology Review*, **27**, 425-457.

**Garety, P.A., Hemsley, D.R. & Wessely, S.** (1991). Reasoning in deluded schizophrenic and paranoid patients. *Journal of Nervous and Mental Disease* **179**, 194-201.

**Garety, P.A. & Freeman, D.** (1999). Cognitive approaches to delusions: a critical review of theories and evidence. *British Journal of Clinical Psychology* **38**, 113-154.

**Garety, P.A., Freeman, D., Jolley, S., Dunn, G., Bebbington, P.E., Fowler, D.G., Kuipers, E. & Dudley, R.** (2005). Reasoning, Emotions, and Delusional Conviction in Psychosis *Journal of Abnormal Psychology*, **114**, 373-384.

**Harrington, L., Langdon, R., Siegert, R.J. & McClure, J.** (2005). Schizophrenia, theory of mind, and persecutory delusions. *Cognitive Neuropsychiatry*, **10**, 87-104.

**Harrington, L., Siegert, R.J. & McClure, J.** (2005). Theory of Mind in schizophrenia: a critical review. *Cognitive Neuropsychiatry*,

**10**, 249-286.

**Huq, S.F., Garety, P.A. & Hemsley, D.R.** (1988). Probabilistic judgements in deluded and nondeluded subjects. *Quarterly Journal of Experimental Psychology*,

**40A**, 801-812.

**Langdon, R. ; Mitchie, P.T.; Ward, P.B.; McConaghy, N; Catts, S.V. & Coltheart, M** (1997). Defective self and/or other mentalising in schizophrenia: A cognitive neuropsychological approach. *Cognitive Neuropsychiatry*, **2**, 167-193.

**Moore, R., Blackwood, N., Corcoran, R., Rowse, G., Kinderman, P., Bentall, R. & Howard, R.** (2006). Misunderstanding the Intentions of Others: An Exploratory Study of the Cognitive Etiology of Persecutory Delusions in Very Late-Onset Schizophrenia-Like Psychosis. *American Journal of Geriatric Psychiatry*, **14**, 410-418.

**Peters, E.R. & Garety, P.** (2006). Cognitive functioning in delusions: A longitudinal analysis. *Behaviour Research and Therapy*, **44**, 481-514.

**Peters, E.R., Joseph, S.A. & Garety, P.** (1999). Measurement of delusional ideation in the normal population: Introducing the PDI *Schizophrenia Bulletin*, **25**, 553-576.

**Phillips, L.D. & Edwards, W.** (1966). Conservatism in a simple probability inference task. *Journal of Experimental Psychology*, **72**, 346-354.

**VanDael,F.,Versmissen,D.,Janssen,I.,MyinGermeys,I.,vanOs,J.,&Krabbendam, L.**(2006).Data gathering: Biased in psychosis? *Schizophrenia Bulletin*, **32**, 341-351.

**Wechsler, D.** (1999). *Wechsler Abbreviated Scale of Intelligence*. San Antonio, TX: The Psychological Corporation.

**World Health Organization** (1997). *Schedules for clinical assessment in neuropsychiatry, SCAN* version 2. Geneva: WHO.

**Table One: demographic characteristics of the groups and their performance on the cognitive measures.**

	Currently paranoid (schiz)	Remitted paranoid (schiz)	Currently paranoid (dep)	Non— psychotic dep	Healthy adult	P level
Age	33.95 (8.4)	34.7 (10.3)	36.0 (10.0)	48.4 (11.0)	39.0 (14.0)	<.001
Sex (% male)	67%	62%	55%	33%	42%	<.05
Current IQ	92.2 (10.7)	98.5 (13.2)	98.5 (15.7)	108.2 (13.0)	108.6 (14.5)	<.001
Beads task mean draws to decision	3.1 (2.7)	5.7 (4.6)	3.95 (1.9)	5.1 (2.2)	5.6 (3.2)	<.005
Social task mean draws to decision	3.3 (2.4)	4.8 (4.6)	3.4 (2.1)	5.0 (4.0)	4.9 (3.1)	<.1
Theory of mind stories mean % score	78.4 (17.3)	86.0 (15.1)	82.9 (11.6)	90.7 (11.1)	90.2 (11.1)	<.002
Memory	94.0 (9.2)	92.0 (11.9)	95.4 (6.5)	97.8 (3.8)	98.7	<.006

mean % score					(3.0)	
Non- mental state inference % score	85.0 (12.4)	86.6 (12.5)	86.6 (13.7)	91.7 (11.3)	96.2 (5.9)	<.001
False belief picture sequencing score	4.4 (1.4)	4.4 (1.3)	4.9 (1.3)	5.0 (1.2)	5.3 (1.2)	<.025

Table Two. Correlations between task performance, medication dosage and the subscales of the PDI.

	Medication dosage	PD conviction	PD preoccupation	PD distress
Probabilistic reasoning task performance	.046 (ns)	.054 (ns)	-.035 (ns)	-.16 (ns)
Social probabilistic reasoning task performance	-.095 (ns)	-.129 (ns)	-.15 (ns)	-.11 (ns)
ToM stories performance	-.064 (ns)	.197 (ns)	.337 (p=.01)	.298 (p=.024)
False belief picture sequencing performance	-.176 (ns)	.058 (ns)	-.001 (ns)	-.138 (ns)