



Cochrane
Library

Cochrane Database of Systematic Reviews

Genetic testing for prevention of severe drug-induced skin rash (Protocol)

Alfirevic A, Pirmohamed M, Marinovic B, Jorgensen AL, Harcourt-Smith L

Alfirevic A, Pirmohamed M, Marinovic B, Jorgensen AL, Harcourt-Smith L.
Genetic testing for prevention of severe drug-induced skin rash.
Cochrane Database of Systematic Reviews 2013, Issue 12. Art. No.: CD010891.
DOI: 10.1002/14651858.CD010891.

www.cochranelibrary.com

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
BACKGROUND	1
Figure 1.	2
OBJECTIVES	3
METHODS	3
ACKNOWLEDGEMENTS	5
REFERENCES	6
ADDITIONAL TABLES	9
APPENDICES	11
WHAT'S NEW	13
CONTRIBUTIONS OF AUTHORS	13
DECLARATIONS OF INTEREST	14
SOURCES OF SUPPORT	14

[Intervention Protocol]

Genetic testing for prevention of severe drug-induced skin rash

Ana Alfirevic¹, Munir Pirmohamed¹, Branka Marinovic², Andrea L Jorgensen³, Linda Harcourt-Smith⁴

¹Department of Molecular and Clinical Pharmacology, Institute of Translational Medicine, University of Liverpool, Liverpool, UK.

²Department of Dermatology and Venereology, University Hospital Center Zagreb, School of Medicine University of Zagreb, Zagreb, Croatia. ³Centre for Medical Statistics and Health Evaluation, University of Liverpool, Liverpool, UK. ⁴c/o Cochrane Skin Group, The University of Nottingham, Nottingham, UK

Contact address: Ana Alfirevic, Department of Molecular and Clinical Pharmacology, Institute of Translational Medicine, University of Liverpool, Centre for Personalised Medicine, Block A: Waterhouse Building, 1-5 Brownlow Street, Liverpool, L69 3GE, UK. Ana.Alfirevic@liverpool.ac.uk.

Editorial group: Cochrane Skin Group.

Publication status and date: New, published in Issue 12, 2013.

Citation: Alfirevic A, Pirmohamed M, Marinovic B, Jorgensen AL, Harcourt-Smith L. Genetic testing for prevention of severe drug-induced skin rash. *Cochrane Database of Systematic Reviews* 2013, Issue 12. Art. No.: CD010891. DOI: 10.1002/14651858.CD010891.

Copyright © 2013 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

This is the protocol for a review and there is no abstract. The objectives are as follows:

To assess the effects of prospective pharmacogenetic screening to reduce drug-associated skin reactions in a patient population.

BACKGROUND

We have explained some terms we have used in a glossary. Please see [Table 1](#).

Description of the condition

Some drugs may cause skin rashes that vary in their severity and incidence. Skin reactions caused by drugs, often termed 'drug-induced skin injury' (DISI), are common (carbamazepine-induced skin rash has a 10% incidence rate ([Marson 2007](#))); they present with a range of clinical manifestations ranging from a mild maculopapular skin rash to life-threatening skin rashes such as Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN) ([Pirmohamed 2004](#); [Roujeau 1987](#)). The most severe forms are very rare, but these may result in up to 30% mortality. Less severe


forms of hypersensitivity reactions are troublesome and may prevent people from taking medications that are otherwise effective. The mechanisms involved in the pathogenesis of these drug-induced reactions are still poorly understood; however, immunogenetic and non-immune factors have been implicated. Recent evidence suggests that drug-specific T-cells can be identified in individuals who previously experienced adverse drug reactions (ADRs) to the culprit drug ([Illing 2012](#)).

Description of the intervention

There is increasing evidence from clinical trials that pretreatment genetic testing may reduce the possibility of severe drug-induced hypersensitivity ([Chen 2011](#); [Mallal 2008](#)).

[Figure 1](#) represents a diagram of decision-making informed by genetic testing.

Figure 1. Flowchart of interventions (genetic testing) and outcomes (skin rash) in a patient population prescribed drug X



To date, the strongest association with drug-induced skin injury has been reported with genetic variants in the human leukocyte antigens (HLA) (Amstutz 2013; Chung 2004; Hetherington 2002; Hung 2005; Mallal 2002; McCormack 2011; Ozkaya-Bayazit 2001). Human leukocyte antigens are cell surface proteins involved in presenting antigens to the immune system. They are encoded by most polymorphic genes in the human genome. However, different genetic markers are associated with hypersensitivity in different populations, and the effect size varies in different ethnicities. Also, there is evidence that some common factors could predispose to DISI irrespective of the underlying drug aetiology. In addition, it is possible that different severity phenotypes can share the same predisposing factor.

Table 2 shows reported associations between hypersensitivity reactions, which include skin injury and genetic variants in HLA genes.

How the intervention might work

Two recent clinical trials suggested that pretreatment genetic testing could reduce the possibility of severe hypersensitivity induced

with an anti-AIDS drug, abacavir (Mallal 2008), and an antiepileptic drug, carbamazepine (Chen 2011).

Patients who have a clinical requirement for a particular drug treatment can be stratified on the basis of a genetic test. Those who test positive for the risk marker are not prescribed the culprit drug, while those who test negative are safe to take the medicine of interest. In this way, it may be possible to reduce the incidence of severe drug skin reactions in the genotyped group compared to the randomly assigned group of patients who are not offered genetic testing, but for whom decision on drug choice is based on traditional clinical and biochemical parameters.

Why it is important to do this review

Adverse drug reactions affecting the skin are common; they can have high morbidity and mortality and are a burden for healthcare systems around the world. If we were able to predict them on the basis of a simple genetic test, it should be possible to prevent these reactions with one of the following approaches:

1. by prescribing alternative therapy if available;
2. by informing patients and healthcare providers; or
3. by informing drug developers in order to improve drug design and future drug development.

We aim to assess current research evidence to determine whether prospective pharmacogenetic screening is effective in reducing drug-associated skin reactions.

OBJECTIVES

To assess the effects of prospective pharmacogenetic screening to reduce drug-associated skin reactions in a patient population.

METHODS

Criteria for considering studies for this review

Types of studies

We will include randomised controlled trials (RCTs), which may be single-blind or unblinded.

Types of participants

People (adults and children of either gender) who are prescribed drugs known to cause delayed type hypersensitivity reactions. These include antiepileptic drugs, antiretrovirals, antigout drugs, and antibiotics such as beta-lactams (penicillin, amoxicillin, piperacillin, cephalosporins) and sulphonamides (sulphamethoxazole and trimethoprim).

Types of interventions

We will consider genetic testing for any genetic variants associated with hypersensitivity reactions using all available techniques to determine individual genotypes. The intervention is a randomly allocated genetic test; if the test is positive, a drug that can cause hypersensitivity is avoided.

Types of outcome measures

We will base core outcome measures on several papers describing clinical classification of drug-induced skin reactions, including the recently published paper entitled 'Phenotype standardisation for immune-mediated drug-induced skin injury' (Pirmohamed 2011), as well as earlier papers by the RegiSCAR (European Registry of Severe Cutaneous Adverse Reactions) consortium (Bouvesse 2012; Kardaun 2013; Sekula 2011).

We will assess clinically defined hypersensitivity reaction, immunologically confirmed hypersensitivity reaction (if skin patch testing or lymphocyte proliferation assay data are available), and long-term sequelae (including ophthalmologic, cutaneous, or liver damage, etc).

We have provided a full list of clinically relevant outcomes and distinction between primary and secondary outcomes in [Appendix 1](#).

Primary outcomes

1. The incidence of severe drug-induced skin rash.
2. Long-term sequelae (including ophthalmologic, cutaneous, or liver damage, etc) up to 12 months after the severe drug-induced skin rash.

Secondary outcomes

1. Hospitalisation for drug-induced skin reaction within 3 months of exposure to the drug.
2. Clinical phenotypes of hypersensitivity reactions (organ specific and systemic manifestations) including the following:
 - SJS/TEN (Stevens-Johnson syndrome, toxic epidermal necrolysis);
 - AGEP (acute generalised exanthematous pustulosis); and
 - HSS (hypersensitivity syndrome).

Additional terminology for HSS includes the following: Drug-induced hypersensitivity syndrome (DIHS), Drug reaction with eosinophilia and systemic symptoms (DRESS), and Drug-induced delayed multiorgan hypersensitivity syndrome.

Search methods for identification of studies

We aim to identify all relevant randomised controlled trials (RCTs) regardless of language or publication status (published, unpublished, in press, or in progress).

Electronic searches

We will search the following databases for relevant trials:

- the Cochrane Skin Group Specialised Register;
- the Cochrane Central Register of Controlled Trials (CENTRAL) in *The Cochrane Library*;
- MEDLINE via OVID (from 1946);
- EMBASE via OVID (from 1974); and
- LILACS (Latin American and Caribbean Health Science Information database, from 1982).

We have devised a draft search strategy for randomised controlled trials (RCTs) for MEDLINE (OVID), which is displayed in [Appendix 2](#). This will be used as the basis for search strategies for the other databases listed.

Trials registers

We will search the following trials registers:

- The metaRegister of Controlled Trials (www.controlled-trials.com).
- The US National Institutes of Health Ongoing Trials Register (www.clinicaltrials.gov).
- The Australian New Zealand Clinical Trials Registry (www.anzctr.org.au).
- The World Health Organization International Clinical Trials Registry platform (www.who.int/trialsearch).
- The EU Clinical Trials Register (www.clinicaltrialsregister.eu/).

Searching other resources

References from included studies

We will check the bibliographies of included studies for further references to relevant trials.

Adverse Effects

We will not perform a separate search for adverse effects of the target intervention. However, we will examine data on adverse effects from the included studies we identify.

Data collection and analysis

We plan to include at least one 'Summary of findings' table in our review. In this, we will summarise the primary outcomes for the most important comparison. If we feel there are several major comparisons or that our findings need to be summarised for different populations, we will include further 'Summary of findings' tables.

Selection of studies

Two review authors (AA and AJ) will independently assess studies for inclusion; they will independently screen all the titles and abstracts of publications identified by the searches to assess their eligibility. We will assess the full text of eligible citations for inclusion. We will exclude publications that do not meet the criteria at this stage and prepare a table of 'Characteristics of excluded studies' to clearly differentiate between those studies that are not at all relevant and those that may not fulfil the criteria for inclusion but may be considered relevant by some readers. We will reach consensus on the selection of trials and the final list of studies.

Data extraction and management

Two authors (AA and AJ) will independently extract data from included studies and resolve disagreements by discussion. If consensus is not reached, they will consult a third author (MP). We will collect the following information on study characteristics and methods: study design; inclusion and exclusion criteria; setting; country; language of publication; ethnicity of participants; control population (exposed to the culprit drug or healthy population); description of genotyping techniques used; genotyping quality control, which will include deviation from Hardy-Weinberg equilibrium (Hardy-Weinberg equilibrium is a crucial concept in population genetics; it predicts how gene frequencies will be inherited from generation to generation and is used as a measure of quality of genetic tests) and genotype call rate; age; gender; concomitant medications; time from exposure to culprit drug to skin reaction; type of skin reaction; location of skin lesion; duration; treatment; sequelae; other manifestations indicating systemic involvement; and laboratory tests.

Assessment of risk of bias in included studies

All review authors will assess the risk of bias of three studies as a pilot to ensure we are using consistent methods. AA and AJ will then independently assess the risk of bias in each trial according to the approaches described in the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011). Specifically, we will use The Cochrane Collaboration's tool for assessing risk of bias (Table 8.5.a in the *Cochrane Handbook for Systematic Reviews of Interventions*), which is based on seven domains (sequence, generation, allocation concealment, blinding of participants and personnel, blinding of outcome assessment, incomplete outcome data, selective outcome reporting, and other issues). The tool allows for the risk of bias to be assessed as 'low', 'high', or 'unclear' (indicating lack of information or uncertainty over the potential for bias). An additional author (MP) will assess any disagreements.

Measures of treatment effect

We will use statistical methods in accordance with the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011) to measure treatment effect. We will use mean difference (MD) with 95% confidence interval (CI) for continuous data or indeed standardised mean difference (SMD) with 95% CI where all studies report an outcome using similar scales. We will use risk ratio (RR) with 95% CI for dichotomous data.

Unit of analysis issues

We will take care to avoid a unit of analysis error due to repeated observations on participants, multiple treatments, or re-occurring events. We will also consider issues in cluster randomised trials, such as recruitment bias, baseline comparability of clusters,

and number of clusters, and make sure that appropriate statistical methods are used that take into account weighting, etc.

Dealing with missing data

We will consider the possible different types of missing data. We will deal with missing studies and the associated risk of bias by assessing for publication bias, whilst we will deal with missing outcomes and the associated risk of bias by assessing for selective reporting (see [Assessment of reporting biases](#) section).

In the event of missing summary data, we will contact the study author if data required to calculate outcomes of interest are missing. If no further data are available, we will not include the study in the meta-analysis. However, we will report the limited results from the study in narrative form in the results section and consider whether they are consistent with the results of the meta-analysis for that outcome. Also, where appropriate, we will make assumptions about the missing data (e.g. assuming all missing values to have a particular value, e.g. an adverse event) and conduct sensitivity analyses to test how sensitive the analyses are to our assumptions. We will address the potential implications of the missing data in the Discussion section.

Assessment of heterogeneity

We will assess the extent of heterogeneity using the I^2 statistic. We will use the following thresholds for the interpretation of the I^2 statistic:

- 0% to 40% = might not be important;
- 30% to 60% = moderate heterogeneity;
- 50% to 90% = may represent substantial heterogeneity; and
- 75% to 100% = considerable heterogeneity.

If we do a meta-analysis, we shall report the I^2 statistic and interpret the two data together.

Assessment of reporting biases

We will assess biases, including publication bias, 'time-lag' bias, outcome reporting bias, language bias, and citation bias. We will use a funnel plot, Begg test, and Egger test to evaluate publication bias (Begg 1989; Egger 1998). We will only carry out tests for funnel plot asymmetry when there are at least 10 studies included in the meta-analysis, because when there are fewer studies, the power of the tests is too low to distinguish chance from real asymmetry.

Data synthesis

If there is no significant clinical heterogeneity, we will synthesise the results in meta-analyses. We will synthesise data according to type of intervention (e.g. genotype test). We will use a random-effects model.

Where events are rare, a random-effects approach may be inappropriate. Where events are rare, extra care will be taken to adopt appropriate methods of meta-analysis as recommended in the *Cochrane Handbook for Systematic Reviews of Interventions* (section 16.9), since many meta-analysis methods are suboptimal where events are rare through results being biased, confidence intervals being too wide, or power being too low. Choice of method will be guided by control group risk, likely treatment effect size, and consideration of balance in numbers of treated and control participants in the constituent studies. Where the control groups differ, e.g. they are drawn from a healthy population or from a population of people treated with the culprit drug but without any adverse effects, we will conduct separate analysis. We will use Review Manager to undertake the meta-analyses.

Subgroup analysis and investigation of heterogeneity

Where there is substantial heterogeneity, we will explore the causes by way of subgroup analyses. Indeed, some HLA genetic variants are associated with a high risk of drug-induced skin reactions, so interventions would only be given to the appropriate subpopulation.

We will consider the following:

- participant factors (age, ethnicity, ADR classification, and comparability of participant groups); and
- trial design issues (genotyping methodology and quality control, blinding, drugs included, and drug dosage and duration of use).

Sensitivity analysis

We will evaluate the robustness of the results of the meta-analyses by removing trials of low methodological quality as defined by their risk of bias.

ACKNOWLEDGEMENTS

The Cochrane Skin Group editorial base wishes to thank Luigi Naldi who was the Cochrane Dermatology Editor for this protocol; Jo Leonardi-Bee and Ching-Chi Chi who were the Statistical and Methods Editors, respectively; the clinical referee, Olivier Chosidow; and the consumer referee, Jack Tweed.

REFERENCES

Additional references

Amstutz 2013

Amstutz U, Ross CJ, Castro-Pastrana LI, Rieder MJ, Shear NH, Hayden MR, et al. HLA-A*31:01 and HLA-B*15:02 as genetic markers for carbamazepine hypersensitivity in children. *Clinical Pharmacology & Therapeutics* 2013;**94**(1): 142–9. [MEDLINE: 23588310]

Begg 1989

Begg CB, Berlin JA. Publication bias and dissemination of clinical research. *Journal of the National Cancer Institute* 1989;**81**(2):107–15. [MEDLINE: 2642556]

Bouvrès 2012

Bouvrès S, Valeyrie-Allanore L, Ortonne N, Konstantinou MP, Kardaun SH, Bagot M, et al. Toxic epidermal necrolysis, DRESS, AGEP: do overlap cases exist?. *Orphanet Journal of Rare Diseases* 2012;**7**:72. [MEDLINE: 23009177]

Carr 2013

Carr DF, Chaponda M, Jorgensen AL, Castro EC, van Oosterhout JJ, Khoo SH, et al. Association of human leukocyte antigen alleles and nevirapine hypersensitivity in a Malawian HIV-infected population. *Clinical Infectious Diseases* 2013;**56**(9):1330–9. [MEDLINE: 23362284]

Chantarangsu 2009

Chantarangsu S, Mushiroda T, Mahasirimongkol S, Kiertiburanakul S, Sungkanuparph S, Manosuthi W, et al. HLA-B*3505 allele is a strong predictor for nevirapine-induced skin adverse drug reactions in HIV-infected Thai patients. *Pharmacogenetics and Genomics* 2009;**19**(2): 139–46.

Chen 2011

Chen P, Lin JJ, Lu CS, Ong CT, Hsieh PF, Yang CC, et al. Carbamazepine-induced toxic effects and HLA-B*1502 screening in Taiwan. *New England Journal of Medicine* 2011;**364**(12):1126–33. [MEDLINE: 21428768]

Cheung 2013

Cheung YK, Cheng SH, Chan EJ, Lo SV, Ng MH, Kwan P. HLA-B alleles associated with severe cutaneous reactions to antiepileptic drugs in Han Chinese. *Epilepsia* 2013;**54**(7): 1307–14. [MEDLINE: 23692434]

Chong 2013

Chong KW, Chan DW, Cheung YB, Ching LK, Hie SL, Thomas T, et al. Association of carbamazepine-induced severe cutaneous drug reactions and HLA-B*1502 allele status, and dose and treatment duration in paediatric neurology patients in Singapore. *Archives of Disease in Childhood* 2013 Nov 13 [Epub ahead of print]. [MEDLINE: 24225276]

Chung 2004

Chung WH, Hung SI, Hong HS, Hsieh MS, Yang LC, Ho HC, et al. Medical genetics: a marker for Stevens-Johnson syndrome. *Nature* 2004;**428**(6982):486. [MEDLINE: 15057820]

Ding 2010

Ding WY, Lee CK, Choon SE. Cutaneous adverse drug reactions seen in a tertiary hospital in Johor, Malaysia. *International Journal of Dermatology* 2010;**49**(7):834–41. [MEDLINE: 20618508]

Egger 1998

Egger M, Smith GD. Bias in location and selection of studies. *BMJ* 1998;**316**(7124):61–66. [MEDLINE: 9451274]

Gao 2012

Gao S, Gui XE, Liang K, Liu Z, Hu J, Dong B. HLA-dependent hypersensitivity reaction to nevirapine in Chinese Han HIV-infected patients. *AIDS Research and Human Retroviruses* 2012;**28**(6):540–3. [MEDLINE: 21902584]

Gatanaga 2007

Gatanaga H, Yazaki H, Tanuma J, Honda M, Genka I, Teruya K, et al. HLA-Cw8 primarily associated with hypersensitivity to nevirapine. *AIDS* 2007;**21**(2):264–5. [MEDLINE: 17197830]

Hetherington 2002

Hetherington S, Hughes AR, Mosteller M, Shortino D, Baker KL, Spreen W, et al. Genetic variations in HLA-B region and hypersensitivity reactions to abacavir. *Lancet* 2002;**359**(9312):1121–2. [MEDLINE: 11943262]

Higgins 2011

Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration. Available from www.cochrane-handbook.org.

Hughes 2004

Hughes DA, Vilar FJ, Ward CC, Alfirevic A, Park BK, Pirmohamed M. Cost-effectiveness analysis of HLA B*5701 genotyping in preventing abacavir hypersensitivity. *Pharmacogenetics* 2004;**14**(6):335–42. [MEDLINE: 15247625]

Hughes 2004a

Hughes AR, Mosteller M, Bansal AT, Davies K, Haneline SA, Lai EH, et al. Association of genetic variations in HLA-B region with hypersensitivity to abacavir in some, but not all, populations. *Pharmacogenomics* 2004;**5**(2):203–11. [MEDLINE: 15016610]

Hung 2005

Hung SI, Chung WH, Liou LB, Chu CC, Lin M, Huang HP, et al. HLA-B*5801 allele as a genetic marker for severe cutaneous adverse reactions caused by allopurinol. *Proceedings of the National Academy of Sciences of the United States of America* 2005;**102**(11):4134–9. [MEDLINE: 15743917]

Hung 2006

Hung SI, Chung WH, Jee SH, Chen WC, Chang YT, Lee WR, et al. Genetic susceptibility to carbamazepine-induced cutaneous adverse drug reactions. *Pharmacogenetics and Genomics* 2006;**16**(4):297–306. [MEDLINE: 16538176]

Hung 2010

Hung SI, Chung WH, Liu ZS, Chen CH, Hsieh MS, Hui RC, et al. Common risk allele in aromatic antiepileptic-drug induced Stevens-Johnson syndrome and toxic epidermal necrolysis in Han Chinese. *Pharmacogenomics* 2010;**11**(3): 349–56. [MEDLINE: 20235791]

Illing 2012

Illing PT, Vivian JP, Dudek NL, Kostenko L, Chen Z, Bharadwaj M, et al. Immune self-reactivity triggered by drug-modified HLA-peptide repertoire. *Nature* 2012;**486**(7404):554–8. [MEDLINE: 22722860]

Kaniwa 2008

Kaniwa N, Saito Y, Aihara M, Matsunaga K, Tohkin M, Kurose K, et al. HLA-B locus in Japanese patients with anti-epileptics and allopurinol-related Stevens-Johnson syndrome and toxic epidermal necrolysis. *Pharmacogenomics* 2008;**9**(11):1617–22. [MEDLINE: 19018717]

Kardaun 2013

Kardaun SH, Sekula P, Valeyrie-Allanore L, Liss Y, Chu CY, Creamer D, et al. Drug reaction with eosinophilia and systemic symptoms (DRESS): an original multisystem adverse drug reaction. Results from the prospective RegiSCAR study. *British Journal of Dermatology* 2013;**169**(5):1071–80. [MEDLINE: 23855313]

Kazeem 2009

Kazeem GR, Cox C, Aponte J, Messenheimer J, Brazell C, Nelsen AC, et al. High-resolution HLA genotyping and severe cutaneous adverse reactions in lamotrigine-treated patients. *Pharmacogenetics and Genomics* 2009;**19**(9):661–5. [MEDLINE: 19668019]

Kim 2005

Kim SH, Choi JH, Lee KW, Kim SH, Shin ES, Oh HB, et al. The human leucocyte antigen-DRB1*1302-DQB1*0609-DPB1*0201 haplotype may be a strong genetic marker for aspirin-induced urticaria. *Clinical and Experimental Allergy* 2005;**35**(3):339–44. [MEDLINE: 15784113]

Kulkantrakorn 2012

Kulkantrakorn K, Tassaneeyakul W, Tiamkao S, Jantararungtong T, Prabmechai N, Vannaprasaht S, et al. HLA-B*1502 strongly predicts carbamazepine-induced Stevens-Johnson syndrome and toxic epidermal necrolysis in Thai patients with neuropathic pain. *Pain Practice* 2012;**12**(3):202–8. [MEDLINE: 21676164]

Li 2007

Li H, Dai Y, Huang H, Li L, Leng S, Cheng J, et al. HLA-B*1301 as a biomarker for genetic susceptibility to hypersensitivity dermatitis induced by trichloroethylene among workers in China. *Environmental Health Perspectives* 2007;**115**(11):1553–6. [MEDLINE: 18007983]

Likanonsakul 2009

Likanonsakul S, Rattanatham T, Feangvad S, Uttayamakul S, Prasithsirikul W, Tunthanathip P, et al. HLA-Cw*04 allele associated with nevirapine-induced rash in HIV-infected Thai patients. *AIDS Research and Therapy* 2009;**6**: 22. [MEDLINE: 19845952]

Littera 2006

Littera R, Carcassi C, Masala A, Piano P, Serra P, Ortu F, et al. HLA-dependent hypersensitivity to nevirapine in Sardinian HIV patients. *AIDS* 2006;**20**(12):1621–6. [MEDLINE: 16868443]

Locharernkul 2008

Locharernkul C, Loplumlert J, Limotai C, Korkij W, Desudchit T, Tongkobpetch S, et al. Carbamazepine and phenytoin induced Stevens-Johnson syndrome is associated with HLA-B*1502 allele in Thai population. *Epilepsia* 2008;**49**(12):2087–91. [MEDLINE: 18637831]

Mallal 2002

Mallal S, Nolan D, Witt C, Masel G, Martin AM, Moore C, et al. Association between presence of HLA-B*5701, HLA-DR7, and HLA-DQ3 and hypersensitivity to HIV-1 reverse-transcriptase inhibitor abacavir. *Lancet* 2002;**359**(9308):727–32. [MEDLINE: 11888582]

Mallal 2008

Mallal S, Phillips E, Carosi G, Molina JM, Workman C, Tomazic J, et al. HLA-B*5701 screening for hypersensitivity to abacavir. *New England Journal of Medicine* 2008;**358**(6): 568–79. [MEDLINE: 18256392]

Man 2007

Man CB, Kwan P, Baum L, Yu E, Lau KM, Cheng AS, et al. Association between HLA-B*1502 allele and antiepileptic drug-induced cutaneous reactions in Han Chinese. *Epilepsia* 2007;**48**(5):1015–8. [MEDLINE: 17509004]

Marson 2007

Marson AG, Al-Kharusi AM, Alwaidh M, Appleton R, Baker GA, Chadwick DW, et al. The SANAD study of effectiveness of carbamazepine, gabapentin, lamotrigine, oxcarbazepine, or topiramate for treatment of partial epilepsy: an unblinded randomised controlled trial. *Lancet* 2007;**369**(9566):1000–15. [MEDLINE: 17382827]

Martin 2004

Martin AM, Nolan D, Gaudieri S, Almeida CA, Nolan R, James I, et al. Predisposition to abacavir hypersensitivity conferred by HLA-B*5701 and a haplotypic Hsp70-Hom variant. *Proceedings of the National Academy of Sciences of the United States of America* 2004;**101**(12):4180–5. [MEDLINE: 15024131]

Martin 2005

Martin AM, Nolan D, James I, Cameron P, Keller J, Moore C, et al. Predisposition to nevirapine hypersensitivity associated with HLA-DRB1*0101 and abrogated by low CD4 T-cell counts. *AIDS* 2005;**19**(1):97–9.

McCormack 2011

McCormack M, Alfirevic A, Bourgeois S, Farrell JJ, Kasperaviciute D, Carrington M, et al. HLA-A*3101 and carbamazepine-induced hypersensitivity reactions in Europeans. *New England Journal of Medicine* 2011;**364**(12):1134–43. [MEDLINE: 21428769]

Mehta 2009

Mehta TY, Prajapati LM, Mittal B, Joshi CG, Sheth JJ, Patel DB, et al. Association of HLA-B*1502 allele and carbamazepine-induced Stevens-Johnson syndrome among

- Indians. *Indian Journal of Dermatology, Venereology and Leprology* 2009;**75**(6):579–82. [MEDLINE: 19915237]
- Ozeki 2011**
Ozeki T, Mushiroda T, Yowang A, Takahashi A, Kubo M, Shirakata Y, et al. Genome-wide association study identifies HLA-A*3101 allele as a genetic risk factor for carbamazepine-induced cutaneous adverse drug reactions in Japanese population. *Human Molecular Genetics* 2011;**20**(5):1034–41. [MEDLINE: 21149285]
- Ozkaya-Bayazit 2001**
Ozkaya-Bayazit E, Akar U. Fixed drug eruption induced by trimethoprim-sulfamethoxazole: evidence for a link to HLA-A30 B13 Cw6 haplotype. *Journal of the American Academy of Dermatology* 2001;**45**(5):712–7. [MEDLINE: 11606921]
- Palikhe 2008**
Palikhe NS, Kim SH, Park HS. What do we know about the genetics of aspirin intolerance?. *Journal of Clinical Pharmacy and Therapeutics* 2008;**33**(5):465–72. [MEDLINE: 18834360]
- Pellicano 1997**
Pellicano R, Lomuto M, Ciavarella G, Di Giorgio G, Gasparini P. Fixed drug eruptions with feprazone are linked to HLA-B22. *Journal of the American Academy of Dermatology* 1997;**36**(5 Pt 1):782–4. [MEDLINE: 9146544]
- Pirmohamed 2004**
Pirmohamed M, James S, Meakin S, Green C, Scott AK, Walley TJ, et al. Adverse drug reactions as cause of admission to hospital: prospective analysis of 18 820 patients. *BMJ* 2004;**329**(7456):15–9. [MEDLINE: 15231615]
- Pirmohamed 2011**
Pirmohamed M, Friedmann PS, Molokhia M, Loke YK, Smith C, Phillips E, et al. Phenotype standardization for immune-mediated drug-induced skin injury. *Clinical Pharmacology & Therapeutics* 2011;**89**(6):896–901. [MEDLINE: 21562486]
- Quiralte 1999**
Quiralte J, Sánchez-García F, Torres MJ, Blanco C, Castillo R, Ortega N, et al. Association of HLA-DR11 with the anaphylactoid reaction caused by nonsteroidal anti-inflammatory drugs. *Journal of Allergy and Clinical Immunology* 1999;**103**(4):685–9. [MEDLINE: 10200020]
- Rodríguez-Pérez 1994**
Rodríguez-Pérez M, González-Domínguez J, Matarán L, García-Pérez S, Salvatierra D. Association of HLA-DR5 with mucocutaneous lesions in patients with rheumatoid arthritis receiving gold sodium thiomalate. *Journal of Rheumatology* 1994;**21**(1):41–3. [MEDLINE: 8151585]
- Romano 1998**
Romano A, De Santis A, Romito A, Di Fonso M, Venuti A, Gasbarrini GB, et al. Delayed hypersensitivity to aminopenicillins is related to major histocompatibility complex genes. *Annals of Allergy, Asthma & Immunology* 1998;**80**(5):433–7. [MEDLINE: 9609616]
- Roujeau 1986**
Roujeau JC, Bracq C, Huynh NT, Chausselet E, Raffin C, Duédari N. HLA phenotypes and bullous cutaneous reactions to drugs. *Tissue Antigens* 1986;**28**(4):251–4. [MEDLINE: 3544335]
- Roujeau 1987**
Roujeau JC, Huynh TN, Bracq C, Guillaume JC, Revuz J, Touraine R. Genetic susceptibility to toxic epidermal necrolysis. *Archives of Dermatology* 1987;**123**(9):1171–3. [MEDLINE: 3477129]
- Saag 2008**
Saag M, Balu R, Phillips E, Brachman P, Martorell C, Burman W, et al. High sensitivity of human leukocyte antigen-b*5701 as a marker for immunologically confirmed abacavir hypersensitivity in white and black patients. *Clinical Infectious Diseases* 2008;**46**(7):1111–8. [MEDLINE: 18444831]
- Sekula 2011**
Sekula P, Liss Y, Davidovici B, Dunant A, Roujeau JC, Kardaun S, et al. Evaluation of SCORTEN on a cohort of patients with Stevens-Johnson syndrome and toxic epidermal necrolysis included in the RegiSCAR study. *Journal of Burn Care & Research* 2011;**32**(2):237–45. [MEDLINE: 21228709]
- Sieben 2002**
Sieben S, Kawakubo Y, Al Masaoudi T, Merk HF, Blömeke B. Delayed-type hypersensitivity reaction to paraphenylenediamine is mediated by 2 different pathways of antigen recognition by specific alpha beta human T-cell clones. *Journal of Allergy and Clinical Immunology* 2002;**109**(6):1005–11. [MEDLINE: 12063532]
- Tangamornsuksan 2013**
Tangamornsuksan W, Chaikunapruk N, Somkrua R, Lohitnavy M, Tassaneeyakul W. Relationship between the HLA-B*1502 allele and carbamazepine-induced Stevens-Johnson syndrome and toxic epidermal necrolysis: a systematic review and meta-analysis. *JAMA Dermatology* 2013;**149**(9):1025–32. [MEDLINE: 23884208]
- Tassaneeyakul 2009a**
Tassaneeyakul W, Jantararungtong T, Chen P, Lin PY, Tiamkao S, Khunarkornsiri U, et al. Strong association between HLA-B*5801 and allopurinol-induced Stevens-Johnson syndrome and toxic epidermal necrolysis in a Thai population. *Pharmacogenetics and Genomics* 2009;**19**(9):704–9. [MEDLINE: 19696695]
- Tassaneeyakul 2010**
Tassaneeyakul W, Tiamkao S, Jantararungtong T, Chen P, Lin SY, Chen WH, et al. Association between HLA-B*1502 and carbamazepine-induced severe cutaneous adverse drug reactions in a Thai population. *Epilepsia* 2010;**51**(5):926–30. [MEDLINE: 20345939]
- Torres 2008**
Torres MJ, Mayorga C, Cornejo-García JA, Lopez S, Chaves P, Rondon C, et al. Monitoring non-immediate

allergic reactions to iodine contrast media. *Clinical and Experimental Immunology* 2008;**152**(2):233–8. [MEDLINE: 18341616]

Vitezica 2008

Vitezica ZG, Milpied B, Lonjou C, Borot N, Ledger TN, Lefebvre A, et al. HLA-DRB1*01 associated with cutaneous hypersensitivity induced by nevirapine and efavirenz. *AIDS* 2008;**22**(4):540–1. [MEDLINE: 18301070]

Watanabe 2010

Watanabe H, Tohyama M, Kamijima M, Nakajima T, Yoshida T, Hashimoto K, et al. Occupational trichloroethylene hypersensitivity syndrome with

human herpesvirus-6 and cytomegalovirus reactivation. *Dermatology* 2010;**221**(1):17–22. [MEDLINE: 20407216]

References to other published versions of this review

Tassaneeyakul 2009

Tassaneeyakul W, Jantararungtong T, Chen P, Lin PY, Tiamkao S, Khunarkornsiri U, et al. Strong association between HLA-B*5801 and allopurinol-induced Stevens-Johnson syndrome and toxic epidermal necrolysis in a Thai population. *Pharmacogenetics and Genomics* 2009;**19**(9): 704–9. [MEDLINE: 19696695]

* Indicates the major publication for the study

ADDITIONAL TABLES

Table 1. Glossary of terms

Term	Explanation
Allele	One of two or more alternative forms of a gene at corresponding sites (loci) on homologous chromosomes
Hardy-Weinberg equilibrium	This states that allele and genotype frequencies in a population will remain constant from generation to generation in the absence of other evolutionary influences
HLA	Human leukocyte antigen: a group of protein molecules located on bone marrow and other cells that can provoke an immune response
Hypersensitivity	A state of altered reactivity in which the body reacts with an exaggerated immune response to a foreign substance, such as a drug
Polymorphic	A variation in the DNA that is too common to be due merely to new mutation. A polymorphism must have a frequency of at least 1% in a population
Maculopapular rash	A rash with both macules (flat and coloured like a freckle) and papules (a small raised spot)

Table 2. Associations between drug-induced skin injury and genetic variants in the HLA genes

Drugs associated with skin injury	Class of drug	HLA allele	Population	Reference
Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN)				
Allopurinol	Antiuric acid	B*5801	Han Chinese	Hung 2005
-	-	-	Thai	Tassaneeyakul 2009a

Table 2. Associations between drug-induced skin injury and genetic variants in the HLA genes (Continued)

-	-	-	Japanese	Kaniwa 2008
-	-	-	Malay	Ding 2010
Carbamazepine	Antiepileptic	B*1502	Han Chinese	Cheung 2013 ; Chung 2004 ; Chong 2013 ; Hung 2006 ; Man 2007
-	-	-	Thai	Kulkantrakorn 2012 ; Locharernkul 2008 ; Tassaneeyakul 2010 ; Tangamornsuksan 2013
-	-	-	Malay	Ding 2010
-	-	-	Indian	Mehta 2009
-	-	A*3101	White	Amstutz 2013 ; McCormack 2011 ;
-	-	A*3101	Japanese	Ozeki 2011
Phenytoin	Antiepileptic	B*1502	Han Chinese	Hung 2010 ; Man 2007
-	-	-	Thai	Locharernkul 2008 ;
Oxicam	Non-steroidal anti-inflammatory drug (NSAID)	A2, B12	White	Roujeau 1987
Sulphamethoxazole	Antibiotic	A29, B12, DR7	White	Roujeau 1986
Hypersensitivity syndrome (DIHS or DRESS)				
Abacavir	Antiretroviral	B*5701	White	Hetherington 2002 ; Hughes 2004 ; Mallal 2002 ; Mallal 2008 ; Martin 2004
-	-	-	African Americans	Hughes 2004a ; Saag 2008
Aminopenicillins	Antibiotic	A2, Drw52	White	Romano 1998
Nevirapine	Antiretroviral	DRB1*01	White - Australian	Martin 2005
-	-	DRB1*01	White - French	Vitezica 2008

Table 2. Associations between drug-induced skin injury and genetic variants in the HLA genes (Continued)

-	-	Cw8, B14	White - Italian	Littera 2006
-	-	Cw8	Japanese	Gatanaga 2007
-	-	B*3505	Thai	Chantarangsu 2009
-	-	Cw4	Thai	Likanonsakul 2009
-	-	C*0404	Black African	Carr 2013
-	-	Cw*04	Chinese	Gao 2012
Aspirin	NSAIDS	DRB1*1302, DQB1*0609	-	Kim 2005 ; Palikhe 2008
-	NSAIDS	DR11	-	Quiralte 1999
Iodine contrast media	-	DR	White - Spanish	Torres 2008
Paraphenylenediamine	Hair dye	DP	White - German	Sieben 2002
Gold sodium thiomalate	Treatment of rheumatoid arthritis	DR5	White - Spanish	Rodriguez-Pérez 1994
Lamotrigine	Antiepileptic	B*5801, A*6801	White	Kazeem 2009
Trichloroethylene	Industrial solvent, dry cleaning	B*1301	Japanese	Li 2007 ; Watanabe 2010
Fixed drug eruptions				
Co-trimoxazole	Antibiotic	A30, B13, Cw6	White - Turkish	Ozkaya-Bayazit 2001
Feprazone	Analgesic	B22	-	Pellicano 1997

APPENDICES

Appendix 1. Outcomes adapted from Pirmohamed 2011

Primary outcome:

- Drug-induced skin reaction (yes, no)

Secondary outcome:

- Clinical phenotype:

SJS/TEN (Stevens-Johnson syndrome, toxic epidermal necrolysis)

- Skin detachment 1% to 10% (SJS), 10% to 30% (overlap syndrome), and > 30% (TEN)
- Severe, often hemorrhagic, erosions of mucous membranes
- Other manifestations indicating systemic involvement (e.g. fever, liver chemistry elevations, intestinal and pulmonary manifestations, or the presence of lymphopenia)
- Severe pain and tenderness in the skin
- Target lesions, representing the degree of epidermal necrosis

AGEP (acute generalised exanthematous pustulosis)

- Acute widespread edematous erythema followed by a sterile pustular eruption. Often the pustules are first localised in the neck, groin, and axillae, and later become widely disseminated
- Fever (temperature > 38 °C)
- Neutrophilia with or without a mild eosinophilia

HSS (Hypersensitivity syndrome)

- Additional terminology includes Drug-induced hypersensitivity syndrome (DIHS), Drug reaction with eosinophilia and systemic symptoms (DRESS), Drug-induced delayed multiorgan hypersensitivity syndrome
- Variable skin manifestations; exanthema are most common
- Increased liver function tests, hepatitis, cholestasis
- Colitis
- Nephritis
- Pneumonitis
- Aseptic meningitis, encephalitis, inappropriate antidiuretic hormone syndrome
- Myocarditis
- Myositis
- Lymphocytic thyroiditis
- Eosinophilia, atypical lymphocytes, agranulocytosis, thrombocytopenia, haemolytic anaemia, aplastic anaemia
- Lymphadenopathy, pseudolymphoma

Appendix 2. MEDLINE (OVID) search strategy

1. exp Exanthema/
2. exanthema.ti,ab.
3. (rash or rashes).ti,ab.
4. drug induced skin injury.ti,ab.
5. exp Drug Hypersensitivity/ge, pc [Genetics, Prevention & Control]
6. (drug adj2 hypersensitivit\$).ti,ab.
7. hypersensitiv\$ reaction\$.ti,ab.
8. hypersensitivity syndrome.ti,ab.
9. drug eruption\$.ti,ab.
10. exp Drug Toxicity/ge, pc [Genetics, Prevention & Control]
11. drug toxic\$.ti,ab.
12. adverse drug reaction\$.ti,ab.
13. toxic epidermal necrolysis.ti,ab. or exp Epidermal Necrolysis, Toxic/

14. stevens johnson syndrome.ti,ab. or exp Stevens-Johnson Syndrome/
15. exp Acute Generalized Exanthematous Pustulosis/
16. Acute Generalized Exanthematous Pustulosis.ti,ab.
17. erythema multiforme.ti,ab. or exp Erythema Multiforme/
18. dress syndrome.ti,ab.
19. "Drug Reaction with Eosinophilia and Systemic Symptoms".ti,ab.
20. "Drug Rash with Eosinophilia and Systemic Symptoms".ti,ab.
21. or/1-20
22. exp Genetic Testing/
23. genetic test\$.ti,ab.
24. exp Pharmacogenetics/
25. (pharmacogenomic\$ or pharmacogenetic\$).ti,ab.
26. screening.ti,ab.
27. patch test\$.ti,ab.
28. exp Patch Tests/
29. exp HLA Antigens/
30. hla allele\$.ti,ab.
31. exp Polymorphism, Genetic/
32. genetic polymorphism\$.ti,ab.
33. exp Genetic Variation/
34. (genetic variant\$ or genetic variation\$).ti,ab.
35. or/22-34
36. randomized controlled trial.pt.
37. controlled clinical trial.pt.
38. randomized.ab.
39. placebo.ab.
40. clinical trials as topic.sh.
41. randomly.ab.
42. trial.ti.
43. 36 or 37 or 38 or 39 or 40 or 41 or 42
44. exp animals/ not humans.sh.
45. 43 not 44
46. 21 and 35 and 45

WHAT'S NEW

Date	Event	Description
19 May 2014	Amended	The Declaration of interest section was updated (clinical referee's statement added)

CONTRIBUTIONS OF AUTHORS

AA was the contact person with the editorial base.

AA co-ordinated the contributions from the co-authors and wrote the final draft of the protocol.

AA, AJ, and MP worked on the methods sections.

AA, MP, and BM drafted the clinical sections of the background and responded to the clinical comments of the referees.

AJ responded to the methodology and statistics comments of the referees.

MP, AJ, BM, and LHS contributed to writing the protocol.

LHS was the consumer co-author and checked the protocol for readability and clarity. She also ensured that the outcomes are relevant to consumers.

AA is the guarantor of the final review.

Disclaimer

The views and opinions expressed therein are those of the authors and do not necessarily reflect those of the NIHR, NHS or the Department of Health, UK.

DECLARATIONS OF INTEREST

Ana Alfirevic: none declared.

Munir Pirmohamed: none declared.

Branka Marinovic: none declared.

Andrea L Jorgensen: none declared.

Linda Harcourt-Smith: none declared.

Olivier Chosidow, clinical referee, works in a department that is a referral center for toxic and auto-immune blistering diseases, and he has participated in the RegiSCAR (European Registry of Severe Cutaneous Adverse Reactions) group.

SOURCES OF SUPPORT

Internal sources

- No financial support, Other.

External sources

- The National Institute for Health Research (NIHR), UK.

The NIHR, UK, is the largest single funder of the Cochrane Skin Group.