

Letters to the editor

OVERVIEW

Please submit letters for the editor's consideration within three weeks of receipt of *Clinical Medicine*. Letters should ideally be limited to 350 words, and sent by email to: clinicalmedicine@rcplondon.ac.uk

Vitamin D – what is normal according to latest research and how should we deal with it?

Editor – This interesting article summarised the public health concern presented by vitamin D deficiency. The observational data that links vitamin D deficiency to many chronic diseases demonstrates a potential impact extending far beyond traditionally recognised endocrine effects on bone health. The benefits of vitamin D supplementation, in the absence of randomised controlled trial data, are currently yet to be fully defined.

The author advocates 'a targeted, rather than population-based screening approach to vitamin D testing', but adds that 'empirical supplementation of at risk groups should be considered'.¹ Any strategy that includes blanket vitamin D supplementation should be approached with caution: certain individuals in the population will have a genetic predisposition to vitamin D toxicity, unmasked only through higher vitamin D intake.

In the 1950s, routine supplementation of formula milk products with vitamin D₃, in an effort to prevent rickets, led to a sudden increase in infants presenting with idiopathic infantile hypercalcaemia, in some cases proving fatal.² Almost 60 years later, the discovery of loss-of-function mutations in *CYP24A1*, which encodes vitamin D 24-hydroxylase – an enzyme with a key role in vitamin D catabolism – finally linked these events.³ Furthermore, an adult cohort who present with calcium-containing renal stones and nephrocalcinosis are recognised to carry *CYP24A1* mutations.⁴ At present, the population frequency of *CYP24A1* variants has not been established.⁵ The rarity of these presentations often leads to delayed diagnosis, a feature that will hopefully change with increased awareness and easier access to diagnostic testing.

As research into vitamin D supplementation expands, we must be careful to tailor our treatment towards each individual, and remember that Vitamin D supplements are not the answer for everyone. ■

Conflicts of interests

The author has no conflicts of interests.

FAY JOANNE HILL

Core medical trainee, Newcastle-Upon-Tyne Hospitals,
Newcastle, UK

References

- 1 Gittoes NJL. Vitamin D - what is normal according to latest research and how should we deal with it? *Clin Med* 2016;171–4.
- 2 British Paediatric Association. Hypercalcaemia in infants and vitamin D. *BMJ* 1956;2:149.
- 3 Schlingmann KP, Kaufmann M, Weber S *et al*. Mutations in *CYP24A1* and idiopathic infantile hypercalcemia. *N Engl J Med* 2011;365:410–21.
- 4 Dinour D, Beckerman P, Ganon L *et al*. Loss-of-function mutations of *CYP24A1*, the vitamin D 24-hydroxylase gene, cause long-standing hypercalciuric nephrolithiasis and nephrocalcinosis. *J Urol* 2013;190:552–7.
- 5 Sayers J, Hynes AM, Rice SJ, Hogg P, Sayer JA. Searching for *CYP24A1* mutations in cohorts of patients with calcium nephrolithiasis. *OA Nephrology* 2013;1:1–6.

Viral arthritis

Editor – I am grateful to Marks and Marks¹ for their excellent overview of viral arthritis, which provided many useful clinical pointers. Viral arthritis acquired abroad will continue to increase in line with the ever-expanding number of travelers visiting or returning to the UK, especially from tropical climes.² As noted in the review, aetiology of viral arthritis includes alphaviruses such as Chikungunya and also flaviviruses such as Dengue. There is another topical flavivirus that is relevant for the generalist faced with a returned traveler with a fever, rash and arthralgia/arthritis: Zika.

Zika virus has been described as a 'new global health threat' for 2016.³ While the majority of those infected with Zika will be asymptomatic, there are major concerns about a possible association between Zika infection in pregnant women and congenital microcephaly.⁴ Twenty one cases of Zika have been diagnosed in the UK since April 2015 in travellers returning from affected countries (predominantly the Americas and the Caribbean).⁵ Returning travellers presenting with a maculopapular rash and/or fever with at least one of the following symptoms (not explained by other medical conditions) meet the World Health Organization criteria of a suspected case: conjunctivitis, arthralgia, or arthritis.⁶ Other symptoms may include itching/pruritus, headache, myalgia, lower back pain, and retro-orbital pain – all of which are compatible with a diagnosis of Dengue fever or Chikungunya.⁵

Public Health England provides comprehensive guidance on Zika virus infection, including a clinical algorithm of investigation of suspected cases and exposed pregnant women.⁵ Management of people with Zika virus infection is supportive, with intravenous fluids if necessary and paracetamol for fever and arthralgia/arthritis. It should be noted that non-steroid anti-inflammatory agents are best avoided until the diagnosis of Dengue has been excluded, because of bleeding risk.

Letters to the editor

With expanding global travel, emerging and tropical infections are becoming ever more pertinent to and prevalent in day-to-day primary and secondary care practice in the UK. Clinicians should remain vigilant to the 1% of acute arthritis cases of viral aetiology, which include Zika virus infection. ■

TOM WINGFIELD

*Infectious diseases specialist registrar and academic
clinical lecturer,
Institute of Infection and Global Health, Liverpool University,
Liverpool, UK*

References

- 1 Marks M, Marks JL. Viral arthritis. *Clin Med* 2016;16:129–34.
- 2 Coltart CEM, Behrens RH. The new health threats of exotic and global travel. *Br J Gen Pract* 2012;62:512–3.
- 3 Zika virus: a new global threat for 2016. *Lancet* 2016;387:96.
- 4 Ahmad SS, Amin TN, Ustianowski A. Zika virus: management of infection and risk. *BMJ* 2016;352:i1062.
- 5 Public Health England. *Zika virus*. Health protection – guidance. PHE, 2016. Available online at www.gov.uk/guidance/zika-virus [Accessed 13 May 2016].
- 6 World Health Organization. *Zika virus disease*. WHO, 2015. Available online at www.who.int/csr/disease/zika/case-definition/en/ [Accessed 13 May 2016].



Museum and Garden

Royal College of Physicians Museum and Medicinal Garden

Visit the Royal College of Physicians (RCP) and discover 500 years of medical history in an acclaimed architectural setting.

The RCP is the oldest medical college in England, and since its foundation in 1518 it has built up magnificent collections of books, manuscripts, portraits, silver, and medical artefacts.

Our celebrated Grade I listed headquarters in Regent's Park is the creation of British architect Sir Denys Lasdun and our garden contains over 1,300 plants, each used as medicine around the world over the past 3,000 years.

Guided tours
We offer guided tours of the building and garden, and welcome bookings for groups of six or more. Refreshments can be booked and tours tailored for special interest groups. Guided tour charges are £5 per person (refreshments extra).

Royal College of Physicians, 11 St Andrews Place,
Regent's Park, London NW1 4LE.
Open: Monday– Friday, 9am– 5pm, free entry

Library, Archive and Museum Services
Tel: +44 (0)20 3075 1543
Email: history@rcplondon.ac.uk

Free entry



Royal College
of Physicians