

First Measurement of the Muon Neutrino Charged Current Single Pion Production Cross Section on Water with the T2K Near Detector

K. Abe,⁴⁷ C. Andreopoulos,^{45, 26} M. Antonova,²¹ S. Aoki,²³ A. Ariga,¹ S. Assylbekov,⁷ D. Autiero,²⁸ S. Ban,²⁴ M. Barbi,³⁹ G.J. Barker,⁵⁵ G. Barr,³⁵ P. Bartet-Friburg,³⁶ M. Batkiewicz,¹² F. Bay,¹⁰ V. Berardi,¹⁷ S. Berkman,³ S. Bhadra,⁵⁹ S. Bienstock,³⁶ A. Blondel,¹¹ S. Bolognesi,⁵ S. Bordoni,¹⁴ S.B. Boyd,⁵⁵ D. Brailsford,^{25, 16} A. Bravar,¹¹ C. Bronner,²² M. Buizza Avanzini,⁹ R.G. Calland,²² T. Campbell,⁷ S. Cao,²⁴ J. Caravaca Rodríguez,¹⁴ S.L. Cartwright,⁴³ R. Castillo,¹⁴ M.G. Catanesi,¹⁷ A. Cervera,¹⁵ D. Cherdack,⁷ N. Chikuma,⁴⁶ G. Christodoulou,²⁶ A. Clifton,⁷ J. Coleman,²⁶ G. Collazuol,¹⁹ D. Coplowe,³⁵ L. Cremonesi,³⁸ A. Dabrowska,¹² G. De Rosa,¹⁸ T. Dealtry,²⁵ P.F. Denner,⁵⁵ S.R. Dennis,²⁶ C. Densham,⁴⁵ D. Dewhurst,³⁵ F. Di Lodovico,³⁸ S. Di Luise,¹⁰ S. Dolan,³⁵ O. Drapier,⁹ K.E. Duffy,³⁵ J. Dumarchez,³⁶ S. Dytman,³⁷ M. Dziewiecki,⁵⁴ S. Emery-Schrenk,⁵ A. Ereditato,¹ T. Feusels,³ A.J. Finch,²⁵ G.A. Fiorentini,⁵⁹ M. Friend,^{13, *} Y. Fujii,^{13, *} D. Fukuda,³³ Y. Fukuda,³⁰ A.P. Furmanski,⁵⁵ V. Galymov,²⁸ A. Garcia,¹⁴ S.G. Giffin,³⁹ C. Giganti,³⁶ K. Gilje,³² F. Gizzarelli,⁵ M. Gonin,⁹ N. Grant,²⁵ D.R. Hadley,⁵⁵ L. Haegel,¹¹ M.D. Haigh,⁵⁵ P. Hamilton,¹⁶ D. Hansen,³⁷ J. Harada,³⁴ T. Hara,²³ M. Hartz,^{22, 51} T. Hasegawa,^{13, *} N.C. Hastings,³⁹ T. Hayashino,²⁴ Y. Hayato,^{47, 22} R.L. Helmer,⁵¹ M. Hierholzer,¹ A. Hillairet,⁵² A. Himmel,⁸ T. Hiraki,²⁴ S. Hirota,²⁴ M. Hogan,⁷ J. Holeczek,⁴⁴ S. Horikawa,¹⁰ F. Hosomi,⁴⁶ K. Huang,²⁴ A.K. Ichikawa,²⁴ K. Ieki,²⁴ M. Ikeda,⁴⁷ J. Imber,⁹ J. Insler,²⁷ R.A. Intonti,¹⁷ T.J. Irvine,⁴⁸ T. Ishida,^{13, *} T. Ishii,^{13, *} E. Iwai,¹³ K. Iwamoto,⁴⁰ A. Izmaylov,^{15, 21} A. Jacob,³⁵ B. Jamieson,⁵⁷ M. Jiang,²⁴ S. Johnson,⁶ J.H. Jo,³² P. Jonsson,¹⁶ C.K. Jung,^{32, †} M. Kabirnezhad,³¹ A.C. Kaboth,^{41, 45} T. Kajita,^{48, †} H. Kakuno,⁴⁹ J. Kameda,⁴⁷ D. Karlen,^{52, 51} I. Karpikov,²¹ T. Katori,³⁸ E. Kearns,^{2, 22, †} M. Khabibullin,²¹ A. Khotjantsev,²¹ D. Kielczewska,^{53, †} T. Kikawa,²⁴ H. Kim,³⁴ J. Kim,³ S. King,³⁸ J. Kisiel,⁴⁴ A. Knight,⁵⁵ A. Knox,²⁵ T. Kobayashi,^{13, *} L. Koch,⁴² T. Koga,⁴⁶ A. Konaka,⁵¹ K. Kondo,²⁴ A. Kopylov,²¹ L.L. Kormos,²⁵ A. Korzenev,¹¹ Y. Koshio,^{33, †} W. Kropp,⁴ Y. Kudenko,^{21, §} R. Kurjata,⁵⁴ T. Kutter,²⁷ J. Lagoda,³¹ I. Lamont,²⁵ E. Larkin,⁵⁵ P. Lasorak,^{38, 38} M. Laveder,¹⁹ M. Lawe,²⁵ M. Lazos,²⁶ T. Lindner,⁵¹ Z.J. Liptak,⁶ R.P. Litchfield,¹⁶ X. Li,³² A. Longhin,¹⁹ J.P. Lopez,⁶ T. Lou,⁴⁶ L. Ludovici,²⁰ X. Lu,³⁵ L. Magaletti,¹⁷ K. Mahn,²⁹ M. Malek,⁴³ S. Manly,⁴⁰ A.D. Marino,⁶ J. Marteau,²⁸ J.F. Martin,⁵⁰ P. Martins,³⁸ S. Martynenko,³² T. Maruyama,^{13, *} V. Matveev,²¹ K. Mavrokoridis,²⁶ W.Y. Ma,¹⁶ E. Mazzucato,⁵ M. McCarthy,⁵⁹ N. McCauley,²⁶ K.S. McFarland,⁴⁰ C. McGrew,³² A. Mefodiev,²¹ C. Metelko,²⁶ M. Mezzetto,¹⁹ P. Mijakowski,³¹ C.A. Miller,⁵¹ A. Minamino,²⁴ O. Mineev,²¹ S. Mine,⁴ A. Missert,⁶ M. Miura,^{47, †} S. Moriyama,^{47, †} Th.A. Mueller,⁹ S. Murphy,¹⁰ J. Myslik,⁵² T. Nakadaira,^{13, *} M. Nakahata,^{47, 22} K.G. Nakamura,²⁴ K. Nakamura,^{22, 13, *} K.D. Nakamura,²⁴ S. Nakayama,^{47, †} T. Nakaya,^{24, 22} K. Nakayoshi,^{13, *} C. Nantais,⁵⁰ C. Nielsen,³ M. Nirkko,¹ K. Nishikawa,^{13, *} Y. Nishimura,⁴⁸ P. Novella,¹⁵ J. Nowak,²⁵ H.M. O’Keefe,²⁵ R. Ohta,^{13, *} K. Okumura,^{48, 22} T. Okusawa,³⁴ W. Orszyszczak,⁵³ S.M. Oser,³ T. Ovsyannikova,²¹ R.A. Owen,³⁸ Y. Oyama,^{13, *} V. Palladino,¹⁸ J.L. Palomino,³² V. Paolone,³⁷ N.D. Patel,²⁴ M. Pavin,³⁶ D. Payne,²⁶ J.D. Perkin,⁴³ Y. Petrov,³ L. Pickard,⁴³ L. Pickering,¹⁶ E.S. Pinzon Guerra,⁵⁹ C. Pistillo,¹ B. Popov,^{36, ¶} M. Posiadala-Zezula,⁵³ J.-M. Poutissou,⁵¹ R. Poutissou,⁵¹ P. Przewlocki,³¹ B. Quilain,²⁴ T. Radermacher,⁴² E. Radicioni,¹⁷ P.N. Ratoff,²⁵ M. Ravonel,¹¹ M.A.M. Rayner,¹¹ A. Redij,¹ E. Reinherz-Aronis,⁷ C. Riccio,¹⁸ P. Rojas,⁷ E. Rondio,³¹ S. Roth,⁴² A. Rubbia,¹⁰ A. Rychter,⁵⁴ R. Sacco,³⁸ K. Sakashita,^{13, *} F. Sánchez,¹⁴ F. Sato,¹³ E. Scantamburlo,¹¹ K. Scholberg,^{8, †} S. Schoppmann,⁴² J. Schwehr,⁷ M. Scott,⁵¹ Y. Seiya,³⁴ T. Sekiguchi,^{13, *} H. Sekiya,^{47, 22, †} D. Sgalaberna,¹¹ R. Shah,^{45, 35} A. Shaikhiev,²¹ F. Shaker,⁵⁷ D. Shaw,²⁵ M. Shiozawa,^{47, 22} T. Shirahige,³³ S. Short,³⁸ M. Smy,⁴ J.T. Sobczyk,⁵⁸ H. Sobel,^{4, 22} M. Sorel,¹⁵ L. Southwell,²⁵ P. Stamoulis,¹⁵ J. Steinmann,⁴² T. Stewart,⁴⁵ P. Stowell,⁴³ Y. Suda,⁴⁶ S. Suvorov,²¹ A. Suzuki,²³ K. Suzuki,²⁴ S.Y. Suzuki,^{13, *} Y. Suzuki,²² R. Tacik,^{39, 51} M. Tada,^{13, *} S. Takahashi,²⁴ A. Takeda,⁴⁷ Y. Takeuchi,^{23, 22} H.K. Tanaka,^{47, †} H.A. Tanaka,^{50, 51, **} D. Terhorst,⁴² R. Terri,³⁸ T. Thakore,²⁷ L.F. Thompson,⁴³ S. Tobayama,³ W. Toki,⁷ T. Tomura,⁴⁷ C. Touramanis,²⁶ T. Tsukamoto,^{13, *} M. Tzanov,²⁷ Y. Uchida,¹⁶ A. Vacheret,¹⁶ M. Vagins,^{22, 4} Z. Vallari,³² G. Vasseur,⁵ T. Wachala,¹² K. Wakamatsu,³⁴ C.W. Walter,^{8, †} D. Wark,^{45, 35} W. Warzycha,⁵³ M.O. Wascko,^{16, 13} A. Weber,^{45, 35} R. Wendell,^{24, †} R.J. Wilkes,⁵⁶ M.J. Wilking,³² C. Wilkinson,¹ J.R. Wilson,³⁸ R.J. Wilson,⁷ Y. Yamada,^{13, *} K. Yamamoto,³⁴ M. Yamamoto,²⁴ C. Yanagisawa,^{32, ††} T. Yano,²³ S. Yen,⁵¹ N. Yershov,²¹ M. Yokoyama,^{46, †} J. Yoo,²⁷ K. Yoshida,²⁴ T. Yuan,⁶ M. Yu,⁵⁹ A. Zalewska,¹² J. Zalipska,³¹ L. Zambelli,^{13, *} K. Zaremba,⁵⁴ M. Ziembicki,⁵⁴ E.D. Zimmerman,⁶ M. Zito,⁵ and J. Żmuda⁵⁸

(The T2K Collaboration)

¹University of Bern, Albert Einstein Center for Fundamental Physics,
Laboratory for High Energy Physics (LHEP), Bern, Switzerland

²Boston University, Department of Physics, Boston, Massachusetts, U.S.A.

³University of British Columbia, Department of Physics and Astronomy, Vancouver, British Columbia, Canada

- ⁴University of California, Irvine, Department of Physics and Astronomy, Irvine, California, U.S.A.
⁵IRFU, CEA Saclay, Gif-sur-Yvette, France
- ⁶University of Colorado at Boulder, Department of Physics, Boulder, Colorado, U.S.A.
⁷Colorado State University, Department of Physics, Fort Collins, Colorado, U.S.A.
⁸Duke University, Department of Physics, Durham, North Carolina, U.S.A.
⁹Ecole Polytechnique, IN2P3-CNRS, Laboratoire Leprince-Ringuet, Palaiseau, France
¹⁰ETH Zurich, Institute for Particle Physics, Zurich, Switzerland
¹¹University of Geneva, Section de Physique, DPNC, Geneva, Switzerland
¹²H. Niewodniczanski Institute of Nuclear Physics PAN, Cracow, Poland
¹³High Energy Accelerator Research Organization (KEK), Tsukuba, Ibaraki, Japan
¹⁴Institut de Fisica d'Altes Energies (IFAE), The Barcelona Institute of Science and Technology, Campus UAB, Bellaterra (Barcelona) Spain
¹⁵IFIC (CSIC & University of Valencia), Valencia, Spain
¹⁶Imperial College London, Department of Physics, London, United Kingdom
- ¹⁷INFN Sezione di Bari and Università e Politecnico di Bari, Dipartimento Interuniversitario di Fisica, Bari, Italy
¹⁸INFN Sezione di Napoli and Università di Napoli, Dipartimento di Fisica, Napoli, Italy
¹⁹INFN Sezione di Padova and Università di Padova, Dipartimento di Fisica, Padova, Italy
²⁰INFN Sezione di Roma and Università di Roma "La Sapienza", Roma, Italy
²¹Institute for Nuclear Research of the Russian Academy of Sciences, Moscow, Russia
²²Kavli Institute for the Physics and Mathematics of the Universe (WPI), The University of Tokyo Institutes for Advanced Study, University of Tokyo, Kashiwa, Chiba, Japan
²³Kobe University, Kobe, Japan
²⁴Kyoto University, Department of Physics, Kyoto, Japan
²⁵Lancaster University, Physics Department, Lancaster, United Kingdom
²⁶University of Liverpool, Department of Physics, Liverpool, United Kingdom
- ²⁷Louisiana State University, Department of Physics and Astronomy, Baton Rouge, Louisiana, U.S.A.
²⁸Université de Lyon, Université Claude Bernard Lyon 1, IPN Lyon (IN2P3), Villeurbanne, France
²⁹Michigan State University, Department of Physics and Astronomy, East Lansing, Michigan, U.S.A.
³⁰Miyagi University of Education, Department of Physics, Sendai, Japan
³¹National Centre for Nuclear Research, Warsaw, Poland
- ³²State University of New York at Stony Brook, Department of Physics and Astronomy, Stony Brook, New York, U.S.A.
³³Okayama University, Department of Physics, Okayama, Japan
³⁴Osaka City University, Department of Physics, Osaka, Japan
³⁵Oxford University, Department of Physics, Oxford, United Kingdom
³⁶UPMC, Université Paris Diderot, CNRS/IN2P3, Laboratoire de Physique Nucléaire et de Hautes Energies (LPNHE), Paris, France
- ³⁷University of Pittsburgh, Department of Physics and Astronomy, Pittsburgh, Pennsylvania, U.S.A.
³⁸Queen Mary University of London, School of Physics and Astronomy, London, United Kingdom
³⁹University of Regina, Department of Physics, Regina, Saskatchewan, Canada
⁴⁰University of Rochester, Department of Physics and Astronomy, Rochester, New York, U.S.A.
⁴¹Royal Holloway University of London, Department of Physics, Egham, Surrey, United Kingdom
⁴²RWTH Aachen University, III. Physikalisches Institut, Aachen, Germany
⁴³University of Sheffield, Department of Physics and Astronomy, Sheffield, United Kingdom
⁴⁴University of Silesia, Institute of Physics, Katowice, Poland
- ⁴⁵STFC, Rutherford Appleton Laboratory, Harwell Oxford, and Daresbury Laboratory, Warrington, United Kingdom
⁴⁶University of Tokyo, Department of Physics, Tokyo, Japan
⁴⁷University of Tokyo, Institute for Cosmic Ray Research, Kamioka Observatory, Kamioka, Japan
⁴⁸University of Tokyo, Institute for Cosmic Ray Research, Research Center for Cosmic Neutrinos, Kashiwa, Japan
⁴⁹Tokyo Metropolitan University, Department of Physics, Tokyo, Japan
⁵⁰University of Toronto, Department of Physics, Toronto, Ontario, Canada
⁵¹TRIUMF, Vancouver, British Columbia, Canada
- ⁵²University of Victoria, Department of Physics and Astronomy, Victoria, British Columbia, Canada
⁵³University of Warsaw, Faculty of Physics, Warsaw, Poland
⁵⁴Warsaw University of Technology, Institute of Radioelectronics, Warsaw, Poland
⁵⁵University of Warwick, Department of Physics, Coventry, United Kingdom
⁵⁶University of Washington, Department of Physics, Seattle, Washington, U.S.A.
⁵⁷University of Winnipeg, Department of Physics, Winnipeg, Manitoba, Canada
⁵⁸Wroclaw University, Faculty of Physics and Astronomy, Wroclaw, Poland
⁵⁹York University, Department of Physics and Astronomy, Toronto, Ontario, Canada

(Dated: May 26, 2016)

The T2K off-axis near detector, ND280, is used to make the first differential cross section measurements of muon neutrino charged current single positive pion production on a water target at energies ~ 0.8 GeV. The differential measurements are presented as a function of the muon and

pion kinematics, in the restricted phase-space defined by $p_{\pi^+} > 200 \text{ MeV}/c$, $p_{\mu} > 200 \text{ MeV}/c$, $\cos(\theta_{\pi^+}) > 0.3$ and $\cos(\theta_{\mu}) > 0.3$. The total flux integrated ν_{μ} charged current single positive pion production cross section on water in the restricted phase-space is measured to be $\langle\sigma\rangle_{\phi} = 4.25 \pm 0.48$ (stat) ± 1.56 (syst) $\times 10^{-40} \text{ cm}^2/\text{nucleon}$. The total cross section is consistent with the NEUT prediction ($5.03 \times 10^{-40} \text{ cm}^2/\text{nucleon}$) and 2σ lower than the GENIE prediction ($7.68 \times 10^{-40} \text{ cm}^2/\text{nucleon}$). The differential cross sections are in good agreement with the NEUT generator. The GENIE simulation reproduces well the shapes of the distributions, but over-estimates the overall cross section normalization.

PACS numbers: 14.60.Pq, 14.60.Lm, 25.30.Pt, 29.40.Ka, 29.40.Mc

I. INTRODUCTION

The T2K long baseline neutrino experiment [1] has the primary goal to precisely measure neutrino oscillation parameters through measurements of ν_e appearance and ν_{μ} disappearance from a ν_{μ} beam. As neutrinos are chargeless and colorless, neutrino oscillation experiments rely on the detection of charged particles coming from charged current (CC) and neutral current (NC) interactions to infer neutrino properties, e.g. CC quasi-elastic (QE) interactions allow the calculation of the neutrino energy from the lepton kinematics. The knowledge of ν_{μ} and ν_e interaction cross sections is then fundamental to infer neutrino properties correctly. ν_{μ} CC resonant interactions are part of the signal and sometimes of the background of oscillation experiments, and a better understanding of this channel could be beneficial not only to T2K, but to the neutrino community in general, as there are discrepancies between models and experimental data.

Both the MiniBooNE [2] and MINERvA [3] collaborations provided measurements of the CC single positive pion production ($\text{CC}1\pi^+$) cross sections in mineral oil and plastic scintillator, respectively. The $\text{CC}1\pi^+$ cross section is described by the particles leaving the nucleus, i.e. one muon, one positive pion and any number of nucleons. There are large discrepancies between the MiniBooNE and MINERvA experiments, and the historic ANL [4] and BNL [5] bubble chamber results, which could be due to nuclear effects that if not modelled correctly can modify the effective measured cross-section. The MiniBooNE and MINERvA results show also significant normalization and shape discrepancies between each other [3], and currently no theoretical model can explain all the pion production data available. Additional pion production data can help to constrain the pion production models and give valuable information on the nucleon- Δ axial form factor [6, 7].

We present the first $\text{CC}1\pi^+$ differential cross section measurements on water. A $\text{CC}1\pi^+$ measurement on water will have a strong impact on the T2K oscillation analysis, as current results suffer from large cross section systematic uncertainties related to the differences in targets between near and far detectors (carbon versus water) [8]. These data will also be beneficial to future atmospheric and long-baseline experiments, that plan to use a water target, such as the Hyper-Kamiokande experiment [9].

II. T2K EXPERIMENT

The T2K long baseline neutrino experiment uses the J-PARC facility in Tokai, Japan, to produce 30 GeV protons, which produce charged pions by colliding with a graphite target and consequently result in a high purity ν_{μ} beam. The beam center axis is directed 2.5° off-axis towards Super-Kamiokande [10] at 295 km from J-PARC. Two near detectors are located at 280 m from the target, the on-axis near detector (INGRID [11]) and the off-axis near detector (ND280).

Neutrino Beam Flux—The predicted neutrino beam flux [12] peaks at 0.6 GeV and its fractional composition is 92.6% ν_{μ} , 6.2% $\bar{\nu}_{\mu}$, 1.1% ν_e , 0.1% $\bar{\nu}_e$. The proton interactions with the graphite target are simulated with the FLUKA2008 package [13], The propagation of secondary and tertiary pions and kaons and their decays to neutrinos is simulated with GEANT3 [14]. The hadron interactions are modeled with GCALOR [15] and tuned to hadron production data from external experiments, such as the CERN NA61/SHINE experiment [16, 17].

Neutrino Interaction Model—Based on the prediction of the neutrino flux, the NEUT [18] (version 5.1.4.2) event generator is used to simulate neutrino interactions in ND280.

For Changed Current Quasi Elastic (CCQE) and Neutral Current Quasi Elastic (NCQE) interactions, NEUT uses the Llewellyn Smith model [19] integrated with the relativistic Fermi gas (RFG) model by Smith and Morniz to describe the nucleons within the nucleus [20]. The outgoing nucleon is also required to have larger momentum than the Fermi surface momentum (Pauli blocking), which is 217 MeV/c for carbon and 225 MeV/c oxygen.

NEUT uses the Rein-Sehgal model for resonant interactions [21], considering 18 resonances with masses below 2 GeV/c² and their interference terms. In addition 20%

* also at J-PARC, Tokai, Japan

† affiliated member at Kavli IPMU (WPI), the University of Tokyo, Japan

‡ deceased

§ also at National Research Nuclear University "MEPhI" and Moscow Institute of Physics and Technology, Moscow, Russia

¶ also at JINR, Dubna, Russia

** also at Institute of Particle Physics, Canada

†† also at BMCC/CUNY, Science Department, New York, New York, U.S.A.

of the Δ resonances undergo pion-less Δ decay, in which the Δ is absorbed by the nuclear medium without emitting any pions: $\Delta + N \rightarrow N' + N''$. The NEUT pion production model is tuned using neutrino interaction data from the MiniBooNE experiments [2, 22], as explained in Ref [8]. In particular, the axial mass for resonant $CC1\pi^+$ interactions is set to 1.41 GeV, and the overall $CC1\pi^+$ normalization for energies less than 2.5 GeV is further increased by 15% compared to predictions when the axial mass is set to 1.41 GeV.

Coherent pion production is simulated for both NC and CC interactions using the Rein-Sehgal model [23], including the PCAC (Partially Conserved Axial vector Current) lepton mass correction for CC interactions [24].

DIS (deep inelastic scattering) processes are simulated using GRV98 parton distribution functions [25] and corrections following the Bodek and Yang model [26] to improve the agreement with experiments in the low- Q^2 region. To avoid double counting with the single pion resonant production, only multiple pion production processes are considered for the invariant mass of the recoiling hadron system $W < 2 \text{ GeV}/c^2$. PYTHIA/JETSET [27] is used for hadronisation at energies above 2 GeV, and an internal NEUT method is used at lower energies.

After the simulation of the initial neutrino-nucleon interaction, final state interactions are simulated with the cascade model [28]. Each particle is propagated inside the nucleus with steps determined by the mean free path. The mean free path depends on the position inside the nucleus and the momentum of the particle. At each step, an interaction is generated according to the probability calculated from each cross section such as charge exchange, absorption or scattering. If an interaction occurs, the resulting particles are used for stepping through the rest of the nucleus. This process continues until all particles are either absorbed in the nucleus or escape it. Data from several pion scattering experiments are used to tune this model [8].

Additional information on the models used to simulate the neutrino interactions and the hadron transport in the nuclear medium can be found in references [8, 18].

The results in this paper are also compared to the GENIE generator [29], as it provides a general framework valid over a large range of experiments, targets and neutrino energy. GENIE uses essentially the same models as NEUT for the neutrino interactions simulation, but they differ in the implementation and value of some of the parameters, such as M_A^{RES} which is set to 1.12 GeV. Single pion production (before final state interactions) comes from resonant and coherent processes in NEUT, whereas GENIE also considers DIS contributions to it. Although GENIE considers a lower value of M_A^{RES} , the predicted single positive pion production cross-section is larger than in NEUT, because DIS processes are allowed to contribute to this state.

Near detector—ND280 is a complex of different sub-detectors enclosed in the refurbished UA1/NOMAD magnet. The origin of the ND280 coordinate system is at

the center of the magnet and the 0.2 T magnetic field is along the $+x$ direction. The z axis is along the nominal neutrino beam axis, and x and y axes are horizontal and vertical, respectively.

The ND280 Tracker region contains two fine-grained detectors (FGDs [30]) which are used as the neutrino interaction target, sandwiched between three gaseous time projection chambers (TPCs [31]) which are used to track charged particles. The most upstream FGD (FGD1) primarily consists of polystyrene scintillator bars with layers oriented alternately in the x and y directions allowing 3D tracking of charged particles. The downstream FGD (FGD2) has a similar structure, but the polystyrene bars are interleaved with water layers, creating a modular structure of water layer + x layer + y layer + water layer, and so on (see Figure 1). The areal density of an xy module and a water module are respectively $2146.3 \pm 14.4 \text{ mg/cm}^2$ and $2792.6 \pm 13.4 \text{ mg/cm}^2$. This structure allows the measurement of neutrino interactions on water.

The electromagnetic calorimeters (ECals [32]), made of layers of lead and scintillator bars, surround the Tracker region (Barrel-ECals) with one module downstream of it (Ds-ECal). Upstream of the Tracker there is a π^0 detector (PØD [33]), consisting of scintillator, water and brass layers.

Magnet return yokes surround the entire detector to make the magnetic field uniform and contain it inside the detector. Plastic scintillators in the yoke form the side muon range detectors (SMRDs [34]).

The analysis here presented uses FGD2 as the active interaction target, where in a signal event the neutrino interacts with a nucleus in the water layer, and the charged lepton coming from a CC interaction is tracked in the downstream scintillator layers. These results are based on data taken from November 2010 to May 2013. The good quality data collected during this period corresponds to 5.6×10^{20} protons on target (p.o.t.).

III. SELECTION OF ν_μ $CC1\pi^+$ INTERACTIONS IN WATER

Muon neutrino interactions are selected by using the highest momentum negative track starting in the fiducial volume (FV) of the FGD2. The FGD2 FV begins 58 mm inward from the lateral edges of the FGD2 and 7.5 mm inward from the upstream FGD2 edge (as shown in Figure 1). These tracks are required to enter the TPC3 (located immediately downstream of FGD2) and deposit energy compatible with a muon-like track. Additional tracks matched between the FDG and TPC associated with the same muon candidate vertex are tagged as either protons or positive, negative or neutral pions by looking at the trajectory and energy deposit in the TPCs, and at electromagnetic showers in the ECals. More details on the ν_μ CC inclusive and multi-pion selections can be found in References [8, 35], respectively, where the only




FIG. 1. Schematic view of FGD2 and its fiducial volume (FV) delimited by the red line. The first upstream scintillator layer is not included in the FV.

differences are that in this analysis interactions in FGD2 are selected, rather than in FGD1, and the ECals are used to tag neutral pions. $CC1\pi^+$ -like events are selected by requiring one muon, one positive pion, no other additional pions and any number of nucleons.

Because of the structure of the FGD2 (see Figure 1), interaction vertices occurring in a water module will be reconstructed in the x layer downstream of it. A water-enhanced sample can be selected by requiring the vertex to be in the x layer, whilst a scintillator sample can be selected by requiring the reconstructed vertex to be in the y layer.

The signal sample of this analysis is composed of 1402 selected $CC1\pi^+$ water-enhanced events in the full phase-space, with 30.9% purity of true ν_μ $CC1\pi^+$ interactions on water. To avoid relying on the simulation to describe regions of efficiency < 0.1 , the analysis presented restricts the kinematic phase-space to the region defined by $p_\mu > 200$ MeV/ c , $p_{\pi^+} > 200$ MeV/ c , $\cos(\theta_\mu) > 0.3$ and $\cos(\theta_{\pi^+}) > 0.3$. With these restrictions in the phase-space of the signal definition, the signal efficiency goes from 13.3% to 30.7%.

The signal sample is selected with a purity of 39.9% in the restricted phase-space. Multi-pion interactions can be mis-identified as $CC1\pi^+$ interactions when one or more pions are absorbed by the detector or simply not reconstructed; 29.2% of the signal sample is composed of this background. $CC0\pi$ interactions (3.9%) come into the selection when the proton is mis-identified as a π^+ . The total background from $CC1\pi^+$ interactions occurring

TABLE I. Percentage of true NEUT topologies ($CC1\pi^+$ and $CCnon1\pi^+$) in the restricted phase-space, and number of data events in different modules of the FGD2 for the signal ($CC1\pi^+$ water-enhanced) and two external samples ($CC1\pi^+$ scintillator and $CC1\pi^+n\pi$ water-enhanced).

True NEUT topology	Selected samples		
	$CC1\pi^+$ water	$CC1\pi^+$ scintillator	$CC1\pi^+n\pi$ water
$CC1\pi^+$ water	39.9 %	5.9 %	7.7 %
$CC1\pi^+$ scint	25.7 %	54.6 %	4.8 %
CC non $1\pi^+$ water	18.5 %	8.3 %	49.0 %
CC non $1\pi^+$ scint	14.6 %	28.7 %	36.5 %
non ν_μ CC	0.6 %	0.9 %	1.2 %
Out of FV	0.5 %	1.7 %	0.9 %
Data in reduced phase-space	1275	431	885
Data in full phase-space	1402	491	944

in the scintillator amounts to 25.7% of the signal sample, including interactions occurring in the y layer whose vertex is reconstructed in the x layer. Non- ν_μ CC interactions (0.6%) include both NC and CC interactions due to the $\bar{\nu}$ and ν_e components in the beam. They mainly come into the selection when a π^- from a NC interaction is mis-identified as the muon candidate.

These backgrounds are constrained with two external samples. A sample of selected ν_μ CC events with one π^+ and at least one, but maximum 3, negative or neutral pions ($CC1\pi^+n\pi$ water-enhanced sample) is used to constrain the non- $CC1\pi^+$ interactions, which include the CC non $1\pi^+$, non ν_μ CC and out of FV backgrounds. A sample of selected ν_μ $CC1\pi^+$ events in the y layers of the FGD2 ($CC1\pi^+$ scintillator sample) is used to constrain the background coming from the interactions in the scintillator. Table I shows the composition of the signal and external samples according to the NEUT generator. Distributions of the pion kinematics (p_{π^+} and $\cos(\theta_{\pi^+})$), the muon kinematics (p_μ and $\cos(\theta_\mu)$), the cosine of the angle between the muon and the pion ($\cos(\theta_{(\mu,\pi^+)})$) and reconstructed neutrino energy (E_ν^{rec}) in the selected sample are shown in Figure 2. The reconstructed neutrino energy is found by applying 4-momentum conservation and assuming the target nucleon is at rest and the remaining final-state particle is a nucleon:

$$E_\nu^{rec} = \frac{m_\mu^2 + m_\pi^2 - 2m_N E_f + 2(\mathbf{p}_\mu \cdot \mathbf{p}_\pi)}{(2E_f - \mathbf{p}_\mu \cos(\theta_\mu) - \mathbf{p}_\pi \cos(\theta_{\pi^+}) - m_N)} \quad (1)$$

where m_μ , m_π and m_N are the masses of the muon, the pion and the nucleon respectively; $E_f = E_\mu + E_\pi$; p_x , \mathbf{p}_x and θ_x are the 4-momentum, 3-momentum and angle with the neutrino direction of the particle considered ($x = \mu, \pi^+$). Distributions of the pion momentum in the external samples are found in Figure 3.

Systematic uncertainties—The TPC and FGD detector systematic uncertainties are the same as the ones described in References [8, 35]. The ECal particle identification systematic uncertainties are evaluated with high




FIG. 2. Reconstructed pion kinematics (top), muon kinematics (middle), $\cos(\theta_{(\mu, \pi+)})$ (bottom left) and neutrino energy (bottom right) distributions of the events in the ν_μ CC $1\pi^+$ water-enhanced sample. The NEUT Monte Carlo prediction is separated into the ν_μ CC $1\pi^+$ interactions in water, ν_μ CC $1\pi^+$ interactions in scintillator, ν_μ CCnon $1\pi^+$ interactions in water, ν_μ CCnon $1\pi^+$ interactions in scintillator, non ν_μ CC interactions, and interactions outside of the FGD2 FV. The last bin in the p_{π^+} , p_μ and E_ν^{rec} distributions contains all the over-flow events.

purity samples of electrons and muons, as explained in Reference [36].

The isolated ECal reconstruction systematic uncertainty is evaluated with a control sample of both isolated and non-isolated ECal objects, due to the difficulties of finding a control sample with just isolated ECal objects. The efficiency is found to be 0.303 ± 0.003 in simulation and 0.315 ± 0.009 in data for the Barrel-ECal, and 0.826 ± 0.002 in simulation and 0.839 ± 0.007 for the Ds-ECal. These efficiencies are used to correct the sim-

ulation efficiency for tagging isolated-ECal objects only, which is 0.352 for the Barrel-ECal and 0.163 for the Ds-ECal.

The FGD water modules mass uncertainty is 0.55%. The FGD layer migration uncertainties have been evaluated in detail for this analysis. These migrations are divided into forward (i.e. when the reconstructed vertex is a layer downstream of the true vertex) and backward migrations (i.e. when the reconstructed vertex is a layer upstream of the true vertex). The forward migrations




FIG. 3. Reconstructed pion momentum distributions of the events in the two external samples: $CC1\pi^+$ scintillator (left) and $CC1\pi^+n\pi$ water-enhanced (right). The last bin contains all the over-flow events.

come from a hit reconstruction inefficiency. Their overall uncertainty is estimated to be 3.3% with a control sample of cosmic muons passing through both FGDs. The backward migrations come from low energy backward going particles that are fitted with the muon track and move the vertex one or more layers upstream. These latter migrations are estimated using the $CC0\pi$ and CC multi-pion samples in FGD2: a normalization uncertainty of 30% is assigned to them.

The flux uncertainties are evaluated with beamline and hadron production measurements. The hadron production uncertainties dominate the neutrino flux uncertainties, with a smaller contribution from the neutrino beam direction and proton beam uncertainties. The systematic uncertainty for the ν_μ flux at ND280 varies from 10% and 15% depending on the neutrino energy [12].

The uncertainties related to the cross section model (final state interactions, CCQE model, pion production model and nuclear model) are constrained using external data and comparisons between different existing models. A summary of these uncertainties can be found in Reference [8].

IV. UNFOLDING METHOD

The Bayesian unfolding technique by d'Agostini [37] has been successfully used by past T2K cross section measurements to extract the cross sections (see References [35, 38]). The first estimate of the true distribution is found by applying the unsmearing matrix $P(t_j|r_i)$ (found with Bayes' theorem) to the data distribution:

$$\hat{N}_{t_j} = \frac{1}{\epsilon_j} \sum_i P(t_j|r_i) (N_{r_i} - \sum_{\text{all backgrounds } k} \alpha_k B_{r_i,k}), \quad (2)$$

where t_j (r_i) indicates the true (reconstructed) bin for each observable, N_{r_i} is the number of reconstructed events in bin r_i , $B_{r_i,k}$ is the number of predicted events

in bin r_i of background type k , α_k is a normalization constant derived from the external samples, and ϵ_j is the true efficiency in bin t_j .

Eq. 2 uses a background subtraction where the coefficients α_k are 1 if that part of the background is not constrained by any external sample, or otherwise calculated as:

$$\alpha_k = \frac{C_{\text{data},k}}{C_{\text{MC},k}}, \quad (3)$$

where $C_{\text{data},k}$ is the total number of events in external sample k in data and $C_{\text{MC},k}$ is the total number of events in external sample k in MC. In this analysis the background is divided into 2 groups: the $CC1\pi^+$ interactions in scintillator or in the scintillator-like component of the water modules, that are constrained with the $CC1\pi^+$ scintillator sample; the non $CC1\pi^+$ background which is constrained with the $CC1\pi^+n\pi$ water-enhanced sample.

The FGD2 water modules are composed of oxygen (73.83%), carbon (15.05%), hydrogen (10.48%), silicon (0.39%), and magnesium (0.25%). The carbon, silicon and magnesium come from the polycarbonate structure that enclose the liquid water. They compose the scintillator-like component of the water-modules and can be subtracted out with the x -layer as they have similar composition.

The effect of the systematic uncertainties on the cross section measurements is evaluated by using pseudo-experiments for each error source to calculate the covariance matrix:

$$V_{ij}^s = \frac{1}{N} \sum_{s_n=1}^N (\sigma_i^{s_n} - \sigma_i^{\text{nom}})(\sigma_j^{s_n} - \sigma_j^{\text{nom}}), \quad (4)$$

where $\sigma_i^{s_n}$ is the differential cross section in bin i evaluated with throw n of the uncertainty s , and σ_i^{nom} is the nominal differential cross section in bin i . Statistical and systematic uncertainties are evaluated by varying the contents of each bin according to Poisson and Gaussian statistics, respectively.




FIG. 4. Unfolded ν_μ CC $1\pi^+$ differential cross sections as a function of pion kinematics (top), muon kinematics (center), $\cos(\theta_{(\mu,\pi^+)})$ (bottom left) and E_V^{rec} (bottom right) in the reduced phase-space of $p_{\pi^+} > 200$ MeV/c, $p_\mu > 200$ MeV/c, $\cos(\theta_{\pi^+}) > 0.3$ and $\cos(\theta_\mu) > 0.3$. For the E_V^{rec} , the $\sigma(E)$ is presented as a model dependent result. The inner (outer) error bars show the statistical (total) uncertainty on the data. The dashed (solid) line shows the NEUT, version 5.1.4.2, (GENIE, version 2.6.4) prediction.

V. CROSS-SECTION RESULTS

For a given variable X , the flux integrated differential cross section for bin t_k is defined as:

$$\left\langle \frac{\partial \sigma}{\partial X} \right\rangle_{t_k} = \frac{N_{t_k}^{unfolded}}{T \Phi \Delta X_{t_k}}, \quad (5)$$

where $N_{t_k}^{unfolded}$ is the estimated number of events in bin t_k (as given by Eq. 2), T is the number of target

nucleons, Φ is the ν_μ flux per unit area and integrated over neutrino energy (as detailed in Reference [12]), and ΔX_{t_k} is the width of bin t_k . Even though single pion resonant production has a threshold at 480 MeV, no cut is applied to the ν_μ flux, as the CC $1\pi^+$ signal definition includes processes with different thresholds as well.

The number of target nucleons is computed considering only the oxygen and hydrogen in the FGD2 water modules, as the carbon, silicon and magnesium components are removed by the Bayesian unfolding with background

subtraction. The total number of target nucleons is found to be:

$$T = N_A \cdot V_{FV} \cdot \rho \sum_{a=O,H} f_a \frac{A_a}{M_a} = 2.55 \cdot 10^{29},$$

where $N_A = 6.022 \cdot 10^{23} \text{ mol}^{-1}$ is the Avogadro number, V_{FV} is the volume of the modules considered inside the FV, $\rho = \rho_{area}/\Delta z$ is the total density of the water modules of the FGD2 ($\rho_{area} = 2798.7 \pm 5.4 \text{ mg/cm}^2$ is the total areal density and $\Delta z = 2.79 \text{ cm}$ is the width of each water module); a runs over the elements present in the water modules; f_a is the mass fraction; A_a represents the averaged number of nucleons per nucleus; and M_a is the atomic mass.

Figure 4 shows the differential cross section as a function of pion kinematics (top), muon kinematics (center), $\cos(\theta_{(\mu,\pi+)})$ (bottom left), and E_{ν}^{rec} (bottom right). For the E_{ν}^{rec} the $\sigma(E)$ is presented as a model dependent result, as the E_{ν}^{rec} is unfolded to the true neutrino energy as predicted by NEUT. The NEUT and GENIE (version 2.6.4 [29]) predictions are also shown in the plots. The differential cross sections obtained are compatible with the NEUT prediction, but a small suppression is seen at $0.5 < p_{\pi^+} < 0.7 \text{ GeV}$ and $\cos(\theta_{\pi^+}) > 0.95$. This might be linked to the model for CC coherent interactions used in NEUT: NEUT greatly over-estimates the amount of coherent interactions especially at low E_{π^+} [39]. The GENIE simulation reproduces well the shapes of the distributions, but over-estimates the overall cross section normalization.

The total flux integrated cross section is computed as:

$$\langle \sigma \rangle_{\Phi} = \frac{N_{\text{total}}}{T \cdot \Phi}. \quad (6)$$

The total flux integrated ν_{μ} CC single positive pion production cross section on water in the restricted phase-space is measured to be $\langle \sigma \rangle_{\Phi} = 4.25 \pm 0.48 \text{ (stat)} \pm 1.56 \text{ (syst)} \times 10^{-40} \text{ cm}^2/\text{nucleon}$. This result is compatible with the NEUT prediction of $5.03 \times 10^{-40} \text{ cm}^2/\text{nucleon}$, and about 2σ away from the GENIE prediction $7.68 \times 10^{-40} \text{ cm}^2/\text{nucleon}$. The dominant systematic uncertainties on this result are those related to the cross section model (23.9%) and flux parameters (25.5%). The flux uncertainties are larger than the initial uncertainty size because of the background subtraction procedure. Final state interactions and detector systematic uncertainties contribute with 5.3% and 10.8%, respectively. The data and MC statistical errors are estimated as 10.7% and 3.3%, respectively. Figure 5 shows the total ν_{μ} CC1 π^+ cross section on water in the reduced phase-space of $p_{\pi^+} > 200 \text{ MeV}/c$, $p_{\mu} > 200 \text{ MeV}/c$, $\cos(\theta_{\pi^+}) > 0.3$ and $\cos(\theta_{\mu}) > 0.3$, with the T2K ν_{μ} flux and the NEUT and GENIE predictions.

The data related to this measurement can be found together with the cross section results obtained when unfolding the muon kinematics and neutrino energy distributions in Reference [40].




FIG. 5. Total ν_{μ} CC1 π^+ cross section on water in the reduced phase-space of $p_{\pi^+} > 200 \text{ MeV}/c$, $p_{\mu} > 200 \text{ MeV}/c$, $\cos(\theta_{\pi^+}) > 0.3$ and $\cos(\theta_{\mu}) > 0.3$. The T2K data point is placed at the ν_{μ} flux mean energy. The vertical error represents the total uncertainty, and the horizontal bar represents 68% of the flux each side of the mean. The T2K flux distribution is shown in gray. The NEUT (version 5.1.4.2) and GENIE (version 2.6.4) predictions are the total ν_{μ} CC1 π^+ predictions as a function of neutrino energy. The NEUT and GENIE averages are the flux-averaged predictions.

Conclusion—The T2K off-axis near detector ND280 is used to extract the first ν_{μ} CC1 π^+ differential cross sections on water as a function of the pion kinematics and muon-pion angle. These results will be beneficial to the T2K experiment and the neutrino community in general, as a better understanding of neutrino induced pion production on water at energy below 2 GeV would result in a higher sensitivity to the measurement of oscillation parameters. The cross section is evaluated in the restricted phase-space defined by $p_{\mu} > 200 \text{ MeV}/c$, $p_{\pi^+} > 200 \text{ MeV}/c$, $\cos(\theta_{\mu}) > 0.3$ and $\cos(\theta_{\pi^+}) > 0.3$. The results are in good agreement with the NEUT generator and a general suppression is seen compared to the GENIE generator. The total ν_{μ} CC1 π^+ cross section on water is found to be $\langle \sigma \rangle_{\Phi} = 4.25 \pm 0.48 \text{ (stat)} \pm 1.56 \text{ (syst)} \times 10^{-40} \text{ cm}^2/\text{nucleon}$, which is in good agreement with the NEUT prediction and is 2σ lower than the GENIE prediction.

Acknowledgements—We thank the J-PARC staff for superb accelerator performance and the CERN NA61 Collaboration for providing valuable particle production data. We acknowledge the support of MEXT, Japan; NSERC (Grant No. SAPPJ-2014-00031), NRC and CFI, Canada; CEA and CNRS/IN2P3, France; DFG, Germany; INFN, Italy; National Science Centre (NCN), Poland; RSF, RFBR and MES, Russia; MINECO and ERDF funds, Spain; SNSF and SERI, Switzerland; STFC, UK; and DOE, USA. We also thank CERN for the UA1/NOMAD magnet, DESY for the HERA-B magnet mover system, NII for SINET4, the WestGrid and SciNet consortia in Compute Canada, and GridPP in the United Kingdom. In addition, participation of individual

researchers and institutions has been further supported by funds from ERC (FP7), H2020 Grant No. RISE-

GA644294-JENNIFER, EU; JSPS, Japan; Royal Society, UK; and the DOE Early Career program, USA.

-
- [1] K. Abe *et al.* (T2K Collaboration), Nucl. Instrum. Methods **A659**, 106 (2011).
- [2] A. A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), Phys. Rev. D **83**, 052007 (2011).
- [3] B. Eberly *et al.* (Minerva Collaboration), Physical Review D **92**, 092008 (2015).
- [4] G. M. Radecky *et al.* (ANL Collaboration), Phys. Rev. D **25**, 1161 (1982).
- [5] T. Kitagaki *et al.* (BNL Collaboration), Phys. Rev. D **34**, 2554 (1986).
- [6] O. Lalakulich and U. Mosel, Phys. Rev. C **88**, 017601 (2013).
- [7] M. Martini and M. Ericson, Phys. Rev. C **90**, 025501 (2014).
- [8] K. Abe *et al.* (T2K Collaboration), Phys. Rev. D **91**, 072010 (2015).
- [9] K. Abe *et al.* (Hyper-Kamiokande Collaboration), Progress of Theoretical and Experimental Physics **2015**, 053C02 (2015).
- [10] S. Fukuda *et al.* (Super-Kamiokande Collaboration), Nucl. Instrum. Methods **A501**, 418 (2003).
- [11] M. Otani *et al.*, Nucl. Instrum. Methods **A623**, 368 (2010).
- [12] K. Abe *et al.* (T2K Collaboration), Phys. Rev. D **87**, 012001 (2013).
- [13] G. Battistoni, F. Cerutti, A. Fasso, A. Ferrari, S. Muraro, J. Ranft, S. Roesler, and P. Sala, in *AIP Conference Proceedings*, Vol. 896 (2007) pp. 31–49.
- [14] R. Brun, F. Carminati, and S. Giani, CERN-W5013 (1994).
- [15] C. Zeitnitz and T. A. Gabriel, In Proc. of International Conference on Calorimetry in High Energy Physics (1993).
- [16] N. Abgrall *et al.* (NA61/SHINE Collaboration), Phys. Rev. C **84**, 034604 (2011).
- [17] N. Abgrall *et al.* (NA61/SHINE Collaboration), Phys. Rev. C **85**, 035210 (2012).
- [18] Y. Hayato, Nucl. Phys. Proc. Supp. **B112**, 171 (2002).
- [19] C. L. Smith, Physics Reports **3**, 261 (1972).
- [20] R. Smith and E. Moniz, Nuclear Physics B **43**, 605 (1972).
- [21] D. Rein and L. M. Sehgal, Annals of Physics **133**, 79 (1981).
- [22] A. A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), Phys. Rev. D **83**, 052009 (2011).
- [23] D. Rein and L. M. Sehgal, Nuclear Physics B **223**, 29 (1983).
- [24] D. Rein and L. Sehgal, Physics Letters B **657**, 207 (2007).
- [25] M. Glück, E. Reya, and A. Vogt, The European Physical Journal C-Particles and Fields **5**, 461 (1998).
- [26] A. Bodek and U. Yang, arXiv preprint hep-ex/0308007 (2003).
- [27] T. Sjöstrand, Computer Physics Communications **82**, 74 (1994).
- [28] L. Salcedo, E. Oset, M. Vicente-Vacas, and C. Garcia-Rocio, Nuclear Physics A **484**, 557 (1988).
- [29] C. Andreopoulos *et al.*, Nucl. Instrum. Methods **A614**, 87 (2010).
- [30] P. Amaudruz *et al.*, Nucl. Instrum. Methods **A696**, 1 (2012).
- [31] N. Abgrall *et al.*, Nucl. Instrum. Methods **A637**, 25 (2011).
- [32] D. Allan *et al.*, JINST **8**, P10019 (2013).
- [33] S. Assylbekov *et al.*, Nucl. Instrum. Methods **A686**, 48 (2012).
- [34] S. Aoki *et al.*, Nucl. Instrum. Methods **A698**, 135 (2013).
- [35] K. Abe *et al.* (T2K Collaboration), Phys. Rev. D **87**, 092003 (2013).
- [36] K. Abe *et al.* (T2K Collaboration), Phys. Rev. D **89**, 092003 (2014).
- [37] G. D’Agostini, Nucl. Instrum. Methods **A362**, 487 (1995).
- [38] K. Abe *et al.* ((T2K Collaboration)), Phys. Rev. Lett. **113**, 241803 (2014).
- [39] K. Abe *et al.* (T2K Collaboration), arXiv preprint arXiv:1604.04406 (2016).
- [40] K. Abe *et al.*, “T2K public data,” <http://t2k-experiment.org/results/nd280data-numu-cc1pi-xs-on-h2o-2015>.