

1 **Title: Comment on “The extent of forest in dryland biomes”**

2
3 **Authors:** Daniel M. Griffith^{1*}, Caroline E.R. Lehmann^{2,3}, Caroline A.E. Strömberg⁴, Catherine
4 L. Parr^{5,6,7}, R. Toby Pennington^{8,9}, Mahesh Sankaran^{10,11}, Jayashree Ratnam¹⁰, Christopher J.
5 Still¹, Rebecca L. Powell¹², Niall P. Hanan¹³, Jesse B. Nippert¹⁴, Colin P. Osborne^{15,16}, Stephen
6 Good¹⁷, T. Michael Anderson¹⁸, Ricardo M. Holdo¹⁹, Joseph W. Veldman^{20,21}, Giselda
7 Durigan²², Kyle W. Tomlinson²³, William A. Hoffmann²⁴, Sally Archibald^{3,25}, and William J.
8 Bond^{26,27}

9
10 **Affiliations:**

11 ¹Forest Ecosystems and Society, Oregon State University, Corvallis, Oregon, U.S.A.

12 ²School of GeoSciences, University of Edinburgh, Edinburgh EH9 EFF, U.K.

13 ³Centre for African Ecology, School of Animal, Plant and Environmental Sciences, University of
14 the Witwatersrand, Johannesburg, Private Bag X3, WITS, 2050, South Africa

15 ⁴Department of Biology & Burke Museum of Natural History & Culture, University of
16 Washington, Seattle, Washington, U.S.A.

17 ⁵Department of Earth, Ocean and Ecological Sciences, School of Environmental Sciences,
18 University of Liverpool, Liverpool

19 ⁶School of Animal, Plant and Environmental Sciences, University of the Witwatersrand,
20 Johannesburg, South Africa

21 ⁷Department of Zoology & Entomology, University of Pretoria, Pretoria 0002, South Africa

22 ⁸Royal Botanic Garden Edinburgh, Edinburgh EH3 5LR, UK

23 ⁹Department of Geography, University of Exeter, EX4 4RJ, UK

24 ¹⁰National Centre for Biological Sciences, Tata Institute of Fundamental Research, GKVK
25 Campus, Bellary Road, Bangalore, India.

26 ¹¹School of Biology, University of Leeds, Leeds LS29JT, UK.

27 ¹²Department of Geography and the Environment, University of Denver, Denver, Colorado,
28 U.S.A.

29 ¹³Plant and Environmental Sciences, New Mexico State University, Las Cruces, New Mexico,
30 U.S.A.

31 ¹⁴Division of Biology, Kansas State University, Manhattan, Kansas, U.S.A

32 ¹⁵Department of Animal and Plant Sciences, University of Sheffield, Sheffield S10 2TN, U.K.

33 ¹⁶Grantham Centre for Sustainable Futures, University of Sheffield, Sheffield S10 2TN, U.K.

34 ¹⁷Department of Biological & Ecological Engineering, Oregon State University, Corvallis,
35 Oregon, U.S.A.

36 ¹⁸Department of Biology, Wake Forest University, Winston-Salem, North Carolina, U.S.A.

37 ¹⁹Odum School of Ecology, University of Georgia, Athens, Georgia, U.S.A.

38 ²⁰Department of Ecosystem Science and Management, Texas A&M University, College Station

39 Texas, USA

40 ²¹Instituto Boliviano de Investigación Forestal, Casilla 6204, Santa Cruz, Bolivia

41 ²²Assis State Forest, Forestry Institute of São Paulo State, Assis, SP, Brazil

42 ²³Center for Integrative Conservation, Xishuangbanna Tropical Botanical Garden, Chinese
43 Academy of Sciences, Menglun 666303, Yunnan, People's Republic of China

44 ²⁴Department of Plant and Microbial Biology, North Carolina State University, Raleigh, North
45 Carolina, U.S.A.

46 ²⁵Natural Resources and the Environment, Council for Scientific and Industrial Research,
47 Pretoria, South Africa

48 ²⁶South African Environmental Observation Network, National Research Foundation,
49 Claremont, South Africa

50 ²⁷Department of Biological Sciences, University of Cape Town, Rondebosch, South Africa

51 *To whom correspondence should be addressed: griffith.dan@gmail.com

52

53 **Abstract (60/60 words):**

54 Bastin et al (Reports, 12 May 2017, p. 635) infer forest as more globally extensive than
55 previously estimated using tree cover data. However, their forest definition does not reflect
56 ecosystem function or biotic composition. These structural and climatic definitions inflate forest
57 estimates across the tropics and undermine conservation goals leading to inappropriate
58 management policies and practices in tropical grassy ecosystems.

59

60 **Main text (824/1000 words; 15/15 references; 2/2 Figures):**

61 Bastin *et al* (1) used high-resolution Google Earth images to estimate tree canopy cover in
62 213,795 (0.5 ha) globally distributed plots. Extrapolation of these plot-level data produced a
63 forest cover classification where they concluded “dry forests” cover ~ 40% more of the global
64 land area than previously estimated, increasing global forest cover estimates by 9%. However,
65 their calculation of forest extent is based on a structural definition adopted by the Food and
66 Agriculture Organization of the United Nations (FAO), where areas greater than 0.5 hectares and
67 with more than 10% tree cover are considered forest (1). As a consequence of applying the FAO
68 forest definition, Bastin *et al* (1) misclassify as dry forest many tropical regions that are in fact
69 savannas. Savannas differ from forests in having a continuous grassy ground layer which
70 supports fire and grazing mammals. These disturbances select for functionally distinct plant traits
71 and species that are different from forests in their biodiversity and ecosystem services (2, 3).

72 Bastin *et al* (1) refer to plots with 10-40% tree cover as open forest and over 40% as closed
73 forest. These “forest” classes clearly overlap with savannas which can range in tree cover from 0
74 – 80% (4). Tree cover has been previously demonstrated as an unreliable metric by which to
75 differentiate forest and savanna (3), and sites classified by Bastin *et al* (1) as forest include iconic
76 savannas such as Kruger National Park (Fig 1). Additionally, the FAO “forest” definition applied
77 by Bastin *et al* (1) includes sites where tree cover is “temporarily under 10% but is expected to
78 recover,” an unclear guideline implying degradation rather than accounting for known temporal
79 variability in savanna tree cover (5–7). Consequently, the majority of “new” forest identified
80 here resulted from the misclassification of tropical savannas as “forests” (Bastin *et al* Fig
81 S12)(8).

82
83 Implications of misclassification of savanna as forest include support for afforestation,
84 modification of mammalian grazer and browser regimes, and fire suppression policies (9), as fire
85 and large herbivores are generally considered to be at odds with the integrity of forest
86 ecosystems (2, 10, 11). In contrast, it is the loss of these processes in many savannas that results
87 in their degradation (8). Over millions of years, fires and herbivores have driven the evolution of
88 herbaceous plants with belowground buds, underground trees and trees with thick insulating
89 bark, traits which make savanna species functionally distinct from forest species (5, 9).

90 Afforestation and fire suppression policies in savannas risk destroying a wealth of specialized
91 and endemic savanna biodiversity that underpin unique ecological processes, and compromising
92 ecosystem functions such as carbon cycling and water and energy exchange (5, 6, 9, 11, 12).

93 Further, afforestation strategies negatively impact grassy ecosystem function by altering the
94 hydrology and/or trophic structure (2, 8) of entire landscapes. Many of the sites identified by
95 Bastin *et al* (1) as forest fall within areas identified as opportunities for “forest and landscape
96 restoration” (6), increasing the very real risk that misclassification could misdirect afforestation
97 policies (8).

98
99 Further underlying the misclassification of savanna is an assumption that biomes can be
100 delineated using a single simple metric of climate (*i.e.*, aridity index). Using a threshold aridity
101 index (0.65) belies the rich ecological complexity in identification and characterization of
102 biomes, the subject of debate for a century (reviewed in 11). Historical contingencies in the

103 distribution and evolution of plant lineages and their associated functional traits generate critical
104 biogeographic variation in the limits of biomes and their dynamics in response to climate (*e.g.*,
105 savannas across continents) (14). Because of this complexity, the climate threshold in Bastin *et*
106 *al* (1) also misclassifies some wet Neotropical forests (in Amazonian Ecuador and Peru, and on
107 the Pacific coast of Ecuador and Colombia) as dry forest (15). Recent evidence overwhelmingly
108 shows that definitions of forest based solely on tree cover or climate thresholds ignores key
109 functional difference between closed and open canopy vegetation types (2, 3, 6, 8).

110
111 Many of the ecosystems identified by Bastin *et al* (1) are not forest but savannas (3, 5) where
112 low tree cover is the result of natural processes (4, 5, 8, 9). Their aim was “to accurately
113 determine how much forest and tree cover remains in dryland biomes” (p.635). This aim implies
114 that dryland systems were once widely forested, which is incorrect. In Figure 2, we map
115 locations derived from (5) providing fossil evidence that many “forest” sites in Bastin *et al* (1)
116 have supported tree-grass mosaic vegetation over millennia. Conservation policies should reflect
117 savanna antiquity and not equate low tree cover with degradation. Moreover, although we have
118 focused on savannas, the inflation of forest extent could equally hamper conservation in other
119 threatened forests. An example is the dry forests of Latin America, which lack adequate
120 protected areas to safeguard their unique and geographically heterogeneous flora (15). While the
121 data collected by Bastin *et al* (1) are impressive and potentially useful, the use of the FAO forest
122 definition is damaging to conservation goals across the tropics.

123
124 **Acknowledgments:** CJS and DMG were supported by National Science Foundation award
125 1342703; CAES by NSF award 1253713. CLP is funded by a Royal Society-DFID Africa
126 Capacity Building Research grant. RTP acknowledges funding from the Scottish Government’s
127 Rural and Environmental Science and Analytical Services (RESAS) Division, NERC
128 NE/I028122/1 and NERC/CONFAP NE/N000587/1. DMG and CERL led the drafting of the
129 manuscript with contributions from all authors. CAES created the fossil map. NPH, JR, CLP,
130 and JWV provided images.

131

132 **References:**

133

134 1. J.-F. Bastin *et al.*, The extent of forest in dryland biomes. *Science*. **356**, 635–638 (2017).

135 2. C. E. R. Lehmann, C. L. Parr, Tropical grassy biomes: linking ecology, human use and
136 conservation. *Philos. Trans. R. Soc. B Biol. Sci.* **371**, 20160329 (2016).

137 3. J. Ratnam *et al.*, When is a “forest” a savanna, and why does it matter? *Glob. Ecol.*
138 *Biogeogr.* **20**, 653–660 (2011).

139 4. M. Sankaran *et al.*, Determinants of woody cover in African savannas. *Nature*. **438**, 846–849
140 (2005).

141 5. C. A. E. Strömberg, Evolution of Grasses and Grassland Ecosystems. *Annu. Rev. Earth*
142 *Planet. Sci.* **39**, 517–544 (2011).

143 6. J. W. Veldman *et al.*, Where Tree Planting and Forest Expansion are Bad for Biodiversity
144 and Ecosystem Services. *BioScience*. **65**, 1011–1018 (2015).

145 7. N. P. Hanan, A. T. Tredennick, L. Prihodko, G. Bucini, J. Dohn, Analysis of stable states in
146 global savannas: is the CART pulling the horse?: Analysis of stable states in global
147 savannas. *Glob. Ecol. Biogeogr.* **23**, 259–263 (2014).

148 8. C. L. Parr, C. E. R. Lehmann, W. J. Bond, W. A. Hoffmann, A. N. Andersen, Tropical
149 grassy biomes: misunderstood, neglected, and under threat. *Trends Ecol. Evol.* **29**, 205–213
150 (2014).

151 9. W. J. Bond, Ancient grasslands at risk. *Science*. **351**, 120–122 (2016).

152 10. S. Archibald, C. E. R. Lehmann, J. L. Gomez-Dans, R. A. Bradstock, Defining pyromes and
153 global syndromes of fire regimes. *Proc. Natl. Acad. Sci.* **110**, 6442–6447 (2013).

154 11. G. P. Asner, A. J. Elmore, L. P. Olander, R. E. Martin, A. T. Harris, Grazing systems,
155 ecosystem responses, and global change. *Annu. Rev. Environ. Resour.* **29**, 261–299 (2004).

156 12. T. D. Searchinger *et al.*, High carbon and biodiversity costs from converting Africa's wet
157 savannas to cropland. *Nat. Clim. Change*. **5**, 481–486 (2015).


158 13. S. I. Higgins, R. Buitenwerf, G. R. Moncrieff, Defining functional biomes and monitoring
159 their change globally. *Glob. Change Biol.* **22**, 3583–3593 (2016).

160 14. C. E. R. Lehmann *et al.*, Savanna Vegetation-Fire-Climate Relationships Differ Among
161 Continents. *Science*. **343**, 548–552 (2014).

162 15. DRYFLOR *et al.*, Plant diversity patterns in neotropical dry forests and their conservation
163 implications. *Science*. **353**, 1383–1387 (2016).


164

165 **Figures:**



166

167 **Figure 1.** Examples of savannas, with continuous grass layers and discontinuous tree canopies,
168 that are misclassified as forests by the FAO 10% tree cover threshold in Bastin *et al* (1). (A)
169 *Acacia*-grass mixture from Australia, functionally a savanna according to contemporary
170 ecological understanding. This is Fig S3 from Bastin *et al* (1). (B) *Combretum* savanna in Kruger
171 National Park, South Africa. Photo credit: CLP. (C) South-Sahel site in Lakamané, Mali. This
172 site has ~12.4% tree cover, is heavily grazed and experiences frequent fires. Photo credit: NPH.
173 (D) Savanna from Isalo National Park, Madagascar. Photo credit: CERL. (E) Savanna (cerrado)
174 in eastern lowland Bolivia. This site is within the “dry subhumid” zone in Bastin *et al* and
175 experiences frequent fires. Photo credit: JWV. (F) Long-term monitoring plot in an *Anogeissus*-
176 *Terminalia-Chloroxylon* savanna in Amrabad Tiger Reserve, southern India. Photo credit: JR.



177

178

179 **Figure 2.** Forest distribution in “drylands,” from Bastin *et al* (2017), where points (red dots)
180 highlight areas where fossil evidence (*e.g.*, fossil floras and faunas, stable carbon isotopes) has
181 demonstrated past occurrence (>0.5 million years ago, but mainly 4-22 million years ago) of
182 grass-dominated habitats and their faunas across continents (5). Although savanna extent has
183 shifted with changing climates and disturbance regimes and exact compositions have changed
184 during the last 22 million years, it is abundantly clear these regions have deep evolutionary roots
185 as mixed tree-grass systems (5). Note: ocean points represent paleovegetation data reconstructed
186 from marine cores.

187