



Ongoing and planned activities to improve the management of patients with Type 1 diabetes across Africa; implications for the future

Journal:	<i>Hospital Practice</i>
Manuscript ID	Draft
Manuscript Type:	Review Article
Sub-Manuscript Type:	Invited submission
Therapeutic Area:	Diabetes, Type 1 Diabetes
Keywords – Click here to find your MeSH keywords.:	Type 1 diabetes, healthcare policies, insulin, primary healthcare, glucose monitoring, patient co-payments, Africa

SCHOLARONE™
Manuscripts

Abstract

Background: Currently there are 19 million people in Africa with diabetes, mainly Type 2 diabetes (T2DM) (95%), estimated to grow to 47 million people by 2045. However, there are concerns with early diagnosis of patients with Type 1 diabetes (T1DM) with often patients presenting late with complications. Concerns also to access and affordability of insulin and affordability of monitoring equipment and test strips with typically high patient co-payments with potentially catastrophic consequences for families. This negatively impacts on the quality of care adversely affecting morbidity and mortality. There are also issues of patient education and psychosocial support affecting patients' quality of life. These challenges need to be debated and potential future activities discussed to improve the future care of patients with T1DM across Africa. Methodology: Documentation of the current situation across Africa including the epidemiology and economics of diabetes as well as facilities and available treatments within public healthcare system for patients with T1DM. In addition, ongoing activities to improve future care. Subsequently, contextualise the findings with input from senior level government, academic and other professionals from across Africa to provide future guidance. Results: Whilst the prevalence rates for T1DM are considerably lower than T2DM, there are ongoing concerns with late diagnosis of T1DM as well as the routine provision of insulin and monitoring equipment across Africa. High patient co-payments exacerbate the situation. There are ongoing developments to address this including the instigation of universal health care and partnerships with non-governmental organisations, patient organisations and pharmaceutical companies to improve future care. Their impact remains to be seen. A range of activities have been documented for all key stakeholder groups to improve future care. Conclusion: There are concerns with the management of patients with T1DM across Africa. A number of activities have been suggested to address this and will be monitored.

1. Introduction

The focus on the management of patients with diabetes in Africa has increased considerably in recent years driven by growing prevalence rates as well as increasing costs to healthcare systems (1-3). Alongside this, there is increasing morbidity and mortality from non-communicable diseases (NCDs) including diabetes as well as cardiovascular diseases and hypertension (4-7), as well as and increasing comorbidity with HIV and TB especially among children (8, 9), focusing key stakeholder attention on the care of patients with diabetes. In South Africa for instance as a result of rising

1
2
3 prevalence rates, diabetes is currently the second most common cause among recorded deaths (10,
4 11). Diabetes, including Type 1 diabetes (T1DM), is also known to adversely affect patients' quality of
5
6 life due to the diabetes related complications (12-18).
7
8

9 Current estimates suggest that approximately 19 million people in Africa have diabetes with an
10 estimated 47 million likely to have diabetes by 2045 (19, 20), with almost a third of patients needing
11 insulin for their diabetes (21). High prevalence rates are seen particularly among North African
12 countries (22, 23). Other authors publishing in 2016 suggested that up to 29.3million people in Africa
13 had diabetes (17). In 2015, approximately 321,000 deaths in sub-Saharan Africa (SSA) were
14 attributable to diabetes, with most of these occurring in people 60 or under (1, 24). This increased to
15 366,200 across Africa in 2019 (20). Concerns with insulin costs, insulin resistance, and the
16 inaccessibility of insulins are key contributing factors to current mortality rates (21).
17
18
19
20
21
22
23
24

25 High prevalence rates for diabetes have resulted in estimates that approximately US\$10.81 billion
26 was spent on direct medical costs in 2015 for treating patients with diabetes in SSA, with out-of-
27 pocket expenditure likely to exceed 50% of overall health expenditure in a number of African countries
28 (1). Total costs increased to US\$13billion across Africa in 2019 for patients aged between 20 to 79
29 years (20). These costs are exacerbated by the fact that patients with diabetes in hospitals have the
30 longest stay due to diverse complications including stroke and amputations (12, 25). The total costs
31 for treating patients with diabetes in SSA including direct medical costs are likely to increase to
32 US\$59.32billion by 2030 (1), most of which will be for treating patients with type 2 diabetes (T2DM).
33 Costs will be enhanced by the lack of health technologies to screen, diagnose and effectively manage
34 patients with diabetes. Having said this, less than 10% of the total worldwide costs for treating
35 patients with diabetes are currently from the African, North African, South East Asian and Middle East
36 Regions combined (26). The cost of treating patients with T1DM in Africa is typically higher compared
37 with T2DM patients at a ratio between 1.80 to 5.66 (27), with others calculating ratios of between 1.5
38 to 4.4 although taking into account more continents (27, 28)
39
40
41
42
43
44
45
46
47
48
49
50
51
52

53 There have also been cost estimates for treating diabetes among individual Africa countries. In 2015,
54 Quaye et al estimated the mean annual financial cost of managing one diabetic case at hospital
55 clinics in Ghana was US \$372.65, with medicines accounting for 71% of the total financial cost (29). In
56 2015, Pei estimated the total cost of diabetes for 40 households in Ghana, including patients with both
57
58
59
60

1
2
3 T1DM and T2DM, was approximately 15,000 Cedi/month of which 66.5% was direct costs (30). In
4 view of the expenditures involved, 52.9% of the households surveyed in Ghana have experienced
5 catastrophic health expenditure when family members are diagnosed with diabetes (30). In Mali,
6 patients with diabetes typically spend an average of US\$230 per year on their treatment without
7 adding in the cost for managing any complications, with the cost borne by the State in Mali for the
8 treatment of diabetes without complications currently estimated at US\$20million per year (21, 31) (see
9 Table 2 for further details). Suleiman *et al.* (2014) estimated the annual direct costs of treating
10 patients with T1DM in Nigeria was NGN18 billion (US\$112.54million), with costs of NGN100,032.89
11 (US\$625.21) per patient (32). In their review, Mutyambizi *et al.* (2018) estimated the national direct
12 costs of treating patients with diabetes in Nigeria (T1DM and T2DM) was between I\$3.5 to I\$4.5
13 billion per annum, which compares with estimated national costs (direct and indirect) in Morocco of
14 between I\$5.9 to I\$8.2 billion per annum (27).

15
16
17
18
19
20
21
22
23
24
25
26
27 There are concerns that governments and health systems across Africa are struggling to cope with
28 the demand and cost of services to treat patients with diabetes and its complications unnecessarily
29 increasing morbidity and mortality (1, 33). This includes concerns that children in Africa are at high
30 risk of being diagnosed with T1DM only after complications occur, or when they are critically ill, due to
31 a lack of knowledge among parents, caregivers and school teachers (34). This is beginning to change
32 with several initiatives now in place across Africa to improve the management of patients with
33 diabetes. These include patient led strategies to strengthen healthcare systems, strategies to improve
34 guideline adherence where there are concerns with sub-optimal management, assist patients with
35 appropriate lifestyle changes including diet and exercise, provide free medicines and diagnostic
36 equipment as well as instigate quality targets (35-54). There are also projects involving pharmacists
37 educating patients about the signs and symptoms and treatments to improve self-management for
38 those with chronic NCDs (55). Further details regarding ongoing strategies across Africa to improve
39 the care of patients with diabetes are summarised in the recent paper of Godman *et al* (56).

40
41
42
43
44
45
46
47
48
49
50
51
52
53 The vast majority of patients with diabetes in Africa have T2DM, with higher prevalence rates of
54 patients with T2DM versus T1DM seen in Africa compared with other parts of the world (1, 56-60). In
55 their study published in 2008, Majaliwa *et al.* reported that incidence rates for patients with T1DM in
56 Africa ranged from 1.5/100000 in Tanzania to 20/100000 in Morocco, with 9.8% of patients in Ethiopia
57
58
59
60

1
2
3 attending diabetes clinics having T1DM (61). Higher rates are seen in children in SSA where T1DM is
4 one of the most common endocrine and metabolic conditions (62, 63). Reported prevalence rates for
5 patients with T1DM across Africa were low in 2011, ranging from 4 per 100,000 in Mozambique up to
6
7 12 per 100,000 in Zambia (58). More recent incidence and prevalence rates are seen in Table 1.
8
9

10 Typically, published figures demonstrate an increasing prevalence of T1DM across Africa (45, 57, 58,
11 64-67), although this is not universal (57). Current incidence and prevalence figures may be
12
13 underestimates if parents fail to understand the features of childhood diabetes and do not take their
14 children for diagnosis and treatment at healthcare facilities (30, 59, 63, 68), with estimates that up to
15
16 half or more of patients with T1DM are not diagnosed (34, 69).
17
18
19

20 Prevalence rates for patients with T1DM in Africa are likely to be higher if the age of diagnosis is
21 extended up to 26 years as opposed to 0 to 14 years (1), enhanced by the age at diagnosis of T1DM
22 typically higher among the black population in Africa compared with the white population (70). In
23
24 Rwanda, recent studies suggest prevalence rates are over three times higher when young adults up
25
26 to 26 years are included in epidemiological studies versus those only including children up to 14 years
27
28 of age (71, 72). Similarly in Ethiopia, there was a rapid increase in the incidence of T1DM in children
29
30 in their mid-teens up to the age of thirty compared with those under 15 years of age (73).
31
32
33

34 Insert Table 1

35 In 2015, the International Diabetes Federation (IDF) believed there were 7600 new cases each year
36
37 of diabetes among children in Africa (17), with generally higher rates among North African countries
38
39 (17). Typically in Africa, children with diabetes are insulin-dependent and most likely autoimmune
40
41 (74), with auto-immune beta-cell destruction having an important role in the pathogenesis of T1DM
42
43 among African patients (75). There are a number of HLA class II alleles associated with T1DM among
44
45 the black population in Southern Africa, which is important for care delivery (76). It is estimated that
46
47 T1DM accounts for over 90% of children with diabetes (17, 63). The latest estimates (2019) from the
48
49 IDF suggest there are 25,800 children and adolescents (0 – 19 years) with T1DM in Africa, with
50
51 10,300 newly diagnosed cases each year (77). However, still a significant number of patients remain
52
53 undiagnosed (34, 69, 77). There are also ongoing debates regarding the role of malnutrition in the
54
55 development of T1DM among African children, as well as malnutrition-related diabetes, enhancing the
56
57 poor prognosis for T1DM patients in Africa especially those from resource-poor settings (59, 73).
58
59

60 Despite its impact on the health of patients including early mortality, there are still concerns that

1
2
3
4 patients with T1DM are not receiving evidence based care in Africa at the expense of ongoing
5 strategies to reduce morbidity and mortality associated with T2DM (45, 56). Studies published in 2005
6 documented high mortality rates for children with T1DM (78, 79). For instance, in Mozambique the life
7 expectancy of children under 15 years with T1DM was 3.5 years (1). Common causes of death in
8 African children with diabetes include diabetic ketoacidosis (DKA), severe hypoglycaemia, infection
9 and eventually microvascular complications (17, 69, 80). Overall, DKA is the most common cause of
10 death in children and adolescents with T1DM, accounting for approximately half of all deaths in
11 patients younger than 24 years with diabetes (59, 81, 82). Typically at diagnosis of T1DM in Africa, up
12 to 80% or more of children and adolescents newly diagnosed with T1DM have DKA (17, 34, 69, 80,
13 83-85). Independent risk factors associated with DKA in children include their age, parents'
14 knowledge of the signs and symptoms of DKA, health care professional (HCP) education, erratic
15 supply and poor adherence rates to prescribed medicines as well as infections before DKA (80, 86-
16 88). High mortality rates due to DKA in developing countries including African countries are due to
17 higher rates of infection, less developed medical service as well as delays in seeking care with over
18 reliance on traditional medicines and healers (69, 80, 81, 89, 90). Severe hypoglycaemia is also
19 common among children in SSA, and is associated with high mortality (17, 65, 91) as well as a
20 substantial impact on both productivity and healthcare utilisation (92). There are also concerns that
21 lack of correction dosing for hyperglycaemia, lack of carbohydrate counting and lack of access to
22 rapid-acting insulin among children with T1DM results in poor glycaemic control, which negatively
23 impacts on the quality of life of patients (93). Having said this, good management of nocturnal and
24 daytime non-severe hypoglycemic events that minimize hypoglycemia while maintaining good
25 glycemic control may positively impact upon the psychological wellbeing of people with diabetes,
26 reducing healthcare costs and improving productivity (94). Women with T1DM in their reproductive
27 age should also be identified as a high-risk group where access to specialized prepregnancy clinics
28 should be made available so that their T1DM can be intensively managed throughout pregnancy via a
29 multidisciplinary team. The use of multiple insulin injection regimens, including the use of insulin
30 aspart during pregnancy, is both well tolerated and effective, and may offer some benefits with
31 respect to postprandial glycaemic control. This is also a key consideration for the future as care for
32 patients with T1DM improves (95).

33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
There are also concerns with the extent of complications among patients with T1DM across Africa

1
2
3 including retinopathy, nephropathy, neuropathy and hypertension as well as infections including
4 malaria and tuberculosis (1, 17, 59, 69, 96-98). Improved diet, family support offering psychosocial
5 support, as well as family support by assisting with correct and constant administration of medicines,
6 including insulin, can help to reduce the complications of T1DM (65). Psychosocial support is
7 important with T1DM adversely affecting the quality of life of patients resulting in the need for all key
8 stakeholder groups to work together to improve the care and lives of patients with T1DM (13, 14, 99,
9 100). This includes the need to develop coping strategies and address issues of stigma (99), which
10 are especially crucial if coping strategies involve avoiding injections (99). There is also a need to
11 address concerns with the availability of dialysis services across Africa given current concerns (101,
12 102).

13
14 While mortality and morbidity rates have improved among African countries in recent years in patients
15 with T1DM, the gains seen in Africa are far less than those seen in developed countries where there
16 have been substantially greater declines in morbidity and mortality in recent years (45). For instance
17 in Gabon, Damiens *et al.* (2019) estimated mortality at 7% approximately four years after T1DM
18 diagnosis (57). Others have documented higher rates suggesting that up to 50% of patients with
19 T1DM in Africa die in the first five years after diagnosis (57). In addition, among some African
20 countries, the life expectancy of a child with T1DM can be as low as one year (103). There are a
21 number of elements of care that adversely affect improvements in the morbidity and mortality of
22 African patients with T1DM (45). These include failure to seek help, unavailability of regular screening
23 programmes, inability to make a formal diagnosis, inadequate and inappropriate care exacerbated by
24 the lack of equipment, facilities and professionals, e.g. Ghana currently has less than fifteen
25 endocrinologists, poor knowledge among patients and parents, frequent episodes of hypoglycaemia
26 exacerbated by a lack of insulin and the cost of insulin as well as monitoring sticks and syringes, poor
27 storage of insulin, poor adherence to treatments and poor eating habits as well as cultural aspects of
28 care (1, 17, 21, 30, 45, 47, 48, 63, 65, 104-113). The WHO in 2015 found that in SSA only 51% of the
29 countries had metformin routinely available and only 40% had insulin readily available, well below the
30 80% target (1, 114). In addition, many African countries currently have inadequate facilities to manage
31 both the microvascular and macrovascular complications of diabetes (1, 115).

32
33 Overall in many African countries, patients with T1DM have been at a disadvantage with a greater
34 focus on infectious diseases; however, as mentioned, this is starting to change (56, 80). We are also
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 seeing programmes introduced among African countries to help transition patients from paediatric to
4 adult services, and this is likely to grow as care is improved for children with T1DM in Africa (116).
5
6

7
8 A key area to address to improve the management of patients with T1DM across Africa along with
9 healthcare strengthening is access to, and regular availability, of insulins as well as increased
10 availability of blood glucose monitoring devices (1, 50, 59, 69, 106). Previous studies have found that
11 <25% of T1DM patients in the Democratic Republic of Congo had regular access to insulin <25% of
12 the time and in Tanzania different types of insulin were available in public health facilities in only 8%–
13 17% of the time with one version of insulin was only available 34% of the time (59, 117). Poor
14 availability and use of insulin are not helped by high costs and co-payments. Other key areas to
15 address include patient empowerment on the knowledge about their disease and treatment to help
16 reduce non-adherence to insulins and rates of hypoglycaemia (55, 118, 119).
17
18
19
20
21
22
23
24
25

26 Table 2 provides more details regarding the economics and availability of insulin and glucose
27 monitoring among African countries highlighting current challenges. Typically in Africa, the direct
28 costs of treating diabetes approaches US\$300/ patient corresponding to the annual income of many
29 families (65). We are beginning to see access programmes and other approaches among African
30 countries to address concerns with insulin availability as well as a lack of education and facilities, and
31 these initiatives are likely to grow alongside moves to enhance universal healthcare across Africa (46,
32 69, 120-123). This also includes addressing concerns with the availability of insulin analogues for
33 insulin resistant cases exacerbated by cost considerations (107, 124, 125). We are also seeing
34 groups such as the IDF improve access to HbA1c testing among African countries including
35 Cameroon and Guinea (49, 50). For instance in the Cameroon, under the Changing Diabetes in
36 Children (CdiC) programme, all children and adolescents attending relevant centres are provided with
37 free insulin as well as glucose monitors and strips (48). Diabetes control has been reported to
38 improve with the length of T1DM once patients had been through any denial phases and had
39 accepted the condition and its implications on lifestyle requirements (48). Recently, the partnership
40 between the Ministry of Health in Rwanda and Abbott through the Team Type 1 Foundation has
41 resulted in the donation of glucose meters and nearly 12 million test strips to improve the diagnosis
42 and management of patients with T1DM in Rwanda for the next four years (53), and this will be
43 closely monitored to provide future guidance,
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 The WHO lists soluble insulins and intermediate insulins in the 2019 Essential Medicines List (126),
4 and these should be readily available to help patients with T1DM. However, there are concerns with
5 the quality of low-cost generic insulins as these are more complex to produce than oral generic tablets
6
7 (17, 106).
8

9
10 Insert Table 2

11
12 Consequently, the principal objective of this paper is to debate ongoing challenges and potential
13 future activities that could be instigated to improve the care of patients with T1DM in Africa. This
14 includes issues surrounding medicine availability and storage along with monitoring equipment and
15 other measures to improve the care of patients with T1DM. There have been a number of systematic
16 reviews and guidelines looking at key issues surrounding the management of patients with diabetes in
17
18 Africa including the epidemiology, economics and initiatives to strengthen healthcare systems (1, 2,
19 61, 62, 98, 109, 127-131). However to date, there have only been a limited number of publications
20 that have comprehensively focused on all aspects of care covering epidemiology, diagnosis,
21 education, and treatment including access to appropriate medicines and monitoring equipment as well
22 as adherence rates. There have also only been a limited number of publications assessing the
23 influence and impact of different interventions to improve the care of patients with T1DM across
24 Africa. In view of this, we sought to document and discuss ongoing and potential activities that could
25 be introduced for all key stakeholder groups across Africa to improve the future care of patients with
26 T1DM. This includes contextualising the findings from a wide range of co-authors from across Africa
27 and wider to provide future guidance.
28
29

30 **2. Methodology**

31 We did not perform a systematic review for the reasons already given. However, there is an urgent
32 need to improve the management of patients with T1DM in the public healthcare system across Africa
33 building on recent publications. Consequently, we conducted a narrative review of the literature. In
34 addition, in view of the paucity of publications researching the influence and impact of different
35 interventions to improve the care of patients with T1DM, we used senior level personnel from
36 governments and their advisers, clinicians, academia, rational medicine use advisers, HTA personnel,
37 as well as patient organisation personnel, from across Africa and wider to suggest potential activities
38 that could be introduced to improve the future care of patients with T1DM in Africa. This included both
39 short- and long-term initiatives building on an understanding of the current situation in their country.
40
41 The advice given to improve the care of patients with T1DM was based on the considerable
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 knowledge and experience of the co-authors as well as publications known to them. The co-authors
5
6 have come from a wide range of backgrounds and countries. We have successfully used this
7
8 approach in previous publications to stimulate debate in other priority healthcare areas and situations
9
10 to provide future guidance (56, 132-144).

11
12 We did not split the African countries into low- or middle-income countries as the issues surrounding
13
14 the management of patients with T1DM appears to apply across Africa. Consequently, countries can
15
16 learn from each other. We also did not review each paper to assess whether the presence of
17
18 diabetes, especially T1DM, had been defined according to WHO and other internationally recognised
19
20 diagnostic criteria in view of the objectives of this paper (130, 145).

21 22 **3. Findings and suggested activities**

23
24 T1DM is typically diagnosed within primary healthcare centres (PHCs) in the public healthcare system
25
26 among a number of African countries; alternatively, within district and tertiary hospitals. Generally,
27
28 most patients with DKA are diagnosed at secondary and tertiary hospitals.

29
30
31 Table 3 contains details of treatments and facilities routinely available within the public healthcare
32
33 systems among African countries to manage T1DM as well as its complications. As mentioned, the
34
35 WHO Essential Medicines List (WHO/ EML) lists soluble insulins and intermediate insulins (126).
36
37 However in their latest guidance on the management of patients with diabetes including T1DM, the
38
39 WHO suggests long-acting insulin analogues can be used to control blood glucose levels in patients
40
41 with T1DM who have frequent severe hypoglycaemia with human insulin (146). This will be an issue
42
43 in many African countries given considerable differences in costs and existing concerns with the
44
45 routine availability of NPH insulins (107, 147).

46
47 Insert Table 3

48
49 Boxes 1 to 6 contain suggested activities that can be undertaken to improve the care and outcomes of
50
51 patients with T1DM in Africa, Activities for healthcare professional groups and patients/ patient
52
53 organisations are broken down into short and longer term activities.

54
55 Alongside these activities, African governments need to see how they can procure insulins and
56
57 monitoring technologies at more affordable prices for patients where there are concerns with high co-
58
59 payment levels. The private sector can play its role by lowering prices and/ or enhancing access
60

1
2
3 programmes in Africa building on existing initiatives (46, 103). In addition in their procurement
4 negotiations, African countries need to use their own and surrounding countries' existing price data in
5 theirs and surrounding countries to negotiate improved prices with the potential for developing
6 mechanisms to foster competition and secure supplies of insulin with only a limited number of
7 companies supplying them with insulin (109). In time as more biosimilars become available,
8 increased competition should lead to lower prices and greater availability across Africa (building on
9 Table 3) (107). Improved forecasting and distribution systems should help reduce stock-outs building
10 on recent initiatives in South Africa (4).

11
12
13
14
15
16
17
18 Insert Boxes 1 to 6

19 20 21 **4. Discussion and next steps**

22 There is a growing prevalence of patients with T1DM across Africa. Increased morbidity and mortality
23 can be expected in view of the number of patients not diagnosed or misdiagnosed, concerns with the
24 routine availability of insulins even those within country's EML, and high co-payments for insulins in a
25 number of African countries. High co-payments for glucose monitoring equipment and testing strips
26 across many African countries add to the patient's and families' financial burden. The consequences
27 are potentially catastrophic for families. There are also concerns regarding the lack of HCP support
28 including education and psychosocial support. Specialists are also needed to help improve the
29 management of patients with T1DM across Africa. This includes help with patients coming to terms
30 with their disease as well as help with adherence to the medicines prescribed to reduce the
31 complications of T1DM including DKA, hypoglycaemia and microvascular complications.

32 Encouragingly, we are seeing patient organisations and NGOs including the IDF playing an increasing
33 role among African countries to improve the care of patients with T1DM. This is likely to grow. A
34 greater focus is also ongoing among countries to improve the management of patients with NCDs.

35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Boxes 1 to 6 provide suggestions for governments as well as all key stakeholder groups to improve
the care of patients with T1DM both in the short and longer term. We will be monitoring this progress
in the future as countries seek to improve the care of their patients with T1DM. Diagnosis and
management rates for patients with diabetes including T1DM need to be appreciably improved across
Africa to achieve the World Health Organisation's (WHO) 25×25 Global Action Plan (2013) as well as
Sustainable Development Goal (SDG) 3 goal to reduce premature deaths due to NCDs across
countries by a third by 2030 (148-152). With respect to increasing the routine availability of insulins

1
2
3
4 alongside reducing co-payments where concerns, it is better for countries to focus initially on
5
6 increasing the availability of NPH insulins as opposed to more expensive insulin analogues apart from
7
8 insulin resistant patients. This is because a number of published studies from academic and HTA
9
10 units have not demonstrated significant differences between NPH and the more expensive long acting
11
12 insulins in terms of reducing hypoglycaemia episodes or on patient's quality of life (153-160).
13
14 However, this finding was not universal (161). More recently, long acting insulins have been
15
16 recommended as second line in patients who are allergic to conventional insulins or have failed to
17
18 have their T1DM controlled with conventional insulins (146, 162, 163). First though the availability of
19
20 WHO EML recommended insulins (126) needs to become routine at affordable prices to patients;
21
22 preferably free-of-charge. In addition, it is also important to increase the availability and affordability of
23
24 glucose monitoring equipment and strips to further improve future care.

25
26 We are aware of a number of limitations with our study. This includes the fact that we did not
27
28 undertake a systematic review for the reasons already mentioned. We are also aware that there have
29
30 been a paucity of studies assessing the influence and impact of interventions to improve the care of
31
32 patients with T1DM among African countries. However, we have incorporated the views of a wide
33
34 variety of senior level personnel from across Africa with experience of managing, organising,
35
36 researching or educating others about the care of patients with T1DM to help address such concerns.
37
38 Consequently, we believe our findings and suggestions are robust.

39 **5. Conclusion**

40
41 The improved management of patients with T1DM across Africa is starting to be addressed as
42
43 countries develop and refine their action plans to improve the care of patients with NCDs as part of
44
45 SDG goals, with further suggestions contained in Boxes 1 to 6. However, there are still considerable
46
47 concerns and all key stakeholder groups need to work together to appreciably improve the future care
48
49 and self care of patients with T1DM across Africa.

50
51 We will be monitoring developments regarding the future management of T1DM among African
52
53 countries, including the impact of interventions to improve their care. In addition, it is important to
54
55 rapidly publish any findings so that African countries can learn from each other.
56
57
58
59
60

Tables

Table 1 - Epidemiology of Type 1 diabetes across Africa

Country	Epidemiology
General (45, 62)	<ul style="list-style-type: none"> Incidence rates of 4.3 per 1000000 among African children aged between 0 to 14 years and 10.3 per 1000000 in those aged between 0 and 19 years Prevalence rates of 10.3 per 1000000 among African children aged between 0 to 14 years and 25.8 per 1000000 in those aged between 0 and 19 years These incidence and prevalence rates are appreciably lower than those seen in other regions of the world with incidence rates as high as 25.1 per 1000000 among children in Europe aged between 0 to 14 years and up to 31.1 per 1000000 in those aged between 0 and 19 years There is an increase in pregnant adolescents girls presenting with T1DM in SSA
Botswana	<ul style="list-style-type: none"> The IDF currently estimates that the national prevalence rates for diabetes among adults is 5.5% (77) The number of children and adolescents with T1DM in Botswana was recently (2019) estimated at 99 (164)
Cameroon	<ul style="list-style-type: none"> The number of children and adolescents with T1DM in Cameroon was recently (2019) estimated at 743 (164)
Ethiopia	<ul style="list-style-type: none"> The incidence rate of diabetes among children in Ethiopia in 2013 aged between 0 and 14 years was 0.3 per 100,000 in 2013 (26, 165) In their study, Balcha <i>et al.</i> (2018) found an appreciably higher rate of T1DM in urban areas (incidence rate of 11.9/ 100,000 for men and 6.1/ 100,000 for women) vs. rural areas (2.2/ 100,000 for men and 1.0/ 100,000 for women) (73) Currently in Ethiopia, it is estimated (2019) that 2127 children and adolescents have T1DM (164)
Gabon	<ul style="list-style-type: none"> Reported incidence rates of 3.4/ 100,000 inhabitants/ year in 2015 and 2.9 in 2017 (57) It is estimated that 89 children (2019) in Gabon currently have T1DM (164)
Ghana	<ul style="list-style-type: none"> The overall prevalence of diabetes in adults in Ghana was 3.3% in 2014; principally T2DM (30), with higher rates seen in urban towns (30, 68) This has risen to 3.6% in 2017 (518,400 cases of diabetes in Ghana principally T2DM) (166) Currently in Ghana, it is estimated (2019) 1209 children and adolescents have T1DM (164)
Kenya	<ul style="list-style-type: none"> Currently in Kenya, it is estimated (2019) 1694 children and adolescents have T1DM (164)
Morocco	<ul style="list-style-type: none"> The overall prevalence of diabetes in adults in Morocco is 7.3%, much of which will be patients with T2DM as opposed to T1DM (167-169)
Namibia	<ul style="list-style-type: none"> Adekanmbi <i>et al.</i> (2019) recently estimated the prevalence of diabetes in Namibia at 5.1% including patients with T1DM and T2DM (170) Currently it is estimated (2019) in Namibia that 127 children and adolescents have T1DM (164)
Nigeria	<ul style="list-style-type: none"> Hospital based studies published in 2010 and 2013 showed prevalence rates of T1DM up to 3.1/1000 among children aged 15 years or below (90, 171)

	<ul style="list-style-type: none"> • However, there is appreciable variation depending on the locality ranging from 0.1/1000 in the South-East of Nigeria , 0.33/1000 in Sokoto (extreme North-West of Nigeria), 2.3/1000 in Kano (North-West), and 10.1/1000 in Jos, North-Central (84, 171-173). • The IDF currently estimates (2019) that there are 2954 children and adolescents in Nigeria with T1DM (164)
Rwanda	<ul style="list-style-type: none"> • The overall prevalence of diabetes in Rwanda is approximately 3% of the population, mainly T2DM (174) • Currently (2019) there appear to be over 630 children and adolescents in Rwanda with T1DM (164)
Somalia	<ul style="list-style-type: none"> • The IDF currently estimates (2019) that there are 57 children and adolescents in Somalia with T1DM (164)
South Africa	<ul style="list-style-type: none"> • In South Africa in 2015, approximately 2.3 million people had diabetes principally T2DM (67), with estimates of 1599 children and adolescents with T1DM in 2019 (164) • However there is a high rate of misdiagnosis in both sectors in South Africa as well as high rates of DKA (34) • Diabetes Type 1A class found to be more prevalent among young black South Africans than their white/ Indian peers (175) <p>There is increase</p>
Sudan	<ul style="list-style-type: none"> • Incidence estimates were 10.3/100,000 in 1995 (176) • Recently, the IDF currently estimated (2019) there are 208 children and adolescents in South Sudan with T1DM (164)
Tanzania	<ul style="list-style-type: none"> • Incidence rate among children in Tanzania aged between 0 and 14 years in 2013 was 0.9 per 100,000 in 2013 (26, 165) • More recently, Jasem <i>et al.</i> (2019) calculated an annual incidence of between 1.8 to 1.9 per 100,000 children aged under 15 years, with a peak incidence at 10 to 14 years (165) • The IDF currently estimates (2019) there are 1984 children and adolescents in the Republic of Tanzania with T1DM (164)
Zambia	<ul style="list-style-type: none"> • According to the WHO, the overall prevalence of diabetes in Zambia is 4.2% with no breakdown between T1DM and T2DM (177) • The number of children and adolescents with T1DM in Zambia is currently estimated at 557 (164)
Zimbabwe	<ul style="list-style-type: none"> • Among a total of 29,514 study participants, the overall pooled prevalence of diabetes in Zimbabwe including patients with T1DM before 1980 was 0.44% (95% CI 0.0-1.9%), and after 1980 the pooled prevalence was 5.7% (95% CI 3.3-8.6%) (178) • The IDF currently estimates (2019) there are 464 children and adolescents in Zimbabwe with T1DM (164)

NB: DKA= diabetic ketoacidosis; IDF: International Diabetes Federation; T1DM: Type 1 Diabetes mellitus; T2DM: Type 2 Diabetes mellitus; WHO: World Health Organization

Table 2 – Economics of diabetes across Africa especially T1DM, insulin availability and glucose strips

Country	Summary details
Sub-Saharan Africa	<ul style="list-style-type: none"> Ogle <i>et al.</i> (2016) ascertained that the median cost of blood glucose strips among seven African nations was US\$0.50, and the minimal reasonable yearly costs to families with a child or adolescent with diabetes ranged from 74% to 377% of per-capita gross income (1, 129) The regular availability of monitors and test strips is rare for most patients in African countries with the cost of monitoring beyond the financial means of most families, and many clinics currently cannot routinely measure glucose levels (17, 129, 179) Only a limited number of governments among those documented by Ogle <i>et al.</i> (2016) fully provided insulin (129) The mean availability of insulins in the public sector ranged from 55% to 80% among 13 LMICs including four African countries (107). Median government procurement prices for human insulins was US\$5 among the 13 countries (107) Wide variations have been seen in the prices of a 10-ml, 100-IU/ml vial equivalent of insulin ranging from US\$1.0 to US\$ 11.5 (median US\$ 4.7) among countries in the WHO Regional Office for Africa (109)
East Africa	Many patients on cost and other grounds still reuse syringes, which is discouraged as this can increase lipohypertropy (128)
Burkina Faso	Estimated published minimal annual direct costs of treating T1DM patients and supplies is US\$1185 (129), equating to 177% per capita income
Egypt	<ul style="list-style-type: none"> In their study, Zhang <i>et al.</i> (2010) documented Egypt's healthcare expenditure for patients with diabetes including both T1DM and T2DM reached US\$ 116 per person - 16% of total healthcare expenditure (23, 180) Expenditures are currently likely to be considerably more than this, with greater expenditure per person among patients with T1DM versus T2DM
Gabon	<ul style="list-style-type: none"> Damiens <i>et al.</i> in their study (2019) found that in 76% of patients with T1DM the cost of the medicines was an obstacle to the adherence to the medicines prescribed (57), with 69% saying they do not have a supply of blood sugar test strips to adapt insulin doses In view of the financial effects of the disease, if patients are alive after 4 years post diagnosis they typically have poor metabolic balances and recurrent DKA resulting in premature death (57)
Ghana	<ul style="list-style-type: none"> The total cost of diabetes for 40 households in Ghana including patients with both T1DM and T2DM was approximately 15,000 Cedi/month in 2015 of which 66.5% was direct costs (30) Overall, 52.9% of the households surveyed in Ghana experienced catastrophic health expenditure (30)
Kenya	<ul style="list-style-type: none"> Ngwiri <i>et al.</i> in their study (2015) found that 69% of children there were concerns with the unavailability of mixed formulation insulins (65) In their study, Pastakia <i>et al.</i> (2018) found that 85% of patients with diabetes including both T1DM and T2DM being treated at Webuye District Hospital (rural diabetes clinic) received regular HbA1c testing (181). This is welcomed with the average cost of HbA1c measurement at US10 – 25 with

	<p>50% of residents typically living on less than 1US\$/ day (6, 17, 182)</p> <ul style="list-style-type: none"> • The management of patients with diabetes in Kenya including T1DM patients is a key Government objective to reduce morbidity and mortality rates (54) • In their recent study, Shannon <i>et al.</i> (2019) found that despite a ceiling price ceiling of US\$5 per vial of insulin, many patients could not afford the additive expense of care including monitoring, other medicines and travel to the clinics. Encouragingly though, there was a greater number of stock-outs among facilities not taking part in the access scheme (46)
Mozambique	<ul style="list-style-type: none"> • The government currently buys insulin at approximately US\$4.30/ vial. Since the average person requires 13 vials a year, the government must prioritise whether it provides insulin for one person or essential medicines for nearly 40 others (47) • In addition in Mozambique, previously only a limited number of facilities have the necessary equipment to undertake blood glucose analysis (47); however, this is now being addressed with the help of the IDF
Namibia	<ul style="list-style-type: none"> • There is universal access to cost-free diabetic care in Namibia's public sector, which can be accessed from primary health care to tertiary institutions (5) • A wide range of insulins including soluble and long acting insulins are available to the public via PHCs financed by the government of the Republic of Namibia • A wide range of insulin preparations can also be accessed in the private sector as well as through a cost sharing system, Average co-payments range from US\$3.4 to US\$5.1 per vial. Without medical aid insurance, the cost of insulin preparations would range from US\$11.3 to US\$17.1 per vial • The challenges of managing patients with T1DM include: long distances between homes and health care facilities for people in the rural areas where the cost of transport and lost wages needs to be factored in as well as a lack of appropriate storage systems in rural settings
Nigeria	<ul style="list-style-type: none"> • Medication costs accounted for 72.3% of the total direct cost of patients with diabetes followed by the cost of laboratory investigations (17.6%), with costs typically out-of-pocket (183). Other studies have suggested that the costs of medicines can account for up to 90% of total costs (27) • Ogbera <i>et al.</i> (2012) estimated the monthly mean costs of insulin was 5212.8 Nigerian Naira (US\$33.1) equating to 29% of the minimum wage on insulin alone for patients with T1DM (184) • In their study, Onyiruka <i>et al.</i> (2012) estimated the mean monthly direct cost of routine diabetes care was 10, 950 Nigerian Naira (US\$73) with the insulin responsible for 51.1% of this cost (185)
Rwanda	<ul style="list-style-type: none"> • Rwanda's community-based health insurance system covered approximately 80% of the population in 2015–2016 with growth ongoing as Rwanda strives towards universal health coverage (121, 186). The healthcare system also covers country's poorest citizens aiming to enhance access to healthcare for all including medicines • The Rwanda Diabetic Association is active in many areas across the country to improve the care of patients with diabetes including T1DM. Activities include running diabetes

	<p>awareness programmes, providing relevant information on diabetes management including medical treatment, training HCPs, as well as creating a network between key diabetes care stakeholders and a registry of diabetic patients in Rwanda. Activities also include providing information to children and young adults on insulin (187)</p> <ul style="list-style-type: none"> • Other initiatives to improve the care of patients with T1DM in Rwanda include the Team Type 1 Foundation and Abbott to donate glucose meters as well as nearly 12 million test strips (53)
South Africa	<ul style="list-style-type: none"> • The cost of hospitalisation due to diabetes in South Africa ranges from I\$ 80 to I\$ 461 (27), with the treatment of DKA put a further strain on the system due to late presentations (34) • There are concerns glucose testing strips are not available to every patient and children are being treated with premixed insulin without access to rapid-acting insulin in public health facilities (93)
Sudan	<ul style="list-style-type: none"> • Encouragingly, the Audit Report of Global Fund Grants for the Republic of Sudan (2019) stated that the national medicines supply fund (NMSF), which is the public supply agency in Sudan, has robust processes and controls (188) • This is illustrated by the NMSF had never experienced stock-outs of insulin from 2013 to 2018. In addition, the NMSF has continued to supply insulin at fixed prices to the country because it has continued to use a fixed exchange rate of SDG18 for US\$1.00 for procurement despite the high inflation rate in Sudan, e.g. 63.3%, 2018. Consequently, public healthcare facilities in Sudan have managed to have a continuous supply of insulin preparations at prices lower than private sector (189): <ul style="list-style-type: none"> ○ Insulin (Mixtard®): Public retail price US\$0.82 Private retail price US\$4.76 ○ Insulin (Actrapid®) Public retail price US\$0.82, Private retail price US\$4.76 ○ Insulin Glargine: Public retail price US\$4.52, Private retail price US\$8.40. • However, the cost of T1DM is still a burden to families as a significant percent of household out-of-pocket expenses are used on medicines and medical consumables. • A study published in 2005 reported that the direct cost of diabetes care required 65% of the health expenditure of the whole family with the costs of insulin constituting approximately one-third of the total diabetes care expenditure (190) • A more recent study showed that a four-fold increase in the cost of managing diabetes in Sudan occurred between 2005 and 2013 (191), with costs expected to rise further
Zambia	<ul style="list-style-type: none"> • In Zambia, there are initiatives in place for patients to receive free or subsidized insulins through the national health insurance scheme operationalised in 2020 (47) • However, even at an average price of US\$2.00/ vial for insulin this is still a financial burden in countries where many patients live on less than US\$1.00/ day (47) • Overall insulin (soluble) was available in 38% of facilities recently surveyed by Mutale <i>et al.</i> (2018) and 18% in rural facilities (33)

	<ul style="list-style-type: none"> • Kalungia <i>et al.</i> (2017) also confirmed that stock-outs were high for short acting (83%) and long acting (69%) preparations among the surveyed healthcare facilities (192) • Kaiser <i>et al.</i> (2019) also found variable availability of insulins (short acting, intermediate acting and 30/70) ranging from a low of 4.6% among 99 surveyed pharmacists in three adjacent provinces in Zambia up to a 46.4% (193). Median prices of insulins also varied among the different pharmacies, e.g. between US\$11.33 to US\$24.79 for intermediate insulin and US\$16.80 - \$US28.82 for insulin 30/70 (193) • The routine availability of glucose strips at public healthcare facilities is still irregular with issues of affordability • Socioeconomic circumstances continue to hamper the use of advanced technologies to manage T1DM in Zambia. Major barriers to address include not only poverty and poor access to diabetes care, but also lack of knowledge among caregivers of T1DM patients which often leads to increased risk of misdiagnosis and late detection of T1DM (99).
--	--

NB: HCP: Healthcare Professional; T1DM = Type 1 diabetes; T2DM = Type 2 diabetes; PHCs = Primary healthcare centres

Table 3 – Facilities and treatments available in the public healthcare system among African countries to manage T1DM and its complications

Country	Treatments and Facilities
Botswana	<ul style="list-style-type: none"> • Random blood glucose within public healthcare facilities is part and parcel of the initial evaluation when patients first present with suspected diabetes in Botswana • Insulins routinely available among public healthcare facilities in Botswana include: <ul style="list-style-type: none"> • Short-acting regular human insulins <ul style="list-style-type: none"> ○ Actrapid ○ Rapid-acting analogue insulins ○ Aspart (NovoRapid) ○ Glulisine [Apidra] • Intermediate-acting (basal) human insulins <ul style="list-style-type: none"> ○ NPH - neutral protamine Hagedorn • Long-acting (basal) analogue insulins <ul style="list-style-type: none"> ○ Glargine [Lantus] • Pre-mixed human (biphasic) insulins <ul style="list-style-type: none"> ○ 30% Regular + 70% NPH [Actraphane] • Pre-mixed analogue insulins <ul style="list-style-type: none"> ○ Rapid-acting plus basal (NPH) ○ Biphasic aspart [NovoMix-] ○ Biphasic lispro [Humalog Mix25] <p>NB: Most of these drugs are available in specialized diabetes clinics. In non-specialised clinics, only a few insulins are available on specialist prescription forms (special order- NovoMix, NovoRapid).</p> <ul style="list-style-type: none"> • Treatments available to help manage the complications of T1DM within the public healthcare system include: <ul style="list-style-type: none"> ○ Medicines, e.g. blood pressure medicines – routinely available free of charge within public healthcare facilities ○ Procedures: retinal photocoagulation, renal replacement therapy by dialysis (both haemodialysis and peritoneal dialysis are available in tertiary healthcare facilities with the costs covered by the Government of Botswana), and renal replacement therapy by transplantation (currently performed outside of Botswana with the costs covered by government); coronary artery disease treatment is also available for diabetes patients where pertinent

	<ul style="list-style-type: none"> ○ Basic technologies: Blood glucose measurement, Urine strips for glucose and ketone measurements, HbA1c
Cameroon	<ul style="list-style-type: none"> • Patients are routinely seen by nurses and general practitioners in the Cameroon, with specialist care only available at regional or reference hospitals • All care is currently out of pocket. Micro insurance schemes do not cover medical costs for diabetes or related conditions. • Random blood sugar, fasting blood sugar, and urinalysis are routine tests to assess the care of patients with T1DM. HbA1c is not available. • Available options for insulin include INSULET RAPID (DCI: INSULINE GLARGINE 1000 UI/mL) ; INSULET MIX (DCI: INSULINE GLARGINE 1000 UI/mL); and INSULET NPH (DCI : INSULINE GLARGINE 100 UI/mL) • Chronic disease support groups known as Non-Communicable Disease clinics are common and provide monthly services to patients mostly for checkups
Ghana	<ul style="list-style-type: none"> • Insulins currently available in the Ghana EML for prescribing in PHCs include Insulin Aspart, Insulin Detemir, Insulin Glargine, Insulin Lispro, Insulin Soluble, Insulin Isophane (NPH) and Premixed (mixture of soluble insulin and intermediate NPH insulin in a 30:70 ratio) • The only insulins available under the National Health Insurance Scheme (NHIS) include insulin soluble, NPH and Premixed (30/70) while the rest are purchased out of pocket when prescribed (194) • Whilst some basic laboratory tests such as fasting blood glucose are provided under the NHIS, other relevant routine tests such as HbA1c and other supplies such as glucometer test strips and insulin syringes and needles currently have to be paid for out-of-pocket • Patients in PHCs are typically reviewed every 2-3 months and given refills of their insulin based on current availability at PHCs. However, not all health centres have dietitians and patients have to typically rely on their physician or nurse for food/nutrition advice • Complicated cases of T1DM requiring advanced specialist care including dialysis, renal replacement therapy are referred to one of the few available tertiary hospitals. Advanced care is not covered by the NHIS and therefore service are paid for out of pocket
Kingdom of Eswatini (Swaziland)	<p>Insulins listed in the EML of the Kingdom include:</p> <ol style="list-style-type: none"> 1. Soluble insulin 100units/ml injection 2. Isophane 100units/ml injection 3. Zinc insulin 100units/ml injection 4. Insulin+Isophane (30/70) 100units/ ml injection
Namibia	<ul style="list-style-type: none"> • A wide range of insulins including soluble and long acting insulins are available to the public via PHCs financed by the government of the Republic of Namibia.
Nigeria	<ul style="list-style-type: none"> • Short acting or meal time insulins, premixed insulin and insulin glargine are in the Nigerian EML. However, there are problems with availability and affordability of insulin glargine • The Diabetic Association of Nigeria has been involved with the development of practice guidelines in Nigeria to improve the care of patients with diabetes. The association has also had agreements with pharmaceutical companies to reduce the cost of their medicines to increase access and usage • There is typically routine monitoring of blood glucose using glucometer and laboratory testing in patients treated in secondary and tertiary hospitals. Routine monitoring of HbA1c is typically too expensive for most patients due to high co-payment levels in Nigeria • There are typically yearly referrals to eye clinics for assessment and treatment as well as routine treatment for systemic hypertension and dyslipidaemia when this occurs. Dialysis facilities are available but can be problematic in view of the costs involved

South Africa	<ul style="list-style-type: none"> • Screening of diabetes occurs at all levels of care depending on the patients' clinical conditions and their complaints • Random and fasting blood glucose within any healthcare facility is part and parcel of the initial evaluation • Insulins routinely available with public healthcare facilities include: <ul style="list-style-type: none"> ○ Insulin, short acting Regular human insulin ○ Insulin, intermediate acting ○ Insulin, biphasic mixtures of regular human insulin and intermediate acting insulin in different proportions, e.g. 30/70 (30% regular insulin and 70% intermediate acting insulin). • Treatment is typically initiated by medical doctors at any level of care within the public system depending on the availability of the physicians in PHC facilities on the day of consultation, with maintenance medicines sent to the PHC facilities monthly from the supporting district and regional hospitals • Stable patients are typically reviewed every six months • Basic technologies include blood glucose measurement, urine strips for glucose and ketone measurements, HbA1c, retinal examinations, renal function tests, neurological examinations, continuous blood sugar monitoring • Non-pharmacological management discussed include: diet, physical activity, lifestyle advice, behavioural therapy • Dialysis is available at the provincial and tertiary hospitals • However, there have been concerns with suboptimal control of hyperglycaemia in Africans despite being on treatment leading to calls for greater monitoring to reduce complications (195) • This has resulted in projects including TIP – a collaboration between the University of Pretoria, the Department of Health nationally and locally, and the City of Tshwane - Tshwane Insulin Project (TIP) initiated in 2018 and funded for over five years by Eli Lilly and Company. The TIP recognises concerns with initiating and optimising the use of insulin in patients with diabetes, with improving the management in primary care a key goal (196, 197) – building on concerns with the lack of HCPs trained in insulin use as well as patient-related factors including psychological fears with diabetes and insulin in South Africa • Under the TIP, there will be a major role for HCPs in primary care including primary care nurses and clinic physicians to help diagnose, initiate and monitor the use of insulins as well as test the feasibility of telemedicine and phone apps to improve the care of patients especially in more rural areas (196)
Sudan	<p>Insulin preparations listed on the Sudan NEML are (198):</p> <ul style="list-style-type: none"> • Short acting (soluble) Insulin • Glargine • Insulin Lispro • Insulin mixed • Intermediate-acting insulin <p>Based on the actual need as specified by the practitioners, NMSF covers only three of the insulin preparations included in the EML as follow:</p> <ul style="list-style-type: none"> • Insulin (Mixtard®) • Insulin (Actrapid®) • Insulin Glargine
Zambia	<ul style="list-style-type: none"> • Insulin preparations listed on the Zambia EML are: <ul style="list-style-type: none"> ○ Short acting (soluble) Insulin ○ Isophane Insulin, Biphasic Insulin ○ Actraphane or mixtard (30/70) ○ Long acting insulins <p>In Zambia, healthcare interventions available to prevent and manage complications include:</p>

	<ul style="list-style-type: none"> ○ Available routine monitoring of blood levels including BP measurement/treatment across all levels of public healthcare ○ At level 2 and 3 (district and tertiary hospitals), monitoring of patients regarding the development of retinopathy, neuropathy, etc., typically takes place ○ Dialysis for patients with nephropathy is mostly available at tertiary level hospitals ○ Referral and treatment of diabetes ketoacidosis takes place at level 2 and 3 hospitals ● Diabetes clinics manned by specialist physician established at level 2 and 3 hospitals
Zimbabwe	<ul style="list-style-type: none"> ● Listed in the Zimbabwe EML include: <ul style="list-style-type: none"> ○ Bolus insulins, e.g. rapid and short acting insulins including ASPART and ACTRAPID ○ Intermediate and long acting insulin analogues including biphasic insulins (30/70), PROTAPHANE as well as insulin glargine and degludec ● T1DM is typically treated starting in district hospitals. Routine laboratory testing facilities are available but there are concerns with the lack of consumables. In addition, the cost of tests is a deterrent for most diabetic patients in public institutions compromising their chronic care ● Complications associated with T1DM are assessed as part of routine care by physicians at eye and diabetic clinics including those in tertiary level facilities ● Dialysis is available at some provincial hospitals as well as in all 5 central hospitals in the country

NB: EML = Essential Medicines List; T1DM = Type 1 diabetes; PHCs = Primary healthcare centres

Box 1 – National/ regional governments and authorities

Leadership and governance

- Commit at the highest level to make quality T1DM healthcare services and medication based on current guidelines within countries available for all T1DM patients. In the long term, this should be free at the point of delivery irrespective of the ability of patients or their families
- Instigate initiatives to highlight to key government and other officials that T1DM is a public health concern across Africa. As part of these initiatives, include primary healthcare and disease prevention as part of a national health strategy to reduce morbidity and mortality due to NCDs
- Institutionalise routine screening for potential patients with T1DM in all healthcare facilities including both symptomatic and asymptomatic patients, and make screening free of charge. A key area will be primary healthcare centres within public healthcare systems as opposed to hospitals and specialist clinics
- Provide incentives for increasing the number of endocrinologists within a country where pertinent. In the short term, this could include using International aid to boost the number of endocrinologists within a country
- Introduce/ expand effective chronic disease management systems of care for patients with T1DM to improve the efficiency of care delivery
- Strengthen policy and systems towards T1DM management including encouraging adherence to WHO guidance as well as regular updates on country guidelines and NCD policies that incorporate T1DM. In addition, encourage operational research to monitor improvements in care delivery. As part of this, seek to provide direct economic, medical and social support for families with a child with T1DM
- Restructure where pertinent inventory management and procurement systems for insulin and its accessory devices nationally and among PHCs to ensure routine availability and accessibility to all types of insulin necessary for the effective management of patients with T1DM

Health care workforce

- Increase recruitment/ retention of general practitioners, family physicians with an interest in diabetes, dieticians, specialist pharmacists, and clinical psychologist with a special focus on PHCs for comprehensive clinical management in line with a growing focus on PHCs to diagnose and manage patients with T1DM in Africa. This alongside greater recruitment and retention of endocrinologists in Africa. As a result, improve and attain a good distribution of a relevant healthcare workforce across the spectrum
- Task shifting to enable pharmacists, nurses and community healthcare workers help run diabetes clinics and educational initiatives in both urban and rural settings
- Introduce a syllabus for HCPs on NCDs including T1DM so that all staff including clinical and non clinical staff can create a common language and framework for managing patients with T1DM
- Following this, regular continuous training and performance development for of all HCPs to improve their diagnosis and care of patients with T1DM including prevention and treatment of potential complications. This includes information of the mechanism of action of medicines and potential adverse events – and as part of this look to make available charts on how to manage T1DM including self-monitoring and potential complications

Medical products and technologies

- Improve the availability and supply of products and technologies used for T1DM management including routine/ consistent availability of insulins on the WHO EML as well as making fasting blood sugar levels and HbA1c testing more affordable for patients. This can be through collaborations with NGOs especially around the free provision of glucometers and test strips
- In the long term seek to include routine availability of long acting analogues with the increasing availability of biosimilars
- Explore solar- dependent refrigeration for insulin preparations at the household level especially in rural settings
- Consider non-invasive methods of glucose monitoring to improve compliance to self-care modalities
- Utilise cost-effective interventions
- Commit in the longer term for governments and health authorities to use HTA as part of decision-making processes to guide investment and funding decisions based on the efficacy, effectiveness, safety and cost-effectiveness of different technologies

Financing

- Improve healthcare financing towards T1DM management to reduce current morbidity and mortality rates. This can include expansion of health insurance towards universal access in a number of African countries
- This includes reducing or negating out-of-pocket payments and co-payment for insulin as well as for monitoring including glucose strips
- Improve medicine and product quality, access and pricing mechanisms to control for substandard products as well as improved prices given the price variations in insulins and other medicines to manage patients with T1DM across Africa and within countries
- Incentivise pharmacists as well other HCPs to establish medicines management centres in rural settings
- **Research and Innovation**
- Invest in epidemiological research where this is currently lacking including potentially the establishment of a national surveillance system/ databases for patients with T1DM to improve monitoring and evaluation of care delivery. This can include the introduction of IT systems/ electronic medical records to enable monitoring of care at the patient level
- Invest in HTA research where this is lacking to enable greater critique of evidence based potential alternatives within finite resources to improve efficient use of resources
- Invest in research surrounding possible innovative solutions including cost-effective solutions
- Encourage greater collaboration and linkage between research institutions, academia and policy makers

NB: EML = Essential Medicines List; HCPs = Healthcare professionals; HTA = Health Technology Assessment; T1DM = Type 1 diabetes; PHCs = Primary healthcare centres; NCDs = Non-communicable diseases; WHO = World Health Organisation

Box 2 – Physicians (hospital and ambulatory care)**Short term**

- Including greater opportunities for training and re-training of ambulatory care physicians in the management of patients with T1DM starting in pharmacy and medical schools and progressing to in-service training. This can also include greater knowledge surrounding key issues such as adherence to the medicines prescribed for both T1DM and its complications as well as the need for greater psychological support to improve patient care
- Work with key agencies to develop local evidence based guidelines for the management of patients with T1DM. These can subsequently be used as a basis for future quality improvement programmes
- Encourage more specialist doctors in fields such as endocrinology to contribute to better of patients with T1DM across Africa
- Work with others to seek to improve the availability of medicines for all T1DM patients
- Work with others to address ongoing bottlenecks in the pharmaceutical supply chain of T1DM medicines and products as well as working with others to help ensure quality and affordable medicines and technologies

Longer term

- Seek to attain a good distribution of physicians across the spectrum of health service provision, and geographical areas (rural and urban) settings
- Regular updating of guidelines when any changes in available medicines and technologies

NB: T1DM – Type 1 diabetes

Box 3 – Pharmacists (hospital and ambulatory care)**Short term**

- Continuous training and re-training of pharmacists regarding the management and education of patients with T1DM starting in pharmacy school
- Improve the participation and role of community pharmacists in providing care to patients with T1DM including counselling especially dealing with key issues such as re-fill prescriptions, insulins and needles as well as glucose monitoring. Include patient education to empower them to address key issues such as adherence, diet and lifestyles as well as concerns with the impact of T1DM on their quality of life
- Enhance the role of pharmacists in pharmacovigilance, medication error reporting and quality assurance of medicines and products used in T1DM. This can include the employment of Doctor of Pharmacy graduates (where these exist) in the public healthcare sector to improve patient-centred care
- Integrate and accredit community pharmacy services within national health insurance schemes. Services can be enhanced by introducing additional pharmacist training on the care of patients with diabetes. Look to include mandatory continuous performance development programmes
- Establishment and strengthening of medicines information services to all healthcare professionals as well as patients
- Encourage proactive data collection, analysis and feedback to both patients and healthcare providers at the facility level in terms of agreed diabetes management monitoring indicators

Longer term

- Conduct research and development of novel treatment modalities, medicines and products, including pharmacogenetics of T1DM in the population
- Provide incentives and opportunities for operational research on NCDs such as T1DM
- Potentially seek to establish pharmacist-led diabetic services at all levels of healthcare to address concerns with available facilities and personnel

NB: T1DM – Type 1 diabetes

Box 4 – Healthcare professionals in ambulatory and hospital care including nurse practitioners**Short term**

- Enhance competences and skills of the healthcare workforce to better diagnose and manage patients with T1DM through improved training including in-service training. Look to include the development and introduction of dedicated diabetic nurse educators
- Seek to train ambulatory and acute hospital care staff regarding any relevant advances and technologies used in the care of patients with T1DM
- Routinely provide relevant equipment for their roles

Longer term

- Undertake quality improvement research to address gaps in strategies and technical aspects to improve future service delivery

NB: T1DM – Type 1 diabetes

Box 5. Health technology (other than pharmaceutical therapy)**Short term**

- Continuous training and re-training of clinical technologists regarding the management of patients with diabetes
- Enhance the role of clinical technologists in surveillance, quality assurance of results and products used in T1DM. This can include the employment of clinical technology graduates (where these exist) in the public healthcare sector to improve patient-centred care
- Integrate and accredit community laboratory services within national health insurance scheme, enhanced by introduced of additional training on the care of patients with diabetes. This can ultimately include mandatory continuous performance development programmes

Longer term

- Involvement of biomedical engineers for development of newer diagnostic techniques
- Conduct research and development of novel diagnostic modalities that are affordable and can improve the management of patients with T1DM efficiently (such as continuous glucose monitoring system)
- Provide incentives and opportunities for operational research on NCDs such as T1DM
- Involvement of experts in Health Technology Assessment for measuring the impact of different health technologies on the future management of patients with T1DM

NB: NCDs = Non-communicable diseases; T1DM – Type 1 diabetes

Box 6 – Suggested activities among patients/ support services**Short term**

- Encourage the development and expansion of patient support networks to enhance the identification and subsequent management of patients with T1DM including the importance of diet and physical activities as well as the need for psychological management for patients with T1DM. For instance, in Zambia, the Diabetes Association of Zambia has been key in encouraging and engaging key stakeholder groups on diabetes matters, with the Diabetes Association in Nigeria playing a key role in guideline development and securing lower prices for medicines
- Undertake health education to communities to increase awareness, access to care and social support for patients with T1DM. This can include establishment of youth groups for children and adolescents in every community. Such groups can promote self care and reduce stigmatization associated with T1DM through group sharing of experiences

Longer term

- Instigate comprehensive adolescent health initiatives as part of patient education programmes that help to reduce the complications of T1DM including DKA, neuropathy, nephropathy and CV diseases
- Developing pertinent initiatives among T1DM patients and their families, particularly among adolescents, to ensure acceptance of diabetes and ant diabetes care plan as part of their lives - viewing diabetes as manageable and as their responsibility. As part of this, work with patients and patient organisations to gain valuable information on diabetes and diabetes management to improve future care (199)

- Encourage greater patient engagement and involvement in the health technology assessment process

NB: DKA= diabetic ketoacidosis; T1DM – Type 1 diabetes

References

1. Atun R, Davies JI, Gale EAM, Barnighausen T, Beran D, Kengne AP, et al. Diabetes in sub-Saharan Africa: from clinical care to health policy. *The lancet Diabetes & endocrinology*. 2017;5(8):622-67.
2. Mapa-Tassou C, Katte JC, Mba Maadjhou C, Mbanya JC. Economic Impact of Diabetes in Africa. *Current diabetes reports*. 2019;19(2):5.
3. WHO. GLOBAL REPORT ON DIABETES. 2016. Available at URL: http://apps.who.int/iris/bitstream/10665/204871/1/9789241565257_eng.pdf?ua=1.
4. Meyer JC, Schellack N, Stokes J, Lancaster R, Zeeman H, Defty D, et al. Ongoing Initiatives to Improve the Quality and Efficiency of Medicine Use within the Public Healthcare System in South Africa; A Preliminary Study. *Frontiers in pharmacology*. 2017;8:751.
5. Nashilongo MM, Singu B, Kalemeera F, Mubita M, Naikaku E, Baker A, et al. Assessing Adherence to Antihypertensive Therapy in Primary Health Care in Namibia: Findings and Implications. *Cardiovascular drugs and therapy*. 2017;31(5-6):565-78.
6. Mbui JM, Oluka MN, Guantai EM, Sinei KA, Achieng L, Baker A, et al. Prescription patterns and adequacy of blood pressure control among adult hypertensive patients in Kenya; findings and implications. *Expert review of clinical pharmacology*. 2017;10(11):1263-71.
7. Mwitwa JC, Francis JM, Omech B, Botsile E, Oyewo A, Mokgwathi M, et al. Glycaemic, blood pressure and low-density lipoprotein-cholesterol control among patients with diabetes mellitus in a specialised clinic in Botswana: a cross-sectional study. *BMJ open*. 2019;9(7):e026807.
8. Webb EA, Hesselting AC, Schaaf HS, Gie RP, Lombard CJ, Spitaels A, et al. High prevalence of Mycobacterium tuberculosis infection and disease in children and adolescents with type 1 diabetes mellitus. *The international journal of tuberculosis and lung disease*. 2009;13(7):868-74.
9. Reid MJA, Tsimba BM, Kirk B. HIV and diabetes in Africa. *African Journal of Diabetes Medicine*. 2012; 20 (2): 28-32.
10. Statistics South Africa. Mortality and causes of death in South Africa, 2016: Findings from death notification. Available at URL: <http://www.statssa.gov.za/publications/P03093/P030932016.pdf>.
11. Moosa A, Bezuidenhout S, Meyer JC, Godman B. Knowledge regarding medicines management of type-2 diabetes among patients attending a Community Health Centre in South Africa. *Journal of Pharmaceutical Health Services Research*. 2019, 10; 13–21.
12. Fasanmade OA, Dagogo-Jack S. Diabetes Care in Nigeria. *Ann Glob Health*. 2015;81(6):821-9.
13. Rwegerera GM, Moshomo T, Gaenamong M, Oyewo TA, Gollakota S, Rivera YP, et al. Health-related quality of life and associated factors among patients with diabetes mellitus in Botswana. *Alexandria Journal of Medicine*. 2018;54(2):111-8.
14. da Mata AR, Alvares J, Diniz LM, da Silva MR, Alvernaz dos Santos BR, Guerra Junior AA, et al. Quality of life of patients with Diabetes Mellitus Types 1 and 2 from a referral health centre in Minas Gerais, Brazil. *Expert review of clinical pharmacology*. 2016;9(5):739-46.
15. Hayes A, Arima H, Woodward M, Chalmers J, Poulter N, Hamet P, et al. Changes in Quality of Life Associated with Complications of Diabetes: Results from the ADVANCE Study. *Value in Health*. 2016;19(1):36-41.
16. Romero-Naranjo F, Espinosa-Uquillas C, Barrera-Guarderas F, Gordillo Altamirano F. Which Factors may reduce the Health-Related Quality of Life of Ecuadorian Patients with Diabetes? *Puerto Rico health sciences journal*. 2019;38:102-8.
17. Piloya-Were T, Sunni M, Ogle GD, Moran A. Childhood diabetes in Africa. *Current opinion in endocrinology, diabetes, and obesity*. 2016;23(4):306-11.
18. Bos M, Agyemang C. Prevalence and complications of diabetes mellitus in Northern Africa, a systematic review. *BMC public health*. 2013;13:387-.
19. International Diabetes Federation. IDF Africa Members. 2019. Available at URL: <https://idf.org/our-network/regions-members/africa/members/25-south-africa.html>.
20. IDF. IDF DIABETES ATLAS Ninth Edition Africa. 2019. Available at URL: https://diabetesatlas.org/upload/resources/2019/2019_regional_factsheet.pdf.
21. Elliot J, Besançon S, Sachs J. Almost 100 years after it was discovered, let's make access to insulin a reality in Africa! 2016. Available at URL: <https://ideas4development.org/en/almost-100-years-after-it-was-discovered-lets-make-access-to-insulin-a-reality-in-africa/>.

22. Majeed A, El-Sayed AA, Khoja T, Alshamsan R, Millett C, Rawaf S. Diabetes in the Middle-East and North Africa: an update. *Diabetes research and clinical practice*. 2014;103(2):218-22.
23. Sherif S, Sumpio BE. Economic development and diabetes prevalence in MENA countries: Egypt and Saudi Arabia comparison. *World journal of diabetes*. 2015;6(2):304-11.
24. International Diabetes Federation. IDF Diabetes Atlas 7th edition. Available at URL: <https://www.idf.org/e-library/epidemiology-research/diabetes-atlas/13-diabetes-atlas-seventh-edition.html>.
25. Ogbera AO, Fasanmade O, Ohwovoriole AE, Adediran O. An assessment of the disease burden of foot ulcers in patients with diabetes mellitus attending a teaching hospital in Lagos, Nigeria. *The international journal of lower extremity wounds*. 2006;5(4):244-9.
26. da Rocha Fernandes J, Ogurtsova K, Linnenkamp U, Guariguata L, Seuring T, Zhang P, et al. IDF Diabetes Atlas estimates of 2014 global health expenditures on diabetes. *Diabetes research and clinical practice*. 2016;117:48-54.
27. Mutyambizi C, Pavlova M, Chola L, Hongoro C, Groot W. Cost of diabetes mellitus in Africa: a systematic review of existing literature. *Global Health*. 2018;14(1):3.
28. Ng CS, Lee JYC, Toh MP, Ko Y. Cost-of-illness studies of diabetes mellitus: a systematic review. *Diabetes research and clinical practice*. 2014;105(2):151-63.
29. Quaye EA, Amporful EO, Akweongo P, Aikins MK. Analysis of the Financial Cost of Diabetes Mellitus in Four Cocoa Clinics of Ghana. *Value in health regional issues*. 2015;7:49-53.
30. Pei F. Managing Diabetes in Urban Ghana: Is it Affordable? 2015. Available at URL: https://dukespace.lib.duke.edu/dspace/bitstream/handle/10161/10014/Pei_duke_0066N_12921.pdf?sequence=1&isAllowed=y.
31. Alouki K, Delisle H, Besancon S, Balde N, Sidibe-Traore A, Drabo J, et al. Simple calculator to estimate the medical cost of diabetes in sub-Saharan Africa. *World J Diabetes*. 2015;6(16):1312-22.
32. Suleiman IA, Festus JA. Cost of illness among diabetes mellitus patients in Niger Delta, Nigeria. *J Pharm Health Serv Res*. 2015;6(1):53-60.
33. Mutale W, Bosomprah S, Shankalala P, Mweemba O, Chilengi R, Kapambwe S, et al. Assessing capacity and readiness to manage NCDs in primary care setting: Gaps and opportunities based on adapted WHO PEN tool in Zambia. *PLoS one*. 2018;13(8):e0200994-e.
34. Reddy Y, Gannie Y, Pillay K. Characteristics of children presenting with newly diagnosed type1 diabetes. *SAJCH*. 2013; 7(2).
35. Ministry of Health and Social Services, Primary Health Care Directorate, Namibia. National Multisectoral Strategic Plan For Prevention and Control of Non-Communicable Diseases (NCDs) in Namibia 2017/18 – 2021/22. 2017. Available at URL: <https://www.iccp-portal.org/system/files/plans/NAMIBIA%20NATIONAL%20MULTISECTORAL%20STRATEGIC%20PLAN%20FOR%20PREVENTION%20AND%20CONTROL%20OF%20NCDs.pdf>.
36. Mukanu MM, Zulu JM, Mweemba C, Mutale W. Responding to non-communicable diseases in Zambia: a policy analysis. *Health Research Policy and Systems*. 2017;15(1):34.
37. Federal Ministry of Health, SIDCAIN, World Diabetes Foundation. Diabetes and NCDs in Nigeria - Perspectives, Challenges & the Way Forward. 2017. Available at URL: <http://sidcain.org/download/ABUJA%20Proceedings%20Reviewed%20Nov%202017.pdf>.
38. NDOH. National Department of Health Republic of South Africa. Adherence Guidelines for HIV, TB and NCDs. Policy and service delivery guidelines to care, adherence to treatment and retention in care. February 2016. Available at URL: <https://www.nacosa.org.za/wp-content/uploads/2016/11/Integrated-Adherence-Guidelines-NDOH.pdf>
39. KINGDOM OF ESWATINI - MINISTRY OF HEALTH. 2017 NATIONAL PREVENTION AND CONTROL OF NON COMMUNICABLE DISEASES (NCDs). 2017 Annual Program Report.
40. Igbojiaku OJ, Harbor OC, Ross A. Compliance with diabetes guidelines at a regional hospital in KwaZulu-Natal, South Africa. *African Journal of Primary Health Care & Family Medicine*. 2013;5(1):447.
41. SEMDSA 2017 Guidelines for the Management of Type 2 diabetes mellitus. SEMDSA Type 2 Diabetes Guidelines Expert Committee. *JEMDSA* 2017; 22(1)(Supplement 1): S1-S196.
42. Awodele O, Osuolale JA. Medication adherence in type 2 diabetes patients: study of patients in Alimosho General Hospital, Igando, Lagos, Nigeria. *Afr Health Sci*. 2015;15(2):513-22.
43. Botswana National Multisectoral Strategy for the Prevention and Control of Non-Communicable Diseases. 2017. Available at URL: <https://www.iccp-portal.org/system/files/plans/20171129%20Botswana%20multisec%20NCDs%20strategy%20-%20final.pdf>.

- 1
- 2
- 3
- 4 44. Khogali SS, Ali WA, Mohamed SY, Abdelrahim HE, Mirghani AA, Ali RH et al. Knowledge, attitude and practice of Sudanese individuals with type 2 diabetes about medication used in treatment of diabetes, hypertension and dyslipidaemia: a matter of debate or matter of concern? *J Public Health Emerg.* 2018;2:23.
- 5
- 6
- 7 45. Bahendeka SK. Diabetes in sub-Saharan Africa: let us not forget type 1. *The Lancet Diabetes & endocrinology.* 2017;5(8):575-7.
- 8
- 9 46. Shannon GD, Haghparast-Bidgoli H, Chelagat W, Kibachio J, Skordis-Worrall J. Innovating to increase access to diabetes care in Kenya: an evaluation of Novo Nordisk's base of the pyramid project. *Glob Health Action.* 2019;12(1):1605704.
- 10
- 11 47. International Insulin Foundation – Fact Sheet on diabetes in sub-Saharan Africa. Available at URL: <http://www.access2insulin.org/uploads/4/9/1/0/4910107/factsheet.pdf>.
- 12
- 13 48. Djonou C, Tankeu AT, Dehayem MY, Tchetchoua DN, Mbanja JC, Sobngwi E. Glycemic control and correlates in a group of sub Saharan type 1 diabetes adolescents. *BMC research notes.* 2019;12(1):50.
- 14
- 15 49. Balde N, Camara A, Sobngwi-Tambekou J, Balti EV, Tchatchoua A, Fezeu L, et al. Improving access to HbA1c in sub-Saharan Africa (IA3) cohort: cohort profile. *The Pan African medical journal.* 2017;27:275-.
- 16
- 17 50. Park PH, Pastakia SD. Access to Hemoglobin A1c in Rural Africa: A Difficult Reality with Severe Consequences. *Journal of diabetes research.* 2018;2018:6093595.
- 18
- 19 51. Republic of Zambia. Ministry of Health. Zambia National Health Strategic Plan 2017–2021. Lusaka; 2017. Available at URL: <https://www.moh.gov.zm/docs/ZambiaNHSP.pdf>.
- 20
- 21 52. Tapela NM, Tshisimogo G, Shatera BP, Letsatsi V, Gaborone M, Madidimalo T, et al. Integrating noncommunicable disease services into primary health care, Botswana. *Bulletin of the World Health Organization.* 2019;97(2):142-53.
- 22
- 23 53. WHO Africa. The Team Type 1 Foundation and Abbott partner to donate glucose meters and nearly 12 million test strips to help people living with diabetes in Rwanda. 2020. Available at URL: <https://www.afro.who.int/news/team-type-1-foundation-and-abbott-partner-donate-glucose-meters-and-nearly-12-million-test>.
- 24
- 25 54. Kenya Ministry of Health. Transforming Health: Accelerating attainment of Universal Health Coverage. The KENYA HEALTH SECTOR STRATEGIC AND INVESTMENT PLAN – KHSSP July 2012 –June 2017. Available at URL: <http://e-cavi.com/wp-content/uploads/2014/11/kenya-health-sector-strategic-investment-plan-2013-to-2017.pdf>.
- 26
- 27 55. Rampamba EM, Meyer JC, Helberg EA, Godman B. Empowering Hypertensive Patients in South Africa to Improve Their Disease Management: A Pharmacist-Led Intervention. *J Res Pharm Pract.* 2019;8(4):208-13.
- 28
- 29 56. Godman B, Basu D, Pillay Y, Mwita JC, Rweggerera GM, Anand Paramadhas BD et al. Review of ongoing activities and challenges to improve the care of patients with Type 2 diabetes across Africa and the implications for the future. 2020. Accepted for publication in *Frontiers in Pharmacology.*
- 30
- 31 57. Pambou Damiens A, Ganga-Zandzou PS, Tsoucka Ibounde E, Kayemba-Kay's S, Baye E, Biloghe P, et al. Type 1 diabetes mellitus in Gabon: A study of epidemiological aspects. *International journal of pediatrics & adolescent medicine.* 2019;6(3):87-91.
- 32
- 33 58. Hall V, Thomsen RW, Henriksen O, Lohse N. Diabetes in Sub Saharan Africa 1999-2011: Epidemiology and public health implications. a systematic review. *BMC public health.* 2011;11:564-.
- 34
- 35 59. Pastakia SD, Pekny CR, Manyara SM, Fischer L. Diabetes in sub-Saharan Africa - from policy to practice to progress: targeting the existing gaps for future care for diabetes. *Diabetes Metab Syndr Obes.* 2017;10:247-63.
- 36
- 37 60. Uloko AE, Ofoegbu EN, Chinenye S, Fasanmade OA, Fasanmade AA, Ogbera AO, et al. Profile of Nigerians with diabetes mellitus - Diabcare Nigeria study group (2008): Results of a multicenter study. *Indian J Endocrinol Metab.* 2012;16(4):558-64.
- 38
- 39 61. Majaliwa ES, Elusiyan BE, Adesiyun OO, Laigong P, Adeniran AK, Kandi CM, et al. Type 1 diabetes mellitus in the African population: epidemiology and management challenges. *Acta bio-medica.* 2008;79(3):255-9.
- 40
- 41 62. Patterson CC, Karuranga S, Salpea P, Saeedi P, Dahlquist G, Soltesz G, et al. Worldwide estimates of incidence, prevalence and mortality of type 1 diabetes in children and adolescents: Results from the International Diabetes Federation Diabetes Atlas, 9th edition. *Diabetes research and clinical practice.* 2019;157:107842.
- 42
- 43 63. Chikani UN, Bisi-Onyemaechi AI, Oguonu T, Ugege SM, Ogugua C. Childhood diabetes: a myth or reality?- perception of the public from a low-income country: a cross-sectional study. *BMC public health.* 2018;18(1):852.
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

- 1
- 2
- 3 64. Padoa CJ. The epidemiology and pathogenesis of type 1 diabetes mellitus in Africa. *Journal of Endocrinology, Metabolism and Diabetes of South Africa*. 2011;16(3):130-6.
- 4 65. Ngwiri T, Were F, Predieri B, Ngugi P, Iughetti L. Glycemic Control in Kenyan Children and
- 5 Adolescents with Type 1 Diabetes Mellitus. *International journal of endocrinology*. 2015;2015:761759.
- 6 66. International Diabetes Federation. *IDF Diabetes Atlas*. 2017; 8th edition. Available at URL:
- 7 <http://www.diabetesatlas.org>.
- 8 67. Mutyambizi C, Booysen F, Stokes A, Pavlova M, Groot W. Lifestyle and socio-economic
- 9 inequalities in diabetes prevalence in South Africa: A decomposition analysis. *PLoS one*.
- 10 2019;14(1):e0211208.
- 11 68. Amoah AG, Owusu SK, Adjei S. Diabetes in Ghana: a community based prevalence study in
- 12 Greater Accra. *Diabetes research and clinical practice*. 2002;56(3):197-205.
- 13 69. Niba LL. Determinants of outcome of children with type 1 diabetes in the North West Region
- 14 of Cameroon. 2016. Available at URL: https://edoc.ub.uni-muenchen.de/20100/7/Niba_Loveline.pdf.
- 15 70. Padoa CJ, Rheeder P, Pirie FJ, Motala AA, van Dyk JC, Crowther NJ. Identification of a
- 16 subgroup of black South Africans with type 1 diabetes who are older at diagnosis but have lower
- 17 levels of glutamic acid decarboxylase and islet antigen 2 autoantibodies. *Diabetic medicine : a journal*
- 18 *of the British Diabetic Association*. 2019.
- 19 71. Marshall SL, Edidin D, Arena VC, Becker DJ, Bunker CH, Gishoma C, et al. Prevalence and
- 20 incidence of clinically recognized cases of Type 1 diabetes in children and adolescents in Rwanda,
- 21 Africa. *Diabetic medicine*. 2015;32(9):1186-92.
- 22 72. Marshall SL, Edidin DV, Arena VC, Becker DJ, Bunker CH, Gishoma C, et al. Glucose control
- 23 in Rwandan youth with type 1 diabetes following establishment of systematic, HbA1c based, care and
- 24 education. *Diabetes research and clinical practice*. 2015;107(1):113-22.
- 25 73. Balcha SA, Phillips DIW, Trimble ER. Type 1 Diabetes in a Resource-Poor Setting:
- 26 Malnutrition Related, Malnutrition Modified, or Just Diabetes? *Current diabetes reports*.
- 27 2018;18(7):47.
- 28 74. Pirie FJ, Pegoraro R, Motala AA, Rauff S, Rom L, Govender T, et al. Toll-like receptor 3 gene
- 29 polymorphisms in South African Blacks with type 1 diabetes. *Tissue Antigens*. 2005;66(2):125-30.
- 30 75. Panz VR, Kalk WJ, Zouvanis M, Joffe BI. Distribution of autoantibodies to glutamic acid
- 31 decarboxylase across the spectrum of diabetes mellitus seen in South Africa. *Diabetic medicine*.
- 32 2000;17(7):524-7.
- 33 76. Pirie FJ, Hammond MG, Motala AA, Omar MA. HLA class II antigens in South African Blacks
- 34 with type I diabetes. *Tissue Antigens*. 2001;57(4):348-52.
- 35 77. International Diabetes Federation. *IDF Atlas Ninth Edition*. 2019. Available at URL:
- 36 https://diabetesatlas.org/upload/resources/material/20200106_152211_IDFATLAS9e-final-web.pdf.
- 37 78. Gill GV, Huddle KR, Monkoe G. Long-term (20 years) outcome and mortality of Type 1
- 38 diabetic patients in Soweto, South Africa. *Diabetic medicine*. 2005;22(12):1642-6.
- 39 79. Beran D, Yudkin JS, de Courten M. Access to care for patients with insulin-requiring diabetes
- 40 in developing countries: case studies of Mozambique and Zambia. *Diabetes Care*. 2005;28(9):2136-
- 41 40.
- 42 80. Atkilt HS, Turago MG, Tegegne BS. Clinical Characteristics of Diabetic Ketoacidosis in
- 43 Children with Newly Diagnosed Type 1 Diabetes in Addis Ababa, Ethiopia: A Cross-Sectional Study.
- 44 *PLoS one*. 2017;12(1):e0169666-e.
- 45 81. Kitabchi AE, Umpierrez GE, Miles JM, Fisher JN. Hyperglycemic crises in adult patients with
- 46 diabetes. *Diabetes Care*. 2009;32(7):1335-43.
- 47 82. Azevedo M, Alla S. Diabetes in sub-saharan Africa: kenya, mali, mozambique, Nigeria, South
- 48 Africa and zambia. *International journal of diabetes in developing countries*. 2008;28(4):101-8.
- 49 83. Onyiriuka AN, Ifebi E. Ketoacidosis at diagnosis of type 1 diabetes in children and
- 50 adolescents: frequency and clinical characteristics. *Journal of diabetes and metabolic disorders*.
- 51 2013;12(1):47.
- 52 84. Ibekwe M U, Ibekwe R C. Pattern of Type 1 Diabetes Mellitus in Abakaliki, Southeastern,
- 53 Nigeria. *Pediatr Oncall J*. 2011;8: 59-62.
- 54 85. Murunga AN, Owira PM. Diabetic ketoacidosis: an overlooked child killer in sub-Saharan
- 55 Africa? *Trop Med Int Health*. 2013;18(11):1357-64.
- 56 86. Ndebele NFM, Naidoo M. The management of diabetic ketoacidosis at a rural regional
- 57 hospital in KwaZulu-Natal. *Afr J Prim Health Care Fam Med*. 2018;10(1):e1-e6.
- 58 87. Majaliwa ES, Mohn A, Chiavaroli V, Ramaiya K, Swai AB, Chiarelli F. Management of diabetic
- 59 ketoacidosis in children and adolescents in sub-Saharan Africa: a review. *East Afr Med J*.
- 60 2010;87(4):167-73.

- 1
2
3 88. Iovane B, Cangelosi AM, Bonaccini I, Di Mauro D, Scarabello C, Panigari A, et al. Diabetic ketoacidosis at the onset of Type 1 diabetes in young children Is it time to launch a tailored campaign for DKA prevention in children <5 years? *Acta bio-medica*. 2018;89(1):67-71.
- 4
5
6 89. Van Zyl DG. Diagnosis and treatment of diabetic ketoacidosis. *South African Family Practice*. 2008;50(1):35-9.
- 7
8 90. Adeleke SI, Asani MO, Belonwu RO, Gwarzo GD, Farouk ZL. Childhood diabetes mellitus in Kano, North West, Nigeria. *Nigerian journal of medicine*. 2010;19(2):145-7.
- 9
10 91. Mukama LJ, Moran A, Nyindo M, Philemon R, Msuya L. Improved glycemic control and acute complications among children with type 1 diabetes mellitus in Moshi, Tanzania. *Pediatric diabetes*. 2013;14(3):211-6.
- 11
12
13 92. Emral R, Pathan F, Cortés CAY, El-Hefnawy MH, Goh S-Y, Gómez AM, et al. Self-reported hypoglycemia in insulin-treated patients with diabetes: Results from an international survey on 7289 patients from nine countries. *Diabetes research and clinical practice*. 2017;134:17-28.
- 14
15
16 93. Kalweit KL, Briers N, Olorunju S. The success of various management techniques used in South African children with type 1 diabetes mellitus. *South African medical journal*. 2015;105(5):400-4.
- 17
18
19 94. Fulcher G, Singer J, Castañeda R, Fraige Filho F, Maffei L, Snyman J, et al. The psychosocial and financial impact of non-severe hypoglycemic events on people with diabetes: two international surveys. *Journal of medical economics*. 2014;17(10):751-61.
- 20
21
22 95. Kinsley B. Achieving better outcomes in pregnancies complicated by type 1 and type 2 diabetes mellitus. *Clinical therapeutics*. 2007;29 Suppl D:S153-S60.
- 23
24
25 96. Muddu M, Mutebi E, Ssinabulya I, Kizito S, Mondo CK. Hypertension among newly diagnosed diabetic patients at Mulago National Referral Hospital in Uganda: a cross sectional study. *Cardiovascular journal of Africa*. 2018;29(4):218-24.
- 26
27
28 97. Ahmed SA, Badi S, Tahir H, Ahmed MH, Almobarak AO. Knowledge and practice of diabetic foot care in Sudan: A cross sectional survey. *Diabetes & metabolic syndrome*. 2019;13(4):2431-5.
- 29
30
31 98. Wagnew F, Eshetie S, Kibret GD, Zegeye A, Dessie G, Mulugeta H, et al. Diabetic nephropathy and hypertension in diabetes patients of sub-Saharan countries: a systematic review and meta-analysis. *BMC research notes*. 2018;11(1):565.
- 32
33
34 99. Hapunda G, Abubakar A, van de Vijver F, Pouwer F. Living with type 1 diabetes is challenging for Zambian adolescents: qualitative data on stress, coping with stress and quality of care and life. *BMC endocrine disorders*. 2015;15:20-.
- 35
36
37 100. Nieuwesteeg A, Pouwer F, van der Kamp R, van Bakel H, Aanstoot H-J, Hartman E. Quality of life of children with type 1 diabetes: a systematic review. *Current diabetes reviews*. 2012;8(6):434-43.
- 38
39
40 101. Oluyombo R, Okunola OO, Olanrewaju TO, Soje MO, Obajolowo OO, Ayorinde MA. Challenges of hemodialysis in a new renal care center: call for sustainability and improved outcome. *International journal of nephrology and renovascular disease*. 2014;7:347-52.
- 41
42
43 102. Makhele L, Matlala M, Sibanda M, Martin AP, Godman B. A Cost Analysis of Haemodialysis and Peritoneal Dialysis for the Management of End-Stage Renal Failure At an Academic Hospital in Pretoria, South Africa. *PharmacoEconomics - open*. 2019.
- 44
45
46 103. Beran D, Hirsch IB, Yudkin JS. Why Are We Failing to Address the Issue of Access to Insulin? A National and Global Perspective. *Diabetes Care*. 2018;41(6):1125-31.
- 47
48
49 104. MANSFIELD J. TYPE 1 DIABETES IN GHANA: A GLOBAL COMMUNITY DISCUSSION. 2018. Available at URL: <https://beyondtype1.org/type-1-diabetes-ghana/>.
- 50
51
52 105. Pirie FJ, Jairam V, Paruk IM, Connolly C, Motala AA. High frequency of hypoglycaemia in patients with type 1 diabetes mellitus attending a tertiary diabetes clinic in Durban, South Africa. *Diabetes research and clinical practice*. 2019;155:107783.
- 53
54
55 106. Beran D, Ewan M, Laing R. Access to insulin - Current Challenges & Constraints. October 2015. Available at URL: <https://apps.who.int/medicinedocs/documents/s22269en/s22269en.pdf>.
- 56
57
58 107. Ewen M, Joosse HJ, Beran D, Laing R. Insulin prices, availability and affordability in 13 low-income and middle-income countries. *BMJ global health*. 2019;4(3):e001410.
- 59
60
61 108. Beran D, Ewen M, Laing R. Constraints and challenges in access to insulin: a global perspective. *The lancet Diabetes & endocrinology*. 2016;4(3):275-85.
- 62
63
64 109. Beran D, Laing RO, Kaplan W, Knox R, Sharma A, Wirtz VJ, et al. A perspective on global access to insulin: a descriptive study of the market, trade flows and prices. *Diabetic medicine*. 2019;36(6):726-33.
- 65
66
67 110. Tiruneh GG, Abebe N, Dessie G. Self-reported hypoglycemia in adult diabetic patients in East Gojjam, Northwest Ethiopia: institution based cross-sectional study. *BMC Endocr Disord*. 2019;19(1):17.

- 1
2
3 111. Kabsay H, Fantahun B, Nedi T, Demoz GT. Evaluation of Hypoglycemia and Associated
4 Factors among Patients with Type 1 Diabetes on Follow-Up Care at St. Paul's Hospital Millennium
5 Medical College, Addis Ababa, Ethiopia. *Journal of diabetes research*. 2019;2019:9037374.
- 6 112. Kratzer J. Structural barriers to coping with type 1 diabetes mellitus in Ghana: experiences of
7 diabetic youth and their families. *Ghana medical journal*. 2012;46(2 Suppl):39-45.
- 8 113. T1International. Ghana. Available at URL: <https://www.t1international.com/Ghana/>.
- 9 114. World Health Organisation (WHO). Global action Plan for the prevention and control of
10 noncommunicable diseases. 2013. Available at URL:
11 https://apps.who.int/iris/bitstream/handle/10665/94384/9789241506236_eng.pdf;jsessionid=7E6647E
12 [D415CFBF876A7CC201127FB76?sequence=1](https://apps.who.int/iris/bitstream/handle/10665/94384/9789241506236_eng.pdf;jsessionid=7E6647ED415CFBF876A7CC201127FB76?sequence=1).
- 13 115. Kapwata T, Manda S. Geographic assessment of access to health care in patients with
14 cardiovascular disease in South Africa. *BMC health services research*. 2018;18(1):197-.
- 15 116. Essaddam L, Kallali W, Jemel M, Kandara H, Kammoun I, Hsairi M, et al. Implementation of
16 effective transition from pediatric to adult diabetes care: epidemiological and clinical characteristics-a
17 pioneering experience in North Africa. *Acta diabetologica*. 2018;55(11):1163-9.
- 18 117. Peck R, Mghamba J, Vanobberghen F, Kavishe B, Rugarabamu V, Smeeth L, et al.
19 Preparedness of Tanzanian health facilities for outpatient primary care of hypertension and diabetes:
20 a cross-sectional survey. *The Lancet Global health*. 2014;2(5):e285-e92.
- 21 118. Pillay DK, Ross AJ, Campbell L. A review of hypoglycaemia in a South African family practice
22 setting. *African journal of primary health care & family medicine*. 2016;8(1):e1-e6.
- 23 119. Kaplan H, Amodb A, van Zyk FH, Reddy J, van Tondere A, Tsybale E. Incidence of
24 hypoglycaemia in the South African population with diabetes: results from the IDMPs Wave 7 study.
25 *Journal of Endocrinology, Metabolism and Diabetes of South Africa* 2019; 24(2):58–64.
- 26 120. Sandoz - A Novartis Division. Kenya is first country to launch 'Novartis Access', expanding
27 affordable treatment options against chronic diseases. 2015. Available at URL:
28 [https://www.sandoz.com/news/media-releases/kenya-first-country-launch-novartis-access-expanding-](https://www.sandoz.com/news/media-releases/kenya-first-country-launch-novartis-access-expanding-affordable-treatment)
29 [affordable-treatment](https://www.sandoz.com/news/media-releases/kenya-first-country-launch-novartis-access-expanding-affordable-treatment).
- 30 121. Kabeza CB, Harst L, Schwarz PEH, Timpel P. Assessment of Rwandan diabetic patients'
31 needs and expectations to develop their first diabetes self-management smartphone application
32 (Kir'App). *Therapeutic advances in endocrinology and metabolism*. 2019;10:2042018819845318-.
- 33 122. Novo Nordisk. NOVO NORDISK EXPANDS PROGRAMME TO REACH 20,000 CHILDREN
34 WITH DIABETES IN DEVELOPING COUNTRIES. 2016. Available at URL:
35 <http://www.multivu.com/players/uk/7962351-novo-nordisk-programme-children-diabetes/>.
- 36 123. Roche. Changing diabetes care for children in Africa. 2019. Available at URL:
37 https://www.roche.com/sustainability/access-to-healthcare/ath_diabetes.htm.
- 38 124. Heise T. Getting closer to physiologic insulin secretion. *Clinical therapeutics*. 2007;29 Suppl
39 D:S161-S5.
- 40 125. Danne T. Flexibility of rapid-acting insulin analogues in children and adolescents with
41 diabetes mellitus. *Clinical therapeutics*. 2007;29 Suppl D:S145-S52.
- 42 126. WHO. World Health Organization Model List of Essential Medicines 2019. Available at URL:
43 [https://apps.who.int/iris/bitstream/handle/10665/325771/WHO-MVP-EMP-IAU-2019.06-](https://apps.who.int/iris/bitstream/handle/10665/325771/WHO-MVP-EMP-IAU-2019.06-eng.pdf?ua=21st%20List%202019)
44 [eng.pdf?ua=21st List 2019](https://apps.who.int/iris/bitstream/handle/10665/325771/WHO-MVP-EMP-IAU-2019.06-eng.pdf?ua=21st List 2019).
- 45 127. Jaam M, Ibrahim MIM, Kheir N, Awaisu A. Factors associated with medication adherence
46 among patients with diabetes in the Middle East and North Africa region: A systematic mixed studies
47 review. *Diabetes research and clinical practice*. 2017;129:1-15.
- 48 128. Silver B, Ramaiya K, Andrew SB, Fredrick O, Bajaj S, Kalra S, et al. EADSG Guidelines:
49 Insulin Therapy in Diabetes. *Diabetes therapy*. 2018;9(2):449-92.
- 50 129. Ogle GD, Kim H, Middlehurst AC, Silink M, Jenkins AJ. Financial costs for families of children
51 with Type 1 diabetes in lower-income countries. *Diabetic medicine*. 2016;33(6):820-6.
- 52 130. Manne-Goehler J, Geldsetzer P, Agoudavi K, Andall-Brereton G, Aryal KK, Bicaba BW, et al.
53 Health system performance for people with diabetes in 28 low- and middle-income countries: A cross-
54 sectional study of nationally representative surveys. *PLoS Med*. 2019;16(3):e1002751.
- 55 131. Owolabi MO, Yaria JO, Daivadanam M, Makanjuola AI, Parker G, Oldenburg B, et al. Gaps in
56 Guidelines for the Management of Diabetes in Low- and Middle-Income Versus High-Income
57 Countries-A Systematic Review. *Diabetes care*. 2018;41(5):1097-105.
- 58 132. Haque M, McKimm J, Godman B, Abu Bakar M, Sartelli M. Initiatives to reduce postoperative
59 surgical site infections of the head and neck cancer surgery with a special emphasis on developing
60 countries. *Expert review of anticancer therapy*. 2019;19(1):81-92.
133. Godman B, Wettermark B, van Woerkom M, Fraeyman J, Alvarez-Madrado S, Berg C, et al.
Multiple policies to enhance prescribing efficiency for established medicines in Europe with a

- particular focus on demand-side measures: findings and future implications. *Frontiers in pharmacology*. 2014;5:106.
134. Godman B, Malmstrom RE, Diogene E, Gray A, Jayathissa S, Timoney A, et al. Are new models needed to optimize the utilization of new medicines to sustain healthcare systems? *Expert review of clinical pharmacology*. 2015;8(1):77-94.
135. Godman B, Bucsecs A, Vella Bonanno P, Oortwijn W, Rothe CC, Ferrario A, et al. Barriers for Access to New Medicines: Searching for the Balance Between Rising Costs and Limited Budgets. *Front Public Health*. 2018;6:328.
136. Godman B, Malmstrom RE, Diogene E, Jayathissa S, McTaggart S, Cars T, et al. Dabigatran - a continuing exemplar case history demonstrating the need for comprehensive models to optimize the utilization of new drugs. *Frontiers in pharmacology*. 2014;5:109.
137. Ermisch M, Bucsecs A, Vella Bonanno P, Arickx F, Bybau A, Bochenek T, et al. Payers' Views of the Changes Arising through the Possible Adoption of Adaptive Pathways. *Frontiers in pharmacology*. 2016;7:305.
138. Campbell SM, Godman B, Diogene E, Furst J, Gustafsson LL, MacBride-Stewart S, et al. Quality indicators as a tool in improving the introduction of new medicines. *Basic & clinical pharmacology & toxicology*. 2015;116(2):146-57.
139. Godman B, Shrank W, Andersen M, Berg C, Bishop I, Burkhardt T, et al. Policies to enhance prescribing efficiency in Europe: findings and future implications. *Frontiers in pharmacology*. 2010;1:141.
140. Moorkens E, Vulto AG, Huys I, Dylst P, Godman B, Keuerleber S, et al. Policies for biosimilar uptake in Europe: An overview. *PLoS one*. 2017;12(12):e0190147.
141. Bochenek T, Abilova V, Alkan A, Asanin B, de Miguel Beriain I, Besovic Z, et al. Systemic Measures and Legislative and Organizational Frameworks Aimed at Preventing or Mitigating Drug Shortages in 28 European and Western Asian Countries. *Frontiers in pharmacology*. 2017;8:942.
142. Vella Bonanno P, Bucsecs A, Simoens S, Martin AP, Oortwijn W, Gulbinovic J, et al. Proposal for a regulation on health technology assessment in Europe - opinions of policy makers, payers and academics from the field of HTA. *Expert review of pharmacoeconomics & outcomes research*. 2019;19(3):251-61.
143. Godman B, Haque M, McKimm J, Abu Bakar M, Sneddon J, Wale J, et al. Ongoing strategies to improve the management of upper respiratory tract infections and reduce inappropriate antibiotic use particularly among lower and middle-income countries: findings and implications for the future. *Current medical research and opinion*. 2019:1-27.
144. Godman B, Grobler C, Van-De-Lisle M, Wale J, Barbosa WB, Masele A, et al. Pharmacotherapeutic interventions for bipolar disorder type II: addressing multiple symptoms and approaches with a particular emphasis on strategies in lower and middle-income countries. *Expert opinion on pharmacotherapy*. 2019;20(18):2237-55.
145. World Health Organisation. Definition and diagnosis of diabetes mellitus and intermediate hyperglycemia: report of a WHO/IDF consultation. Available at URL: https://apps.who.int/iris/bitstream/handle/10665/43588/9241594934_eng.pdf;jsessionid=75A0C3B74F6572D012AB00D9F8A23D4E?sequence=1.
146. WHO. Guidelines on second-and third-line medicines and type of insulin for the control of blood glucose levels in non-pregnant adults with diabetes mellitus. 2018. Available at URL: <https://apps.who.int/iris/bitstream/handle/10665/272433/9789241550284-eng.pdf?ua=1>.
147. Beran D, Ewen M, Lipska K, Hirsch IB, Yudkin JS. Availability and Affordability of Essential Medicines: Implications for Global Diabetes Treatment. *Current diabetes reports*. 2018;18(8):48.
148. Roth GA, Johnson C, Abajobir A, Abd-Allah F, Abera SF, Abyu G, et al. Global, Regional, and National Burden of Cardiovascular Diseases for 10 Causes, 1990 to 2015. *Journal of the American College of Cardiology*. 2017;70(1):1-25.
149. Joseph P, Leong D, McKee M, Anand SS, Schwalm JD, Teo K, et al. Reducing the Global Burden of Cardiovascular Disease, Part 1: The Epidemiology and Risk Factors. *Circulation research*. 2017;121(6):677-94.
150. Measuring progress and projecting attainment on the basis of past trends of the health-related Sustainable Development Goals in 188 countries: an analysis from the Global Burden of Disease Study 2016. *Lancet*. 2017;390(10100):1423-59.
151. UN. Sustainable Development Goal 3 - Ensure healthy lives and promote well-being for all at all ages. 2019. Available at URL: <https://sustainabledevelopment.un.org/sdg3>.
152. Morton S, Pencheon D, Squires N. Sustainable Development Goals (SDGs), and their implementation: A national global framework for health, development and equity needs a systems approach at every level. *British medical bulletin*. 2017;124(1):81-90.

- 1
2
3 153. Caires de Souza AL, de Assis Acurcio F, Guerra Junior AA, Rezende Macedo do Nascimento
4 RC, Godman B, Diniz LM. Insulin glargine in a Brazilian state: should the government disinvest? An
5 assessment based on a systematic review. *Applied health economics and health policy*.
6 2014;12(1):19-32.
- 7 154. de Souza AL, Acurcio Fde A, Guerra Junior AA, do Nascimento RC, Godman B, Diniz LM.
8 Authors' reply to Dr. Malerbi: "Insulin glargine in a Brazilian state: should the government disinvest?".
9 *Applied health economics and health policy*. 2014;12(5):561-3.
- 10 155. Almeida P, Silva TBC, de Assis Acurcio F, Guerra Junior AA, Araujo VE, Diniz LM, et al.
11 Quality of Life of Patients with Type 1 Diabetes Mellitus Using Insulin Analog Glargine Compared with
12 NPH Insulin: A Systematic Review and Policy Implications. *The patient*. 2018;11(4):377-89.
- 13 156. Marra LP, Araujo VE, Silva TB, Diniz LM, Guerra Junior AA, Acurcio FA, et al. Clinical
14 Effectiveness and Safety of Analog Glargine in Type 1 Diabetes: A Systematic Review and Meta-
15 Analysis. *Diabetes therapy*. 2016;7(2):241-58.
- 16 157. Institut für Qualität und Wirtschaftlichkeit im Gesundheitswesen [Institute of Quality and
17 Efficiency in Health Care]: Long-acting insulin analogues in the treatment of diabetes mellitus type 1.
18 2010. Available at URL: [Http://www.iqwig.de/download/A05-01_Executive-Summary_Long-](http://www.iqwig.de/download/A05-01_Executive-Summary_Long-acting_insulin_analogues_in_diabetes_mellitus_type_1.pdf)
19 [acting_insulin_analogues_in_diabetes_mellitus_type_1.pdf](http://www.iqwig.de/download/A05-01_Executive-Summary_Long-acting_insulin_analogues_in_diabetes_mellitus_type_1.pdf)
- 20 158. Holden SE, Poole CD, Morgan CL, Currie CJ. Evaluation of the incremental cost to the
21 National Health Service of prescribing analogue insulin. *BMJ open*. 2011;1(2):e000258.
- 22 159. Horvath K, Jeitler K, Berghold A, Ebrahim SH, Gratzner TW, Plank J, et al. Long-acting insulin
23 analogues versus NPH insulin (human isophane insulin) for type 2 diabetes mellitus. *The Cochrane*
24 *database of systematic reviews*. 2007(2):Cd005613.
- 25 160. Singh SR, Ahmad F, Lal A, Yu C, Bai Z, Bennett H. Efficacy and safety of insulin analogues
26 for the management of diabetes mellitus: a meta-analysis. *CMAJ*. 2009;180(4):385-97.
- 27 161. Monami M, Marchionni N, Mannucci E. Long-acting insulin analogues vs. NPH human insulin
28 in type 1 diabetes. A meta-analysis. *Diabetes, obesity & metabolism*. 2009;11(4):372-8.
- 29 162. NICE. Type 1 diabetes in adults: diagnosis and management NICE guideline [NG17]. 2016.
30 Available at URL: <https://www.nice.org.uk/guidance/ng17>.
- 31 163. Moodie P. More from PHARMAC on long-acting insulin analogues: insulin glargine now
32 funded. *The New Zealand medical journal*. 2006;119(1236):U2040-U.
- 33 164. International Diabetes Federation. IDF Ninth Edition - Country Summary Table. 2019.
34 Available at URL: <https://diabetesatlas.org/en/resources/>.
- 35 165. Jasem D, Majaliwa ES, Ramaiya K, Najem S, Swai ABM, Ludvigsson J. Incidence,
36 prevalence and clinical manifestations at onset of juvenile diabetes in Tanzania. *Diabetes research*
37 *and clinical practice*. 2019;156:107817.
- 38 166. International Diabetes Federation. IDF Africa Members - Ghana. 2017. Available at URL:
39 <https://idf.org/our-network/regions-members/africa/members/11-ghana.html>.
- 40 167. IDF. Middle East and North Africa. 2019. Available at URL: [https://idf.org/our-network/regions-](https://idf.org/our-network/regions-members/middle-east-and-north-africa/members/41-morocco.html)
41 [members/middle-east-and-north-africa/members/41-morocco.html](https://idf.org/our-network/regions-members/middle-east-and-north-africa/members/41-morocco.html).
- 42 168. Dinar Y, Belahsen R. Diabetes Mellitus in Morocco: Situation and Challenges of Diabetes
43 Care. *Journal of Scientific Research & Reports*. 2014; 3(19): 2477-2485.
- 44 169. Chetoui A, Kaoutar K, Kardoudi A, Boutahar K, Chigr F, Najimi M. Epidemiology of diabetes in
45 Morocco: review of data, analysis and perspectives. *International Journal of Scientific & Engineering*
46 *Research* 2018; 9 (12): 1310 - 16.
- 47 170. Adekanmbi VT, Uthman OA, Erqou S, Echouffo-Tcheugui JB, Harhay MN, Harhay MO.
48 Epidemiology of prediabetes and diabetes in Namibia, Africa: A multilevel analysis. *Journal of*
49 *diabetes*. 2019;11(2):161-72.
- 50 171. Ugege O, Ibitoye PK, Jiya NM. Childhood diabetes mellitus in sokoto, north-western Nigeria:
51 A ten year review. *Sahel Med J*. 2013;16:97-101.
- 52 172. Umar UI. Pattern of presentation of Type 1 diabetic patients in Kano, Nigeria. *Niger J Basic*
53 *Clin Sci* 2016;13:85-8.
- 54 173. John C, Abok II, Yilgwan C. Clinical profile of childhood type 1 diabetes in Jos, Nigeria.
55 *African Journal of Diabetes Medicine*. 2013; 21 (1): 11-13.
- 56 174. Nsabimana J de Dieu. Diabetes prevalence in Rwanda at 3% – MoH. 2019. Available at URL:
57 <https://www.newtimes.co.rw/news/diabetes-prevalence-rwanda-3-moh>.
- 58 175. Ekpebegh CO, Longo-Mbenza B. Clinical, immunologic and insulin secretory characteristics
59 of young black South African patients with diabetes: Hospital based single centre study. *Diabetes*
60 *research and clinical practice*. 2013;99(3):380-4.
176. Elamin A, Ghalib M, Eltayeb B, Tuvemo T. High incidence of type 1 diabetes mellitus in
Sudanese children, 1991-1995. *Annals of Saudi medicine*. 1997;17(4):478-80.

177. World Health Organisation. Country Profile - Zambia. 2019. Available at URL: https://www.who.int/diabetes/country-profiles/zmb_en.pdf?ua=1.
178. Mutowo M, Gowda U, Mangwiro JC, Lorgelly P, Owen A, Renzaho A. Prevalence of diabetes in Zimbabwe: a systematic review with meta-analysis. *International journal of public health*. 2015;60(1):1-11.
179. Ogle GD, Middlehurst AC, Silink M. The IDF Life for a Child Program Index of diabetes care for children and youth. *Pediatric diabetes*. 2016;17(5):374-84.
180. Zhang P, Zhang X, Brown J, Vistisen D, Sicree R, Shaw J, et al. Global healthcare expenditure on diabetes for 2010 and 2030. *Diabetes research and clinical practice*. 2010;87(3):293-301.
181. Pastakia SD, Nuche-Berenguer B, Pekny CR, Njuguna B, O'Hara EG, Cheng SY, et al. Retrospective assessment of the quality of diabetes care in a rural diabetes clinic in Western Kenya. *BMC endocrine disorders*. 2018;18(1):97-.
182. Matheka DM, Kilonzo JM, Munguti CM, Mwangi PW. Pattern, knowledge and practices of HbA1C testing among diabetic patients in a Kenyan tertiary referral hospital. *Globalization and health*. 2013;9:55-.
183. Fadare J, Olamoyegun M, Gbadegesin BA. Medication adherence and direct treatment cost among diabetes patients attending a tertiary healthcare facility in Ogbomoso, Nigeria. *Malawi Med J*. 2015;27(2):65-70.
184. Ogbera AO, Kuku SF. Insulin use, prescription patterns, regimens and costs.-a narrative from a developing country. *Diabetology & metabolic syndrome*. 2012;4(1):50-.
185. Onyiriuka N, Ezomo OO, Onyiriuka RC. Cost of treating Insulin-requiring diabetes in children and adolescents. *Journal of Institute of Medicine*. 2012;34(1):2-8.
186. Yamey G, Fewer S, Beyeler N. Achieving a "Grand Convergence" in Global Health by 2035: Rwanda Shows the Way Comment on "Improving the World's Health Through the Post-2015 Development Agenda: Perspectives From Rwanda". *International journal of health policy and management*. 2015;4(11):789-91.
187. Rwanda Diabetes Association. 2019. Available at URL: <https://rwandadiabetes.com/>.
188. Global Fund Grants in the Republic of Sudan - GF-OIG-19-010. Geneva, Switzerland. 2019. Available from URL: https://www.theglobalfund.org/media/8433/oig_gf-oig-19-010_report_en.pdf?u=637066540330000000.
189. Mohamed Ali, GK. 3,121 days in the driving seat of the National Medical Supplies Fund: What lessons to be shared? Book manuscript under proofreading for publication.
190. Elrayah H, Eltom M, Bedri A, Belal A, Rosling H, Ostenson C-G. Economic burden on families of childhood type 1 diabetes in urban Sudan. *Diabetes research and clinical practice*. 2005;70(2):159-65.
191. Eliadarous H. Exploring the impact of diabetes in Sudan: Out-of-pocket expenditure and social consequences of diabetes on patients and their families. Stockholm, Sweden. 2017. Available at URL: https://openarchive.ki.se/xmlui/bitstream/handle/10616/45956/Thesis_Hind_Eliadarous.pdf?sequence=1&isAllowed=y.
192. Kalungia CA, Mwale M, Sondashi IS, Mweetwa B, Yassa P, Kadimba G. Availability of Essential Antihypertensive and Antidiabetic Medicines in Public Health Facilities in Lusaka District, Zambia. *Medical Journal of Zambia* 2017; 44 (3): 140-8.
193. Kaiser AH, Hehman L, Forsberg BC, Simangolwa WM, Sundewall J. Availability, prices and affordability of essential medicines for treatment of diabetes and hypertension in private pharmacies in Zambia. *PloS one*. 2019;14(12):e0226169-e.
194. Ghana National Health Insurance Medicine list 2018. Available at URL: <http://www.nhis.gov.gh/files/2018%20NHIS%20ML.pdf>.
195. Kalk WJ, Raal J, Joffe BI. The prevalence and incidence of and risk factors for, micro-albuminuria among urban Africans with type 1 diabetes in South Africa: An inter-ethnic study. *International Journal of Diabetes Mellitus*. 2010; 2; 148–153.
196. Diabetes South Africa. The Tshwane Insulin Project. 2018. Available at URL: <https://www.diabetessa.org.za/the-tshwane-insulin-project/>.
197. Ntuli C. Tuks sets up diabetes treatment programme. 2019. Available at URL: <https://www.iol.co.za/pretoria-news/tuks-sets-up-diabetes-treatment-programme-37175398>.
198. Federal Ministry of Health, Sudan. National Essential Medicines List, 2014. Available from URL: https://www.humanitarianresponse.info/sites/www.humanitarianresponse.info/files/documents/files/sudan_national_essential_medicines_list_.pdf.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

199. Jonker D, Deacon E, van Rensburg E, Segal D. Illness perception of adolescents with well-controlled type 1 diabetes mellitus. *Health Psychol Open*. 2018;5(2):2055102918799968-.

For Review Only