

# Limit theorems for continuous-state branching processes with immigration

Clément Foucart\* & Chunhua Ma † & Linglong Yuan‡

October 1, 2020

## Abstract

We prove and extend some results stated by Mark Pinsky in *Limit theorems for continuous state branching processes with immigration* [Bull. Amer. Math. Soc. **78** (1972), 242–244]. Consider a continuous-state branching process with immigration  $(Y_t, t \geq 0)$  with branching mechanism  $\Psi$  and immigration mechanism  $\Phi$  (CBI $(\Psi, \Phi)$  for short). We shed some light on two different asymptotic regimes occurring when  $\int_0 \frac{\Phi(u)}{|\Psi(u)|} du < \infty$  or  $\int_0 \frac{\Phi(u)}{|\Psi(u)|} du = \infty$ . We first observe that when  $\int_0 \frac{\Phi(u)}{|\Psi(u)|} du < \infty$ , supercritical CBIs have a growth rate dictated by the branching dynamics, namely there is a renormalization  $\tau(t)$ , only depending on  $\Psi$ , such that  $(\tau(t)Y_t, t \geq 0)$  converges almost-surely to a finite random variable. When  $\int_0 \frac{\Phi(u)}{|\Psi(u)|} du = \infty$ , it is shown that the immigration overwhelms the branching dynamics and that no linear renormalization of the process can exist. Asymptotics in the second regime are studied in details for all non-critical CBI processes via a non-linear time-dependent renormalization in law. Three regimes of weak convergence are then exhibited, and a misprint in Pinsky’s paper is corrected. CBI processes with critical branching mechanisms subject to a regular variation assumption are also studied.

**Key words.** Continuous-state branching processes, Immigration, Grey martingale, Limit distribution, Non-linear renormalization, Regularly varying functions.

*MSC (2010):* primary 60J80; secondary 60G70; 60G55

## 1 Introduction

Continuous-state branching processes with immigration (CBI for short) have been defined by Kawazu and Watanabe [KW71]. They are scaling limits of Galton-Watson Markov chains with immigration, see e.g. [KW71, Theorem 2.2]. Recent years have seen renewed interest in this class of Markov processes. They appear for instance as strong solutions of some stochastic differential equations with jumps, see Dawson and Li [DL12], and in a more applied point of view, form an important subclass of the so-called affine processes, which are known in the financial mathematics setting for modelling interest rates, see [DFS03].

---

\*LAGA Université Sorbonne Paris Nord E-mail: foucart@math.univ-paris13.fr

†School of Mathematical Sciences and LPMC, Nankai University E-mail: mach@nankai.edu.cn

‡Department of Mathematical Sciences, University of Liverpool; Department of Mathematical Sciences, Xi’an Jiaotong-Liverpool University E-mail: yuanlinglongcn@gmail.com

We mention for instance the works in this direction of Jiao et al. [JMS17] and Barczy et al. [BAKP18] where certain CBI processes are studied from a statistical point of view.

The asymptotic behaviors of Galton-Watson processes with immigration have been extensively studied since the seventies. We refer to the works of Cohn [Coh77], Heathcote [Hea65], Heyde [Hey70], Pakes [Pak79] and Seneta [Sen70] and to their references. Transience and Recurrence of CBIs have been characterized by Duhalde et al. Fine properties of the stationary distributions of CBI processes, when they exist, have also been recently established in Chazal et al. [CLP18] and Keller-Ressel and Mijatovic [KRM12]. In the case where no stationary distribution exists, less attention has been paid to the limit theorems of CBI processes. It will certainly not be surprising that the results found in the seventies for Galton-Watson processes with immigration, have counterparts in the continuous-state and continuous-time framework. A year after Kawazu and Watanabe's founding work, Pinsky thus published a short note [Pin72] without proof, on the limits of CBIs. We believe however of interest to write down some details and resume in this article the study of limit theorems for CBIs initiated by Pinsky.

We start by proving an almost-sure convergence for CBI processes by adapting Grey's approach [Gre74] to the framework with immigration (Theorem 2). We then provide a general non-linear renormalization in law (Theorem 6). To the best of our knowledge this latter renormalization does not appear in the literature about Galton-Watson processes with immigration. We explain now our main results. Denote respectively by  $\Psi$  and  $\Phi$  the branching and immigration mechanisms, we will recall their definitions in the next section. In the case of supercritical branching, we show the existence of two distinct almost-sure asymptotic regimes according to the convergence/divergence of the integral  $\int_0 \frac{\Phi(u)}{|\Psi(u)|} du$ .

When this integral converges, i.e  $\int_0 \frac{\Phi(u)}{|\Psi(u)|} du < \infty$ , the branching dynamics takes precedence over immigration and directs the divergence of the process towards infinity. More precisely, under the classical  $L \ln L$  moment assumption (also called Kesten-Stigum condition) over the branching Lévy measure, the CBI process grows at the same exponential rate as the pure branching process. On the other hand, when it diverges, i.e  $\int_0 \frac{\Phi(u)}{|\Psi(u)|} du = \infty$ , immigration is so substantial that the branching, although supercritical, is somehow overtaken. So that, typically the process grows faster than the pure branching process on its event of non-extinction.

A similar dichotomy occurs more generally for non-critical CBI processes when we consider their longterm behaviour in law. Our main contribution is to design a non-linear time-dependent renormalization in law of non-critical CBI( $\Psi, \Phi$ )-processes satisfying  $\int_0 \frac{\Phi(u)}{|\Psi(u)|} du = \infty$ . We shall find, see Theorem 6, a deterministic function  $\lambda \mapsto r_t(\lambda)$  only depending on  $\Psi$  and  $\Phi$  such that

$$r_t(1/Y_t) \xrightarrow[t \rightarrow \infty]{} e_1 \text{ in law} \quad (1.1)$$

where  $e_1$  is a standard exponential random variable. The latter renormalization is actually equivalent to the following property, see Corollary 1. Given two independent CBI( $\Psi, \Phi$ ) processes  $(Y_t, t \geq 0)$  and  $(\tilde{Y}_t, t \geq 0)$  such that  $\int_0 \frac{\Phi(u)}{|\Psi(u)|} du = \infty$ , one has

$$\frac{Y_t}{\tilde{Y}_t} \xrightarrow[t \rightarrow \infty]{} \Lambda \text{ in law} \quad (1.2)$$

where  $\mathbb{P}(\Lambda = 0) = \mathbb{P}(\Lambda = \infty) = \frac{1}{2}$ . As a consequence of (1.2), we shall see that no function  $(\eta(t), t \geq 0)$  exists such that  $\eta(t)Y_t$  converges in law towards a nondegenerate random variable (i.e whose support is not contained in  $\{0, \infty\}$ ). This has been shown by Cohn [Coh77] in the setting of discrete time and space. The meaning of the limit in law (1.1)

will be made more explicit by introducing further assumptions on the rate of divergence of the integral  $\int_0^{\frac{\Phi(u)}{|\Psi(u)|}} du$ , namely on the speed at which  $\int_{\varepsilon}^{\frac{\Phi(u)}{|\Psi(u)|}} du$  goes to  $\infty$  as  $\varepsilon$  goes to 0. In the same vein as Pakes, see [Pak79], we design three regimes of divergence: Slow (S), Log (L) and Fast (F). Each corresponds to a specific renormalization and a specific limiting law. The faster the integral diverges, the more the branching dynamics is overtaken by immigration. This is reflected by the different renormalizations occurring in the three regimes. In particular, in the fast case (F) the branching mechanism plays no role in the renormalization. Pinsky's result [Pin72, Theorem 2] which corresponds to the subcritical case under condition (S) has now a proof, see Remark 14–ii), and a misprint in his statement is corrected.

**Notation:** We denote respectively by  $\xrightarrow{d}$  and  $\xrightarrow{p}$  the convergence in law and the convergence in probability. We use the relation symbol  $\sim$  when the ratio of the two terms on the two sides of it converges to 1 (if any of the two terms is random, the convergence holds almost surely). The probability measure and its expectation are denoted by  $\mathbb{P}$  and  $\mathbb{E}$ . For any  $x \geq 0$ ,  $\mathbb{P}_x$  denotes the law of a CBI process started from  $x$ . The integrability of a function  $f$  in a neighbourhood of 0 is denoted by  $\int_0 f(x) dx < \infty$  (similarly  $\int^{\infty} f(x) dx < \infty$  denotes the integrability of  $f$  in a neighbourhood of  $\infty$ ). Last, we denote functions, either deterministic or random, vanishing in the limit by  $o(1)$ .

The paper is organized as follows. First we recall in Section 2 the definition of a CBI process and some of its most fundamental properties. Our main results are stated in Section 3. We first establish in Section 3.1 the almost-sure convergence results in the supercritical case. Then, Section 3.2 is devoted to the study of convergence in law in the non-critical case when  $\int_0^{\frac{\Phi(u)}{|\Psi(u)|}} du = \infty$ . We define the three regimes (S), (L), (F) in Section 3.3. The last section treats some critical branching mechanisms fulfilling certain regular variation properties.

## 2 Preliminaries

We recall hereafter the definition of a CBI process and some of its most fundamental properties. Our main references are Chapter 3 of Li's book [Li11] and Chapter 12 of Kyprianou's book [Kyp14]. We recall that a random variable is nondegenerate if its support is not contained in  $\{0, \infty\}$ . We say a random variable is proper if it is finite almost surely.

Write  $\pi$  and  $\nu$  for two  $\sigma$ -finite nonnegative measures on  $(0, \infty)$  satisfying respectively  $\int_0^{\infty} (z \wedge z^2) \pi(dz) < \infty$  and  $\int_0^{\infty} (1 \wedge z) \pi(dz) < \infty$ . Consider a triple  $(\sigma, b, \beta)$  such that  $\sigma \geq 0, b \in \mathbb{R}$  and  $\beta \geq 0$ . Let  $\Psi$  be the Laplace exponent of a spectrally positive Lévy process with finite mean<sup>1</sup> and whose characteristic triple is  $(b, \sigma, \pi)$ . Let  $\Phi$  be the Laplace exponent of a subordinator with drift  $\beta$  and Lévy measure  $\nu$ . They are specified by the Lévy-Khinchine formula

$$\Psi(q) = bq + \frac{1}{2}\sigma^2 q^2 + \int_0^{\infty} (e^{-qu} - 1 + qu)\pi(du), \quad q \geq 0,$$

So  $\Psi$  is convex (i.e.,  $\Psi''(q) \geq 0, \forall q \geq 0$ ) with  $\Psi(0) = 0$ . Similarly,

$$\Phi(q) = \beta q + \int_0^{\infty} (1 - e^{-qu})\nu(du), \quad q \geq 0,$$

So  $\Phi$  is a concave continuous, strictly increasing function with  $\Phi(0) = 0$ .

<sup>1</sup>We assume  $|\Psi'(0+)| < \infty$ , so that in particular the CBI process does not explode.

A CBI process with branching and immigration mechanisms  $\Psi$  and  $\Phi$ , is a strong Markov process  $(Y_t, t \geq 0)$  taking values in  $[0, \infty)$  whose transition kernels are characterized by their Laplace transforms. So for  $\lambda \geq 0$  and  $x \in \mathbb{R}_+$ ,

$$\mathbb{E}_x[e^{-\lambda Y_t}] = \exp\left(-xv_t(\lambda) - \int_0^t \Phi(v_s(\lambda))ds\right), \quad (2.3)$$

where the map  $t \mapsto v_t(\lambda)$  is the solution to the differential equation

$$\frac{\partial}{\partial t}v_t(\lambda) = -\Psi(v_t(\lambda)), \quad v_0(\lambda) = \lambda. \quad (2.4)$$

Note that  $v_{t+s}(\lambda) = v_t(v_s(\lambda))$  from the Markov property. The above display implies, by a change of variable that

$$r_t(\lambda) := \int_0^t \Phi(v_s(\lambda))ds = \int_{v_t(\lambda)}^\lambda \frac{\Phi(u)}{\Psi(u)}du. \quad (2.5)$$

Then (2.3) can also be written as

$$\mathbb{E}_x[e^{-\lambda Y_t}] = \exp(-xv_t(\lambda) - r_t(\lambda)). \quad (2.6)$$

Note also that for any  $t \geq 0$  and any  $n \in \mathbb{N}$ ,  $Y_t = Y_t^1 + \dots + Y_t^n$  in law where  $((Y_t^i)_{t \geq 0}, 1 \leq i \leq n)$  are i.i.d copies of a CBI  $(\Psi, \frac{1}{n}\Phi)$  process. So that in particular,  $Y_t$  has an infinite divisible law on  $\mathbb{R}_+$  and  $\lambda \mapsto r_t(\lambda)$  is the Laplace exponent of a subordinator (with no killing term). For any  $t \geq 0$ , we set  $r_t(\infty) = \int_{\bar{v}_t}^\infty \frac{\Phi(u)}{\Psi(u)}du \in [0, \infty]$ , where  $\bar{v}_t := \lim_{\lambda \rightarrow \infty} \uparrow v_t(\lambda)$ , with the convention that if  $\int^\infty \frac{\Phi(u)}{\Psi(u)}du = \infty$  then  $r_t(\infty) = \infty$  for all  $t > 0$ . From (2.5), we easily check that  $r_t(\infty) < \infty$  as soon as  $\int^\infty \frac{\Phi(u)}{\Psi(u)}du < \infty$ . Letting  $\lambda$  tend to  $\infty$  in (2.6) readily entails that  $r_t(\infty) < \infty$  if and only if  $\mathbb{P}_x(Y_t = 0) > 0$ . We refer the reader interested in the zero-set of CBIs to [FUB14].

Existence and unicity of CBI processes have been established in [KW71, Theorem 1.1]. Recently Dawson and Li [DL12], see also [?], have shown that any CBI is the strong solution of a certain stochastic differential equation (SDE) with jumps.

Suppose that  $(\Omega, \mathcal{F}_t, \mathbb{P})$  is a filtered probability space satisfying the usual hypotheses. Let  $\{B_t\}_{t \geq 0}$  be an  $(\mathcal{F}_t)$ -Brownian motion. Let  $N_0(ds, dz, du)$  and  $N_1(ds, du)$  denote two  $(\mathcal{F}_t)$ -Poisson random measures on  $(0, \infty)^3$  and  $(0, \infty)^2$  with intensities  $ds\pi(dz)du$  and  $ds\nu(dz)$ . We assume that the Brownian motion and the Poisson random measures are independent of each other. Let  $\tilde{N}_0(ds, dz, du)$  be the corresponding compensated measure of  $N_0$ , namely  $\tilde{N}_0(ds, dz, du) := N_0(ds, dz, du) - ds\pi(dz)du$ . The following SDE

$$Y_t = Y_0 + \sigma \int_0^t \sqrt{Y_s} dB_s + \int_0^t \int_0^\infty \int_0^{Y_{s-}} z \tilde{N}_0(ds, dz, du) + \int_0^t (\beta - bY_s) ds + \int_0^t \int_0^\infty z N_1(ds, dz). \quad (2.7)$$

admits a unique strong solution whose law is that of a CBI with branching mechanism  $\Psi$  and immigration mechanism  $\Phi$ . When there is no immigration, that is to say  $\Phi \equiv 0$ , the drift  $\beta$  and the Poisson random measure  $N_1$  vanish and the process  $(Y_t, t \geq 0)$  solution to (2.7) is a continuous-state branching process (CB process for short) with branching mechanism  $\Psi$ . When  $\Psi \equiv 0$ , only the immigration part remains and  $(Y_t, t \geq 0)$  is a subordinator with Laplace exponent  $\Phi$ . Lastly, we recall that a (sub)-critical CB  $(\Psi)$  process conditioned on the non-extinction is a CBI  $(\Psi, \Phi)$  process with  $\Phi = \Psi' - \Psi'(0+)$ , see Lambert [Lam07], Li [Li00, Theorem 4.1] and Fittipaldi and Fontbona [FF12].

Recall the form of the function  $\Psi$  and notice that  $b = \Psi'(0+)$ . A CBI process is said to be critical, subcritical or supercritical according as  $b = 0$ ,  $b > 0$  or  $b < 0$ . Note that  $\Psi$  has at most two roots. Introduce

$$\rho = \inf\{z > 0, \Psi(z) \geq 0\}, \quad \inf \emptyset = \infty.$$

We see that  $\rho = 0$  if  $b \geq 0$  and  $\rho > 0$  if  $b < 0$ . In particular  $\rho = \infty$  if and only if  $-\Psi$  is the Laplace exponent of a subordinator. By (2.4), if  $0 < \lambda < \rho$  (resp.  $\lambda > \rho$ ), then  $v_t(\lambda) \in [\lambda, \rho]$  is increasing (resp.  $v_t(\lambda) \in [\rho, \lambda]$  is decreasing) in  $t$ . Then (2.4) implies

$$\int_{v_t(\lambda)}^{\lambda} \frac{dz}{\Psi(z)} = t, \quad \forall t \in [0, \infty), \quad \forall \lambda \in (0, \infty) \setminus \{\rho\}, \quad (2.8)$$

Recall  $\bar{v}_t := \lim_{\lambda \rightarrow \infty} \uparrow v_t(\lambda) \in [0, \infty]$  and set  $\bar{v} := \lim_{t \rightarrow \infty} \downarrow \bar{v}_t \in [0, \infty]$ . Grey shows in [Gre74] that

$$\bar{v}_t < \infty \text{ for all } t > 0 \text{ if and only if } \int^{\infty} \frac{dq}{\Psi(q)} < \infty \text{ (Grey's condition)}. \quad (2.9)$$

Note that  $\rho \leq \bar{v}$ , and if  $\bar{v} < \infty$  then  $\bar{v} = \rho$ . We refer to Section 3.2 of [Li11] for proofs of the following technical statements; see also [Gre74]. We gather in the next lemma analytical results on the map  $\lambda \mapsto v_t(\lambda)$  and its inverse (whenever it exists).

**Lemma 1.** *The map  $\lambda \mapsto v_t(\lambda)$  is strictly increasing on  $[0, \infty)$ . For any  $t \geq 0$ , let  $\lambda \mapsto v_{-t}(\lambda)$  be the inverse map of  $\lambda \mapsto v_t(\lambda)$ . This is a strictly increasing function, well-defined on  $[0, \bar{v}_t)$  which satisfies for all  $s, t \geq 0$  and  $0 \leq \lambda < \bar{v}_{s+t}$*

$$v_{-(s+t)}(\lambda) = v_{-s}(v_{-t}(\lambda)).$$

For  $0 \leq \lambda < \bar{v}_t$ , such that  $\Psi(\lambda) \neq 0$ , by (2.8) one has

$$\int_{\lambda}^{v_{-t}(\lambda)} \frac{dz}{\Psi(z)} = \int_{v_t(v_{-t}(\lambda))}^{v_{-t}(\lambda)} \frac{dz}{\Psi(z)} = t. \quad (2.10)$$

In particular, in the supercritical case, i.e.  $b \in (-\infty, 0)$ , for  $\lambda \in (0, \rho)$

$$\frac{\partial v_{-t}(\lambda)}{\partial t} = \Psi(v_{-t}(\lambda)), \quad v_0(\lambda) = \lambda. \quad (2.11)$$

The map  $t \mapsto v_{-t}(\lambda)$  is decreasing and by letting  $t \rightarrow \infty$  in (2.10) we see that  $v_{-t}(\lambda) \xrightarrow[t \rightarrow \infty]{} 0$ .

Moreover,  $v_{-(t+u)}(\lambda)/v_{-t}(\lambda) \xrightarrow[t \rightarrow \infty]{} e^{bu}$  for any  $u \geq 0$ .

The following theorem has been announced by Pinsky [Pin72] and provides some first information on the growth rate. It has been established in the (sub)critical case by Li, see [Li11, Theorem 3.20, P.66] and Keller-Ressel and Mijatović see [KRM12, Appendix].

**Theorem 1** (Pinsky [Pin72], Li [Li11]). *Let  $(Y_t, t \geq 0)$  be a CBI process with  $|\Psi'(0+)| < \infty$ . Set  $\tau(t) = e^{bt}$  if  $b < 0$  and  $\tau(t) = 1$  if  $b \geq 0$ . The process  $(\tau(t)Y_t, t \geq 0)$  converges in law, as  $t \rightarrow \infty$ , towards a proper random variable if and only if*

$$\int_0^{\infty} \frac{\Phi(u)}{|\Psi(u)|} du < \infty. \quad (2.12)$$

If  $\int_0^{\infty} \frac{\Phi(u)}{|\Psi(u)|} du = \infty$ , then for all  $z \geq 0$ ,  $\mathbb{P}_x(\tau(t)Y_t \leq z) \xrightarrow[t \rightarrow \infty]{} 0$ , that is to say  $(\tau(t)Y_t, t \geq 0)$  converges to  $\infty$  in probability.

Our paper aims to enrich the above Theorem 1 by studying almost-sure limits in the supercritical case and finding new results on the growth rates when  $\Psi$  is non-critical and  $\int_0^{\frac{\Phi(u)}{|\Psi(u)|}} du = \infty$ . It is clear that a supercritical CBI process  $(Y_t, t \geq 0)$  is transient i.e.  $Y_t \xrightarrow[t \rightarrow \infty]{} \infty$  a.s. The properties of transience and recurrence for subcritical and critical CBI processes have been studied in Duhalde et al. [DFM14]. It is established in [DFM14, Theorem 3] that a (sub)critical CBI( $\Psi, \Phi$ ) process is recurrent or transient according as

$$\mathcal{E} := \int_0 \frac{dx}{\Psi(x)} \exp\left(-\int_x^1 \frac{\Phi(u)}{\Psi(u)} du\right) = \infty \text{ or } < \infty. \quad (2.13)$$

We see that the integral  $\int_0 \frac{\Phi(u)}{|\Psi(u)|} du$  plays a crucial role in this integral test. In particular, it is worth noticing that in the (sub)critical case, the divergence of  $\int_0 \frac{\Phi(u)}{|\Psi(u)|} du$  is necessary for the CBI process to be transient but not sufficient.

The notion of regularly varying functions will be used in several places. Recall that a function  $R$  is regularly varying at  $\infty$  (resp. at 0) with index  $\theta \in \mathbb{R}$  if for any  $\lambda > 0$

$$\frac{R(\lambda x)}{R(x)} \rightarrow \lambda^\theta \text{ as } x \rightarrow \infty \text{ (resp. } 0). \quad (2.14)$$

The function  $R$  is said to be *slowly varying* if  $\theta = 0$  and if  $R$  is regularly varying with index  $\theta$ , then  $R$  has the form  $R(x) = x^\theta L(x)$  for all  $x \geq 0$  with  $L$  a slowly varying function. We stress that those functions occur naturally in the study as for instance, in the supercritical case, i.e.  $b \in (-\infty, 0)$ , Lemma 1 ensures that the function  $t \mapsto v_{-\ln(t)}(\lambda)$  is regularly varying at  $\infty$  with index  $b$ . We refer the reader to Bingham et al. [BGT87] for a reference on those functions.

## 3 Results

### 3.1 Almost-sure limits

This section deals with the so-called Seneta-Heyde norming for CBI processes. We refer to Seneta [Sen70] and Heyde [Hey70] for the seminal papers in the discrete setting; see also Lambert [Lam07]. When no immigration is taken into account, namely  $\Phi \equiv 0$ , this study has been done by Grey [Gre74] and Bingham [Bin76]. We refer the reader for instance to the end of Chapter 12 of Kyprianou's book [Kyp14].

Recall (2.9),  $\bar{v} \in [0, \infty]^2$ ,  $t \mapsto v_{-t}(\lambda)$  and its Equation (2.11).

**Theorem 2.** *Let  $(Y_t, t \geq 0)$  be a CBI( $\Psi, \Phi$ ) with a supercritical branching mechanism  $\Psi$  i.e.  $b < 0$ . Let  $\lambda < \bar{v}$ . Then,*

i) *if  $\int_0 \frac{\Phi(u)}{|\Psi(u)|} du < \infty$  then  $v_{-t}(\lambda)Y_t \xrightarrow[t \rightarrow \infty]{} W^\lambda \mathbb{P}_x$ -a.s., where  $W^\lambda$  is a non-degenerate proper random variable with Laplace exponent*

$$\mathbb{E}_x[e^{-\theta W^\lambda}] = \exp\left(-xv_{\ln(\frac{\theta}{\bar{v}})}(\lambda) + \int_0^{v_{\ln(\frac{\theta}{\bar{v}})}(\lambda)} \frac{\Phi(u)}{\Psi(u)} du\right), \quad (3.15)$$

ii) *if  $\int_0 \frac{\Phi(u)}{|\Psi(u)|} du = \infty$  then  $v_{-t}(\lambda)Y_t \xrightarrow[t \rightarrow \infty]{} \infty \mathbb{P}_x$ -a.s.*

---

<sup>2</sup> $\bar{v}$  is finite if and only if Grey's condition holds.

*Remark 3.* i) Without immigration, namely when  $\Phi \equiv 0$ , Theorem 2-i) matches with [Gre74, Theorem 2] for supercritical CBs. See also Duquesne and Labbé [DL13, Lemma 2.2] for the expression of the Laplace transform (3.15) with  $\Phi \equiv 0$ .

ii) According to Grey [Gre74], if  $\int^\infty (x \ln x) \pi(dx) < \infty$  then  $v_{-t}(\lambda) \underset{t \rightarrow \infty}{\sim} K_\lambda e^{bt}$  for some constant  $K_\lambda > 0$ . So that in this case, as mentioned in the introduction, the CBI process grows almost-surely exponentially fast.

iii) In the non-critical case, the condition  $\int_0^\infty \frac{\Phi(u)}{|\Psi(u)|} du < \infty$  is equivalent to  $\int^\infty \ln(u) \nu(du) = \infty$  where  $\nu$  is the immigration measure  $\nu$ , see the forthcoming Remark 11

We will adapt Grey's martingales, known for CB processes, see [Gre74], to the setting of CBIs. It is worth mentioning that for a supercritical CB process (with finite mean), Grey found the same almost-sure renormalization  $v_{-t}(\lambda)$  on the event of non-extinction.

As explained in the introduction Theorem 2 reflects the fact that two regimes occur according to the convergence/divergence of the integral  $\int_0^\infty \frac{\Phi(u)}{|\Psi(u)|} du$ . In case i), the branching dictates the growth of the process, in case ii) it is overwhelmed by the immigration.

*Proof.* Consider  $(Y_t, t \geq 0)$  a CBI  $(\Psi, \Phi)$ . Recall (2.6). Fix  $\lambda \in (0, \bar{v})$ . We show that the process  $(M_t^\lambda, t \geq 0)$  defined by

$$M_t^\lambda = \exp\left(-v_{-t}(\lambda)Y_t + \int_\lambda^{v_{-t}(\lambda)} \frac{\Phi(u)}{\Psi(u)} du\right)$$

is a positive martingale. For every  $t \geq 0$ ,  $\mathbb{E}[M_t^\lambda] \leq \exp\left(\int_\lambda^{v_{-t}(\lambda)} \frac{\Phi(u)}{\Psi(u)} du\right) < \infty$ . The random variables  $M_t^\lambda$  are thus integrable and

$$\begin{aligned} \mathbb{E}[M_{t+s}^\lambda | \mathcal{F}_t] &= \mathbb{E}\left[\exp\left(-v_{-(t+s)}(\lambda)Y_{t+s} + \int_\lambda^{v_{-(t+s)}(\lambda)} \frac{\Phi(u)}{\Psi(u)} du\right) \middle| \mathcal{F}_t\right] \\ &= \exp\left(\int_\lambda^{v_{-(t+s)}(\lambda)} \frac{\Phi(u)}{\Psi(u)} du\right) \mathbb{E}[\exp(-v_{-(t+s)}(\lambda)Y_{t+s}) | Y_t] \quad (\text{by the Markov property}) \\ &= \exp\left(\int_\lambda^{v_{-(t+s)}(\lambda)} \frac{\Phi(u)}{\Psi(u)} du\right) \exp\left(-Y_t v_s(v_{-(t+s)}(\lambda)) - \int_{v_s(v_{-(t+s)}(\lambda))}^{v_{-(t+s)}(\lambda)} \frac{\Phi(u)}{\Psi(u)} du\right) \\ &= \exp\left(\int_\lambda^{v_{-(t+s)}(\lambda)} \frac{\Phi(u)}{\Psi(u)} du\right) \exp(-Y_t v_{-t}(\lambda)) \exp\left(\int_{v_{-(t+s)}(\lambda)}^{v_{-t}(\lambda)} \frac{\Phi(u)}{\Psi(u)} du\right) \\ &= \exp\left(\int_\lambda^{v_{-t}(\lambda)} \frac{\Phi(u)}{\Psi(u)} du\right) \exp(-Y_t v_{-t}(\lambda)) = M_t^\lambda, \end{aligned}$$

where the third equality follows from (2.6) and the fourth equality follows from the fact that  $v_s(v_{-(t+s)}(\lambda)) = v_s(v_{-s} \circ v_{-t}(\lambda)) = v_{-t}(\lambda)$ . In particular, we see that the process  $(e^{-v_{-t}(\lambda)Y_t}, t \geq 0)$  is a positive supermartingale. This entails that the process  $(v_{-t}(\lambda)Y_t, t \geq 0)$  converges, as  $t$  goes to infinity,  $\mathbb{P}_x$ -almost-surely in  $\mathbb{R}_+ := [0, \infty]$ . It remains to study its limit. We denote it by  $W^\lambda$  and shall see that it is infinite almost-surely in case ii).

Since  $\Psi$  is supercritical, one has  $\rho > 0$ , and  $\Psi(u) < 0$  for  $0 < u < \rho$ . By Lemma 1, we know that

$$v_{-t}(\lambda) \xrightarrow[t \rightarrow \infty]{} 0.$$

For fixed  $\theta > 0$ , we choose  $t$  large enough such that  $v_{-t}(\lambda), \theta v_{-t}(\lambda) \in (0, \rho)$ . One has

$$\mathbb{E}_x[e^{-\theta v_{-t}(\lambda)Y_t}] = \exp\left(-xv_t(\theta v_{-t}(\lambda)) - \int_{v_t(\theta v_{-t}(\lambda))}^{\theta v_{-t}(\lambda)} \frac{\Phi(u)}{\Psi(u)} du\right). \quad (3.16)$$

We have by (2.8) and (2.10),

$$\int_{\lambda}^{v_t(\theta v_{-t}(\lambda))} \frac{dz}{\Psi(z)} = \int_{\lambda}^{v_{-t}(\lambda)} \frac{dz}{\Psi(z)} + \int_{v_{-t}(\lambda)}^{\theta v_{-t}(\lambda)} \frac{dz}{\Psi(z)} + \int_{\theta v_{-t}(\lambda)}^{v_t(\theta v_{-t}(\lambda))} \frac{dz}{\Psi(z)} = \int_{v_{-t}(\lambda)}^{\theta v_{-t}(\lambda)} \frac{dz}{\Psi(z)}.$$

Recall that by assumption  $b \in (-\infty, 0)$ . Then

$$\lim_{t \rightarrow \infty} \int_{\lambda}^{v_t(\theta v_{-t}(\lambda))} \frac{dz}{\Psi(z)} = \lim_{t \rightarrow \infty} \int_{v_{-t}(\lambda)}^{\theta v_{-t}(\lambda)} \frac{dz}{b z} = \frac{\ln \theta}{b}. \quad (3.17)$$

Equations (2.4) and (2.10) ensure that

$$v_t(\theta v_{-t}(\lambda)) \xrightarrow{t \rightarrow \infty} v_{-\ln \theta/b}(\lambda). \quad (3.18)$$

Note that the above convergence holds no matter the sign of  $-\ln \theta/b$ . It is not hard to see that  $v_{\ln(-\frac{\theta}{b})}(\lambda) \in (0, \rho)$  as  $\int_0^{\frac{1}{\Psi(z)}} dz = \infty$  and  $\int_{\lambda}^{\rho} \frac{1}{\Psi(z)} dz = \infty$ . In fact, if  $\rho < \infty$ , then  $\Psi(\rho) = 0$ ; if  $\rho = \infty$ , then  $-\Psi$  is the Laplace exponent of a subordinator which implies that  $\int^{\infty} \frac{1}{\Psi(z)} dz = \infty$ . Moreover by (2.10),

$$v_{-\ln(\frac{\theta}{b})}(\lambda) \xrightarrow{\theta \rightarrow 0} 0 \text{ and } v_{-\ln(\frac{\theta}{b})}(\lambda) \xrightarrow{\theta \rightarrow \infty} \rho. \quad (3.19)$$

By taking now the limit as  $t$  goes to  $\infty$  in (3.16), we obtain

$$\mathbb{E}_x[e^{-\theta W^\lambda}] = \exp\left(-x v_{-\ln(\frac{\theta}{b})}(\lambda) + \int_0^{v_{-\ln(\frac{\theta}{b})}(\lambda)} \frac{\Phi(u)}{\Psi(u)} du\right). \quad (3.20)$$

If  $\int_0^{\frac{\Phi(u)}{|\Psi(u)|}} du < \infty$ , then  $W^\lambda < \infty$   $\mathbb{P}_x$ -almost-surely using the first convergence in (3.19). Applying the second convergence in (3.19) and the fact that  $\int^{\rho} \frac{du}{\Psi(u)} = -\infty$ ,

$$\mathbb{P}(W_x^\lambda = 0) = \exp\left(-x\rho + \int_0^{\rho} \frac{\Phi(u)}{\Psi(u)} du\right) = 0.$$

If  $\int_0^{\frac{\Phi(u)}{|\Psi(u)|}} du = \infty$ , then  $\int_0^{\frac{\Phi(u)}{\Psi(u)}} du = -\infty$  as  $\Psi(u) < 0$  for  $0 < u < \rho$ . We immediately have  $\mathbb{E}_x[e^{-\theta W^\lambda}] = 0$  by (3.20) for any  $\theta > 0$ . This implies that  $W^\lambda = \infty$  almost-surely.  $\square$

The next Theorem sheds some light on what limit theorems can be expected in the case  $\int_0^{\frac{\Phi(u)}{|\Psi(u)|}} du = \infty$ . In particular it breaks the hope of finding any law of large numbers.

**Theorem 4.** *Let  $(Y_t, t \geq 0)$  be a supercritical CBI $(\Psi, \Phi)$ . Assume  $\int_0^{\frac{\Phi(u)}{|\Psi(u)|}} du = \infty$ . Then, there exists no deterministic renormalization function  $(\eta(t), t \geq 0)$  such that  $\eta(t)Y_t \xrightarrow{t \rightarrow \infty} V$  almost-surely for some non-degenerate random variable  $V$ .*

*Remark 5.* A simpler proof of Theorem 4, not based on the same arguments as the proof below, is provided in Corollary 1.

*Proof.* We shall use the framework of flow of SDEs as Dawson and Li [DL12]. We recall that by replacing in the SDE (2.7), the Brownian motion  $(B_t, t \geq 0)$  by a white noise  $M(ds, du)$ , see [DL12] and [LM08] for details, one can consider on a same probability space, the SDEs

$$\begin{aligned} Y_t^{(n)}(x) &= x + \sigma \int_n^{n+t} \int_0^{Y_s^{(n)}(x)} M(ds, du) + \int_n^{n+t} \int_0^\infty \int_0^{Y_s^{(n)}(x)} z \tilde{N}_0(ds, dz, du) \\ &\quad + \int_n^{n+t} (\beta - bY_s^{(n)}(x)) ds + \int_n^{n+t} \int_0^\infty z N_1(ds, dz). \end{aligned} \quad (3.21)$$



Those SDEs are known to have pathwise unique solutions. More precisely this provides a sequence of flows of CBI( $\Psi, \Phi$ ) processes  $\{Y_t^{(n)}(x), x \geq 0, t \geq 0, n \geq 0\}$  such that for any  $n \in \mathbb{N}$  and any  $x \geq y \geq 0$ ,  $(Y_t^{(n)}(x) - Y_t^{(n)}(y), t \geq 0)$  is a CB process started from  $x - y$  with branching mechanism  $\Psi$  and is independent of  $\{Y_t^{(n)}(y), t \geq 0\}$ . We denote by  $(Y_t(x), t \geq 0)$  the process  $(Y_t^{(0)}(x), t \geq 0)$  solution to (3.21) for  $n = 0$ . Pathwise uniqueness entails that  $Y_t^{(n)}(Y_n(x)) = Y_{n+t}(x)$  for any  $x \geq 0, t \geq 0$  and  $n \in \mathbb{N}$  almost-surely. Let  $X_t^{(n)}(x) = Y_t^{(n)}(x) - Y_t^{(n)}(0)$ . Then

$$Y_{n+t}(0) = X_t^{(n)}(Y_n(0)) + Y_t^{(n)}(0).$$

By Theorem 2-(ii),  $v_{-t}(\lambda)Y_t^{(n)}(0) \xrightarrow[t \rightarrow \infty]{} \infty$  a.s and applying Theorem 2-(i) to the CB( $\Psi$ ) process  $(X_t^{(n)}(Y_n(0)), t \geq 0)$ , see Remark 3-i), we get  $v_{-t}(\lambda)X_t^{(n)}(Y_n(0)) \xrightarrow[t \rightarrow \infty]{} W^\lambda$  for some finite random variable  $W^\lambda$ . Hence

$$\frac{Y_{n+t}(0)}{Y_t^{(n)}(0)} = 1 + \frac{X_t^{(n)}(Y_n(0))}{Y_t^{(n)}(0)} \xrightarrow[t \rightarrow \infty]{} 1 \quad \text{a.s..} \quad (3.22)$$

Assume that there exists some  $\eta(t) > 0$  such that  $\eta(t)Y_t(0) \xrightarrow[t \rightarrow \infty]{} W_0$  a.s.. Then by (3.22),  $\eta(n+t)Y_t^{(n)}(0) \xrightarrow[t \rightarrow \infty]{} W_0$  a.s., and for  $\ell \in \mathbb{N}$ ,  $\eta(n+\ell)Y_\ell^{(n)}(0) \xrightarrow[\ell \rightarrow \infty]{} W_0$  a.s.. However by iteration, it is not hard to see that for  $\ell \geq 1$ ,

$$\begin{aligned} Y_\ell^{(n)}(0) &= X_{\ell-1}^{(n+1)}(Y_1^{(n)}(0)) + Y_{\ell-1}^{(n+1)}(0) \\ &= \sum_{k=1}^{\ell} X_{\ell-k}^{(n+k)}(Y_1^{(n+k-1)}(0)), \quad (Y_0^{(n+\ell)}(0) = 0) \end{aligned}$$

where  $\{X_t^{(k)}(Y_1^{(k-1)}(0))\}_{k=1}^{\infty}$  is a sequence of independent CB( $\Psi$ ) processes.

By the above iteration,  $Y_\ell^{(n)}(0)$  is measurable with respect to the tail  $\sigma$ -algebra generated by the sequence of independent processes  $(\{X_t^{(k)}(Y_1^{(k-1)}(0)) : t \geq 0\}_{k=n+1}^{\infty}, n \geq 1)$ . Since  $\eta(n+\ell)Y_\ell^{(n)}(0) \xrightarrow[\ell \rightarrow \infty]{} W_0$  a.s., we immediately have  $W_0$  is measurable with respect to the same tail  $\sigma$ -algebra for any  $n \geq 1$ .

Kolmogorov's zero-one law, see e.g. [Dur10, Theorem 2.5.1], asserts that  $W_0$  is a constant or infinite a.s.. Assume that  $W_0$  is a finite positive constant a.s.. Since  $Y_t^{(n)}(0)$  has the same distribution as  $Y_t(0)$ , we immediately have that  $\eta(t)Y_t^{(n)}(0) \xrightarrow[t \rightarrow \infty]{} W_0$  a.s.. Then  $\eta(n+t)/\eta(t) \xrightarrow[t \rightarrow \infty]{} 1$ , which implies that  $\eta(t) \sim L(e^t)$  for some slowly varying function  $L$  at  $\infty$ . However recalling that  $v_{-\ln t}$  is a regularly varying function with index  $b$ , see Lemma 1, we have that  $v_{-t}(\lambda) \sim e^{bt}L^*(e^t)$  as  $t$  goes to  $\infty$  where  $L^*(\cdot)$  is a slowly varying function at  $\infty$  and thus  $v_{-t}(\lambda)/\eta(t) \rightarrow 0$ . This leads to a contradiction. Thus  $W_0$  is 0 or  $\infty$  a.s..  $\square$

The last theorem naturally leads to the problem of finding some renormalization in law of a CBI( $\Psi, \Phi$ ) process in the case  $\int_0 \frac{\Phi(u)}{|\Psi(u)|} du = \infty$ . This is the aim of the next section.

### 3.2 A general limit in law for non-critical CBIs when $\int_0 \frac{\Phi(u)}{|\Psi(u)|} du = \infty$

This section focuses on the study of the long-term behavior of CBI( $\Psi, \Phi$ ) processes satisfying the condition

$$\int_0 \frac{\Phi(u)}{|\Psi(u)|} du = \infty. \quad (3.23)$$

By Theorem 1, in this case  $Y_t$  converges in law to  $\infty$  as  $t$  goes to  $\infty$ . We give a finer description of the behavior of  $Y_t$  through distributional, rather than almost sure, limit theorems. In this section, we prove the main convergence theorem below which provides a non-linear time-dependent renormalization in law of any non-critical CBI process. The three different regimes of convergence in law mentioned in the Introduction are designed in the forthcoming Section 3.3.

Recall the definition of  $r_t(\lambda)$  in (2.5). Recall that  $r_t(\infty) = \infty$  if and only if  $\mathbb{P}_x(Y_t = 0) = 0$  for all  $t > 0$ . In the next theorem, we take the convention  $1/0 = \infty$ .

**Theorem 6.** *Assume (3.23) holds, and  $\Psi$  is non-critical ( $b \neq 0$ ). Then, for all  $x \geq 0$ , we have*

$$r_t(1/Y_t) := \int_{v_t(1/Y_t)}^{1/Y_t} \frac{\Phi(u)}{\Psi(u)} du \xrightarrow{d} e_1, \text{ as } t \rightarrow \infty \text{ under } \mathbb{P}_x \quad (3.24)$$

where  $e_1$  is an exponential random variable with parameter 1.

*Remark 7.* The limiting distribution in (3.24) in law does not depend on the initial state  $x$  of the CBI process. It justifies the perception that in the regime (3.23), the dynamics is governed on the long-term by the immigration part and not by the branching part.

*Proof.* Recall the equations (2.3) and (2.6).

Step 1: We claim that for fixed  $\lambda > 0$ ,  $r_t(\lambda) \rightarrow \infty$  as  $t \rightarrow \infty$ . In fact, in the subcritical case,  $\Psi(u) > 0$  for  $u > 0$ . By (2.8),  $v_t(\lambda) \downarrow 0$  as  $t \rightarrow \infty$ , for any fixed  $\lambda > 0$ . According to (3.23) and the second equality of (2.5), we have that  $r_t(\lambda) \rightarrow \infty$  as  $t \rightarrow \infty$ .

In the supercritical case, still by (2.8),  $v_t(\lambda) \rightarrow \rho$  as  $t \rightarrow \infty$  for fixed  $\lambda > 0$ . Then  $\Phi(v_t(\lambda)) \rightarrow \Phi(\rho) > 0$ , as  $t \rightarrow \infty$ . Together with the first equality of (2.5), we obtain that  $r_t(\lambda) \rightarrow \infty$  as  $t \rightarrow \infty$ .

Step 2: Recall that  $\lambda \mapsto r_t(\lambda)$  is the Laplace exponent of a subordinator with no killing term. For all  $t$ ,  $r_t(0) = 0$ , and  $r_t$  is strictly decreasing in  $\lambda$ . So we can define  $\lambda \mapsto c_t(\lambda)$  as the inverse of  $\lambda \mapsto r_t(\lambda)$ . Fix  $\lambda > 0$ . By Step 1, for any small  $\varepsilon > 0$ , we can find sufficiently large  $t$  such that  $r_t(\varepsilon) > \lambda = r_t(c_t(\lambda))$ , which implies that  $c_t(\lambda) < \varepsilon$ . Thus

$$c_t(\lambda) \xrightarrow{t \rightarrow \infty} 0, \text{ and } v_t(c_t(\lambda)) \xrightarrow{t \rightarrow \infty} 0. \quad (3.25)$$

The second limit follows from (3.23) and the second equality of (2.5) (replacing  $\lambda$  by  $c_t(\lambda)$ ).

Step 3: Notice that we can equivalently show that for any  $\lambda \geq 0$  and  $\theta > 0$ ,

$$\lim_{t \rightarrow \infty} \mathbb{E}_x[e^{-\theta c_t(\lambda) Y_t}] = e^{-\lambda}. \quad (3.26)$$

In fact, if (3.26) holds, then  $c_t(\lambda) Y_t$  converges in distribution to a random variable  $Z$  such that  $\mathbb{P}(Z = \infty) = 1 - \mathbb{P}(Z = 0) = 1 - e^{-\lambda}$ . Therefore, for  $\lambda > 0$ ,

$$\mathbb{P}(r_t(1/Y_t) > \lambda) = \mathbb{P}(1/Y_t > c_t(\lambda)) = \mathbb{P}(c_t(\lambda) Y_t < 1) \xrightarrow{t \rightarrow \infty} e^{-\lambda}, \quad (3.27)$$

which implies (3.24).

Step 4: Note that

$$\mathbb{E}_x[e^{-\theta c_t(\lambda) Y_t}] = \exp(-x v_t(\theta c_t(\lambda)) - r_t(\theta c_t(\lambda))) \quad (3.28)$$

and

$$\begin{aligned} r_t(\theta_{c_t}(\lambda)) &= \int_{v_t(\theta_{c_t}(\lambda))}^{\theta_{c_t}(\lambda)} \frac{\Phi(u)}{\Psi(u)} du \\ &= \int_{c_t(\lambda)}^{\theta_{c_t}(\lambda)} \frac{\Phi(u)}{\Psi(u)} du + \underbrace{\int_{v_t(c_t(\lambda))}^{c_t(\lambda)} \frac{\Phi(u)}{\Psi(u)} du}_{=r_t(c_t(\lambda))=\lambda} + \int_{v_t(\theta_{c_t}(\lambda))}^{v_t(c_t(\lambda))} \frac{\Phi(u)}{\Psi(u)} du. \end{aligned} \quad (3.29)$$

So to obtain (3.26), it suffices to prove that as  $t \rightarrow \infty$ ,

$$v_t(\theta_{c_t}(\lambda)) \rightarrow 0 \quad (3.30)$$

and

$$\int_{c_t(\lambda)}^{\theta_{c_t}(\lambda)} \frac{\Phi(u)}{\Psi(u)} du \rightarrow 0, \quad \int_{v_t(c_t(\lambda))}^{v_t(\theta_{c_t}(\lambda))} \frac{\Phi(u)}{\Psi(u)} du \rightarrow 0. \quad (3.31)$$

By monotonicity of  $\Phi$ , we have

$$\left| \int_{\theta y}^y \frac{\Phi(u)}{\Psi(u)} du \right| \leq \max(\Phi(y), \Phi(\theta y)) \left| \int_{\theta y}^y \frac{1}{\Psi(u)} du \right|. \quad (3.32)$$

On the one hand  $\max(\Phi(y), \Phi(\theta y)) \xrightarrow{y \rightarrow 0} 0$  as  $\Phi$  is continuous and  $\Phi(0) = 0$ . On the other hand, since  $\Psi$  is non-critical, there exists some constant  $h > 0$  such that  $|\Psi(u)| \geq hu$  for  $u$  close enough to 0. This entails that

$$\left| \int_{\theta y}^y \frac{1}{\Psi(u)} du \right| \leq \frac{|\ln \theta|}{h}, \quad (3.33)$$

when  $y$  is small enough. Then by (3.32) and (3.25),

$$\left| \int_{c_t(\lambda)}^{\theta_{c_t}(\lambda)} \frac{\Phi(u)}{\Psi(u)} du \right| \leq \max(\Phi(c_t(\lambda)), \Phi(\theta_{c_t}(\lambda))) \frac{|\ln \theta|}{h} \xrightarrow{t \rightarrow \infty} 0. \quad (3.34)$$

So the first convergence in (3.31) is proved.

Note that

$$\begin{aligned} \int_{v_t(c_t(\lambda))}^{v_t(\theta_{c_t}(\lambda))} \frac{du}{\Psi(u)} &= \int_{v_t(c_t(\lambda))}^{c_t(\lambda)} \frac{du}{\Psi(u)} + \int_{c_t(\lambda)}^{\theta_{c_t}(\lambda)} \frac{du}{\Psi(u)} + \int_{\theta_{c_t}(\lambda)}^{v_t(\theta_{c_t}(\lambda))} \frac{du}{\Psi(u)} \\ &= t + \int_{c_t(\lambda)}^{\theta_{c_t}(\lambda)} \frac{du}{\Psi(u)} - t \\ &= \int_{c_t(\lambda)}^{\theta_{c_t}(\lambda)} \frac{du}{\Psi(u)}, \end{aligned}$$

which entails that  $v_t(\theta_{c_t}(\lambda)) \rightarrow 0$  as  $t \rightarrow \infty$ , since  $v_t(c_t(\lambda)) \rightarrow 0$  by (3.25) and  $\left| \int_{c_t(\lambda)}^{\theta_{c_t}(\lambda)} \frac{du}{\Psi(u)} \right| \leq \ln \theta / h$  by (3.33). So the convergence in (3.30) is proved. Then

$$\begin{aligned} \left| \int_{v_t(c_t(\lambda))}^{v_t(\theta_{c_t}(\lambda))} \frac{\Phi(u)}{\Psi(u)} du \right| &\leq \max\{\Phi(v_t(c_t(\lambda))), \Phi(v_t(\theta_{c_t}(\lambda)))\} \left| \int_{v_t(c_t(\lambda))}^{v_t(\theta_{c_t}(\lambda))} \frac{1}{\Psi(u)} du \right| \\ &= \max\{\Phi(v_t(c_t(\lambda))), \Phi(v_t(\theta_{c_t}(\lambda)))\} \left| \int_{c_t(\lambda)}^{\theta_{c_t}(\lambda)} \frac{1}{\Psi(u)} du \right|, \end{aligned}$$

which goes to 0 similarly as in (3.34). Then the second convergence in (3.31) is proved. So both conditions (3.30) and (3.31) hold true, and we can conclude that (3.24) holds true.  $\square$

*Remark 8.* Note that the steps 1 and 2 also work for the critical case. But step 4 requires  $b \neq 0$ . However the same line of arguments as in this proof will be used in Section 3.4 where we focus on the study of the critical case.

We now provide a corollary leading to a more intuitive probabilistic understanding of Theorem 6. In particular it will provide an alternative proof of Theorem 4. We take the convention  $0/0 = 0 \times \infty = 0$ .

**Corollary 1.** *Assume (3.23) and that  $\Psi$  is non-critical. Let  $(Y_t, t \geq 0)$  and  $(\tilde{Y}_t, t \geq 0)$  be two independent CBI( $\Psi, \Phi$ ) processes started from 0. Then*

$$\mathbb{P}(Y_t/\tilde{Y}_t \xrightarrow[t \rightarrow \infty]{} 0) = \mathbb{P}(Y_t/\tilde{Y}_t \xrightarrow[t \rightarrow \infty]{} \infty) = \frac{1}{2}. \quad (3.35)$$

Moreover, there exists no deterministic renormalization  $(\eta(t), t \geq 0)$  such that  $(\eta(t)Y_t, t \geq 0)$  converges in law towards a non-degenerate random variable.

*Proof.* By a similar argument as in Equation 3.27, since for any  $\lambda > 0$ ,  $c_t(\lambda)Y_t \xrightarrow[t \rightarrow \infty]{} Z$  in law with  $Z$  such that  $\mathbb{P}(Z = \infty) = 1 - \mathbb{P}(Z = 0) = 1 - e^{-\lambda}$ , one has for any  $\theta > 0$  and

$$\mathbb{P}(r_t(\theta/Y_t) > \lambda) = \mathbb{P}(c_t(\lambda)Y_t < \theta) \xrightarrow[t \rightarrow \infty]{} e^{-\lambda} = \mathbb{P}(e_1 > \lambda)$$

with  $e_1$  a standard exponential random variable. Hence for any  $\theta \geq 0$  and for any  $t \geq 0$ , we apply (2.6) to obtain

$$\begin{aligned} \mathbb{E}[e^{-\theta \frac{Y_t}{\tilde{Y}_t}}] &= \tilde{\mathbb{E}}_0[\mathbb{E}_0[e^{-\frac{\theta}{\tilde{Y}_t} Y_t} | \tilde{Y}_t]] = \tilde{\mathbb{E}}[\mathbb{E}[e^{-r_t(\theta/\tilde{Y}_t)} | \tilde{Y}_t]] \\ &= \tilde{\mathbb{E}}[e^{-r_t(\theta/\tilde{Y}_t)}] \xrightarrow[t \rightarrow \infty]{} \mathbb{E}[e^{-e_1}] = \frac{1}{2}. \end{aligned}$$

Therefore  $Y_t/\tilde{Y}_t \xrightarrow[t \rightarrow \infty]{} \Lambda$  in law with  $\mathbb{P}(\Lambda = 0) = \frac{1}{2}$  and  $\mathbb{P}(\Lambda = \infty) = \frac{1}{2}$ .

We show now that there is no renormalization in law. By contradiction, assume that there exists  $(\eta(t), t \geq 0)$  such that  $\eta(t)Y_t \xrightarrow[t \rightarrow \infty]{} V$  in law with  $V$  a non-degenerate random variable. Let  $b > a > 0$  be any two values such that  $\mathbb{P}(V \in [a, b]) > 0$ . Then,

$$\liminf_{t \rightarrow \infty} \mathbb{P}\left(\frac{\eta(t)Y_t}{\eta(t)\tilde{Y}_t} = \frac{Y_t}{\tilde{Y}_t} \in [a/b, b/a]\right) > 0.$$

But this is in contradiction to (3.35). Then necessarily  $V$  is degenerate.  $\square$

*Remark 9.* The statement of Corollary 1 holds true for CBI processes started from arbitrary initial values. Indeed if  $(Y_t(x), t \geq 0)$  and  $(\tilde{Y}_t(y), t \geq 0)$  are two independent CBI( $\Psi, \Phi$ ) started respectively at  $x$  and  $y$ , then for any  $t \geq 0$ ,  $Y_t(x) = X_t(x) + Y_t(0)$  with  $(X_t(x), t \geq 0)$  a CB( $\Psi$ ) started from  $x$  and independent of  $(Y_t(0), t \geq 0)$ . Similarly  $\tilde{Y}_t(y) = \tilde{X}_t(y) + \tilde{Y}_t(0)$  with  $(\tilde{X}_t(y), t \geq 0)$  a CB( $\Psi$ ) started from  $y$  and independent of  $(\tilde{Y}_t(0), t \geq 0)$ . One checks that

$$\frac{Y_t(x)}{\tilde{Y}_t(y)} = \frac{Y_t(0)(1 + X_t(x)/Y_t(0))}{\tilde{Y}_t(0)(1 + \tilde{X}_t(y)/\tilde{Y}_t(0))} \underset{t \rightarrow \infty}{\sim} \frac{Y_t(0)}{\tilde{Y}_t(0)} \text{ a.s.}$$

Indeed if  $\Psi$  is (sub)critical then both  $\tilde{X}_t(x)$  and  $X_t(y)$  converge towards 0 almost-surely. If  $\Psi$  is supercritical then by Theorem 2, for any  $x \geq 0$ ,  $\frac{X_t(x)}{Y_t(0)} = \frac{v_{-t}(\lambda)X_t(x)}{v_{-t}(\lambda)Y_t(0)} \xrightarrow[t \rightarrow \infty]{} 0$  a.s.

### 3.3 Three different regimes

Since the renormalization in Theorem 6 is non-linear and time-dependent, it is rather intricate at a first sight to deduce from it which explicit growth rates are possible. We design here different regimes for which (3.23) holds and the rate can be found explicitly. This establishes and completes [Pin72, Theorem 2].

#### 3.3.1 Definition of the regimes and preliminary calculations

Recall  $\lambda \mapsto c_t(\lambda)$  the inverse of  $\lambda \mapsto r_t(\lambda) = \int_{v_t(\lambda)}^{\lambda} \frac{\Phi(u)}{\Psi(u)} du$ . Theorem 6 indicates that  $Y_t$  should grow at the speed of  $1/c_t(\lambda)$  as  $t \rightarrow \infty$ . However the magnitude of  $c_t(\lambda)$  is rather involved and deserves a careful analysis. We shall simplify (3.24) to more straightforward forms by imposing some additional conditions.

To start with, let us fix some constant  $\lambda_0$  such that  $\lambda_0 \in (0, \infty)$  in the (sub)critical case and  $\lambda_0 \in (0, \rho)$  in the supercritical case. Put

$$\varphi(\lambda) = \int_{\lambda}^{\lambda_0} \frac{du}{|\Psi(u)|}, \quad 0 < \lambda < \lambda_0. \quad (3.36)$$

By assumption  $|\Psi'(0+)| < \infty$  and thus  $\varphi(\lambda) \rightarrow \infty$  as  $\lambda \rightarrow 0$ . The mapping  $\varphi : (0, \lambda_0) \rightarrow (0, \infty)$  is strictly decreasing, and we write  $g$  for its inverse mapping. It is easy to see that  $g$  is a strictly decreasing continuous function on  $(0, \infty)$ , and

$$\lim_{x \rightarrow \infty} g(x) = 0, \quad \lim_{x \rightarrow 0} g(x) = \lambda_0. \quad (3.37)$$

By (2.8), if  $b \geq 0$ , then  $\Psi \geq 0$  and

$$\varphi(v_t(\lambda)) = \int_{v_t(\lambda)}^{\lambda} \frac{du}{\Psi(u)} + \int_{\lambda}^{\lambda_0} \frac{du}{\Psi(u)} = t + \varphi(\lambda). \quad (3.38)$$

Similarly if  $b < 0$ , then  $\Psi(u) \leq 0$  for  $0 \leq u \leq \rho$  and provided that  $v_t(\lambda) \in (0, \lambda_0)$

$$\varphi(v_t(\lambda)) = - \int_{v_t(\lambda)}^{\lambda} \frac{du}{\Psi(u)} - \int_{\lambda}^{\lambda_0} \frac{du}{\Psi(u)} = -t + \varphi(\lambda). \quad (3.39)$$

Applying  $g$  to both sides entails that if  $b \geq 0$

$$v_t(\lambda) = g(\varphi(\lambda) + t), \quad 0 < \lambda < \lambda_0, \quad t > 0 \quad (3.40)$$

and if  $b < 0$

$$v_t(\lambda) = g(\varphi(\lambda) - t), \quad 0 < \lambda < \lambda_0, \quad t > 0. \quad (3.41)$$

Then for any  $x, y > 0$  such that  $\Psi$  never attains zero between  $x, y$ , we obtain by a change of variable that

$$\int_x^y \frac{\Phi(u)}{|\Psi(u)|} du = \int_{g(\varphi(x))}^{g(\varphi(y))} \frac{\Phi(u)}{|\Psi(u)|} du = \int_{\varphi(y)}^{\varphi(x)} \Phi(g(u)) du. \quad (3.42)$$

Inspired by Pinsky [Pin72, Theorem 2], we introduce the following function to characterize the divergence of the integral in (3.23):

$$H(x) := \begin{cases} \frac{1}{|b|} \int_{e^{-x}}^1 \frac{\Phi(u)}{u} du, & \text{if } b \in (-\infty, 0) \cup (0, \infty); \\ \int_{g(x)}^{\lambda_0} \frac{\Phi(u)}{|\Psi(u)|} du, & \text{if } b = 0 \end{cases}, \quad x \geq 0 \quad (3.43)$$

where  $H(0)$  takes the value  $\lim_{x \rightarrow 0} H(x) = 0$  (using (3.37)). It is not hard to check that the condition (3.23) is equivalent to  $H(x) \xrightarrow{x \rightarrow \infty} \infty$ . Based on (3.36), a simple calculation shows that  $H'$  is strictly decreasing and  $H'(x) \rightarrow 0$  as  $x \rightarrow \infty$ . We introduce now different regimes of speed of divergence of the function  $H$  at  $\infty$ . We refer the reader to Bingham et al. [BGT87] for a reference on those functions. The following three different modes of convergence to 0 of  $H'$  correspond to different modes of divergence of  $H$ :

- (S) (slow-divergence)  $xH'(x) \rightarrow 0$  as  $x \rightarrow \infty$  and  $H(x) \rightarrow \infty$  ;
- (L) (log-divergence)  $xH'(x) \rightarrow a$  for some constant  $a > 0$  as  $x \rightarrow \infty$ ;
- (F) (fast-divergence)  $xH'(x) \rightarrow \infty$  as  $x \rightarrow \infty$  and  $H'$  is regularly varying at  $\infty$ .

Note that condition (L) and (F) above entail that  $\lim_{x \rightarrow \infty} H(x) = \infty$ . Since  $H'(x) \rightarrow 0$  as  $x \rightarrow \infty$ , Condition (F) can be given in the following equivalent form:

$$H'(x) = x^{-\delta} \frac{1}{L(x)}, \quad (3.44)$$

where  $L$  is slowly varying at  $\infty$  and  $0 \leq \delta \leq 1$ , and if  $\delta = 0$ ,  $L(x) \rightarrow \infty$  as  $x \rightarrow \infty$ ; if  $\delta = 1$ ,  $L(x) \rightarrow 0$  as  $x \rightarrow \infty$ .

Let  $h$  be the inverse map of  $1/H'$ . Under (3.44) with  $\delta > 0$ ,  $t \mapsto h(t)$  is regularly varying with index  $1/\delta$  at  $\infty$ , i.e.  $h(t) \sim t^{1/\delta} L^*(t)$  for some slowly varying function  $L^*(t)$ . Moreover, since  $H'(x) \rightarrow 0$  as  $x \rightarrow \infty$ , we have

$$\lim_{t \rightarrow \infty} h(t) = \infty. \quad (3.45)$$

It might be difficult to verify the three conditions. We provide a proposition below to study the asymptotic behaviors of  $H$  and  $H'$ . To this purpose, recall the immigration mechanism

$$\Phi(q) = \beta q + \int_0^\infty (1 - e^{-qu}) \nu(du).$$

As observed by Pinsky [Pin72, P. 244], for a non-critical CBI process, a faster rate of divergence in (3.23) implies heavier tails of the Lévy measure  $\nu(du)$ . The following result specifies this idea.

**Proposition 1.** *Assume that  $b \neq 0$  and  $H(x) \rightarrow \infty$  as  $x \rightarrow \infty$ . Denote by  $\bar{\nu}$  the tail of  $\nu$ : for all  $u \geq 0$ ,  $\bar{\nu}(u) = \nu([u, \infty))$ . Then*

$$H(x) \sim \frac{1}{|b|} \int_1^{e^x} \frac{\bar{\nu}(u)}{u} du, \quad \text{as } x \rightarrow \infty.$$

Moreover  $H'(x) = \Phi(e^{-x})/|b|$  for any  $x \geq 0$  and if  $u \mapsto \bar{\nu}(u)$  is slowly varying at  $\infty$  then

$$H'(x) \sim \bar{\nu}(e^x)/|b|, \quad \text{as } x \rightarrow \infty.$$

*Remark 10.* The tail  $u \mapsto \bar{\nu}(u)$  is slowly varying at  $\infty$  if and only if  $\Phi$  is slowly varying at 0, see [BGT87, Theorem 8/1/6, P.333]

*Proof.* Note that  $\Phi(u)/u = \beta + \int_0^\infty e^{-ut} \bar{\nu}(t) dt$ . A simple calculation shows that for  $z \in (0, 1)$

$$\begin{aligned} \int_z^1 \frac{\Phi(u)}{u} du &= \beta(1-z) + \int_0^\infty \frac{\bar{\nu}(u)}{u} (e^{-zu} - e^{-u}) du \\ &= \beta(1-z) + \left( \int_0^1 + \int_1^\infty \right) \frac{\bar{\nu}(u)}{u} (e^{-zu} - e^{-u}) du. \end{aligned}$$

Since  $\int_0^1 u\nu(du) < \infty$ ,  $\int_1^\infty \frac{\bar{\nu}(u)}{u} e^{-u} du < \infty$  and  $\int_{0+} \frac{\Phi(u)}{u} du = \infty$ , we have that as  $z \rightarrow 0$ ,

$$\int_z^1 \frac{\Phi(u)}{u} du \sim \int_1^\infty \frac{\bar{\nu}(u)}{u} e^{-zu} du, \quad (3.46)$$

which implies  $\int_1^x \frac{\bar{\nu}(u)}{u} du \rightarrow \infty$  as  $x \rightarrow \infty$ . It is not hard to see that  $\int_1^x \frac{\bar{\nu}(u)}{u} du$  is also slowly varying at  $\infty$ . It follows from Tauberian theorem (see e.g. Bertoin [Ber96, P.10]) that

$$\int_1^x \frac{\bar{\nu}(u)}{u} du \sim \int_1^\infty \frac{\bar{\nu}(u)}{u} e^{-u/x} du, \quad x \rightarrow \infty. \quad (3.47)$$

The first result follows from (3.46) and (3.47). For the second result, note that  $H'(x) = \Phi(e^{-x})/|b|$ . Without loss of generality, we can assume that the parameter  $\beta$  in  $\Phi$  equals zero. Applying the Tauberian theorem entails that if  $\bar{\nu}(x) \sim \ell(x)$  for some slowly varying function  $\ell$  at  $\infty$ , then  $\frac{\Phi(u)}{u} \sim \frac{1}{u} \ell(1/u)$  as  $u$  goes 0. Hence we have  $H'(x) \sim \bar{\nu}(e^x)/|b|$ ,  $x \rightarrow \infty$ .  $\square$

*Remark 11.* By letting  $z$  to 0 in (3.46), we see that  $\int_0 \frac{\Phi(x)}{x} dx = \infty$  if and only if  $\int^\infty \frac{\bar{\nu}(u)}{u} du = \infty$ , the latter is equivalent to  $\int_1^\infty \ln x \nu(dx) = \infty$ .

Proposition 1 allows us to reformulate the three regimes, in the non-critical case, in terms of the tail of the immigration measure  $\nu$ .

- (S) (slow-divergence)  $\bar{\nu}(x) \ln x \rightarrow 0$  as  $x \rightarrow \infty$  and  $\int_1^\infty \frac{\bar{\nu}(x)}{x} = \infty$ ;
- (L) (log-divergence)  $\bar{\nu}(x) \ln x \rightarrow c$  for some constant  $c > 0$  as  $x \rightarrow \infty$ ;
- (F) (fast-divergence)  $\bar{\nu}(x) \ln x \rightarrow \infty$  as  $x \rightarrow \infty$  and  $\bar{\nu}$  is slowly varying at  $\infty$ .

The constant  $c$  in regime (L) matches with  $a|b|$  where  $a := \lim_{x \rightarrow \infty} xH'(x)$ . We give below some examples of explicit immigration measures  $\nu$  for which the three different regimes may occur in the non-critical cases.

*Example 1.* Let  $b \in (-\infty, 0) \cup (0, \infty)$ .

1. If  $\bar{\nu}(x) \sim \frac{1}{\ln x \ln \ln x}$  as  $x \rightarrow \infty$ , then  $H(x) \sim (\ln \ln x)/|b|$  and  $H'(x) \sim 1/(|b|x \ln x)$ . Condition (S) is satisfied. This example corresponds to Example 3 in [DFM14, Example 3] of null-recurrent CBI.
2. If  $\bar{\nu}(x) \sim c/\ln x$  for some constant  $c > 0$ , as  $x \rightarrow \infty$ , then  $H'(x) \sim c/(|b|x)$ . Condition (L) is satisfied.
3. If  $\bar{\nu}(x) \sim \frac{\ln \ln x}{(\ln x)^\delta}$ , ( $0 < \delta \leq 1$ ) as  $x \rightarrow \infty$ , then  $H'(x) \sim (x^{-\delta} \ln x)/|b|$ . If as  $x \rightarrow \infty$ ,  $\bar{\nu}(x) \sim 1/(\ln \ln x)$ , then  $H'(x) \sim 1/(|b| \ln x)$ . Both cases satisfy Condition (F).

*Remark 12.* Recall the integral test  $\mathcal{E} < \infty$  or  $\mathcal{E} = \infty$  for transience and recurrence of (sub)critical CBIs given in (2.13). Plainly, by a change of variable,  $\mathcal{E} = \int^\infty e^{-H(x)} dx$ . If (F) holds, or (L) is satisfied with  $a > 1$ , then  $\mathcal{E} < \infty$  and the process is transient. In the case (S), or (L) with  $a \leq 1$ ,  $\mathcal{E} = \infty$  and the process is null-recurrent.

We state now an aside result on the growth rate of a subordinator whose Laplace exponent is slowly varying at 0.

**Proposition 2.** *Let  $(I_t, t \geq 0)$  be a subordinator with Laplace exponent  $\Phi$ . Assume that  $\Phi$  is slowly varying at 0, then*

$$t\Phi(1/I_t) \xrightarrow{d} e_1 \text{ as } t \rightarrow \infty.$$

*Proof.* This is reminiscent to the steps 1 and 2 in the proof of Theorem 6. Observe that the stated convergence holds if and only if for all  $\lambda > 0$

$$\mathbb{P}(t\Phi(1/I_t) > \lambda) = \mathbb{P}(I_t < 1/\Phi^{-1}(\lambda/t)) = \mathbb{P}(\Phi^{-1}(\lambda/t)I_t < 1) \xrightarrow[t \rightarrow \infty]{} e^{-\lambda}.$$

The latter will occur if for any  $\theta \geq 0$ ,

$$\mathbb{E}[e^{-\theta\Phi^{-1}(\lambda/t)I_t}] \xrightarrow[t \rightarrow \infty]{} e^{-\lambda}.$$

Since  $\Phi$  is slowly varying at 0 and  $\Phi^{-1}(\lambda/t) \rightarrow 0$  as  $t \rightarrow \infty$ , we have

$$\frac{t}{\lambda}\Phi(\theta\Phi^{-1}(\lambda/t)) = \frac{\Phi(\theta\Phi^{-1}(\lambda/t))}{\Phi(\Phi^{-1}(\lambda/t))} \xrightarrow[t \rightarrow \infty]{} 1.$$

Therefore

$$\mathbb{E}[e^{-\theta\Phi^{-1}(\lambda/t)I_t}] = e^{-t\Phi(\theta\Phi^{-1}(\lambda/t))} \xrightarrow[t \rightarrow \infty]{} e^{-\lambda}$$

which finishes the proof.  $\square$

In the next subsection, we study how the convergence results can be simplified by combining Theorem 6 and the three conditions. These results can be seen as the continuous analogues of those in [Pak79].

### 3.3.2 Subcritical case

**Theorem 13.** *Assume that  $b > 0$ .*

(i) *If Condition (S) holds, let  $m(x) := \exp(\int_{1/x}^1 \frac{\Phi(u)}{\Psi(u)} du)$  for  $x > 0$ . Then*

$$\frac{\ln Y_t}{bt} \xrightarrow{p} 0 \quad \text{and} \quad m(Y_t)/m(e^{bt}) \xrightarrow{d} U, \quad \text{as } t \rightarrow \infty, \quad (3.48)$$

where  $U$  is uniformly distributed on  $[0, 1]$ .

(ii) *If Condition (L) holds, then*

$$\frac{\ln Y_t}{bt} \xrightarrow{d} U_L, \quad \text{as } t \rightarrow \infty, \quad (3.49)$$

where  $\mathbb{P}(U_L \leq \lambda) = \left(\frac{\lambda}{1+\lambda}\right)^a$  for any  $\lambda \geq 0$ .

(iii) *If Condition (F) holds, then*

$$\frac{\ln Y_t}{bt} \xrightarrow{p} \infty \quad \text{and} \quad t\Phi(1/Y_t) \xrightarrow{d} e_1, \quad \text{as } t \rightarrow \infty.$$

If in addition (3.44) holds with  $\delta > 0$ , then we have

$$h(t) = t^{1/\delta}L^*(t) \quad \text{and} \quad \frac{\ln Y_t}{h(bt)} \xrightarrow{d} U_F, \quad \text{as } t \rightarrow \infty, \quad (3.50)$$

where  $L^*$  is some slowly varying function at  $\infty$  and  $U_F$  follows the extreme distribution given by

$$P(U_F \leq \lambda) = \exp(-1/\lambda^\delta), \quad \lambda \geq 0. \quad (3.51)$$



*Remark 14.* i) We observe from Proposition 2, that in the fast regime (F), the branching part plays essentially no role in the growth of the subcritical CBI( $\Psi, \Phi$ ) process, since it has the same growth rate as the immigration subordinator ( $I_t, t \geq 0$ ).

ii) The statement (i) of Theorem 13 has been given in Pinsky [Pin72, Theorem 2] which however contains some incorrectness. The corrected convergence in [Pin72, Theorem 2] reads:

$$P_x(m(X_t)/m(e^{ct}) \geq u^{-1}) \rightarrow u, \forall 0 < u \leq 1 \text{ as } t \rightarrow \infty.$$

We will first prove a lemma.

**Lemma 2.** *Assume that  $b > 0$ . Then  $\varphi(\lambda) \underset{\lambda \rightarrow 0}{\sim} -\frac{1}{b} \ln \lambda$ ,  $-\ln g(x) \underset{x \rightarrow \infty}{\sim} bx$ , and  $\ln v_t(\lambda) \sim -b(t + \varphi(\lambda))$  as  $t \rightarrow \infty$  or/and  $\lambda \rightarrow 0$ .*

*Proof.* Since  $b > 0$ , then  $\Psi(u) > 0$  for any  $u > 0$ . Note that

$$\varphi(\lambda) = \int_{\lambda}^{\lambda_0} \frac{du}{\Psi(u)} \underset{\lambda \rightarrow 0}{\sim} \int_{\lambda}^{\lambda_0} \frac{du}{bu} \underset{\lambda \rightarrow 0}{\sim} -\frac{1}{b} \ln \lambda.$$

Since  $g$  is the inverse function of  $\varphi$ , we have  $-\ln g(x) \underset{x \rightarrow \infty}{\sim} bx$ . Then we establish the last statement by plugging  $x = \varphi(\lambda) + t$  in (3.40).  $\square$

*Proof of Theorem 13* Since under the assumption (3.23), the process  $(Y_t, t \geq 0)$  goes to infinity in probability, by the Skorokhod representation theorem, there is a probability space on which is defined a process with the same one-dimensional laws as the CBI process but who goes to  $\infty$  almost-surely as  $t$  goes to  $\infty$ . When dealing with subcritical or critical processes, we shall also work implicitly with this process.

By definition

$$H(-\ln v_t(1/Y_t)) - H(\ln Y_t) = \frac{1}{b} \int_{v_t(1/Y_t)}^{1/Y_t} \frac{\Phi(u)}{u} du. \quad (3.52)$$

In the sequel, we will frequently use the following two results:

1. Since  $Y_t \xrightarrow{a.s} \infty$ , by Lemma 2 we have that

$$-\ln v_t(1/Y_t) \sim b(t + \varphi(1/Y_t)) \quad \text{and} \quad b\varphi(1/Y_t) \sim \ln Y_t \text{ as } t \rightarrow \infty. \quad (3.53)$$

2. From (3.52) and Theorem 6, together with  $b > 0$ , we get

$$H(-\ln v_t(1/Y_t)) - H(\ln Y_t) = (1 + o(1)) \int_{v_t(1/Y_t)}^{1/Y_t} \frac{\Phi(u)}{\Psi(u)} du \xrightarrow{d} e_1 \text{ as } t \rightarrow \infty. \quad (3.54)$$

(i) We start with the first half of (3.48). Applying first the mean value theorem, and then Condition (S) and (3.53), we see that there exists  $\theta_t$  between  $\ln Y_t$  and  $-\ln v_t(1/Y_t)$  such that

$$\begin{aligned} H(-\ln v_t(1/Y_t)) - H(\ln Y_t) &= \int_{\ln Y_t}^{-\ln v_t(1/Y_t)} \frac{uH'(u)}{u} du \\ &= H'(\theta_t)\theta_t \ln \left( -\frac{\ln v_t(1/Y_t)}{\ln Y_t} \right) \\ &= o(1) \ln \left( \frac{bt}{\ln Y_t} + 1 \right) + o(1). \end{aligned} \quad (3.55)$$

A comparison with (3.54) entails that  $(\ln Y_t)/bt \xrightarrow{p} 0$  as  $t$  goes to  $\infty$ . We now turn to the second half of (3.48). The fact that  $b > 0$  entails that

$$\frac{m(Y_t)}{m(e^{bt})} = \exp\left(\int_{1/Y_t}^{e^{-bt}} \frac{\Phi(u)}{\Psi(u)} du\right) = \exp\{(H(\ln Y_t) - H(bt))(1 + o(1))\}. \quad (3.56)$$

Note that

$$H(\ln Y_t) - H(bt) = H(-\ln v_t(1/Y_t)) - H(bt) - \left(H(-\ln v_t(1/Y_t)) - H(\ln Y_t)\right). \quad (3.57)$$

Since  $b > 0$ , the convexity of  $\Psi$  and equation (2.8) imply that  $v_t(\lambda) \leq e^{-bt}\lambda$  for all  $t \geq 0, \lambda > 0$ . Then  $v_t(1/Y_t) \leq e^{-bt}$  for large enough  $t$  (depending on the path of  $Y$ ) as  $1/Y_t$  goes to 0 almost surely. Using again Condition (S) and (3.53), together with the monotonicity of  $H$ , there exists  $\theta'_t$  between  $bt$  and  $-\ln v_t(1/Y_t)$  such that

$$\begin{aligned} H(-\ln v_t(1/Y_t)) - H(bt) &= H'(\theta'_t)\theta'_t \ln\left(-\frac{\ln v_t(1/Y_t)}{bt}\right) \\ &= o(1) \ln\left(1 + \frac{\ln Y_t}{bt}\right) + o(1). \end{aligned}$$

Since  $(\ln Y_t)/bt \xrightarrow{p} 0$ , the above term converges to 0 in probability as  $t \rightarrow \infty$ . Together with (3.54), (3.56) and (3.57),  $\frac{m(Y_t)}{m(e^{bt})} \xrightarrow{d} \exp(-e_1)$ .

(ii) By Condition (L),

$$H(-\ln v_t(1/Y_t)) - H(\ln Y_t) = \int_{\ln Y_t}^{-\ln v_t(1/Y_t)} H'(u) du \sim a \int_{\ln Y_t}^{-\ln v_t(1/Y_t)} \frac{du}{u} \text{ as } t \rightarrow \infty.$$

Using (3.53), we obtain

$$\int_{\ln Y_t}^{-\ln v_t(1/Y_t)} \frac{du}{u} = \ln\left(-\frac{\ln v_t(1/Y_t)}{\ln Y_t}\right) = \ln\left(\frac{bt}{\ln Y_t} + 1\right) + o(1).$$

Then by (3.54),

$$\frac{\ln Y_t}{bt} \xrightarrow{d} (\exp(e_1/a) - 1)^{-1} \text{ as } t \rightarrow \infty,$$

which implies Theorem 13 (ii).

(iii) The mean value theorem for integrals shows that there is some  $\theta_t$  between  $\ln Y_t$  and  $-\ln v_t(1/Y_t)$  such that

$$\begin{aligned} H(-\ln v_t(1/Y_t)) - H(\ln Y_t) &= \int_{\ln Y_t}^{-\ln v_t(1/Y_t)} \frac{uH'(u)}{u} du \\ &= H'(\theta_t)\theta_t \ln\left(-\frac{\ln v_t(1/Y_t)}{\ln Y_t}\right). \end{aligned}$$

By Condition (F) and (3.54), we have that  $\ln v_t(1/Y_t) \sim -\ln Y_t$  in probability. Together with (3.53), we have

$$\frac{bt}{\ln Y_t} \xrightarrow{p} 0 \text{ as } t \rightarrow \infty. \quad (3.58)$$

Applying (3.39) and (3.42),

$$\int_{v_t(1/Y_t)}^{1/Y_t} \frac{\Phi(u)}{\Psi(u)} du = \int_{\varphi(1/Y_t)}^{\varphi(v_t(1/Y_t))} \Phi(g(u)) du = \int_{\varphi(1/Y_t)}^{t+\varphi(1/Y_t)} \Phi(g(u)) du. \quad (3.59)$$

Note that  $\Phi(g(u)) = bH'(-\ln g(u))$ . By the fact that  $-\ln g(u) \sim bu$  as  $u \rightarrow \infty$  (see Lemma 2), and Condition (F), we have that  $\Phi(g(u))$  is regularly varying at  $\infty$ ; see [Res08, Proposition 0.8-(iv)]. By changing variable and applying a mean value theorem, we develop the above display as follows:

$$\begin{aligned} \int_{v_t(1/Y_t)}^{1/Y_t} \frac{\Phi(u)}{\Psi(u)} du &= \varphi(1/Y_t) \int_1^{t/\varphi(1/Y_t)+1} \Phi(g(\varphi(1/Y_t)u)) du \\ &= t\Phi(g(\varphi(1/Y_t)\theta_t)) \quad (\text{for some } \theta_t \in (1, t/\varphi(1/Y_t) + 1)) \\ &= t\Phi(g(\varphi(1/Y_t))) \cdot \frac{\Phi(g(\varphi(1/Y_t)\theta_t))}{\Phi(g(\varphi(1/Y_t)))} \\ &\sim t\Phi(1/Y_t) \text{ almost-surely } (t \rightarrow \infty) \end{aligned}$$

The validity of the last equivalence is proved as follows: using (3.58), and  $b\varphi(\lambda) \sim -\ln \lambda$  as  $\lambda \rightarrow 0$  (see Lemma 2), we have  $t/\varphi(1/Y_t) \xrightarrow{p} 0$ , hence  $\theta_t \xrightarrow{p} 1$ , as  $t \rightarrow \infty$ . Then, since  $\Phi \circ g$  is regularly varying, the last equivalence holds by locally uniform convergence, see e.g. [BGT87, Theorem 1.2.1]. We then apply Theorem 6 to obtain that  $t\Phi(1/Y_t) \xrightarrow{d} e_1$ , which finishes the proof of the first half of (iii).

Now we focus on the case when  $\delta > 0$  in (3.44). Note that  $H'(x) = \frac{1}{b}\Phi(e^{-x})$ . Since  $h$  is the inverse function of  $1/H'$ , we have  $H'(h(x)) = 1/x$ , for any  $x > 0$ . Then by Karamata's theorem ([Res08, P.23]), the statement on  $h(t)$  in (3.50) holds true.

For  $\lambda > 0$ , we use again  $H'(x) = \frac{1}{b}\Phi(e^{-x})$ , and apply (3.44) and (3.45) to obtain that

$$\frac{H'(\ln Y_t)}{H'(h(bt)\lambda)} = t\Phi(1/Y_t) \frac{H'(h(bt))}{H'(h(bt)\lambda)} \xrightarrow{d} \lambda^\delta e_1 \text{ as } t \rightarrow \infty.$$

Hence,

$$\mathbb{P}(\ln Y_t/h(bt) \leq \lambda) = \mathbb{P}(H'(\ln Y_t)/H'(h(bt)\lambda) \geq 1) \xrightarrow[t \rightarrow \infty]{} \exp(-1/\lambda^\delta).$$

□

### 3.3.3 Supercritical case

Consider a supercritical CBI process  $(Y_t, t \geq 0)$ , i.e.  $b < 0$ . Recall  $v_{-t}(\lambda)$  defined by (2.11) for  $\lambda \in (0, \rho)$ . From Theorem 2-(ii), if (3.23) holds, then  $v_{-t}(\lambda)Y_t \xrightarrow{a.s.} \infty$  as  $t \rightarrow \infty$ . Let  $\rho_t := v_{-t}(\lambda_0)$  for some  $\lambda_0 \in (0, \rho)$ .

**Lemma 3.** *Assume  $b < 0$ . Then  $\ln v_t(\rho_t\lambda) \sim \ln \lambda$ , as  $t \rightarrow \infty$  and  $\lambda \rightarrow 0$ .*

*Proof.* Note that  $\Psi(u) < 0$  for  $u \in (0, \rho)$ . By (3.36),  $\varphi(\lambda) = -\int_\lambda^{\lambda_0} \frac{du}{\Psi(u)}$  for  $0 < \lambda < \lambda_0$ . By (3.18) and (3.19),

$$v_t(\rho_t\lambda) \xrightarrow[t \rightarrow \infty]{} v_{-\ln \lambda/b}(\lambda_0) \quad \text{and} \quad v_{-\ln \lambda/b}(\lambda_0) \xrightarrow[\lambda \rightarrow 0]{} 0.$$

Moreover, recall that  $\lambda \mapsto v_t(\rho_t\lambda)$  is non-decreasing and  $\lim_{t \rightarrow \infty} \rho_t = 0$ . Then for sufficiently large  $t$  and small enough  $\lambda$ , we have  $\rho_t\lambda, v_t(\rho_t\lambda) \in (0, \lambda_0)$ . By (3.39),

$$\int_{v_t(\rho_t\lambda)}^{\lambda_0} (-1/\Psi(u)) du = -t + \varphi(\rho_t\lambda),$$

which implies that

$$v_t(\rho_t\lambda) = g(\varphi(\rho_t\lambda) - t) \quad (3.60)$$

for large  $t$  and small  $\lambda$ .

By (2.10), we obtain

$$\begin{aligned} \varphi(\rho_t\lambda) - t &= \int_{\rho_t\lambda}^{\lambda_0} (-1/\Psi(u))du - \int_{\lambda_0}^{\rho_t} (1/\Psi(u))du \quad (\rho_t = v_{-t}(\lambda_0)) \\ &= \int_{\rho_t\lambda}^{\rho_t} (-1/\Psi(u))du \sim \int_{\rho_t\lambda}^{\rho_t} (-1/bu)du = \frac{\ln \lambda}{b}, \end{aligned}$$

as  $t \rightarrow \infty$  (hence  $\rho_t \rightarrow 0$ ). Similarly as proved in Lemma 2, if  $x \rightarrow \infty$ , we have  $\ln(g(x)) \sim bx$ . Putting  $x = \varphi(\rho_t\lambda) - t \sim \frac{\ln \lambda}{b}$  into (3.60), as  $t \rightarrow \infty$  and  $\lambda \rightarrow 0$ , we get  $\ln v_t(\rho_t\lambda) \sim \ln \lambda$ .  $\square$

**Theorem 15.** *Assume that  $b \in (-\infty, 0)$ .*

(i) *If Condition (S) holds, let  $m(x)$  be the same as in (3.48). Then*

$$-\frac{\ln Y_t}{bt} \xrightarrow{p} 1 \text{ and } m(\rho_t Y_t)/m(e^{-bt}) \xrightarrow{d} U \text{ as } t \rightarrow \infty,$$

where  $U$  is uniformly distributed on  $[0, 1]$ .

(ii) *If Condition (L) holds, then with  $U_L$  defined by (3.49)*

$$-\frac{\ln Y_t}{bt} \xrightarrow{d} 1 + U_L \text{ as } t \rightarrow \infty.$$

(iii) *If Condition (F) holds, then*

$$t\Phi(1/Y_t) \xrightarrow{d} e_1 \text{ as } t \rightarrow \infty.$$

*If in addition (3.44) holds with  $\delta > 0$ , then with  $U_F$  given by (3.51)*

$$h(t) = t^{1/\delta} L^*(t) \text{ and } \frac{\ln Y_t}{h(t)} \xrightarrow{d} U_F \text{ as } t \rightarrow \infty,$$

where  $L^*$  is slowly varying at  $\infty$ .

*Proof.* Recall Lemma 1 that  $\rho_{s+t}/\rho_s \rightarrow e^{bt}$  as  $s \rightarrow \infty$ . Consider the function  $\rho_{\ln s}$  on  $(1, \infty)$ . It is not hard to see that  $\rho_{\ln s}$  is regularly varying with index  $b$  at  $\infty$ . By Bingham *et al.* [BGT87, Proposition 1.3.6 (i)],  $\ln \rho_{\ln s} \sim b \ln s$  ( $s \rightarrow \infty$ ). Consequently

$$\ln \rho_s \sim bs, \quad \text{as } s \rightarrow \infty. \quad (3.61)$$

By Theorem 2-ii), we have  $\rho_t Y_t \rightarrow \infty$  a.s. as  $t$  goes to  $\infty$ . Then by Lemma 3 with  $\lambda = 1/(\rho_t Y_t)$ ,

$$\ln v_t(1/Y_t) \sim -\ln \rho_t Y_t. \quad (3.62)$$

Since  $Y_t \rightarrow \infty$  a.s. as  $t \rightarrow \infty$ , we have  $1/Y_t \leq \rho$  for large enough  $t$  (depending on the path of  $Y$ ) and thus  $v_t(1/Y_t) \geq 1/Y_t$ . Then using the same reasoning as to prove (3.54), we obtain

$$H(\ln Y_t) - H(-\ln v_t(1/Y_t)) \xrightarrow{d} e_1. \quad (3.63)$$

The procedure is the same as that of Theorem 13 and we present the arguments briefly below.

(i) By Condition (S) and (3.62), similarly to (3.55)

$$H(-\ln v_t(1/Y_t)) - H(\ln Y_t) = o(1) \ln \left( -\frac{\ln v_t(1/Y_t)}{\ln Y_t} \right) = o(1) \ln \left( \frac{\ln \rho_t}{\ln Y_t} + 1 \right) + o(1).$$

Combining (3.63) and (3.61), we have that

$$-bt/\ln Y_t \xrightarrow{p} 1, \quad \text{as } t \rightarrow \infty.$$

Using the above display and (3.62), similarly as in (3.55), we obtain

$$H(-bt) - H(\ln Y_t) \xrightarrow{p} 0, \quad H(-\ln v_t(1/Y_t)) - H(\ln \rho_t Y_t) \xrightarrow{p} 0, \quad \text{as } t \rightarrow \infty.$$

Therefore, as  $t \rightarrow \infty$

$$\begin{aligned} H(\ln \rho_t Y_t) - H(-bt) &= H(-\ln v_t(1/Y_t)) - H(\ln Y_t) \\ &\quad - (H(-bt) - H(\ln Y_t)) \\ &\quad - (H(-\ln v_t(1/Y_t)) - H(\ln \rho_t Y_t)) \xrightarrow{d} -e_1. \end{aligned}$$

(ii) In the same manner as in Theorem 13 (ii), we have

$$H(-\ln v_t(1/Y_t)) - H(\ln Y_t) \sim a \int_{\ln Y_t}^{-\ln v_t(1/Y_t)} \frac{du}{u} = a \ln \left( \frac{\ln \rho_t}{\ln Y_t} + 1 \right) + o(1) \text{ as } t \rightarrow \infty.$$

Then by (3.63), we have  $-\ln Y_t/\ln \rho_t \xrightarrow{d} (1 - e^{-e_1/a})^{-1}$  as  $t \rightarrow \infty$ . The result follows using (3.61).

(iii) Recall that  $\rho_t Y_t \rightarrow \infty$  as  $t$  goes to  $\infty$ . For  $t$  large enough (depending on the path of  $Y$ ),  $1/\rho_t Y_t < \lambda_0$  a.s, and by (3.60), we have  $\varphi(v_t(1/Y_t)) = \varphi(v_t(\rho_t/\rho_t Y_t)) = \varphi(1/Y_t) - t$ . Similarly as in (3.59),

$$\int_{v_t(1/Y_t)}^{1/Y_t} \frac{\Phi(u)}{\Psi(u)} du = \int_{\varphi(1/Y_t)-t}^{\varphi(1/Y_t)} \Phi(g(u)) du = \frac{1}{o(1)} \ln \left( \frac{\varphi(1/Y_t)}{\varphi(1/Y_t) - t} \right) \xrightarrow{d} e_1, \text{ as } t \rightarrow \infty$$

which implies that  $t/\varphi(1/Y_t) \xrightarrow{p} 0$  as  $t \rightarrow \infty$ . As in Theorem 13 (iii), we obtain  $t\Phi(1/Y_t) \xrightarrow{d} e_1$  as  $t \rightarrow \infty$ . Then the first half of (iii) is proved. The remaining part when (3.44) holds, is similar as in the subcritical case and therefore omitted.  $\square$

### 3.4 On the critical case

The study of the critical case, i.e.  $b = \Psi'(0+) = 0$ , is more involved as  $v_t(\lambda)$  may have different asymptotics as  $t \rightarrow \infty$  and  $\lambda \rightarrow 0$  according to the behavior of  $\Psi$  near 0. We make the following assumption on the Lévy measure  $\pi$  in the branching mechanism:

Suppose that  $\pi$  satisfies

$$\bar{\pi}(u) \underset{u \rightarrow \infty}{\sim} -\frac{1}{\Gamma(-\alpha)} u^{-1-\alpha} \ell(u), \quad (3.64)$$

where  $\bar{\pi}(u) = \pi(u, \infty)$  for  $u > 0$ ,  $0 < \alpha < 1$  and  $\ell$  is slowly varying at  $\infty$ . By [BGT87, Theorem 8.1.6], the above assumptions are equivalent to

$$\Psi(\lambda) \underset{\lambda \rightarrow 0}{\sim} \lambda^{1+\alpha} \ell(1/\lambda). \quad (3.65)$$

Recall that  $\varphi$  is defined by (3.36) and  $g$  is inverse function of  $\varphi$ . It follows from Karamata's theorem ([Res08, P.17 and P.23]) that

$$\varphi(\lambda) \sim \frac{\lambda^{-\alpha}}{\alpha \ell(1/\lambda)} \quad \text{and} \quad g(1/\lambda) \sim \lambda^{1/\alpha} \ell^*(1/\lambda), \quad \text{as } \lambda \rightarrow 0, \quad (3.66)$$

where  $\ell^*$  is slowly varying at  $\infty$ . We denote by  $\Phi^{-1}$  the inverse function of  $\Phi$ .

**Theorem 16.** *Assume that  $b = 0$  and (3.64) holds.*

(i) *If Condition (S) holds, let  $m(x)$  be defined as in (3.48). Then*

$$\frac{m(Y_t)}{m(1/g(t))} \xrightarrow{d} V \quad \text{as } t \rightarrow \infty,$$

where  $V$  is uniformly distributed on  $[0, 1]$ .

(ii) *If Condition (L) holds, then*

$$g(t)Y_t \xrightarrow{d} V_L \quad \text{as } t \rightarrow \infty,$$

where  $\mathbb{E}[e^{-\lambda V_L}] = (1 + \lambda^\alpha)^{-a}$ ,  $\forall \lambda \geq 0$ .

(iii) *If Condition (F) holds with  $\delta > 0$  in (3.44), then*

$$\varrho_t Y_t \xrightarrow{d} V_F \quad \text{as } t \rightarrow \infty \quad (3.67)$$

where  $\mathbb{E}[e^{-\theta V_F}] = \exp(-\theta^{\delta\alpha})$ , for all  $\theta \geq 0$ , with  $\varrho_t = \Phi^{-1}(1/t) \sim t^{-1/(\delta\alpha)} \bar{\ell}(t)$  as  $t \rightarrow \infty$  for some slowly varying function  $\bar{\ell}$  at  $\infty$ .

*If Condition (F) holds with  $\delta = 0$  in (3.44), then  $t\Phi(1/Y_t) \xrightarrow{d} e_1$  as  $t \rightarrow \infty$ .*

*Remark 17.* When  $\delta \in (0, 1]$ , the convergence (3.67) is equivalent to the following  $t\Phi(1/Y_t) \xrightarrow{d} V_F^{-\delta\alpha}$  as  $t$  goes to  $\infty$ . Indeed for any  $\lambda > 0$ , the property of regular variation of  $\varrho_t$  implies that

$$\begin{aligned} \mathbb{P}(t\Phi(1/Y_t) \geq \lambda) &= \mathbb{P}(\Phi(1/Y_t) \geq \lambda/t) = \mathbb{P}(1/Y_t \geq \varrho_t(\lambda/t)) \\ &= \mathbb{P}(\varrho_{t/\lambda} Y_t \leq 1) \sim \mathbb{P}(\lambda^{1/(\delta\alpha)} \varrho_t Y_t \leq 1) \xrightarrow{t \rightarrow \infty} \mathbb{P}(V_F \leq \lambda^{-1/(\delta\alpha)}) = \mathbb{P}((V_F)^{-\delta\alpha} \geq \lambda). \end{aligned}$$

The random variable  $V_F$  having a stable law,  $(V_F)^{-\delta\alpha}$  is not a standard exponential random variable. Therefore, unlike in the non-critical cases, see Theorem 13-(iii) and Theorem 15-(iii) for which

$$t\Phi(1/Y_t) \xrightarrow{d} e_1 \quad \text{as } t \rightarrow \infty$$

holds true whenever condition (F) holds, no matter what value  $\delta \in [0, 1]$  takes, in the critical case, the above convergence is only true for  $\delta = 0$ .

*Proof.* We start with some observations. By (3.42), we have

$$\int_{v_t(\lambda)}^{\lambda} \frac{\Phi(u)}{\Psi(u)} du = \int_{\varphi(\lambda)}^{t+\varphi(\lambda)} \Phi(g(u)) du, \quad H(x) = \int_{\varphi(\lambda_0)}^x \Phi(g(u)) du.$$

Then  $H'(x) = \Phi(g(x))$ . Note that we will use  $H'(x)$  or  $\Phi(g(x))$  in different contexts.

It was mentioned in Remark 8 that the steps 1 and 2 in the proof of Theorem 6 still hold for the critical case. That is to say; recalling  $r_t(\lambda)$  defined by (2.5) and  $\lambda \mapsto c_t(\lambda)$  its inverse; we have for any fixed  $\lambda > 0$ ,  $c_t(\lambda) \rightarrow 0$  and  $v_t(c_t(\lambda)) \rightarrow 0$  as  $t \rightarrow \infty$ . Therefore

$$\varphi(c_t(\lambda)) \rightarrow \infty \text{ as } t \rightarrow \infty. \quad (3.68)$$

- (i) By the mean value theorem and Condition (S), together with the above display and (3.42)

$$\begin{aligned} \lambda = r_t(c_t(\lambda)) &= \int_{v_t(c_t(\lambda))}^{c_t(\lambda)} \frac{\Phi(u)}{\Psi(u)} du = \int_{\varphi(c_t(\lambda))}^{t+\varphi(c_t(\lambda))} \Phi \circ g(u) du = \int_{\varphi(c_t(\lambda))}^{t+\varphi(c_t(\lambda))} H'(u) du \\ &= \int_{\varphi(c_t(\lambda))}^{t+\varphi(c_t(\lambda))} \frac{uH'(u)}{u} du = \theta_t H'(\theta_t) \ln\left(\frac{t+\varphi(c_t(\lambda))}{\varphi(c_t(\lambda))}\right) = o(1) \ln\left(\frac{t+\varphi(c_t(\lambda))}{\varphi(c_t(\lambda))}\right), \end{aligned}$$

where  $\theta_t$  is a real number in  $[\varphi(c_t(\lambda)), t+\varphi(c_t(\lambda))]$ . Consequently

$$\varphi(c_t(\lambda))/t \rightarrow 0 \text{ as } t \rightarrow \infty.$$

Since  $\varphi$  is regularly varying, see (3.66),

$$\varphi(\theta c_t(\lambda))/\varphi(c_t(\lambda)) = \theta^{-\alpha} \text{ for } \theta > 0 \text{ as } t \rightarrow \infty.$$

Then still by the mean value theorem and Condition (S), we obtain

$$\int_{v_t(c_t(\lambda))}^{v_t(\theta c_t(\lambda))} \frac{\Phi(u)}{\Psi(u)} du = \int_{t+\varphi(\theta c_t(\lambda))}^{t+\varphi(c_t(\lambda))} \Phi \circ g(u) du = o(1) \ln\left(\frac{t+\varphi(c_t(\lambda))}{t+\varphi(\theta c_t(\lambda))}\right), \quad (3.69)$$

which goes to 0 as  $t \rightarrow \infty$ . The same approach applies to obtain  $\int_{\theta c_t(\lambda)}^{c_t(\lambda)} \frac{\Phi(u)}{\Psi(u)} du \rightarrow 0$ . Hence

$$\int_{\theta c_t(\lambda)}^{v_t(\theta c_t(\lambda))} \frac{\Phi(u)}{\Psi(u)} du = \underbrace{\int_{\theta c_t(\lambda)}^{c_t(\lambda)} \frac{\Phi(u)}{\Psi(u)} du}_{\xrightarrow[t \rightarrow \infty]{} 0} + \underbrace{\int_{c_t(\lambda)}^{v_t(c_t(\lambda))} \frac{\Phi(u)}{\Psi(u)} du}_{-r_t(c_t(\lambda)) = -\lambda} + \underbrace{\int_{v_t(c_t(\lambda))}^{v_t(\theta c_t(\lambda))} \frac{\Phi(u)}{\Psi(u)} du}_{\xrightarrow[t \rightarrow \infty]{} 0}.$$

Similarly as Step 4 in the proof of Theorem 6, we have that

$$\int_{v_t(1/Y_t)}^{1/Y_t} \frac{\Phi(u)}{\Psi(u)} du = \int_{\varphi(1/Y_t)}^{t+\varphi(1/Y_t)} \Phi \circ g(u) du \xrightarrow{d} e_1 \text{ as } t \rightarrow \infty. \quad (3.70)$$

Applying the same transformation as in (3.69) to the above display entails that  $\varphi(1/Y_t)/t \xrightarrow{p} 0$  as  $t \rightarrow \infty$ . Using again Condition (S),

$$\int_{g(t)}^{v_t(1/Y_t)} \frac{\Phi(u)}{\Psi(u)} du = \int_{t+\varphi(1/Y_t)}^t \Phi \circ g(u) du = o(1) \ln\left(\frac{t}{t+\varphi(1/Y_t)}\right) \xrightarrow{p} 0. \quad (3.71)$$

Therefore combining the above two displays (3.70) and (3.71) yields

$$-\ln\left(\frac{m(Y_t)}{m(1/g(t))}\right) = \int_{g(t)}^{1/Y_t} \frac{\Phi(u)}{\Psi(u)} du \xrightarrow{d} e_1.$$

This allows one to conclude that (i) holds true.

(ii) By (2.6) and (3.42)

$$\begin{aligned}\mathbb{E}_x[e^{-\lambda g(t)Y_t}] &= \exp\left\{-xv_t(g(t)\lambda) - \int_{v_t(\lambda g(t))}^{\lambda g(t)} \frac{\Phi(u)}{\Psi(u)} du\right\} \\ &= \exp\left\{-xv_t(g(t)\lambda) - \int_{\varphi(\lambda g(t))}^{t+\varphi(\lambda g(t))} \Phi \circ g(u) du\right\}.\end{aligned}$$

We shall study the two terms in the exponential one by one. As  $t \rightarrow \infty$ ,  $g(t) \rightarrow 0$  and consequently  $v_t(g(t)\lambda) \rightarrow 0$ . By Condition (L),

$$\int_{\varphi(\lambda g(t))}^{t+\varphi(\lambda g(t))} \Phi \circ g(u) du \sim a \int_{\varphi(\lambda g(t))}^{t+\varphi(\lambda g(t))} \frac{dx}{x} = a \ln\left(\frac{\varphi(g(t)) + \varphi(\lambda g(t))}{\varphi(\lambda g(t))}\right),$$

which converges to  $a \ln(\lambda^\alpha + 1)$  by (3.66). So the statement in (ii) holds true.

(iii) Since  $uH'(u) = u\Phi(g(u)) \rightarrow \infty$  as  $u \rightarrow \infty$ , by applying the mean value theorem, we get

$$\lambda = r_t(c_t(\lambda)) = \int_{v_t(c_t(\lambda))}^{c_t(\lambda)} \frac{\Phi(u)}{\Psi(u)} du = \int_{\varphi(c_t(\lambda))}^{t+\varphi(c_t(\lambda))} \Phi \circ g(u) du = \frac{1}{o(1)} \ln\left(\frac{t + \varphi(c_t(\lambda))}{\varphi(c_t(\lambda))}\right).$$

This implies that

$$t/\varphi(c_t(\lambda)) \rightarrow 0 \text{ as } t \rightarrow \infty. \quad (3.72)$$

Since  $\varphi$  is regularly varying with index  $-\alpha$ , see (3.66),  $\varphi(\theta c_t(\lambda))/\varphi(c_t(\lambda)) = \theta^{-\alpha}$  for  $\theta > 0$  and thus  $t/\varphi(\theta c_t(\lambda)) \rightarrow 0$  as  $t \rightarrow \infty$ . Then for any  $\theta > 0$ ,

$$\int_{v_t(\theta c_t(\lambda))}^{\theta c_t(\lambda)} \frac{\Phi(u)}{\Psi(u)} du = \int_{\varphi(\theta c_t(\lambda))}^{t+\varphi(\theta c_t(\lambda))} H'(u) du = \varphi(\theta c_t(\lambda)) \int_1^{1+t/\varphi(\theta c_t(\lambda))} H'(\varphi(\theta c_t(\lambda))u) du.$$

By (3.44),  $H'$  (recall  $H' = \Phi \circ g$ ) is regularly varying. Based on the above equality, the mean value theorem yields that for any  $\theta > 0$ ,

$$\int_{v_t(\theta c_t(\lambda))}^{\theta c_t(\lambda)} \frac{\Phi(u)}{\Psi(u)} du \sim t\Phi \circ g(\varphi(\theta c_t(\lambda))) = t\Phi(\theta c_t(\lambda)), \text{ as } t \rightarrow \infty. \quad (3.73)$$

Letting  $\theta = 1$  and  $\lambda = 1$ , by the definition of  $r_t(1), c_t(1)$ , the left term in the equivalence relation above equals 1 and so we have

$$t\Phi(c_t(1)) \rightarrow 1 \text{ as } t \rightarrow \infty. \quad (3.74)$$

If (3.44) holds, then  $\Phi = H' \circ \varphi$  is regularly varying with index  $\delta\alpha$  at 0 by (3.66) and [Res08, Proposition 0.8-(iv)]. Using the above two displays,

$$\int_{v_t(\theta c_t(1))}^{\theta c_t(1)} \frac{\Phi(u)}{\Psi(u)} du \sim t\Phi(c_t(1)) \frac{\Phi(\theta c_t(1))}{\Phi(c_t(1))} \sim \theta^{\delta\alpha}, \quad \text{as } t \rightarrow \infty.$$

As  $c_t(1) \rightarrow 0$ , we also have  $v_t(\theta c_t(1)) \rightarrow 0$  thanks to the above display. Then using (3.28) and (3.29), we conclude that

$$\mathbb{E}_x[e^{-\theta c_t(1)Y_t}] \rightarrow e^{-\theta^{\delta\alpha}}, \text{ as } t \rightarrow \infty.$$

The map  $\Phi^{-1}$  is regularly varying with index  $1/(\delta\alpha)$  at 0 (see for instance [Res08, Proposition 0.8-(v)]). Note that by (3.74),

$$\Phi(c_t(1)) \sim 1/t \text{ as } t \rightarrow \infty.$$



The regular variation of  $\Phi^{-1}$  shows that  $c_t(1) = \Phi^{-1}(\Phi(c_t(1))) \sim \Phi^{-1}(1/t)$ . Thus we have the first result in (iii).

If  $\Phi \circ g$  is slowly varying at  $\infty$ , then  $\Phi \circ g(\varphi(u))$  is slowly varying at 0 by (3.66). It follows from (3.73) that

$$\int_{v_t(\theta c_t(\lambda))}^{\theta c_t(\lambda)} \frac{\Phi(u)}{\Psi(u)} du \sim t \bar{\Phi}(c_t(\lambda)) \frac{\Phi(\theta c_t(\lambda))}{\Phi(c_t(\lambda))} \sim \lambda \text{ as } t \rightarrow \infty,$$

which implies that  $\mathbb{E}_x[e^{-\theta c_t(\lambda) Y_t}] \rightarrow e^{-\lambda}$ . Similar to Step 4 in the proof of Theorem 6, we have that

$$\int_{v_t(1/Y_t)}^{1/Y_t} \frac{\Phi(u)}{\Psi(u)} du = \int_{\varphi(1/Y_t)}^{t+\varphi(1/Y_t)} \Phi \circ g(u) du \xrightarrow{d} e_1 \text{ as } t \rightarrow \infty.$$

Then the second half of (iii) follows the very similar proof of Theorem 13-(iii) and we omit it.  $\square$

*Remark 18.* Recall that in the critical case when  $\Phi = \Psi'$ , the CBI( $\Psi, \Phi$ ) process has the same law as the CB( $\Psi$ ) processes conditioned on the non-extinction. Moreover, we readily check that  $\int_0^{\Psi'(u)} \frac{\Psi'(u)}{\Psi(u)} du = \infty$ . It follows that  $H'(x) = \Phi(g(x)) = \Psi'(g(x))$  for all  $x$ .

We apply now Theorem 1 and our Theorem 16 to the case of stable branching and immigration mechanisms for which explicit calculations can be done.

**Corollary 2** (Stable case). *Assume  $\Psi(q) = dq^{\alpha+1}$  for  $d > 0$  and  $\alpha \in (0, 1]$  and  $\Phi(q) = d'q^\beta$  for  $d' > 0$  and  $\beta \in (0, 1]$ . Then,*

i) if  $\beta/\alpha > 1$  then

$$Y_t \xrightarrow{d} Y_\infty \text{ as } t \rightarrow \infty$$

where  $Y_\infty$  has Laplace transform  $\mathbb{E}[e^{-\lambda Y_\infty}] = e^{-\frac{\lambda^{\beta-\alpha}}{\beta-\alpha}}$  for any  $\lambda \geq 0$ ,

ii) if  $\beta/\alpha = 1$  then

$$t^{-\frac{1}{\alpha}} Y_t \xrightarrow{d} (\alpha d)^{\frac{1}{\alpha}} V_L \text{ as } t \rightarrow \infty$$

where  $V_L$  has Laplace transform  $\mathbb{E}[e^{-\lambda V_L}] = \frac{1}{(1+\lambda^\alpha)^{\frac{d'}{\alpha d}}}$  for any  $\lambda \geq 0$ ,

iii) if  $\beta/\alpha < 1$  then

$$t^{-\frac{1}{\beta}} Y_t \xrightarrow{d} (d')^{1/\beta} V_F \text{ as } t \rightarrow \infty,$$

where  $V_F$  has Laplace transform  $\mathbb{E}[e^{-\lambda V_F}] = e^{-\lambda^\beta}$  for any  $\lambda > 0$ .

*Remark 19.* When  $\beta = \alpha$ , the specific case  $d' = (\alpha + 1)d$  which corresponds to  $\Phi = \Psi'$ , has been studied by Kyprianou and Pardo [KP08, Lemma 3] with other techniques.

*Proof.* First notice that  $\int_0^{\Phi(u)} \frac{\Phi(u)}{\Psi(u)} du = \infty$  iff  $\frac{\beta}{\alpha} \leq 1$ . Statement i) is a direct consequence of Theorem 1. Let  $\lambda_0 = 1$ , we compute  $\varphi(x) = \frac{1}{\alpha d} \left( \frac{1}{x^\alpha} - 1 \right)$  and  $g(x) = \left( \frac{1}{\alpha d x + 1} \right)^{\frac{1}{\alpha}}$  for  $x > 0$ . Moreover  $x \Phi(g(x)) \sim \frac{d'}{(\alpha d)^{\beta/\alpha}} x^{1-\beta/\alpha}$  as  $x$  goes to  $\infty$ . If  $\beta/\alpha = 1$  then Condition (L) is fulfilled for  $a = \frac{d'}{\alpha d}$  and we deduce statement ii) from Theorem 16. If  $\beta/\alpha < 1$  then Condition (F) and (3.44) are fulfilled with  $\delta = \frac{\beta}{\alpha}$ . Theorem 16 also applies.  $\square$

We have focused our study on continuous-state branching processes with immigration whose branching Lévy measure has finite mean, i.e  $\Psi'(0+) = b \in (-\infty, 0)$  or equivalently  $\int_1^\infty z\pi(dz) < \infty$ . The proofs of Theorem 6 and Corollary 1 however do not make use of this assumption. They hold then also true in the case of a non-explosive supercritical CBI( $\Psi, \Phi$ ) with  $\Psi'(0+) = -\infty$ . A similar dichotomy occurs in the long-term behavior whether  $\int_0 \frac{\Phi(u)}{|\Psi(u)|} du < \infty$  or  $\int_0 \frac{\Phi(u)}{|\Psi(u)|} du = \infty$  when  $b = -\infty$ . In the first case, some almost-sure non-linear renormalizations can be found, see [Gre77] and Foucart and Ma [FM19] for the case without immigration. In the case  $\int_0 \frac{\Phi(u)}{|\Psi(u)|} du = \infty$ , similar regimes than those found in Section 3.3 exist, see [BP79] for the case of discrete state-space processes.

**Acknowledgement.** C.F is supported by the French National Research Agency (ANR): LABEX MME-DII (ANR11-LBX-0023-01). C.M is supported by the NSFC of China (11871032) and GXNSF (2018GXNSFAA050031). L.Y is supported by the NSFC of China (11801458).

## References

- [BAKP18] Mátyás Barczy, Mohamed Ben Alaya, Ahmed Kebaier, and Gyula Pap, Asymptotic properties of maximum likelihood estimator for the growth rate for a jump-type cir process based on continuous time observations, *Stochastic Processes and their Applications* **128** (2018), no. 4, 1135 – 1164.
- [Ber96] Jean Bertoin, Lévy processes, Cambridge Tracts in Mathematics, vol. 121, Cambridge University Press, Cambridge, 1996. MR 1406564 (98e:60117)
- [BGT87] N. H. Bingham, C. M. Goldie, and J. L. Teugels, Regular variation, Encyclopedia of mathematics and its applications, Cambridge University Press, Cambridge, Cambridgeshire, New York, 1987, Includes indexes.
- [Bin76] N.H. Bingham, Continuous branching processes and spectral positivity, *Stochastic Processes and their Applications* **4** (1976), no. 3, 217 – 242.
- [BP79] A. D. Barbour and Anthony G. Pakes, Limit theorems for the simple branching process allowing immigration. II. The case of infinite offspring mean, *Adv. in Appl. Probab.* **11** (1979), no. 1, 63–72. MR 517551 (80d:60114b)
- [CLP18] M. Chazal, R. Loeffen, and P. Patie, Smoothness of continuous state branching with immigration semigroups, *J. Math. Anal. Appl.* **459** (2018), no. 2, 619–660.
- [Coh77] Harry Cohn, On the convergence of the supercritical branching processes with immigration, *Journal of Applied Probability* **14** (1977), no. 2, 387–390.
- [DFM14] Xan Duhalde, Clément Foucart, and Chunhua Ma, On the hitting times of continuous-state branching processes with immigration, *Stochastic Process. Appl.* **124** (2014), no. 12, 4182–4201. MR 3264444
- [DFS03] D. Duffie, D. Filipović, and W. Schachermayer, Affine processes and applications in finance, *Ann. Appl. Probab.* **13** (2003), no. 3, 984–1053.
- [DL12] Donald A. Dawson and Zenghu Li, Stochastic equations, flows and measure-valued processes, *Ann. Probab.* **40** (2012), no. 2, 813–857. MR 2952093
- [DL13] Thomas Duquesne and Cyril Labbé, On the Eve property for CSBP, *EJP* (2013).

- [Dur10] R. Durrett, Probability: Theory and examples, 4th ed., Cambridge University Press, USA, 2010.
- [FF12] M. C. Fittipaldi and J. Fontbona, On SDE associated with continuous-state branching processes conditioned to never be extinct, *Electron. Commun. Probab.* **17** (2012), no. 49, 13. MR 2988395
- [FM19] Clément Foucart and Chunhua Ma, Continuous-state branching processes, extremal processes and super-individuals, *Ann. Inst. H. Poincaré Probab. Statist.* **55** (2019), no. 2, 1061–1086.
- [FUB14] Clément Foucart and Gerónimo Uribe Bravo, Local extinction in continuous-state branching processes with immigration, *Bernoulli* **20** (2014), no. 4, 1819–1844. MR 3263091
- [Gre74] D. R. Grey, Asymptotic behaviour of continuous time, continuous state-space branching processes, *J. Appl. Probability* **11** (1974), 669–677. MR 0408016 (53 #11783)
- [Gre77] ———, Almost-sure convergence in Markov branching process with infinite mean, *J. Appl. Probability* **14** (1977), 702–716. MR 0408016 (53 #11783)
- [Hea65] C. R. Heathcote, A branching process allowing immigration, *J. Roy. Statist. Soc. Ser. B* **27** (1965), 138–143. MR 0193680
- [Hey70] C. C. Heyde, Extension of a result of Seneta for the super-critical Galton-Watson process, *Ann. Math. Statist.* **41** (1970), no. 2, 739–742.
- [JMS17] Y. Jiao, C. Ma, and S. Scotti, Alpha-cir model with branching processes in sovereign interest rate modeling, *Finance Stoch* (2017), no. 21, 789–813.
- [KP08] A.E. Kyprianou and J.C. Pardo, Continuous-state branching processes and self-similarity, *Journal of Applied Probability* **45** (2008), no. 4, 1140–1160.
- [KRM12] Martin Keller-Ressel and Aleksandar Mijatović, On the limit distributions of continuous-state branching processes with immigration, *Stochastic Processes and their Applications* **122** (2012), no. 6, 2329 – 2345.
- [KW71] Kiyoshi Kawazu and Shinzo Watanabe, Branching processes with immigration and related limit theorems, *Teor. Verojatnost. i Primenen.* **16** (1971), 34–51. MR 0290475 (44 #7656)
- [Kyp14] Andreas E. Kyprianou, Fluctuations of Lévy processes with applications, second ed., Universitext, Springer, Heidelberg, 2014, Introductory lectures. MR 3155252
- [Lam07] Amaury Lambert, Quasi-stationary distributions and the continuous-state branching process conditioned to be never extinct, *Electron. J. Probab.* **12** (2007), no. 14, 420–446. MR 2299923 (2008b:60183)
- [Li00] Zeng-Hu Li, Asymptotic behaviour of continuous time and state branching processes, *Journal of the Australian Mathematical Society (Series A)* **68** (2000), 68–84.
- [Li11] Z. H. Li, Measure-valued branching Markov processes, Springer, 2011.
- [LM08] Z. Li and C. Ma, Catalytic discrete state branching models and related limit theorems, *J Theor Probab* (2008), no. 21, 936–965.

- [Pak79] Anthony G. Pakes, Limit theorems for the simple branching process allowing immigration. I. The case of finite offspring mean, *Adv. in Appl. Probab.* **11** (1979), no. 1, 31–62. MR 517550 (80d:60114a)
- [Pin72] Mark A. Pinsky, Limit theorems for continuous state branching processes with immigration, *Bull. Amer. Math. Soc.* **78** (1972), 242–244. MR 0295450 (45 #4516)
- [Res08] Sidney I. Resnick, Extreme values, regular variation and point processes, Springer series in operations research and financial engineering, Springer, New York, 2008, Applied probability, vol. 4.
- [Sen70] E. Seneta, On the supercritical Galton-Watson process with immigration, *Math. Biosci.* **7** (1970), 9–14. MR 0270460