

Journal Pre-proofs

Placental Growth Factor Measurements in the Assessment of Women with Suspected Preeclampsia: a Stratified Analysis of the PARROT Trial

Kate E. Duhig, Jenny E. Myers, Chris Gale, Joanna C. Girling, Kate Harding, Andrew Sharp, Nigel A.B. simpson, Derek Tuffnell, Paul T. Seed, Andrew H. Shennan, Lucy C. Chappell

PII: S2210-7789(20)30130-6
DOI: <https://doi.org/10.1016/j.preghy.2020.10.005>
Reference: PREGHY 784


To appear in: *Pregnancy Hypertension: An International Journal of Women's Cardiovascular Health*

Received Date: 10 July 2020
Accepted Date: 8 October 2020


Please cite this article as: Duhig, K.E., Myers, J.E., Gale, C., Girling, J.C., Harding, K., Sharp, A., simpson, N.A.B., Tuffnell, D., Seed, P.T., Shennan, A.H., Chappell, L.C., Placental Growth Factor Measurements in the Assessment of Women with Suspected Preeclampsia: a Stratified Analysis of the PARROT Trial, *Pregnancy Hypertension: An International Journal of Women's Cardiovascular Health* (2020), doi: <https://doi.org/10.1016/j.preghy.2020.10.005>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier B.V. on behalf of International Society for the Study of Hypertension in Pregnancy.



Strobe



Placental Growth Factor Measurements in the Assessment of Women with Suspected Preeclampsia: a Stratified Analysis of the PARROT Trial

Kate E DUHIG, PhD^a Clinical Research Fellow

Jenny E MYERS, PhD^b Professor of Obstetrics

Chris GALE, PhD^c Reader in Neonatal Medicine

Joanna C GIRLING, MA, MRCP^d Consultant Obstetrician & Gynaecologist/ Honorary Senior Lecturer

Kate HARDING, MBBS^e Consultant Obstetrician and Head of Obstetric Services

Andrew SHARP, PhD^f Senior Lecturer in Obstetrics and Gynaecology

Nigel A B SIMPSON, MBBS^g Senior Lecturer in Obstetrics and Gynaecology

Derek TUFFNELL, FRCOG^h Professor of Obstetrics

Mr Paul T SEED, CStat¹, Senior Lecturer in Medical Statistics

Andrew H SHENNAN, MD¹ Professor of Obstetrics

Lucy C CHAPPELL, PhD¹ NIHR Research Professor in Obstetrics

^aDepartment of Women and Children's Health, School of Life Course Sciences, King's College London

(KED, PTS, AHS, LCC). Kate.duhig@kcl.ac.uk; paul.seed@kcl.ac.uk; Andrew.shennan@kcl.ac.uk;

lucy.chappell@kcl.ac.uk

^bThe Division of Developmental Biology and Medicine, University of Manchester.

Jenny.myers@manchester.ac.uk

^c Neonatal Medicine, School of Public Health, Faculty of Medicine, Chelsea and Westminster

Campus, Imperial College London. Christopher.gale@imperial.ac.uk

^dWest Middlesex University Hospital, Chelsea and Westminster Hospital NHS Foundation Trust.

Joanna.girling@chelwest.nhs.uk

^e Guy's and St Thomas' NHS Foundation Trust. Kate.harding@gstt.nhs.uk

^fUniversity of Liverpool and Liverpool Women's Hospital, members of Liverpool Health Partners.

A.sharp@liverpool.ac.uk

^g Department of Women's and Children's Health, Faculty of Medicine and Health, University of

Leeds. n.a.b.simpson@leeds.ac.uk

^h Bradford Institute for Health Research. Derek.Tuffnell@bthft.nhs.uk

This research was supported by grants from the National Institute for Health Research, Research for Patient Benefit Programme (PB-PG-0214-33054) and National Institute for Health Research Professorship (Chappell RP-2014-05-019). The views expressed in this publication are those of the author(s) and not necessarily those of the NHS, the National Institute for Health Research or the Department of Health. PS is funded in part by Tommy's (registered charity number 1060508) and by the Collaboration for Leadership in Applied Health Research and Care South London (National Institute for Health Research). The funders had no involvement in the study design, collection and

analysis of data, data interpretation report writing or the decision to submit the article for publication.

Request for reprints to: Dr Kate Duhig, Department of Women and Children's Health, School of Life Course Sciences, King's College London, Westminster Bridge Road, London, SE1 7EH;

kate.duhig@kcl.ac.uk

The authors report no conflict of interest.

Journal Pre-proofs

Abstract

Objective: Placental growth factor testing decreases time to recognition of preeclampsia and may reduce severe maternal adverse outcomes. This analysis aims to describe the clinical phenotype of women by PIGF concentration, and to determine the mechanism(s) underpinning the reduction in severe maternal adverse outcomes in the PARROT trial, in order to inform how PIGF testing may be optimally used within clinical management algorithms.

Study Design: This was a planned secondary analysis from the PARROT trial that compared revealed PIGF testing and management guidance with usual care in the assessment of women with suspected preterm pre-eclampsia.

Main Outcome Measures: maternal and perinatal outcomes following stratification of women by trial group, and measured PIGF concentration.

Results: 1006 women were included. PIGF <100pg/ml identified women with more marked hypertension, increased adverse maternal outcomes and preterm delivery rates, and higher rates of small for gestational age infants. There was a reduction in adverse maternal outcomes in women whose results were revealed when PIGF levels were 12-100pg/ml compared to usual care (3.8% vs 6.9%; aOR 0.15(95% CI 0.03-0.92). There was no significant difference in gestation at delivery between concealed or revealed groups in any PIGF categories.

Conclusion: Low PIGF concentrations are associated with severe preeclampsia. The reduction in severe adverse maternal outcomes may be mediated through quicker diagnosis and intensive surveillance, as recommended by the management algorithm for those at increased risk. PIGF is particularly beneficial in those who test 12-100pg/ml, as these may be women with silent multi-organ disease who otherwise may go undetected.

Keywords: preeclampsia, PIGF, hypertension in pregnancy, diagnostic testing

Abbreviations: PIGF; placental growth factor, NICE; National Institute of Health and Care Excellence,

Introduction

Preeclampsia complicates around 3% of singleton pregnancies, with hypertension affecting 10% of pregnant women.¹⁻³ Preeclampsia is associated with a high risk of pregnancy complications including iatrogenic preterm birth, maternal and perinatal morbidity, and perinatal mortality.⁴⁻⁷

The placenta plays a central role in the pathogenesis of preeclampsia. Studies of placentally-derived angiogenic factors, such as Placental Growth Factor (PlGF) and soluble fms-like tyrosine kinase-1 (sFlt-1) have led to their development as adjuncts to diagnosis and prognosis.^{8,9} Evidence from prospective cohort studies has shown that angiogenic factors have good test performance for identifying preterm preeclampsia.^{10 11} These studies included women in whom angiogenic factor concentrations were concealed to carers. A recent randomised trial (PARROT) of 1023 women evaluated revealed PlGF measurement with a clinical management algorithm against usual care, forming one of the largest studies of angiogenic factors in the management of suspected preterm preeclampsia. In this trial there was a clinically important reduction in time to diagnosis of preeclampsia with a concurrent reduction seen in severe maternal adverse outcomes with revealed PlGF testing.¹²

The aim of this secondary analysis of the PARROT Trial was to describe clinical phenotypes of pregnancies by measured PlGF concentration. The analysis also assesses how PlGF measurement may have impacted on clinical outcomes, to inform understanding of the mechanism of benefit. We sought to determine how PlGF testing may be optimally used within clinical management algorithms, by evaluating effect of PlGF testing across women categorised by their PlGF level. We focussed statistical testing on mechanistic questions related to how revealing an abnormal result might drive change in processes or pathways of care.

Methods

This was a planned secondary analysis of the PARROT trial, a multicentre stepped wedge cluster randomised controlled trial (ISRCTN 16842031), approved by the London South East Research Ethics Committee (15/LO/2058). Women were recruited from 11 centres with singleton pregnancies and a live fetus from 20⁺⁰ to 36⁺⁶ weeks' gestation with suspected preeclampsia. Suspected preeclampsia was defined as new onset or worsening of existing hypertension, proteinuria, epigastric or right upper quadrant pain, headache with visual disturbances, altered maternal biochemistry or fetal growth restriction. Women were excluded if they had a confirmed diagnosis of preeclampsia at presentation. Randomisation was to intervention or control groups, and this occurred at cluster level, in a stepped wedge design.

Women in the control group received usual care following local hospital practice based on 2010 National Institute of Health and Care Excellence (NICE) guidance on the management of hypertension in pregnancy,¹³ with an additional blood sample taken for concealed PIGF testing. National guidance included treatment with oral labetalol, nifedipine or methyldopa if above the blood pressure target range, a blood pressure target of less than 150/100 mmHg on treatment, twice weekly blood pressure and urine checks in women with gestational hypertension, admission to hospital and delivery at 37 weeks' gestation for those with preeclampsia. Women in the intervention group received revealed PIGF testing integrated into a standard clinical management algorithm based upon national guidance (figure S1). Women were individually consented to participation in the trial. A single PIGF blood sample was taken from each woman at presentation.

All blood samples were processed at each unit on a Triage (Alere, San Diego, CA, now Quidel Cardiovascular Inc., San Diego, CA) instrument. Strict logs of quality control assessment were kept. Staff performing the assay were trained by the trial team and followed a standard operating procedure. The test is CE-Marked and is approved for use in countries recognising the CE Mark. This

clinical study was conducted in the European Union (U.K.) and investigated a clinical application approved under the product's CE Mark.

Outcomes

Outcomes were collected until the primary postnatal discharge of the woman and infant pair from secondary care services. The primary outcome for the PARROT trial was the time from presentation with suspected preeclampsia to having a diagnosis of preeclampsia documented in the clinical notes. Maternal outcomes for this planned secondary analysis were a composite of severe maternal adverse outcomes as defined by the fullPIERS consensus,¹⁴ systolic blood pressure ≥ 160 mmHg, progression to severe preeclampsia (independently adjudicated),¹⁵ placental abruption, mode and onset of delivery, use of medication, proportion of women reaching the diagnostic criteria (irrespective of clinical documentation) for preeclampsia¹⁶. These outcomes matched those used for the primary trial analysis.

Perinatal outcomes included gestation at delivery, preterm birth below 37 weeks' gestation, birthweight and birthweight centile,¹⁷ a composite of severe perinatal adverse outcomes (number of babies with one or more of the following: intraventricular haemorrhage, seizures, retinopathy of prematurity, respiratory distress syndrome, bronchopulmonary dysplasia, necrotising enterocolitis stage 2 and 3,¹⁸ stillbirth, early neonatal death and late neonatal death to 28 days), neonatal unit admission, perinatal death (stillbirths from 24 weeks' gestation to deaths up to seven completed days after birth) and late neonatal death (deaths between 8 and 27 completed days of life).

Sample Size

The sample size was determined for the main PARROT trial.¹² All women who participated in the trial who had a concealed or revealed PIGF result, and outcome data were included in this planned secondary analysis.

Statistical Analysis

Women were stratified by their measured PIGF concentration into the following predetermined groups: PIGF ≥ 100 pg/ml – determined as ‘normal’; PIGF 12-99pg/ml, equivalent to $<5^{\text{th}}$ centile for gestation and determined as ‘low’; PIGF < 12 pg/ml, the lowest limit of detection for the assay and determined as ‘very low’.

These categorical groups were used based on the evidence that ‘low’ PIGF has a high diagnostic accuracy (0.96; 95% confidence interval, 0.89–0.99) and negative predictive value (0.98; 0.93–0.995) of determining preeclampsia requiring delivery in 14 days in prospective observational cohort studies¹¹, and ‘very low’ PIGF is the lowest limit of detection of the assay. We have previously reported that a fixed PIGF threshold of < 100 pg/mL predicted preeclampsia requiring delivery within 14 days or before 37 weeks’ gestation (whichever was sooner) with sensitivity and negative predictive values similar to diagnostic accuracy estimates obtained by using a $< 5^{\text{th}}$ centile cut-off.¹¹ The data were analysed according to their measured PIGF group. To describe clinical phenotype by measured PIGF level, demographic data are presented in the concealed testing group only. We compared how outcomes were influenced by trial arm in each subgroup in order to determine which groups of women benefited in our primary trial, and elucidate how this was achieved.

Outcomes were adjusted for centre and categorical time effects because of the trial design. Effects were estimated using multiple regression including terms for the intervention with fixed effects using dummy variables at each time in each centre. Centre was considered as a categorical variable and fitted as separate dummy variables for each centre. Calendar time was treated as a single categorical time variable. Continuous outcomes were assessed by linear regression. All binary outcomes were analysed using a binomial regression model with a log link. Test performance was evaluated using sensitivity, specificity, positive and negative predictive values and positive and negative likelihood ratios and area under the receiver operating characteristic curves. Mixed effects

log-normal regression curves were generated for the proportion of women diagnosed relative to time from trial entry.

Results

1006 women were included in this secondary analysis: 435 in the usual care group, and 571 in the revealed group (Figure 1). The unequal size of the trial groups was due to the stepped wedge design, such that recruitment increased overall as the trial continued. There was no contamination between trial groups. Among the participants, 236 (23.5%) had PIGF <12pg/ml, 385 (38.3%) had PIGF 12-100pg/ml, and 385 (38.3%) had PIGF >100pg/ml.

Clinical characteristics by PIGF category

In the concealed PIGF <12pg/ml category, 66% of women received a final diagnosis of preeclampsia. The mean highest systolic blood pressure in the 48 hours prior to trial entry in all women was 150 (17) mmHg. The median gestation at delivery was 34.4 weeks, and 59% of the participants delivered within 14 days of enrolment to the trial. Of the babies born to women in the PIGF <12pg/ml category, 46% had a birthweight of <10th centile.

In the concealed PIGF 12-100pg/ml category, 37% of women received a diagnosis of preeclampsia. The mean highest systolic blood pressure in the 48 hours prior to trial entry was 144 (19) mmHg. The median gestation at delivery was 37.4 weeks, with 43% of participants delivered within 14 days of enrolment in the trial. 20% of the babies born to women in the PIGF 12-100pg/ml category had a birthweight of <10th centile.

In the concealed PIGF >100pg/ml category, 12% of women received a final diagnosis of preeclampsia. The mean highest systolic blood pressure in the 48 hours prior to trial entry was 136 (21) mmHg. The median gestation at delivery was 38.2 weeks, with 8% of participants delivered

within 14 days of enrolment in the trial. 9% of the babies born to women in the PIGF >100pg/ml category had a birthweight of <10th centile. Further demographic details and corresponding values for the revealed group are presented in table 1 and table S1.

Diagnosis of Preeclampsia

The proportion of women diagnosed with preeclampsia was not significantly different between the intervention (revealed) and usual care (concealed) in any of the PIGF categories (74% vs 66% for PIGF <12pg/ml, 40% vs 37% for PIGF 12-100pg/ml, and 12% vs 10% for PIGF >100pg/ml) (Figure 2).

Time to diagnosis of preeclampsia was lower in the revealed PIGF testing group (1.9 days) compared to usual care (4.1 days) across all three PIGF groups (time ratio 0.36, 95% CI 0.15–0.87; $p=0.027$).

Within PIGF categories, time to diagnosis in the revealed testing group vs. the concealed testing group was 1.0 vs 2.0 days (adjusted time ratio 0.17 (95% CI 0.03-1.06)) for PIGF <12pg/ml; 2.0 vs 4.6 days (adjusted time ratio 0.66 (95% CI 0.09-4.95)) for PIGF 12-100pg/ml, and 22.8 vs 30.3 days (adjusted time ratio 0.13 (95% CI 0.16-1.07)) for PIGF >100pg/ml (Table 2). Figure S2 shows the mixed-effects lognormal regression curves with the proportion of women diagnosed by time from trial entry and differences in days (A), and weeks (B) with revealed PIGF testing in those women with PIGF <12pg/ml and PIGF 12-100pg/ml, showing shortened time to diagnosis in both PIGF <12pg/ml or PIGF 12-100pg/ml categories.

Maternal Outcomes

Severe maternal adverse outcomes were less frequent with revealed PIGF testing than with usual care overall (22/573 (3.8%) versus 24/446 (5.4%); adjusted OR (aOR) 0.32, 95%CI 0.11-0.95). This was significant in women with PIGF 12-100pg/ml (3.8% vs 6.9%; aOR 0.15 (95% CI 0.03-0.92)) (Table 2, Table S2). There were no significant differences seen in mean systolic or diastolic blood pressure, or the use of magnesium sulfate in the revealed compared to concealed groups in any of the PIGF

categories (table 1). There was an increase seen in the use of antihypertensive medication in the intervention groups versus the usual care group in women with PIGF <12pg/ml (83.1% vs 74.5%; aOR 3.85 (95% CI 1.03 to 8.28). There were no differences seen in the number of antenatal ultrasound scans, vaginal deliveries, or elective or emergency caesarean section rates between the intervention or usual care groups in any of the PIGF categories.

Time to Delivery and Steroid Administration

PIGF categorisation stratified by time to delivery is shown in Figure 3; those with PIGF <12pg/ml, and <100pg/ml had consistently shorter times to delivery when compared to PIGF >100pg/ml.

In women who delivered <35 weeks' gestation, antenatal corticosteroids were given within the seven days prior to delivery in 39% (29/75) of the intervention group vs 16% (6/38) of the control group in women with PIGF <12pg/ml, and in 37.5% (12/32) and 26% (5/19) respectively in women with PIGF 12-100pg/ml.).

Perinatal Outcomes

There was no evidence of a difference significant difference in gestation at delivery, or perinatal adverse outcome rates with the intervention versus usual care in any of the PIGF categories (Table 3, Table S3). The difference in gestational age between the intervention and usual care in the <12pg/ml category was not significant (mean difference -0.03 weeks; -1.72 to 1.66). There were no significant differences in preterm delivery rates (<37 weeks' gestation), or birthweight centiles between the intervention and usual care in any of the PIGF categories.

Discussion

In one of the largest studies of angiogenetic markers for the assessment of women with suspected preterm preeclampsia to date, we have confirmed that in a real-world setting, low and very low PIGF

categories accurately identified women with a phenotype of more severe preeclampsia. Women with low and very low PIGF concentrations have more marked hypertension, a greater number of adverse maternal outcomes, a shorter time to delivery interval and an increased need for preterm delivery, and higher rates of small for gestational age infants when compared with women with normal PIGF concentrations. Women with normal PIGF results have longer time to delivery intervals and rates of small for gestational age infants consistent with the general pregnant population.

PIGF testing does not lead to significantly more cases of preeclampsia being diagnosed, but consistently shortens the time it takes for a clinician to make a diagnosis across all three categories of PIGF. After adjustment for baseline characteristics, gestational age at delivery was not significantly different between the groups. PIGF testing did not appear to cause a significant difference in gestation at delivery by causing or preventing a non-indicated intervention through knowledge of the result.

Despite initial antenatal visit characteristics being very similar across all groups, women with very low PIGF concentrations had the most severe clinical phenotype of preeclampsia at entry to our study. However, whilst women with low PIGF concentrations appear to have an intermediate-risk phenotype of preeclampsia, they remain at increased risk of experiencing severe adverse outcomes compared to those with normal PIGF. One of the aims of stratification was to explore the mechanism(s) underlying the reduction seen in severe maternal adverse events with implementation of revealed PIGF testing. We found that the difference seen in the severe maternal adverse outcome composite was most marked in the PIGF 12-100pg/ml group (aOR 0.15 (95% CI 0.03 to 0.92) and we anticipate that this may offer clinicians an opportunity to identify women at risk of developing severe preeclampsia complications, who may otherwise be considered at lower risk.

The improvement in clinical outcomes in this group may have been mediated by the use of the clinical management algorithm, which recommends increasing antenatal surveillance. Given that the proportion of women receiving a diagnosis is not increased with revealed PIGF, but that a diagnosis is made sooner after presentation, it would be reasonable to hypothesise that the mechanism for this reduction is mediated through increased surveillance and monitoring as recommended by the trial management algorithm. This may be particularly important in the group of women with PIGF 12-100pg/ml who presented with clinical features of gestational hypertension but may also have had sub-clinical multi-organ disease features.

In this study we reiterated that a low or very low PIGF was not an indication for delivery in itself as highlighted by previous studies.¹⁹ In the PIGF <12 mg/ml group, women in the revealed group appear to deliver around one week earlier than those in the masked group (33.4 vs 34.4 weeks), but after pre-specified adjustment for baseline characteristics, this was not significant. It is also possible that within each PIGF category, those who needed earlier delivery were appropriately managed, and those clinically well were monitored, improving outcomes but not significantly changing overall preterm birth rates. However, whilst implementation of revealed PIGF testing does not significantly alter gestation at delivery between the two trial groups overall, we cannot exclude a difference in increasing preterm birth in the very low PIGF category that we were underpowered to demonstrate in this study. The results of the PREPARE study, which aims to determine if the use of sFlt/PIGF and a risk stratification algorithm reduces preterm delivery rates, are awaited.²⁰

Whilst the algorithm did not recommend routine admission for women with low or very low PIGF, and made no recommendations regarding steroid administration or timing of delivery which was left to the discretion of the treating clinicians, we hypothesise that low PIGF may have acted as an early warning sign of impending complications, giving clinicians the opportunity to act accordingly. The finding of the INSPIRE trial, in which women with suspected preeclampsia were individually

randomised to revealed or concealed sFlt-1/PlGF testing demonstrated similar results, that the clinical use of PlGF/sFlt-1 testing enabled more accurate admission rates of high-risk patients without changing admission rates overall, and improved identification of those with preeclampsia.²¹

There was a high prevalence of respiratory distress requiring NNU admission (nearly 30%) among babies with a PlGF <12pg/ml, but this was driven by gestational age at delivery. Of those women who delivered <35 weeks' gestation, revealed PlGF testing was associated with improvements in antenatal steroid administration within the seven days prior to delivery. Overall, 17% more women in the intervention (revealed) group received steroids within the seven days prior to delivery in those delivering <35 weeks' gestation, demonstrating that PlGF may be clinically useful in assisting with the timing of steroid administration.

Given that PARROT was a national, multicentre trial, we would anticipate that the prevalence of disease seen in the trial population would be similar in women presenting with suspected preterm preeclampsia to maternity triage settings throughout the UK. This would support the generalisability of these findings to the wider UK population.

A particular strength of our study is that these analyses focussed on identifying how the use of PlGF impacts on patient management pathways to influence important patient outcomes. This was a large multicentre study evaluating PlGF testing in a pragmatic way to achieve maximum external validity. The Patient Centred Outcomes Research Institute recommends the evaluation of process of care outcomes alongside morbidity outcomes in the evaluation of novel diagnostic tests.²² It is known that effectiveness trials (i.e. in a real-world clinical care setting) can assess the overall performance of an intervention, but that it can be difficult to identify the exact processes that explain the effectiveness of an intervention, due their pragmatic nature.²³ Our cost effectiveness analysis has been previously reported. The resource use data showed that PlGF was overall cost

saving, with an increase in antenatal inpatient costs for those with abnormal PIGF alongside a reduction in outpatient attendances in those with a normal result, suggesting improved risk stratification with PIGF testing.²⁴ As we did not undertake a more detailed process evaluation, the exact components of changes in the antenatal care pathways that contributed to the reduction in severe maternal adverse outcomes may remain unclear. However, this is balanced by the results being considerably more generalisable than if the trial had been undertaken with a very prescriptive management algorithm and multiple checkpoints such that the effect of the intervention might have required these additional components. Finding a significant effect size with a pragmatic algorithm suggests that clinicians found the intervention easy to integrate into their clinical care.

Stratification of the women into six groups based on PIGF concentrations and treatment allocation has created smaller numbers in each comparison group, meaning we may be underpowered to demonstrate important differences in care. This was a planned secondary analysis of an existing trial dataset, and as such the interpretation of the results should be circumspect.

Previous comparative analyses of concealed versus revealed PIGF testing have demonstrated a reduction in perinatal deaths, but these analyses have been between two separate cohort studies with differing inclusion criteria, with a mixture of revealed and concealed testing.²⁵ This trial showed no difference in perinatal deaths with revealed testing; we anticipate that one reason for this is that 55% of the stillbirths in our trial occurred in pre-viable babies (<24 weeks' gestation and <500 grams), where intervention to influence outcome is limited. It may be that in order to prevent viable stillbirth, repeated PIGF testing is needed alongside ultrasound scanning as a means of disease monitoring, in order that interventions (including delivery) can be implemented in a timely manner in those babies at greatest risk of stillbirth. Further research is needed to determine the optimal frequency of repeat testing and to evaluate the impact on perinatal outcomes.

Conclusions

This analysis has shown that the use of revealed PIGF testing with appropriate risk stratification of women, particularly in those with low or very low PIGF, can prevent serious maternal adverse outcomes. PIGF is beneficial in identifying women with a phenotype indicative of placentally-driven disease, particularly those who test 12-100pg/ml, in whom silent multi-organ disease may otherwise go undetected.

Competing interests

None

Journal Pre-proofs

Figure Legends

Figure 1. STROBE Diagram showing participant flow

Figure 2. Final Diagnoses for Women in the PARROT Trial, stratified by PIGF category.

Figure 3: Time to Delivery (Median, IQR) stratified by PIGF concentration for all participants. Red line indicates PIGF <12pg/ml; orange line, PIGF 12-100pg/ml; green line, PIGF >100pg/ml.

Journal Pre-proofs

Figure 1






Figure 2



	Revealed PIGF <12pg/ml N = 130	Concealed PIGF <12pg/ml N = 106	Revealed PIGF 12-100pg/ml N = 212	Concealed PIGF 12-100pg/ml N = 173	Revealed PIGF >100pg/ml N = 229	Concealed PIGF >100pg/ml N = 156
Preeclampsia, n (%)	112 (86.1%)	83 (78.8%)	108 (50.8%)	83 (48%)	33 (14.5%)	29 (18.6%)
Other complications*, n (%)	18 (13.8%)	18 (17.0%)	78 (36.8%)	70 (40.5%)	120 (52.4%)	74 (47.4%)
Normal, n (%)	0 (0.0%)	5 (4.7%)	26 (12.3%)	20 (11.6%)	76 (33.2%)	53 (34.0%)

*Includes chronic hypertension (CHT), gestational hypertension, gestational proteinuria, small for gestational age (SGA), CHT with SGA, chronic kidney disease (CKD)

Figure 3



Interval from test to delivery (days) Median (IQR)	Gestational Age at Testing
	<35 weeks
	PIGF Revealed
Very Low PIGF (<12 pg/ml)	12 (6-22)
Low PIGF (12-100 pg/ml)	26 (16-36)
Normal PIGF (>100 pg/ml)	50 (32-75)
	Usual Care
Very Low PIGF (<12 pg/ml)	17 (7-25)
Low PIGF (12-100 pg/ml)	27 (18-35)
Normal PIGF (>100 pg/ml)	50 (35-76)
	35⁺⁰ to 36⁺⁶ weeks
	PIGF Revealed
Very Low PIGF (<12 pg/ml)	4 (2-8)
Low PIGF (12-100 pg/ml)	13 (7-18)
Normal PIGF (>100 pg/ml)	20 (13-28)
	Usual Care
Very Low PIGF (<12 pg/ml)	8 (5-12)
Low PIGF (12-100 pg/ml)	11 (4-18)
Normal PIGF (>100 pg/ml)	21 (11-28)

Table 1: Clinical Characteristics

	Revealed PIGF <12pg/ml N = 130	Concealed PIGF <12pg/ml N = 106	Revealed PIGF 12-100pg/ml N = 212	Concealed PIGF 12-100pg/ml N = 173	Revealed PIGF >100pg/ml N= 229	Concealed PIGF >100pg/ml N= 156
Age (years) Mean (SD)	31.6 (6.0)	31.0 (6.1)	32.3 (6.0)	32.1 (5.7)	31.7 (5.8)	31.2 (6.4)
Blood pressure at booking (mmHg)						
Systolic mean (SD)	118 (15)	116 (15)	121 (14)	121 (16)	120 (15)	121 (16)
Diastolic mean (SD)	73 (11)	72 (11)	74 (10)	75 (12)	74 (11)	75 (12)
Gestation at enrolment, weeks, median (IQR)	32.3 (28.7,34.3)	34.1 (29.1,35.9)	34.6 (32.4,35.9)	35.1 (33.1,36.1)	32.6 (29.1,34.9)	32.3 (28.9,34.7)
Primiparous (%)	84 (64.6%)	62 (58.5%)	114 (53.8%)	85 (49.1%)	114 (49.8%)	58 (37.2%)
Pre-existing chronic hypertension	20 (15.4%)	15 (14.2%)	29 (13.7%)	28 (16.2%)	37 (16.2%)	25 (16.0%)
Previous preeclampsia (%)	24 (18.5%)	23 (21.7%)	42 (19.8%)	34 (19.7%)	33 (14.4%)	33 (21.2%)
Highest blood pressure in 48 hours prior to study entry (mmHg)						
Systolic mean (SD)	153 (18)	150 (17)	146 (17)	144 (19)	136 (20)	136 (21)
Diastolic mean (SD)	97 (12)	97 (10)	93 (12)	93 (12)	84 (13)	85 (14)
Time to diagnosis of preeclampsia (for those diagnosed) (days)						
Median (IQR)	1.0 (0.3, 4.5)	2.0 (0.3, 9.0)	2.0 (0.9, 8.70)	4.6 (1.0, 14.5)	22.8 (8.4, 39.2)	30.3 (5.9, 65.1)
Effect size (time ratio (95%CI))		0.17 (0.03 – 1.06)		0.66 (0.09-4.95)		0.13 (0.16-1.07)
Number of women diagnosed with preeclampsia n (%)	96 (73.8%)	70 (66.0%)	84 (39.6%)	64 (37.4%)	23 (10.1%)	19 (12.3%)
Severe preeclampsia (ACOG definition) n women (%)	73 (56.2%)	49 (46.2%)	64 (30.2%)	49 (28.3%)	18 (7.9%)	7 (4.5%)

Table 2: Pregnancy Outcomes

	Revealed PIGF <12pg/ml N = 130	Concealed PIGF <12pg/ml N = 106	Revealed PIGF 12-100pg/ml N = 212	Concealed PIGF 12-100pg/ml N = 173	Revealed PIGF >100pg/ml N= 229	Concealed PIGF >100pg/ml N= 156
Maternal adverse outcomes n of women (%) * aOR (95% CI)	8 (6.2%)	6 (5.7%) 0.87 (0.09 to 8.02)	8 (3.8%)	12 (6.9%) 0.15 (0.03 to 0.92)	6 (2.6%)	6 (3.8%) 0.29 (0.02 to 4.34)
Use of antenatal corticosteroids for fetal lung maturity n (%)	98 (75.4%)	54 (50.9%)	67 (31.6%)	51 (29.5%)	35 (15.3%)	22 (14.1%)
Those delivering <35 weeks, % who got steroids in 7 days	29/75 (39%)	6/38 (16%)	12/32 (37.5%)	5/19 (26%)	3/6 (50%)	1/5 (20%)
Gestation at delivery, weeks Mean (SD) Mean Difference (95% CI)	33.4 (3.13)	34.4 (3.72) -0.03 (-1.72 to 1.66)	36.71 (2.48)	37.06 (2.04) -0.40 (-1.25 to 0.45)	38.30 (1.75)	38.23 (2.33) 0.36 (-0.44 to 1.16)
Status at Birth n (%)						
Livebirth	126 (96.9%)	102 (96.2%)	211 (99.5%)	171 (98.8%)	227 (100.0%)	153 (98.1%)
Stillbirth	4 (3.1%)	4 (3.8%)	1 (0.5%)	2 (1.2%)	0 (0.0%)	2 (1.3%)
Miscarriage <24 weeks	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (0.6%)
Birthweight centile by INTERGROWTH Mean (SD) Mean Difference (95% CI)	19.8 (22.4)	25.0 (28.8) 2.2 (-10.8 to 15.2)	41.5 (31.8)	44.1 (32.4) -2.2 (14.0 to 9.5)	57.1 (31.5)	54.8 (30.9) 3.1 (-9.3 to 15.4)
Birthweight centile <10th aOR (95% CI)	54 (42%)	48 (46%) 0.44 (0.15 to 1.27)	47 (22.4%)	35 (20.2%) 0.90 (0.33 to 2.48)	23 (10.1%)	14 (9.0%) 1.85 (0.45 to 7.67)
Neonatal unit admission n (%) aOR (95% CI)	93 (71.5%)	62 (58.5%) 2.37 (0.63-7.92)	73 (34.4%)	54 (31.2%) 2.37 (0.76-7.37)	29 (12.7%)	27 (17.3%)
Perinatal adverse outcome, n of infants (%) ** aOR (95% CI)	49 (37.7%)	27 (25.5%) 1.95 (0.64 to 6.00)	25 (11.8%)	23 (13.5%) 1.62 (0.45 to 5.89)	12 (5.2%)	9 (5.8%) 3.84 (0.29 to 51.31)

*As defined by the fullPIERS consensus¹⁴ (number of women with one or more of the following features; maternal death, eclampsia, Glasgow Coma Scale <13, stroke, transient ischaemic attack, cortical blindness, posterior reversible encephalopathy, retinal detachment, positive inotropic support, infusion of third parenteral antihypertensive, myocardial ischaemia or infarction, blood oxygen saturations <90%, 50% FiO₂ for > 1 hour, intubation (other than for

caesarean section), pulmonary oedema, inotropic support, transfusion of blood products, platelets $<50 \times 10^9$ per litre, hepatic dysfunction, haematoma or rupture, severe acute kidney injury (creatinine $>150 \mu\text{mol/L}$ or $>200 \mu\text{mol/L}$ in chronic kidney disease, dialysis, placental abruption)).

**Number of babies with one or more of the following features: perinatal death, late neonatal death (8-27 completed days of life), necrotising enterocolitis (stage 2 or 3), respiratory distress syndrome, bronchopulmonary dysplasia, seizures, retinopathy of prematurity, intraventricular haemorrhage.

Journal Pre-proof

References

1. Tan MY, Wright D, Syngelaki A, et al. Comparison of diagnostic accuracy of early screening for pre-eclampsia by NICE guidelines and a method combining maternal factors and biomarkers: results of SPREE. *Ultrasound in obstetrics & gynecology : the official journal of the International Society of Ultrasound in Obstetrics and Gynecology* 2018;Epub ahead of print doi: 10.1002/uog.19039.
2. Duley L. The global impact of pre-eclampsia and eclampsia. *Seminars in perinatology* 2009;33:130-7.
3. Redman C. Pre-eclampsia: A complex and variable disease. *Pregnancy hypertension* 2014;4:241-2.
4. Minire A, Mirton M, Imri V, Lauren M, Aferdita M. Maternal complications of preeclampsia. *Medical archives (Sarajevo, Bosnia and Herzegovina)* 2013;67:339-41.
5. Backes CH, Markham K, Moorehead P, Cordero L, Nankervis CA, Giannone PJ. Maternal preeclampsia and neonatal outcomes. *Journal of pregnancy* 2011;2011:214365.
6. Goldenberg RL, Culhane JF, Iams JD, Romero R. Epidemiology and causes of preterm birth. *Lancet (London, England)* 2008;371:75-84.
7. Menzies J, Magee LA, Macnab YC, et al. Current CHS and NHBPEP criteria for severe preeclampsia do not uniformly predict adverse maternal or perinatal outcomes. *Hypertension in pregnancy* 2007;26:447-62.
8. Maynard SE, Min JY, Merchan J, et al. Excess placental soluble fms-like tyrosine kinase 1 (sFlt1) may contribute to endothelial dysfunction, hypertension, and proteinuria in preeclampsia. *J Clin Invest* 2003;111:649-58.
9. Levine RJ, Maynard SE, Qian C, et al. Circulating angiogenic factors and the risk of preeclampsia. *The New England journal of medicine* 2004;350:672-83.
10. Zeisler H, Llurba E, Chantraine F, et al. Predictive Value of the sFlt-1:PIGF Ratio in Women with Suspected Preeclampsia. *The New England journal of medicine* 2016;374:13-22.
11. Chappell LC, Duckworth S, Seed PT, et al. Diagnostic accuracy of placental growth factor in women with suspected preeclampsia: a prospective multicenter study. *Circulation* 2013;128:2121-31.
12. Duhig KE, Myers J, Seed PT, et al. Placental growth factor testing to assess women with suspected pre-eclampsia: a multicentre, pragmatic, stepped-wedge cluster-randomised controlled trial. *Lancet* 2019;393:1807-18.
13. National Institute for Clinical Excellence. *Clinical Guideline 107: Hypertension in Pregnancy, the Management of Hypertensive Disorders During Pregnancy*. 2010.
14. von Dadelszen P, Payne B, Li J, et al. Prediction of adverse maternal outcomes in pre-eclampsia: development and validation of the fullPIERS model. *Lancet (London, England)* 2011;377:219-27.
15. Hypertension in pregnancy. Report of the American College of Obstetricians and Gynecologists' Task Force on Hypertension in Pregnancy. *Obstetrics and gynecology* 2013;122:1122-31.
16. Tranquilli AL, Dekker G, Magee L, et al. The classification, diagnosis and management of the hypertensive disorders of pregnancy: A revised statement from the ISSHP. *Pregnancy hypertension* 2014;4:97-104.
17. Villar J, Cheikh Ismail L, Victora CG, et al. International standards for newborn weight, length, and head circumference by gestational age and sex: the Newborn Cross-Sectional Study of the INTERGROWTH-21st Project. *Lancet* 2014;384:857-68.
18. Bell MJ, Ternberg JL, Feigin RD, et al. Neonatal necrotizing enterocolitis. Therapeutic decisions based upon clinical staging. *Ann Surg* 1978;187:1-7.
19. Ormesher L, Johnstone ED, Shawkat E, et al. A clinical evaluation of placental growth factor in routine practice in high-risk women presenting with suspected pre-eclampsia and/or fetal growth restriction. *Pregnancy hypertension* 2018;14:234-9.

20. Dias MAB, De Oliveira L, Jeyabalan A, et al. PREPARE: protocol for a stepped wedge trial to evaluate whether a risk stratification model can reduce preterm deliveries among women with suspected or confirmed preterm pre-eclampsia. *BMC Pregnancy Childbirth* 2019;19:343.
21. Cerdeira AS, O'Sullivan J, Ohuma EO, et al. Randomized Interventional Study on Prediction of Preeclampsia/Eclampsia in Women With Suspected Preeclampsia: INSPIRE. *Hypertension (Dallas, Tex : 1979)* 2019;74:983-90.
22. Frank L, Basch E, Selby JV. The PCORI perspective on patient-centered outcomes research. *Jama* 2014;312:1513-4.
23. Patsopoulos NA. A pragmatic view on pragmatic trials. *Dialogues Clin Neurosci* 2011;13:217-24.
24. Duhig KE, Seed PT, Myers JE, et al. Placental growth factor testing for suspected pre-eclampsia: a cost-effectiveness analysis. *BJOG : an international journal of obstetrics and gynaecology* 2019;126:1390-8.
25. Sharp A, Chappell LC, Dekker G, et al. Placental Growth Factor informed management of suspected pre-eclampsia or fetal growth restriction: The MAPPLE cohort study. *Pregnancy hypertension* 2018;Epub ahead of print doi: 10.1016/j.preghy.2018.03.013.