

Short Communication**Mechanism of methaemoglobin breakdown by the Lysine-specific gingipain of the periodontal pathogen
*Porphyromonas gingivalis***

John W Smalley^{1*}, Andrew J Birss¹, Borys Szmigielski² and Jan Potempa^{2,3}

¹Unit of Plaque-Related Diseases, School of Dental Science, The University of Liverpool, Liverpool L69 3GN, UK

²Department of Microbiology, Faculty of Biochemistry, Biophysics and Biotechnology, Jagiellonian University, ul. Gronostajowa 7, 30-387 Krakow, Poland, and ³Department of Biochemistry and Molecular Biology, Life Science Building, University of Georgia, Athens, GA 30602, USA

Corresponding author*

e-mail: josmall@liv.ac.uk

Running Title. Methaemoglobin breakdown by *P. gingivalis* K-gingipain.

Key words: haem; haemoglobin; gingipains; Porphyromonas; periodontal disease; protease


Abstract

The R- and K-gingipain proteases of *Porphyromonas gingivalis* are involved in proteolysis of haemoglobin from which the defensive dimeric haem pigment is formed. Whilst oxyhaemoglobin is refractory towards K-gingipain, methaemoglobin is rapidly degraded. Ligation of methaemoglobin with N_3^- , which effectively blocks haem dissociation from the protein, prevented haemoglobin breakdown. Haem-free globin was rapidly degraded by K-gingipain. These data emphasise the need for haemoglobin oxidation which encourages haem dissociation and makes the haem-free globin susceptible to proteolytic attack.

Introduction

Porphyromonas gingivalis is considered to be one of the major causal organisms associated with adult periodontitis (Lamont and Jenkinson, 1998). It elaborates a number of pathogenic factors including the Arginine-(R)- and Lysine-(K)-specific gingipain proteases, which are thought to contribute to periodontal disease progression. Gingipains not only directly degrade components of connective tissue but are also able to significantly interfere with host defence mechanisms and with regulated inflammatory and reparative processes in periodontal tissues (Potempa et al., 2000).

It has been recently shown (Smalley et al., 2004, 2007) that the HRgpA and Kgp act in concert to degrade oxyhaemoglobin for the release of iron protoporphyrin IX and production of the μ -oxo dimeric haem-containing black pigment. Whilst R-specific gingipains show little degradative activity towards haemoglobin, they can attack the oxyhaemoglobin molecule forming methaemoglobin (metHb; HbFe^+OH), the oxidised form of the haem-protein (Smalley et al., 2007). Although the individual α and β globin subunits contain numerous lysine residues, oxyhaemoglobin remains refractory to Kgp alone, resulting in the formation of a haemoglobin haemichrome which is relatively stable towards further Kgp attack. We have also demonstrated that metHb produced from oxyhaemoglobin by treatment with nitrite (Smalley et al., 2007) or with potassium ferricyanide (Smalley, JW, unpublished findings), is also readily attacked by Kgp. Thus, the effect of HRgpA is to “prime” haemoglobin molecules, oxidising them to the metHb form to facilitate complete degradation by Kgp. This also has the effect of rendering haemoglobin iron protoporphyrin IX molecules into the Fe(III) state (Fe(III)PPIX), so that upon proteolytic release from the haem protein, it can react to give the μ -oxo dimer, $[\text{Fe(III)PPIX}]_2\text{O}$, according to the following equation,



Although oxyHb and metHb differ imperceptibly from a structural point of view, these molecules differ in other physicochemical parameters. These include the inability of the metHb form to carry O_2 , and stability of the haem-globin linkage. It has been known for many years that whilst the haemoglobin

holoprotein is relatively stable, loss of the haem moiety to give apohaemoglobin is accompanied by increased protease susceptibility (Antonini and Brunori, 1971a). This would offer an explanation for the ability of Kgp to more easily degrade metHb whether it is generated by pre-treatment of oxyHb with HRgpA or by nitrite (Smalley et al., 2007). Accordingly, in this study we have examined further the effect of Kgp on apohaemoglobin and with azide liganded methaemoglobin (azidometHb), in which the haem-globin linkage is stabilised. We show here that whilst the azide liganded metHb species is less susceptible to attack, the apoprotein is rendered completely susceptible to breakdown by Kgp.

Incubation of metHb with Kgp resulted in the gradual decrease in the Soret band intensity (Fig 1a) indicating disruption of the haemprotein and loss of haem from the molecule. Over this time period there was little change to the intensity of the control metHb Soret band (data not shown). SDS-PAGE of the Kgp-metHb incubation mixture showed that the loss of Soret absorbance was matched by a progressive digestion of the α and β globin chains (Fig 2a). In contrast, in the presence of Kgp, the degree of Soret band loss of the azidometHb was minimal over the first 7 h incubation (Fig 1b), compared to the control. Densitometry confirmed that there was minimal breakdown of the α and β chains of azidometHb compared to metHb (Fig 3). However, when apohaemoglobin was incubated with Kgp there was a much more rapid breakdown of the alpha and beta globin chains (Fig 3), with approximately 70% of the total protein degraded over the first hour compared to approximately 25% for methaemoglobin. Under these enzyme assay conditions Kgp showed a loss of only 1% and 8% of initial activity after 7 and 24 h, respectively, as assessed using the synthetic substrate substrate N- α -acetyl-lysine-p-nitroanilide (data not presented). In this context the longevity of Kgp is biologically relevant for *P. gingivalis* since its haem pigmentation develops over several days and Hb is continuously supplied by lysing erythrocytes at the infected periodontitis sites.

Thus, these present results confirm previous observations (Smalley et al., 2007) that K-gingipain is able to rapidly degrade methaemoglobin compared to oxyhaemoglobin. Previous studies have also shown that the

oxidised form of haemoglobin is slightly more susceptible to attack by trypsin (Kimura et al., 1978). It has also been demonstrated that apo-haemoglobin is more easily degraded by trypsin, chymotrypsin and papain (Murakami and Murachi, 1978). Structural and conformational differences exist between the oxygenated and deoxygenated haemoglobin species which account for the physiological role of this molecule in oxygen carriage (Perutz, 1970). However, oxyhaemoglobin and methaemoglobin differ imperceptibly in their structures to account for the enhanced susceptibility of the latter towards Kgp as demonstrated here. It is known that general denaturation, including α -helix unfolding, leads to enhanced protease susceptibility of haemoglobin (Kimura et al., 1975, 1976). In addition, the presence of haem effectively stabilises the globin structure in both haemoglobin and myoglobin (Crumpton and Polson, 1965; Kawahara et al., 1965), and loss of haem is followed by loss of helical structure and denaturation. This stability is determined by the affinity of haem for the globin protein (Hargrove and Olson, 1996). Both haemoglobin and myoglobin display extremely high affinities for Fe(II) haem, with dissociation constants in the range 10^{-12} to 10^{-15} M (Hargrove et al., 1996). However, when the haem iron is oxidised to the Fe(III) state, the haem-globin affinity is reduced such that it may be transferred to, and bound by, serum albumin which has an association constant for iron(III) protoporphyrin IX in the region of 8×10^7 to 2×10^8 M⁻¹ (Adams and Berman, 1980; Gatoni et al., 1996). Thus, conversion of oxyhaemoglobin into the oxidised species either by nitrite or by pre-treatment with R-gingipain (Smalley et al., 2007) is accompanied by a greater degree of protein breakdown. Blocking the dissociation of ferric haems from methaemoglobin by adding low spin ligands such as CN⁻ and N₃⁻ (Bunn and Jandl, 1968), would thus lead to the predicted reduction in protease susceptibility. This was found to be the case when azidomethaemoglobin was incubated with Kgp. The time-dependent loss of Soret band intensity of the azidomethaemoglobin was reflected in the decrease in α and β chain breakdown as shown by SDS-PAGE. This resistance to proteolysis was not as a result of enzyme inhibition since Kgp is not inhibited by NaN₃ (Jan Potempa, unpublished data), even at 0.4mM, the concentration of sodium azide used to prepare azidomethaemoglobin.

Conversly, when apohaemoglobin was incubated with Kgp, rapid proteolysis of the globin chains was seen.

What is the significance of the current findings in relationship to haem acquisition by *P. gingivalis*? In moderately deep periodontal pocket sites, mean oxygenation tensions (pO₂) of around 30mmHg have been observed (Hanioka et al., 2000). In addition, periodontal tissue O₂ measurements at such sites reveals haemoglobin oxygen saturation levels in the range 60-70%, but which can be as high as 80% (Hanioka et al., 2000), indicating that these tissues are perfused with oxygenated blood. Thus, following bleeding, for example as a result of micro-ulceration, and haemolysis, periodontal pockets and sub-gingival plaque harbouring *P. gingivalis* may be exposed to oxygenated haemoglobin. Given the refractory nature of oxyhaemoglobin to Kgp proteolysis (Smalley et al., 2007), these present findings emphasise the need for a pre-requisite oxidation step in order for *P. gingivalis* to wrest haem from the oxyhaemoglobin. In this context it is noteworthy that *P. gingivalis* can grow in an oxygenated environment (Diaz and Rogers, 2004).

Methaemoglobin formation, mediated by the action of R-gingipain (Smalley et al., 2007), would serve two important roles. Firstly, it would lower the haem-globin affinity sufficiently to enhance haem dissociation, resulting in collapse of the protein structure, rendering the globin susceptible to proteolytic breakdown and facilitating a fuller release of the iron porphyrin. Secondly, it would poise the iron porphyrin in the Fe(III) state, such that upon proteolytic release, Fe(III)PPIX.OH (haematin) molecules could react together giving the haem dimer, [Fe(III)PPIX]₂O, according to equation 1 above (Smalley et al., 2006).

Acknowledgments

This work was supported in part by grant DE09761 (to J.P.) from National Institutes of Health.

References

- Adams, P.A., and Berman, M.C. (1980). Kinetics and mechanism of the interaction between human serum albumin and monomeric haemin. *Biochem. J.* 191, 95-102.
- Antonini and Brunori (1971a). Solution properties of myoglobin and hemoglobin. In: Hemoglobin and myoglobin in their reactions with ligands, North Holland Publishing Company, Amsterdam, pp. 98-134.
- Antonini and Brunori (1971b) The derivatives of ferric haemoglobin and myoglobin. In: Hemoglobin and myoglobin in their reactions with ligands, North Holland Publishing Company, Amsterdam, pp. 40-54.
- Ascoli, F., Fanelli, M.R., and Antonini, E. (1981). Preparation and properties of apohaemoglobin and reconstituted haemoglobins. *Methods Enzymol.* 76, 72-87.
- Beaven, G.H., Chen, S.H., d'Albis, A., and Gratzer, W.B. (1974). A spectroscopic study of the haemin-human-serum-albumin system. *Eur. J. Biochem.* 41, 539-546.
- Bunn, H.F., and Jandl, J.H. (1968). Exchange of heme among hemoglobins and between hemoglobin and albumin. *J. Biol. Chem.* 243, 465-475.
- Crumton, M.J., and Polson, A. (1965). A comparison of the conformation of sperm whale metmyoglobin with that of apomyoglobin. *J. Mol. Biol.* 11, 722-729.
- Diaz, P.I. and Rogers, A.H. (2004). The effect of oxygen on the growth and physiology of *Porphyromonas gingivalis*. *Oral Microbiol. Immunol.* 19, 88-94.
- Gatoni, M., Boffi, A., Sarti, P., and Chiancone, E. (1996). Stability of the heme-globin linkage in $\alpha\beta$ dimers and isolated chains of human haemoglobin. A study of heme transfer reaction from the immobilized proteins to albumin. *J. Biol. Chem.* 271, 10130-10136.
- Hanioka, T., Tanaka, M., Takaya, K., Matsumori, Y., and Shizukuishi, S. (2000). Pocket oxygen tension in smokers and non-smokers with periodontal disease. *J. Periodontol.* 71, 550-554.

- Hargrove, M.S., and Olson, J.S. (1996). The stability of holomyoglobin is determined by heme affinity. *Biochemistry* 35, 11310-11318.
- Hargrove, M.S., Barrick, D., and Olson, J.S. (1996). The association rate constant for heme binding to globin is independent of protein structure. *Biochemistry* 35, 11293-11299.
- Kawahara, K., Kirshner, A.G., and Tanford, C. (1965). Dissociation of human CO-hemoglobin by urea, guanidine hydrochloride, and other reagents. *Biochemistry* 4, 1203-1213.
- Kimura, H., Tsudzuki, T., and Murachi, T. (1975). Proteolytic degradation of hemoglobin-haptoglobin complex by lysosomal enzymes from rat liver. *J. Biochem.* 77, 909-912.
- Kimura, H., Yamato, S., and Murachi, T. (1976). Different susceptibilities to intracellular proteases of haemoglobin and haemoglobin-haptoglobin complex. *Physiol. Chem. Phys.* 8, 101-106.
- Kimura, H., Yamato, S., and Murachi, T. (1978). Increase in susceptibility of haemoglobin to proteases upon treatment with *p*-mercuribenzoate. *J. Biochem.* 84, 205-211.
- Lamont, R.J., and Jenkinson, H.F. (1998). Life below the gum line: pathogenic mechanisms of *Porphyromonas gingivalis*. *Microbiol. Mol. Biol. Rev.* 62, 1244-1263.
- Murakami, T., and Murachi, T. (1978). Enhanced susceptibility of haemoglobin to proteases upon ascorbic acid oxidation of the heme moiety. *J. Biochem.* 84, 83-91.
- Perutz, M.F. (1970). Stereochemistry and cooperative effects in haemoglobin. *Nature* 228, 726-734.
- Pike, R.N., McGraw, W., Potempa, J., and Travis, J. (1994). Lysine- and arginine-specific proteinases from *Porphyromonas gingivalis*. Isolation, characterization, and evidence for the existence of complexes with hemagglutinins. *J. Biol. Chem.* 269, 406-411.

- Potempa, J., Banbula, A., and Travis, J. (2000). Role of bacterial proteinases in matrix destruction and modulation of host responses. *Periodontol.* 2000 *24*, 153-192.
- Potempa, J., Pike, R., and Travis, J. (1997). Titration and mapping of the active site of cysteine proteinases from *Porphyromonas gingivalis* (gingipains) using peptidyl chloromethanes. *Biol. Chem.* *378*, 223-230.
- Smalley, J.W., Thomas, M.F., Birss, A.J., Withnall, R. and Silver, J. (2004). A combination of both arginine- and lysine-specific gingipain activity of *Porphyromonas gingivalis* is necessary for the generation of the μ -oxo bishaem-containing pigment from haemoglobin. *Biochem. J.* *379*, 833-840.
- Smalley, J.W., Birss, A.J., Szmigielski, B., Potempa, J. (2006). The HA2 haemagglutinin domain of the lysine-specific gingipain (Kgp) of *Porphyromonas gingivalis* promotes μ -oxo bishaem formation from monomeric iron(III) protoporphyrin IX. *Microbiology.* *152*, 1839-1845.
- Smalley, J.W., Birss, A.J., Szmigielski, B., and Potempa, J. (2007). Sequential action of R- and K-specific gingipains of *Porphyromonas gingivalis* in the generation of the haem-containing pigment from oxyhaemoglobin. *Arch. Biochem. Biophys.* *465*, 44-49.

Fig 1




Fig 1

UV-visible absorption spectra of horse methaemoglobin (a) and azidomethaemoglobin (b) recorded during incubation with Kgp at 37°C. Methaemoglobin was prepared by incubation of oxyhaemoglobin with NaNO_2 as previously described (Smalley et al., 2007). Azidomethaemoglobin was prepared as described by (Antonini and Brunori, 1971b). Kgp was isolated and purified from the culture medium of *P. gingivalis* strain HG66 (Pike et al., 1994). Haemoglobin and enzyme concentrations were 4 and $0.4\mu\text{M}$, respectively. Buffer was 0.14M NaCl, 0.1M Tris-HCl, pH 7.5. Spectra were recorded in a 1 cm path length cuvette.

Fig 2




Fig 2

SDS-PAGE of azidomethaemoglobin (a), methaemoglobin (b) and apohaemoglobin (c) during incubation with Kgp. (d) control apohaemoglobin minus Kgp. Incubation conditions as in Fig 1. Haem-free apohaemoglobin was prepared by the acid-acetone method of Ascoli et al. (1981). Enzyme and substrate concentrations were 0.4 and 4 μ M, respectively. SDS-PAGE was carried out on 15% acrylamide gels after sample solubilisation at 100°C for 5 min under reducing conditions as previously described (Smalley et al., 2007). Five μ g of protein were loaded per track.

Fig 3




Fig 3

Loss of intact globin chains of azidomethaemoglobin, methaemoglobin and haem-free globin chains after incubation with Kgp. The α and β chains of each haemoglobin species, separated by SDS-PAGE as in Fig 2, were scanned densitometrically after staining with coomassie blue and integrated using UViBand software. Incubation conditions and protein substrate and enzyme concentrations were the same as described for Fig 2.